

6. Literatur

Alcantara, S., Ruiz, M., De Castro, F., Soriano, E. and Sotelo, C. (2000). Netrin 1 acts as an attractive or as a repulsive cue for distinct migrating neurons during the development of the cerebellar system. *Development* 127, 1359-72.

Altman, J. and Bayer, S. A. (1980a). Development of the brain stem in the rat. I. Thymidine-radiographic study of the time of origin of neurons of the lower medulla. *J Comp Neurol* 194, 1-35.

Altman, J. and Bayer, S. A. (1980b). Development of the brain stem in the rat. IV. Thymidine-radiographic study of the time of origin of neurons in the pontine region. *J Comp Neurol* 194, 905-29.

Altman, J. and Bayer, S. A. (1985). Embryonic development of the rat cerebellum. I. Delineation of the cerebellar primordium and early cell movements. *J Comp Neurol* 231, 1-26.

Altman, J. and Bayer, S. A. (1987a). Development of the precerebellar nuclei in the rat: I. The precerebellar neuroepithelium of the rhombencephalon. *J Comp Neurol* 257, 477-89.

Altman, J. and Bayer, S. A. (1987b). Development of the precerebellar nuclei in the rat: II. The intramural olfactory migratory stream and the neurogenetic organization of the inferior olive. *J Comp Neurol* 257, 490-512.

Altman, J. and Bayer, S. A. (1987c). Development of the precerebellar nuclei in the rat: III. The posterior precerebellar extramural migratory stream and the lateral reticular and external cuneate nuclei. *J Comp Neurol* 257, 513-28.

Altman, J. and Bayer, S. A. (1987d). Development of the precerebellar nuclei in the rat: IV. The anterior precerebellar extramural migratory stream and the nucleus reticularis tegmenti pontis and the basal pontine gray. *J Comp Neurol* 257, 529-52.

Ambrosiani, J., Armengol, J. A., Martinez, S. and Puelles, L. (1996). The avian inferior olive derives from the alar neuroepithelium of the rhombomeres 7 and 8: an analysis by using chick-quail chimeric embryos. *Neuroreport* 7, 1285-8.

Aruga, J., Shimoda, K. and Mikoshiba, K. (2000). A 5' segment of the mouse Zic1 gene contains a region specific enhancer for dorsal hindbrain and spinal cord. *Brain Res Mol Brain Res* 78, 15-25.

Barrow, J. R., Stadler, H. S. and Capecchi, M. R. (2000). Roles of Hoxa1 and Hoxa2 in patterning the early hindbrain of the mouse. *Development* 127, 933-44.

- Bartholoma, A. and Nave, K. A. (1994). NEX-1: a novel brain-specific helix-loop-helix protein with autoregulation and sustained expression in mature cortical neurons. *Mech Dev* 48, 217-28.
- Bel-Vialar, S., Itasaki, N. and Krumlauf, R. (2002). Initiating Hox gene expression: in the early chick neural tube differential sensitivity to FGF and RA signaling subdivides the HoxB genes in two distinct groups. *Development* 129, 5103-15.
- Bell, E., Wingate, R. J. and Lumsden, A. (1999). Homeotic transformation of rhombomere identity after localized Hoxb1 misexpression. *Science* 284, 2168-71.
- Ben-Arie, N., Bellen, H. J., Armstrong, D. L., McCall, A. E., Gordadze, P. R., Guo, Q., Matzuk, M. M. and Zoghbi, H. Y. (1997). Math1 is essential for genesis of cerebellar granule neurons. *Nature* 390, 169-72.
- Berger, S. L. and Kimmel, A. R. (1987). Guide To Molecular Cloning Techniques. San Diego, CA 92101-4495, USA: Academic Press.
- Birmingham, N. A., Hassan, B. A., Wang, V. Y., Fernandez, M., Banfi, S., Bellen, H. J., Fritzsch, B. and Zoghbi, H. Y. (2001). Proprioceptor pathway development is dependent on Math1. *Neuron* 30, 411-22.
- Birnboim, H. C. and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 7, 1513-23.
- Bloch-Gallego, E., Causeret, F., Ezan, F., Backer, S. and Hidalgo-Sanchez, M. (2005). Development of precerebellar nuclei: instructive factors and intracellular mediators in neuronal migration, survival and axon pathfinding. *Brain Res Brain Res Rev* 49, 253-66.
- Borday, C., Wrobel, L., Fortin, G., Champagnat, J., Thaeron-Antono, C. and Thoby-Brisson, M. (2004). Developmental gene control of brainstem function: views from the embryo. *Prog Biophys Mol Biol* 84, 89-106.
- Bourrat, F. and Sotelo, C. (1990). Early development of the rat precerebellar system: migratory routes, selective aggregation and neuritic differentiation of the inferior olive and lateral reticular nucleus neurons. An overview. *Arch Ital Biol* 128, 151-70.
- Bradley, A. and Robertson, E. (1986). Embryo-derived stem cells: a tool for elucidating the developmental genetics of the mouse. *Curr Top Dev Biol* 20, 357-71.
- Briscoe, J. and Ericson, J. (1999). The specification of neuronal identity by graded Sonic Hedgehog signalling. *Semin Cell Dev Biol* 10, 353-62.
- Briscoe, J. and Ericson, J. (2001). Specification of neuronal fates in the ventral neural tube. *Curr Opin Neurobiol* 11, 43-9.

- Briscoe, J., Sussel, L., Serup, P., Hartigan-O'Connor, D., Jessell, T. M., Rubenstein, J. L. and Ericson, J. (1999). Homeobox gene Nkx2.2 and specification of neuronal identity by graded Sonic hedgehog signalling. *Nature* 398, 622-7.
- Cao, S., Liu, J., Song, L. and Ma, X. (2005). The protooncogene c-Maf is an essential transcription factor for IL-10 gene expression in macrophages. *J Immunol* 174, 3484-92.
- Carpenter, E. M., Goddard, J. M., Chisaka, O., Manley, N. R. and Capecchi, M. R. (1993). Loss of Hox-A1 (Hox-1.6) function results in the reorganization of the murine hindbrain. *Development* 118, 1063-75.
- Cheng, L., Samad, O. A., Xu, Y., Mizuguchi, R., Luo, P., Shirasawa, S., Goulding, M. and Ma, Q. (2005). Lbx1 and Tlx3 are opposing switches in determining GABAergic versus glutamatergic transmitter phenotypes. *Nat Neurosci* 8, 1510-5.
- Cheng, T. C., Wallace, M. C., Merlie, J. P. and Olson, E. N. (1993). Separable regulatory elements governing myogenin transcription in mouse embryogenesis. *Science* 261, 215-8.
- Church, G. M. and Gilbert, W. (1984). Genomic sequencing. *Proc Natl Acad Sci U S A* 81, 1991-5.
- Cohen, G. and Sternberg, N. (1989). Genetic analysis of the lytic replicon of bacteriophage P1. I. Isolation and partial characterization. *J Mol Biol* 207, 99-109.
- Corpet, F. (1988). Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res* 16, 10881-90.
- Currie, D. S., Cheng, X., Hsu, C. M. and Monuki, E. S. (2005). Direct and indirect roles of CNS dorsal midline cells in choroid plexus epithelia formation. *Development* 132, 3549-59.
- Dampney, R. A. (1981). Brain stem mechanisms in the control of arterial pressure. *Clin Exp Hypertens* 3, 379-91.
- Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. and McMahon, A. P. (1998). Modification of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre recombinase. *Curr Biol* 8, 1323-6.
- Dauger, S., Pattyn, A., Lofaso, F., Gaultier, C., Goridis, C., Gallego, J. and Brunet, J. F. (2003). Phox2b controls the development of peripheral chemoreceptors and afferent visceral pathways. *Development* 130, 6635-42.
- Ding, L., Takebayashi, H., Watanabe, K., Ohtsuki, T., Tanaka, K. F., Nabeshima, Y., Chisaka, O., Ikenaka, K. and Ono, K. (2005). Short-term lineage analysis of dorsally derived Olig3 cells in the developing spinal cord. *Dev Dyn* 234, 622-32.

- Duus, P. (1990). Neurologisch-topische Diagnostik, 5. Auflage, Thieme.
- Eagleson, G. W. and Harris, W. A. (1990). Mapping of the presumptive brain regions in the neural plate of *Xenopus laevis*. *J Neurobiol* 21, 427-40.
- Einthoven, W., Fahr, G. and De Waart, A. (1950). On the direction and manifest size of the variations of potential in the human heart and on the influence of the position of the heart on the form of the electrocardiogram. *Am Heart J* 40, 163-211.
- Ellenberger, C., Jr., Hanaway, J. and Netsky, M. G. (1969). Embryogenesis of the inferior olfactory nucleus in the rat: a radioautographic study and a re-evaluation of the rhombic lip. *J Comp Neurol* 137, 71-9.
- Eng, S. R., Gratwick, K., Rhee, J. M., Fedtsova, N., Gan, L. and Turner, E. E. (2001). Defects in sensory axon growth precede neuronal death in *Brn3a*-deficient mice. *J Neurosci* 21, 541-9.
- Ericson, J., Morton, S., Kawakami, A., Roelink, H. and Jessell, T. M. (1996). Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity. *Cell* 87, 661-73.
- Ericson, J., Muhr, J., Jessell, T. M. and Edlund, T. (1995a). Sonic hedgehog: a common signal for ventral patterning along the rostrocaudal axis of the neural tube. *Int J Dev Biol* 39, 809-16.
- Ericson, J., Muhr, J., Placzek, M., Lints, T., Jessell, T. M. and Edlund, T. (1995b). Sonic hedgehog induces the differentiation of ventral forebrain neurons: a common signal for ventral patterning within the neural tube. *Cell* 81, 747-56.
- Essick, C. R. (1912). The development of the nuclei pontis and the nucleus arcuatus in man. *Am J Anat* 13:25-54.
- Feinberg, A. P. and Vogelstein, B. (1983). A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* 132, 6-13.
- Filippi, A., Tiso, N., Deflorian, G., Zecchin, E., Bortolussi, M. and Argenton, F. (2005). The basic helix-loop-helix olig3 establishes the neural plate boundary of the trunk and is necessary for development of the dorsal spinal cord. *Proc Natl Acad Sci U S A* 102, 4377-82.
- Fraser, S., Keynes, R. and Lumsden, A. (1990). Segmentation in the chick embryo hindbrain is defined by cell lineage restrictions. *Nature* 344, 431-5.
- Frohman, M. A., Martin, G. R., Cordes, S. P., Halamek, L. P. and Barsh, G. S. (1993). Altered rhombomere-specific gene expression and hyoid bone differentiation in the mouse segmentation mutant, kreisler (kr). *Development* 117, 925-36.

- Fujiwara, K. T., Kataoka, K. and Nishizawa, M. (1993). Two new members of the maf oncogene family, mafK and mafF, encode nuclear b-Zip proteins lacking putative trans-activator domain. *Oncogene* 8, 2371-80.
- Gao, W. Q. and Hatten, M. E. (1994). Immortalizing oncogenes subvert the establishment of granule cell identity in developing cerebellum. *Development* 120, 1059-70.
- Gavrieli, Y., Sherman, Y. and Ben-Sasson, S. A. (1992). Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol* 119, 493-501.
- George.J.; Fitzpatrick, David; Katz, Lawrence.C.; LaMantia, Anthony-Samuel.; McNamara, James.O.; Williams, S. Mark (2001). Neuroscience. 2nd ed., Sunderland (MA): Sinauer Associates, Inc.
- Gross, M. K., Dottori, M. and Goulding, M. (2002). Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. *Neuron* 34, 535-49.
- Gu, H., Zou, Y. R. and Rajewsky, K. (1993). Independent control of immunoglobulin switch recombination at individual switch regions evidenced through Cre-loxP-mediated gene targeting. *Cell* 73, 1155-64.
- Harkmark, W. (1954). Cell migrations from the rhombic lip to the inferior olive, the nucleus raphe and the pons; a morphological and experimental investigation on chick embryos. *J Comp Neurol* 100, 115-209.
- Helms, A. W. and Johnson, J. E. (1998). Progenitors of dorsal commissural interneurons are defined by MATH1 expression. *Development* 125, 919-28.
- Hemmati-Brivanlou, A. and Melton, D. (1997). Vertebrate embryonic cells will become nerve cells unless told otherwise. *Cell* 88, 13-7.
- Herrup, K. (1983). Role of staggerer gene in determining cell number in cerebellar cortex. I. Granule cell death is an indirect consequence of staggerer gene action. *Brain Res* 313, 267-74.
- Hippenmeyer, S., Vrieseling, E., Sigrist, M., Portmann, T., Laengle, C., Ladle, D. R. and Arber, S. (2005). A developmental switch in the response of DRG neurons to ETS transcription factor signaling. *PLoS Biol* 3, e159.
- Hirsch, M. R., Tiveron, M. C., Guillemot, F., Brunet, J. F. and Goridis, C. (1998). Control of noradrenergic differentiation and Phox2a expression by MASH1 in the central and peripheral nervous system. *Development* 125, 599-608.

- His, W. (1891). Die Entwicklung des menschlichen Rautenhirns vom Ende des ersten bis zum Beginn des dritten Monats. I. Verlängertes Mark. Abhandlungen der königlichen sächsischen Gesellschaft der Wissenschaften, *Mathematische-physikalische Klasse*, pp. 1–74.
- Ho, I. C., Hodge, M. R., Rooney, J. W. and Glimcher, L. H. (1996). The proto-oncogene c-maf is responsible for tissue-specific expression of interleukin-4. *Cell* 85, 973-83.
- Ho, I. C., Lo, D. and Glimcher, L. H. (1998). c-maf promotes T helper cell type 2 (Th2) and attenuates Th1 differentiation by both interleukin 4-dependent and -independent mechanisms. *J Exp Med* 188, 1859-66.
- Hoshino, M., Nakamura, S., Mori, K., Kawauchi, T., Terao, M., Nishimura, Y. V., Fukuda, A., Fuse, T., Matsuo, N., Sone, M. et al. (2005). Ptf1a, a bHLH transcriptional gene, defines GABAergic neuronal fates in cerebellum. *Neuron* 47, 201-13.
- Innis, M. A., Gelfand, D. H. and Sninsky, J. J. (1989). PCR Protocols: A Guide To Methods And Applications, (ed., pp. 482. San Diego, CA 92101-4495, USA: Academic Press.
- Inoue, H., Nojima, H. and Okayama, H. (1990). High efficiency transformation of Escherichia coli with plasmids. *Gene* 96, 23-8.
- Irving, C. and Mason, I. (2000). Signalling by FGF8 from the isthmus patterns anterior hindbrain and establishes the anterior limit of Hox gene expression. *Development* 127, 177-86.
- Jan, Y. N. and Jan, L. Y. (1993). HLH proteins, fly neurogenesis, and vertebrate myogenesis. *Cell* 75, 827-30.
- Jessell, T. M. (2000). Neuronal specification in the spinal cord: inductive signals and transcriptional codes. *Nat Rev Genet* 1, 20-9.
- Johnson, J. E., Birren, S. J. and Anderson, D. J. (1990). Two rat homologues of *Drosophila* achaete-scute specifically expressed in neuronal precursors. *Nature* 346, 858-61.
- Joyner, A. L. (1999). Gene Targeting: A Practical Approach. Oxford, United Kingdom: Oxford University Press.
- Ju, M. J., Aroca, P., Luo, J., Puelles, L. and Redies, C. (2004). Molecular profiling indicates avian branchiomotor nuclei invade the hindbrain alar plate. *Neuroscience* 128, 785-96.
- Kandel, E. R., Schwartz, J. H., Jessell, T. M. (1991). Principles of Neural Science. Appleton and Lange, Norwalk, CT.

- Kataoka, K., Fujiwara, K. T., Noda, M. and Nishizawa, M. (1994). MafB, a new Maf family transcription activator that can associate with Maf and Fos but not with Jun. *Mol Cell Biol* 14, 7581-91.
- Kataoka, K., Igarashi, K., Itoh, K., Fujiwara, K. T., Noda, M., Yamamoto, M. and Nishizawa, M. (1995). Small Maf proteins heterodimerize with Fos and may act as competitive repressors of the NF-E2 transcription factor. *Mol Cell Biol* 15, 2180-90.
- Kataoka, K., Nishizawa, M. and Kawai, S. (1993). Structure-function analysis of the maf oncogene product, a member of the b-Zip protein family. *J Virol* 67, 2133-41.
- Kawauchi, D., Taniguchi, H., Watanabe, H., Saito, T. and Murakami, F. (2006). Direct visualization of nucleogenesis by precerebellar neurons: involvement of ventricle-directed, radial fibre-associated migration. *Development* 133, 1113-23.
- Kawauchi, S., Takahashi, S., Nakajima, O., Ogino, H., Morita, M., Nishizawa, M., Yasuda, K. and Yamamoto, M. (1999). Regulation of lens fiber cell differentiation by transcription factor c-Maf. *J Biol Chem* 274, 19254-60.
- Kerppola, T. K. and Curran, T. (1994). A conserved region adjacent to the basic domain is required for recognition of an extended DNA binding site by Maf/Nrl family proteins. *Oncogene* 9, 3149-58.
- Kim, J. I., Li, T., Ho, I. C., Grusby, M. J. and Glimcher, L. H. (1999). Requirement for the c-Maf transcription factor in crystallin gene regulation and lens development. *Proc Natl Acad Sci U S A* 96, 3781-5.
- Kozak, M. (1987). At least six nucleotides preceding the AUG initiator codon enhance translation in mammalian cells. *J Mol Biol* 196, 947-50.
- Kühn, R., Rajewsky, K. and Müller, W. (1991). Generation and analysis of interleukin-4 deficient mice. *Science* 254, 707-10.
- Kuhn, R., Schwenk, F., Aguet, M. and Rajewsky, K. (1995). Inducible gene targeting in mice. *Science* 269, 1427-9.
- Landsberg, R. L., Awatramani, R. B., Hunter, N. L., Farago, A. F., DiPietrantonio, H. J., Rodriguez, C. I. and Dymecki, S. M. (2005). Hindbrain rhombic lip is comprised of discrete progenitor cell populations allocated by Pax6. *Neuron* 48, 933-47.
- Lee, K. J. and Jessell, T. M. (1999). The specification of dorsal cell fates in the vertebrate central nervous system. *Annu Rev Neurosci* 22, 261-94.
- Li, M. Z., Wang, J. S., Jiang, D. J., Xiang, C. X., Wang, F. Y., Zhang, K. H., Williams, P. R. and Chen, Z. F. (2006). Molecular mapping of developing dorsal horn-enriched genes by microarray and dorsal/ventral subtractive screening. *Dev Biol* 292, 555-64.

- Li, S., Qiu, F., Xu, A., Price, S. M. and Xiang, M. (2004). Barhl1 regulates migration and survival of cerebellar granule cells by controlling expression of the neurotrophin-3 gene. *J Neurosci* 24, 3104-14.
- Lu, Q. R., Cai, L., Rowitch, D., Cepko, C. L. and Stiles, C. D. (2001). Ectopic expression of Olig1 promotes oligodendrocyte formation and reduces neuronal survival in developing mouse cortex. *Nat Neurosci* 4, 973-4.
- Lu, Q. R., Sun, T., Zhu, Z., Ma, N., Garcia, M., Stiles, C. D. and Rowitch, D. H. (2002). Common developmental requirement for Olig function indicates a motor neuron/oligodendrocyte connection. *Cell* 109, 75-86.
- Lumsden, A. (1999). Closing in on rhombomere boundaries. *Nat Cell Biol* 1, E83-5.
- Lumsden, A. and Krumlauf, R. (1996). Patterning the vertebrate neuraxis. *Science* 274, 1109-15.
- Machold, R. and Fishell, G. (2005). Math1 is expressed in temporally discrete pools of cerebellar rhombic-lip neural progenitors. *Neuron* 48, 17-24.
- Manzanares, M., Bel-Vialar, S., Ariza-McNaughton, L., Ferretti, E., Marshall, H., Maconochie, M. M., Blasi, F. and Krumlauf, R. (2001). Independent regulation of initiation and maintenance phases of Hoxa3 expression in the vertebrate hindbrain involve auto- and cross-regulatory mechanisms. *Development* 128, 3595-607.
- Manzanares, M., Cordes, S., Ariza-McNaughton, L., Sadl, V., Maruthainar, K., Barsh, G. and Krumlauf, R. (1999). Conserved and distinct roles of kreisler in regulation of the paralogous Hoxa3 and Hoxb3 genes. *Development* 126, 759-69.
- Marin, F. and Puelles, L. (1995). Morphological fate of rhombomeres in quail/chick chimeras: a segmental analysis of hindbrain nuclei. *Eur J Neurosci* 7, 1714-38.
- Massari, M. E. and Murre, C. (2000). Helix-loop-helix proteins: regulators of transcription in eucaryotic organisms. *Mol Cell Biol* 20, 429-40.
- McDonell, M. W., Simon, M. N. and Studier, F. W. (1977). Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J Mol Biol* 110, 119-46.
- McGinnis, W. and Krumlauf, R. (1992). Homeobox genes and axial patterning. *Cell* 68, 283-302.
- McKay, I. J., Muchamore, I., Krumlauf, R., Maden, M., Lumsden, A. and Lewis, J. (1994). The kreisler mouse: a hindbrain segmentation mutant that lacks two rhombomeres. *Development* 120, 2199-211.
- Miale, I. L. and Sidman, R. L. (1961). An autoradiographic analysis of histogenesis in the mouse cerebellum. *Exp Neurol* 4, 277-96.

- Millen, K. J., Millonig, J. H. and Hatten, M. E. (2004). Roof plate and dorsal spinal cord dl1 interneuron development in the dreher mutant mouse. *Dev Biol* 270, 382-92.
- Millhorn, D. E. and Eldridge, F. L. (1986). Role of ventrolateral medulla in regulation of respiratory and cardiovascular systems. *J Appl Physiol* 61, 1249-63.
- Millonig, J. H., Millen, K. J. and Hatten, M. E. (2000). The mouse Dreher gene Lmx1a controls formation of the roof plate in the vertebrate CNS. *Nature* 403, 764-9.
- Mizuguchi, R., Sugimori, M., Takebayashi, H., Kosako, H., Nagao, M., Yoshida, S., Nabeshima, Y., Shimamura, K. and Nakafuku, M. (2001). Combinatorial roles of olig2 and neurogenin2 in the coordinated induction of pan-neuronal and subtype-specific properties of motoneurons. *Neuron* 31, 757-71.
- Moran-Rivard, L., Kagawa, T., Saueressig, H., Gross, M. K., Burrill, J. and Goulding, M. (2001). Evx1 is a postmitotic determinant of v0 interneuron identity in the spinal cord. *Neuron* 29, 385-99.
- Muller, T., Anlag, K., Wildner, H., Britsch, S., Treier, M. and Birchmeier, C. (2005). The bHLH factor Olig3 coordinates the specification of dorsal neurons in the spinal cord. *Genes Dev* 19, 733-43.
- Muller, T., Brohmann, H., Pierani, A., Heppenstall, P. A., Lewin, G. R., Jessell, T. M. and Birchmeier, C. (2002). The homeodomain factor lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. *Neuron* 34, 551-62.
- Nakada, Y., Hunsaker, T. L., Henke, R. M. and Johnson, J. E. (2004). Distinct domains within Mash1 and Math1 are required for function in neuronal differentiation versus neuronal cell-type specification. *Development* 131, 1319-30.
- Nichols, D. H. and Bruce, L. L. (2006). Migratory routes and fates of cells transcribing the Wnt-1 gene in the murine hindbrain. *Dev Dyn* 235, 285-300.
- Nishizawa, M., Kataoka, K., Goto, N., Fujiwara, K. T. and Kawai, S. (1989). v-maf, a viral oncogene that encodes a "leucine zipper" motif. *Proc Natl Acad Sci U S A* 86, 7711-5.
- Novitch, B. G., Chen, A. I. and Jessell, T. M. (2001). Coordinate Regulation of Motor Neuron Subtype Identity and Pan-Neuronal Properties by the bHLH Repressor Olig2. *Neuron* 31, 773-89.
- Nowakowski, R. S., Lewin, S. B. and Miller, M. W. (1989). Bromodeoxyuridine immunohistochemical determination of the lengths of the cell cycle and the DNA-synthetic phase for an anatomically defined population. *J Neurocytol* 18, 311-8.

- Omoteyama, K., Ikeda, H., Imaki, J. and Sakai, M. (2006). Activation of connective tissue growth factor gene by the c-Maf and Lc-Maf transcription factors. *Biochem Biophys Res Commun* 339, 1089-97.
- Ono, K. and Kawamura, K. (1989). Migration of immature neurons along tangentially oriented fibers in the subpial part of the fetal mouse medulla oblongata. *Exp Brain Res* 78, 290-300.
- Pattyn, A., Guillemot, F. and Brunet, J. F. (2006). Delays in neuronal differentiation in Mash1/Ascl1 mutants. *Dev Biol.*
- Pierani, A., Brenner-Morton, S., Chiang, C. and Jessell, T. M. (1999). A sonic hedgehog-independent, retinoid-activated pathway of neurogenesis in the ventral spinal cord. *Cell* 97, 903-15.
- Pierani, A., Moran-Rivard, L., Sunshine, M. J., Littman, D. R., Goulding, M. and Jessell, T. M. (2001). Control of interneuron fate in the developing spinal cord by the progenitor homeodomain protein Dbx1. *Neuron* 29, 367-84.
- Pierce, E. T. (1966). Histogenesis of the nuclei griseum pontis, corporis pontobulbaris and reticularis segmenti pontis (Bechterew) in the mouse. An autoradiographic study. *J Comp Neurol* 126, 219-54.
- Pierce, E. T. (1967). Histogenesis of the dorsal and ventral cochlear nuclei in the mouse. An autoradiographic study. *J Comp Neurol* 131, 27-54.
- Pierce, E. T. (1973). Time of origin of neurons in the brain stem of the mouse. *Prog Brain Res* 40, 53-65.
- Qian, Y., Fritzsch, B., Shirasawa, S., Chen, C. L., Choi, Y. and Ma, Q. (2001). Formation of brainstem (nor)adrenergic centers and first-order relay visceral sensory neurons is dependent on homeodomain protein Rnx/Tlx3. *Genes Dev* 15, 2533-45.
- Rakic, P. (1999). Neurobiology. Discriminating migrations. *Nature* 400, 315-6.
- Ramirez-Solis, R., Rivera-Perez, J., Wallace, J. D., Wims, M., Zheng, H. and Bradley, A. (1992). Genomic DNA microextraction: a method to screen numerous samples. *Anal Biochem* 201, 331-5.
- Ring, B. Z., Cordes, S. P., Overbeek, P. A. and Barsh, G. S. (2000). Regulation of mouse lens fiber cell development and differentiation by the Maf gene. *Development* 127, 307-17.
- Rodriguez, C. I. and Dymecki, S. M. (2000). Origin of the precerebellar system. *Neuron* 27, 475-86.

- Roelink, H., Porter, J. A., Chiang, C., Tanabe, Y., Chang, D. T., Beachy, P. A. and Jessell, T. M. (1995). Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell* 81, 445-55.
- Rossant, J. und Tam, P. (2002). Mouse development – patterning, morphogenesis, and organogenesis, Academic Press.
- Rowitch, D. H., Lu, Q. R., Kessaris, N. and Richardson, W. D. (2002). An 'oligarchy' rules neural development. *Trends Neurosci* 25, 417-22.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. and Arnheim, N. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230, 1350-4.
- Sambrook, J. and Russell, D. W. (2001). Molecular Cloning: A Laboratory Manual. Cold Spring Harbor, New York 11803-2500, USA: Cold Spring Harbor Laboratory Press.
- Sander, M., Paydar, S., Ericson, J., Briscoe, J., Berber, E., German, M., Jessell, T. M. and Rubenstein, J. L. (2000). Ventral neural patterning by Nkx homeobox genes: Nkx6.1 controls somatic motor neuron and ventral interneuron fates. *Genes Dev* 14, 2134-9.
- Sanger, F., Nicklen, S. and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* 74, 5463-7.
- Sasai, Y., Kageyama, R., Tagawa, Y., Shigemoto, R. and Nakanishi, S. (1992). Two mammalian helix-loop-helix factors structurally related to Drosophila hairy and Enhancer of split. *Genes Dev* 6, 2620-34.
- Sauer, B. and Henderson, N. (1990). Targeted insertion of exogenous DNA into the eukaryotic genome by the Cre recombinase. *New Biol* 2, 441-9.
- Schaper, A. (1897). The earliest differentiation in the central nervous system of vertebrates. *Science* 5:430-31.
- Schenkel, J. (2006) Transgene Tiere, Spektrum Akademischer Verlag.
- Schneider-Maunoury, S., Gilardi-Hebenstreit, P. and Charnay, P. (1998). How to build a vertebrate hindbrain. Lessons from genetics. *C R Acad Sci III* 321, 819-34.
- Schneider-Maunoury, S., Topilko, P., Seitandou, T., Levi, G., Cohen-Tannoudji, M., Pournin, S., Babinet, C. and Charnay, P. (1993). Disruption of Krox-20 results in alteration of rhombomeres 3 and 5 in the developing hindbrain. *Cell* 75, 1199-214.

- Schoenwolf, G. C. and Sheard, P. (1990). Fate mapping the avian epiblast with focal injections of a fluorescent-histochemical marker: ectodermal derivatives. *J Exp Zool* 255, 323-39.
- Shojaeian, H., Delhaye-Bouchaud, N. and Mariani, J. (1985). Decreased number of cells in the inferior olfactory nucleus of the developing staggerer mouse. *Brain Res* 353, 141-6.
- Sonhammer, E. L. and Durbin, R. (1995). A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis. *Gene* 167, GC1-10.
- Sorge, J. A. (1988). Bacteriophage lambda cloning vectors. *Biotechnology* 10, 43-60.
- Sternberg, N. and Cohen, G. (1989). Genetic analysis of the lytic replicon of bacteriophage P1. II. Organization of replicon elements. *J Mol Biol* 207, 111-33.
- Tabor, S. and Richardson, C. C. (1987). DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc Natl Acad Sci U S A* 84, 4767-71.
- Takebayashi, H., Nabeshima, Y., Yoshida, S., Chisaka, O. and Ikenaka, K. (2002a). The basic helix-loop-helix factor olig2 is essential for the development of motoneuron and oligodendrocyte lineages. *Curr Biol* 12, 1157-63.
- Takebayashi, H., Ohtsuki, T., Uchida, T., Kawamoto, S., Okubo, K., Ikenaka, K., Takeichi, M., Chisaka, O. and Nabeshima, Y. (2002b). Non-overlapping expression of Olig3 and Olig2 in the embryonic neural tube. *Mech Dev* 113, 169-74.
- Takebayashi, H., Yoshida, S., Sugimori, M., Kosako, H., Kominami, R., Nakafuku, M. and Nabeshima, Y. (2000). Dynamic expression of basic helix-loop-helix Olig family members: implication of Olig2 in neuron and oligodendrocyte differentiation and identification of a new member, Olig3. *Mech Dev* 99, 143-8.
- Tan, K. and Le Douarin, N. M. (1991). Development of the nuclei and cell migration in the medulla oblongata. Application of the quail-chick chimera system. *Anat Embryol (Berl)* 183, 321-43.
- Tanabe, Y. and Jessell, T. M. (1996). Diversity and pattern in the developing spinal cord. *Science* 274, 1115-23.
- Trepel, C. and Racine, R. J. (1999). Blockade and disruption of neocortical long-term potentiation following electroconvulsive shock in the adult, freely moving rat. *Cereb Cortex* 9, 300-5.
- Turner, E. E., Jenne, K. J. and Rosenfeld, M. G. (1994). Brn-3.2: a Brn-3-related transcription factor with distinctive central nervous system expression and regulation by retinoic acid. *Neuron* 12, 205-18.

- Vaage, S. (1969). The segmentation of the primitive neural tube in chick embryos (*Gallus domesticus*). A morphological, histochemical and autoradiographical investigation. *Ergeb Anat Entwicklungsgesch* 41, 3-87.
- Vallstedt, A., Muhr, J., Pattyn, A., Pierani, A., Mendelsohn, M., Sander, M., Jessell, T. M. and Ericson, J. (2001a). Different levels of repressor activity assign redundant and specific roles to Nkx6 genes in motor neuron and interneuron specification. *Neuron* 31, 743-55.
- Vallstedt, A., Muhr, J., Pattyn, A., Pierani, A., Mendelsohn, M., Sander, M., Jessell, T. M. and Ericson, J. (2001b). Different levels of repressor activity assign redundant and specific roles to nkx6 genes in motor neuron and interneuron specification. *Neuron* 31, 743-55.
- Viemari, J. C., Bevengut, M., Burnet, H., Coulon, P., Pequignot, J. M., Tiveron, M. C. and Hilaire, G. (2004). Phox2a gene, A6 neurons, and noradrenaline are essential for development of normal respiratory rhythm in mice. *J Neurosci* 24, 928-37.
- Vogelstein, B. and Gillespie, D. (1979). Preparative and analytical purification of DNA from agarose. *Proc Natl Acad Sci U S A* 76, 615-9.
- von Baer, K. (1828). Über die Entwicklungsgeschichte der Tiere. Königsberg.
- Wang, V. Y., Rose, M. F. and Zoghbi, H. Y. (2005). Math1 expression redefines the rhombic lip derivatives and reveals novel lineages within the brainstem and cerebellum. *Neuron* 48, 31-43.
- Wassarman, K. M., Lewandoski, M., Campbell, K., Joyner, A. L., Rubenstein, J. L., Martinez, S. and Martin, G. R. (1997). Specification of the anterior hindbrain and establishment of a normal mid/hindbrain organizer is dependent on Gbx2 gene function. *Development* 124, 2923-34.
- Wetts, R. and Herrup, K. (1983). Direct correlation between Purkinje and granule cell number in the cerebella of lurcher chimeras and wild-type mice. *Brain Res* 312, 41-7.
- Wilson, S. I. and Edlund, T. (2001). Neural induction: toward a unifying mechanism. *Nat Neurosci* 4 Suppl, 1161-8.
- Wilson, S. I., Rydstrom, A., Trimborn, T., Willert, K., Nusse, R., Jessell, T. M. and Edlund, T. (2001). The status of Wnt signalling regulates neural and epidermal fates in the chick embryo. *Nature* 411, 325-30.
- Wingate, R. (2005). Math-Map(ic)s. *Neuron* 48, 1-4.
- Wingate, R. J. (2001). The rhombic lip and early cerebellar development. *Curr Opin Neurobiol* 11, 82-8.

- Wingate, R. J. and Hatten, M. E. (1999). The role of the rhombic lip in avian cerebellum development. *Development* 126, 4395-404.
- Wolpert, L. (1999). Entwicklungsbiologie. Spektrum Akademischer Verlag GmbH Heidelberg, Berlin.
- Yagi, T., Nada, S., Watanabe, N., Tamemoto, H., Kohmura, N., Ikawa, Y. and Aizawa, S. (1993). A novel negative selection for homologous recombinants using diphtheria toxin A fragment gene. *Anal Biochem* 214, 77-86.
- Yu, D., Ellis, H. M., Lee, E. C., Jenkins, N. A., Copeland, N. G. and Court, D. L. (2000). An efficient recombination system for chromosome engineering in Escherichia coli. *Proc Natl Acad Sci U S A* 97, 5978-83.
- Zanjani, H. S., Mariani, J. and Herrup, K. (1990). Cell loss in the inferior olive of the staggerer mutant mouse is an indirect effect of the gene. *J Neurogenet* 6, 229-41.
- Zhou, Q. and Anderson, D. J. (2002). The bHLH transcription factors OLIG2 and OLIG1 couple neuronal and glial subtype specification. *Cell* 109, 61-73.
- Zhou, Q., Choi, G. and Anderson, D. J. (2001). The bHLH Transcription Factor Olig2 Promotes Oligodendrocyte Differentiation in Collaboration with Nkx2.2. *Neuron* 31, 791-807.
- Zubair, M., Watanabe, E., Fukada, M. and Noda, M. (2002). Genetic labelling of specific axonal pathways in the mouse central nervous system. *Eur J Neurosci* 15, 807-14.