
6 Bibliographie

- [Anonym] (1999). "Merck Index 12". Whitehouse Station, NJ, Merck & Co., Inc.
- [Anonym] (2004). "Breakthrough of the Year - Runners Up". *Science* **306**: 2013-2017.
- Amado, A. M. und RibeiroClaro, P. J. A. (1997). "H/D and D/H exchange rates in alpha-cyclodextrin and alpha-cyclodextrin inclusion compounds - Raman spectroscopic study". *J. Chem. Soc.-Faraday Trans.* **93** (14): 2387-2390.
- Aree, T. (2000). "X-Ray Crystallographic and Neutron Scattering Studies on Hydration Dynamics and Solubility of Methylated Cyclodextrins". Diss. Berlin: FB Biologie, Chemie, Pharmazie, Freie Universität Berlin.
- Aree, T., Hoier, H., Schulz, B., Reck, G. und Saenger, W. (2000a). "Crystal structure of octakis(2,3,6-tri-O-methyl)-gamma-cyclodextrin x 4.5 H₂O: evidence for conformational flexibility of permethylated cyclodextrins". *Carbohydr. Res.* **328** (3): 399-407.
- Aree, T., Hoier, H., Schulz, B., Reck, G. und Saenger, W. R. (2000b). "Novel type of thermostable channel clathrate hydrate formed by heptakis(2,6-di-O-methyl)-beta-cyclodextrin·15H₂O - A paradigm of the hydrophobic effect". *Angew. Chem.-Int. Edit.* **39** (5): 897-899.
- Aree, T., Saenger, W., Leibnitz, P. und Hoier, H. (1999a). "Crystal structure of heptakis(2,6-di-O-methyl)-beta-cyclodextrin dihydrate: a water molecule in an apolar cavity". *Carbohydr. Res.* **315** (3-4): 199-205.
- Aree, T., Saenger, W., Leibnitz, P. und Hoier, H. (1999b). "Erratum to: Crystal structure of heptakis(2,6-di-O-methyl)-beta-cyclodextrin dihydrate: a water molecule in an apolar cavity (vol 315, pg 199, 1999)". *Carbohydr. Res.* **318** (1-4): 201.
- Aree, T., Uson, I., Schulz, B., Reck, G., Hoier, H., Sheldrick, G. M. und Saenger, W. (1999c). "Variation of a theme: Crystal structure with four octakis(2,3,6-tri-O-methyl)-gamma-cyclodextrin molecules hydrated differently by a total of 19.3 water". *J. Am. Chem. Soc.* **121** (14): 3321-3327.
- Atkins, P. W. (1990). "Physikalische Chemie". Weinheim: VCH.
- Badger, R. M. (1940). "The Relation Between the Energy of a Hydrogen Bond and the Frequencies of the O-H Bands". *J. Phys. Chem.* **8** (3): 288-289.
- Badger, R. M. und Bauer, S. H. (1937). "Spectroscopic Studies of the Hydrogen Bond. II. The Shift of the O-H Vibrational Frequency in the Formation of the Hydrogen Bond". *J. Chem. Phys.* **5** (11): 839-851.
- Ball, P. (1999). "H₂O: a Biography of Water". London: Weidenfeld & Nicholson.
- Ball, P. (2001). "H₂O: Biographie des Wassers". München Zürich: Piper Verlag.
- Battino, R. und Clever, H. L. (1966). "The Solubility of Gases in Liquids". *Chem. Rev.* **66** (4): 395 - 463.
- Bender, M. L. und Komiyama, M. (1978). "Cyclodextrin Chemistry". Berlin: Springer Verlag.

- Benedict, W. S., Gailar, N. und Plyer, E. K. (1956). "Rotation-Vibration Spectra of Deuterated Water Vapor". *J. Chem. Phys.* **24**: 1139-1165.
- Ben-Naim, A. und Marcus, Y. (1984). "Solvation thermodynamics of nonionic solutes". *J. Chem. Phys.* **81** (4): 2016-2027.
- Blokzijl, W. und Engberts, J. B. F. N. (1993a). "Hydrophobe Effekte - Ansichten und Tatsachen". *Angew. Chem.* **105**: 1610-1648.
- Blokzijl, W. und Engberts, J. B. F. N. (1993b). "Hydrophobic Effects - Opinions and Facts". *Angew. Chem.-Int. Edit.* **32** (11): 1545-1579.
- Bratos, S. (1975). "Profiles of hydrogen stretching ir bands of molecules with hydrogen bonds: A stochastic theory. I. Weak and medium strength hydrogen bonds". *J. Chem. Phys.* **63** (8): 3499-3509.
- Brooksby, P. A. und Fawcett, W. R. (2000). "Infrared (ATR) study of hydrogen bonding in solutions containing water and ethylene carbonate". *J. Phys. Chem. A* **104** (35): 8307-8314.
- Brügel (1995). "Vibrational spectroscopy of different classes and states of compounds - Infrared and Raman spectroscopy of biomolecules" in: *Infrared and raman spectroscopy: methods and applications*, Hg. B. Schrader: 344-371. Weinheim: VCH.
- Buch, V. und Devlin, J. P. (1999). "A new interpretation of the OH-stretch spectrum of ice". *J. Chem. Phys.* **110** (7): 3437-3443.
- Caira, M. R., Griffith, V. J., Nassimbeni, L. R. und Vanoudtshoorn, B. (1994). "Unusual ${}^1\text{C}_4$ Conformation of a Methylglucose Residue in Crystalline Permethy- β -Cyclodextrin Monohydrate". *J. Chem. Soc. Perk. T 2* (10): 2071-2072.
- Casu, B., Reggiani, M. und Sanderson, G. R. (1979). "Methylated cycloamyloses (cyclodextrins) and their inclusion properties". *Carbohydr. Res.* **76** (1): 59-66.
- Chaplin, M. F. (1999). "A proposal for the structuring of water". *Biophys. Chem.* **83** (3): 211-221.
- Chaplin, M. F. (2003). "Water structure and behavior". Internet-Resource. School of Applied Science - South Bank University. bes.: 16.04.2003. zul. geänd.: 01.04.2003.
- Christen, H. R. (1984). "Chemie". Frankfurt am Main: Diesterweg/Salle.
- Coleman, A. W., Nicolis, I., Keller, N. und Dalbiez, J. P. (1992). "Aggregation of Cyclodextrins: An Explanation of the Abnormal Solubility of β -Cyclodextrin". *J. Includ. Phenom. Mol.* **13**: 139-143.
- Collaborative Computational Project, N. (1994). "The CCP4 Suite: Programs for Protein Crystallography". *Acta Cryst. D* **50**: 760-763.
- Colthup, N. B., Daly, L. H. und Wiberley, S. E. (1990). "Introduction To Infrared And Raman Spectroscopy". San Diego, US: Academic Press, Inc.
- Cramer, F. (1954). "Einschlussverbindungen". Heidelberg: Springer-Verlag.

- D'Arrigo, G., Maisano, G., Mallamace, F., Migliardo, P. und Wanderlingh, F. (1981). "Raman scattering and structure of normal and supercooled water". *J. Chem. Phys.* **75** (9): 4264-4270.
- Davidson, D. (1973). "Clathrate Hydrates" in: *Water: A Comprehensive Treatise*, Hg. F. Franks. **2**: 115-234. New York: Plenum.
- Demtröder, W. (2003). "Laser Spectroscopy: Basic Concepts and Instrumentation". Berlin: Springer-Verlag.
- Dingley, A. J. und Grzesiek, S. (1998). "Direct Observation of Hydrogen Bonds in Nucleic Acid Base Pairs by Internucleotide $^2J_{\text{NN}}$ Couplings". *J. Am. Chem. Soc.* **120** (33): 8293-8297.
- Duchene, D., Hg. (1987). "Cyclodextrin and Their Industrial Uses" Paris: Editions de Santé.
- Dyke, T. R., Mack, K. M. und Muentner, J. S. (1977). "The structure of water dimer from molecular beam electric resonance spectroscopy". *J. Chem. Phys.* **66** (2): 498-510.
- Egyed, O. (1990). "Spectroscopic studies on β -cyclodextrin". *Vib. Spectrosc.* **1**: 225-227.
- Eisenberg, D. und Kauzmann, W. (1969). "The Structure and Properties of Water". Oxford: Oxford University Press.
- Evans, J. C. (1960). "Further studies of unusual effects in the infrared spectra of certain molecules". *Spectrochim. Acta* **16** (9): 994-1000.
- Evans, J. C. und Wright, N. (1960). "A peculiar effect in the infrared spectra of certain molecules". *Spectrochim. Acta* **16** (3): 352-357.
- Fletcher, N. H. (1970). "The Chemical Physics of Ice". Cambridge: Cambridge University Press.
- Folzer, C., Hendricks, R. W. und Narten, A. H. (1971). "Diffraction Pattern and Structure of Liquid Trimethylamine Decahydrate at 5°C". *J. Chem. Phys.* **54** (2): 799-805.
- Frank, H. S. und Evans, M. W. (1945). "Free Volume and Entropy in Condensed Systems III. Entropy in Binary Liquid Mixtures; Partial Molal Entropy in Dilute Solutions; Structure and Thermodynamics in Aqueous Electrolytes". *J. Chem. Phys.* **13** (11): 507-532.
- Frank, H. S. und Wen, W.-Y. (1957). "Ion-solvent interaction. Structural aspects of ion-solvent interaction in aqueous solutions: a suggested picture of water structure". *Discuss. Faraday Soc.* **24**: 133 - 140.
- Freudenberg, K. und Meyer-Delius, M. (1938). "Über die Schardinger-Dextrine aus Stärke". *Ber. Dtsch. Chem. Ges.* **71**: 1596-1600.
- Frömring, K. H. und Szejtli, J. (1994). "Cyclodextrin in Pharmacy". Dordrecht: Kluwer Academic Publishers.
- Gailar, N. M. und Dickey, F. P. (1960). "The vibration-rotation band ν_2 of HDO vapor". *J. Mol. Spectrosc.* **4** (1-6): 1-15.
- Gavira, J. M., Hernanz, A. und Bratu, I. (2003). "Dehydration of β -cyclodextrin: An IR $\nu(\text{OH})$ band profile analysis". *Vib. Spectrosc.* **32**: 137-146.

- Geiger, A., Rahman, A. und Stillinger, F. H. (1979). "Molecular dynamics study of the hydration of Lennard-Jones solutes". *J. Chem. Phys.* **70** (1): 263-276.
- George, W. O. und McIntyre, P. S. (1987). "Infrared spectroscopy". Chichester: John Wiley & Sons.
- Gerthsen, C. (1986). "Physik: ein Lehrbuch zum Gebrauch neben Vorlesungen". Berlin: Springer Vlg.
- Giguere, P. A. (1987). "The bifurcated hydrogen-bond model of water and amorphous ice". *J. Chem. Phys.* **87** (8): 4835-4839.
- Glew, D. N. und Rath, N. S. (1971). "H₂O, HDO, and CH₃OH Infrared Spectra and Correlation with Solvent Basicity and Hydrogen Bonding". *Can. J. Chem.* **49**: 837-856.
- Goutev, N., Nickolov, Z. S., Georgiev, G. und Matsuura, H. (1997). "Hydration of a short chain poly(oxyethylene) (C₁E₂C₁) studied by analysis of the O-H Raman band". *J. Chem. Soc.-Faraday Trans.* **93** (17): 3167-3171.
- Haken, H. und Wolf, H. C. (1994). "Molekülphysik und Quantenchemie: Einführung in die experimentellen und theoretischen Grundlagen". Berlin: Springer.
- Harata, K. (1991). "Recent Advances in the X-Ray Analysis of CD Complexes" in: *Inclusion Compounds: Key Organic Host Systems*, Hg. J. L. Atwood, J. E. D. Davies und D. D. MacNicol. **4**: 311-344. London: Academic Press.
- Harata, K. (1995). "X-Ray Structures of Hexakis(2,6-Di-O-Methyl)-Alpha-Cyclodextrin in 2 Crystal Forms". *Supramol. Chem.* **5** (3): 231-236.
- Harata, K. (1996). "Cristallographic Studies" in: *Comprehensive Supramolecular Chemistry*, Hg. J. L. Atwood, J. E. D. Davies, D. D. MacNicol und F. Vögtle. **3**: Ch. 9, 279-304. Oxford: Pergamon.
- Harata, K. (1998). "Structural aspects of stereodifferentiation in the solid state". *Chem. Rev.* **98** (5): 1803-1827.
- Harata, K., Uekama, K., Odagiri, M. und Hirayama, F. (1987). "Crystal Structures of Hexakis(2,3,6-tri-O-methyl)- α -Cyclodextrin Complexes with (S)-Mandelic and (R)-Mandelic Acid - Chiral Recognition Through the Induced-Fit Conformational Change of the Macrocyclic Ring". *B. Chem. Soc. Jpn.* **60**: 497-502.
- Hare, D. E. und Sorensen, C. M. (1992). "Interoscillator coupling effects on the OH stretching band of liquid water". *J. Chem. Phys.* **96** (1): 13-22.
- Herzberg, G. (1945). "Infrared and Raman Spectra of Polyatomic Molecules". Princeton, N.J.: Van Nostrand-Reinhold.
- Immel, S. und Lichtenthaler, F. W. (1996). "Per-O-methyl α - and β -CD: Cyclodextrins with Inverse Hydrophobicity". *Starch/Stärke* **48**: 225-232.
- Irvine, J. C., Pringsheim, H. und McDonald, J. (1924). "The Constitution of Polysaccharides, Part VIII. The Molecular Structure of β -Hexaamylose". *J. Am. Chem. Soc.* **125**: 942-947.

- Isaacs, E. D., Shukla, A., Platzman, P. M., Hamann, D. R., Barbiellini, B. und Tulk, C. A. (1999a). "Covalency of the Hydrogen Bond in Ice: A Direct X-Ray Measurement". *Phys. Rev. Lett.* **82** (3): 600–603.
- Isaacs, E. D., Shukla, A., Platzman, P. M., Hamann, D. R., Barbiellini, B. und Tulk, C. A. (1999b). "Erratum: Covalency of the Hydrogen Bond in Ice: A Direct X-Ray Measurement [Phys. Rev. Lett. 82, 600 (1999)]". *Phys. Rev. Lett.* **83** (21): 4445.
- Issahary, D., Zolotov, S., Hayon, Y., Fuchs, J. und Sarig, S. (1993). "The Structure of Concentrated Magnesium-Chloride Water Solutions". *Rev. Chem. Eng.* **9** (3-4): 253-265.
- Jeffrey, G. A. (1996). "Hydrate Inclusion Compounds" in: *Comprehensive Supramolecular Chemistry*, Hg. J. L. Atwood, J. E. D. Davies, D. D. MacNicol und F. Vögtle. **6**: 757-788. Oxford: Pergamon.
- Jicsinszky, L., Hashimoto, H., Fenyvesi, E. und Ueno, A. (1996). "Cyclodextrin derivatives" in: *Comprehensive Supramolecular Chemistry*, Hg. J. L. Atwood, J. E. D. Davies, D. D. MacNicol und F. Vögtle. **3**: Ch. 4. Oxford: Pergamon.
- Kaltofen, R. E. (1982). "Tabellenbuch Chemie". Leipzig: VEB Deutscher Verlag für Grundstoffindustrie.
- Kanno, H., Tomikawa, K. und Mishima, O. (1998). "Raman spectra of low- and high-density amorphous ices". *Chem. Phys. Lett.* **293** (5-6): 412-416.
- Koza, M., Schober, H., Tolle, A., Fujara, F. und Hansen, T. (1999). "Formation of ice XII at different conditions". *Nature* **397** (6721): 660-661.
- Laenen, R., Rauscher, C. und Laubereau, A. (1998). "Local substructures of water studied by transient hole-burning spectroscopy in the infrared: Dynamics and temperature dependence". *J. Phys. Chem. B* **102** (46): 9304-9311.
- Lappi, S. E., Smith, B. und Franzen, S. (2004). "Infrared spectra of H₂¹⁶O, H₂¹⁸O and D₂O in the liquid phase by single-pass attenuated total internal reflection spectroscopy". *Spectrochim. Acta A* **60** (11): 2611-2619.
- Latscha, H. P. und Klein, H. A. (1993). "2. Organische Chemie". Berlin: Springer-Verlag.
- Lee, B. und Richards, F. M. (1971). "The interpretation of protein structures: Estimation of static accessibility". *J. Mol. Biol.* **55**: 379 - 400.
- Lehninger, A. L. (1994). "Prinzipien der Biochemie". Heidelberg.
- Libnau, F. O., Toft, J., Christy, A. A. und Kvalheim, O. M. (1994). "Structure of Liquid Water Determined From Infrared Temperature Profiling and Evolutionary Curve Resolution". *J. Am. Chem. Soc.* **116** (18): 8311-8316.
- Lichtenthaler, F. W. und Immel, S. (1996a). "The Lipophilicity Patterns of cyclodextrins and of Non-glucose Cyclooligosaccharides". *J. Inclus. Phenom. Mol.* **25**: 3-16.
- Lichtenthaler, F. W. und Immel, S. (1996b). "On the Hydrophobicity Characteristics of Cyclodextrins: Computer-Aided Visualization of Molecular Lipophilicity Patterns". *Liebigs Ann. Chem.*: 27-37.

- Lobban, C., Finney, J. L. und Kuhs, W. F. (1998). "The structure of a new phase of ice". *Nature* **391** (6664): 268-270.
- Lord, R. C. und Merrifield, R. E. (1953). "Strong Hydrogen Bonds in Crystals". *J. Chem. Phys.* **21** (1): 166-167.
- Loubeyre, P., LeToullec, R., Wolanin, E., Hanfland, M. und Hausermann, D. (1999). "Modulated phases and proton centring in ice observed by X-ray diffraction up to 170 GPa". *Nature* **397** (6719): 503-506.
- Luck, W. A. P. und Fritzsche, M. (1995). "The OH Stretching Mode and a 3 Species Model of Liquid Methanol". *Z. Phys. Chemie-Int. J. Res. Phys. Chem. Chem. Phys.* **191**: 71-86.
- Ludwig, R. (2001). "Wasser: von Clustern in die Flüssigkeit". *Angew. Chem.* **113**: 1856-1876.
- MacLennan, J. M. und Stezowski, J. J. (1980). "Crystal-Structure of Uncomplexed-Hydrated Cyclooctaamylose". *Biochem. Bioph. Res. Co.* **92** (3): 926-932.
- Madan, B. und Sharp, K. (1996). "Heat Capacity Changes Accompanying Hydrophobic and Ionic Solvation: A Monte Carlo and Random Network Model Study". *J. Phys. Chem.* **100** (18): 7713-7721.
- Maeda, Y., Ide, M. und Kitano, H. (1999). "Vibrational spectroscopic study on the structure of water in polymer systems". *J. Mol. Liq.* **80** (2-3): 149-163.
- Manor, P. C. und Saenger, W. (1974). "Topography of cyclodextrin inclusion complexes. III. Crystal and molecular structure of cyclohexaamylose hexahydrate, the water dimer inclusion complex". *J. Am. Chem. Soc.* **96**: 3630-3639.
- Marechal, Y. (1991). "Infrared-Spectra of Water.1. Effect of Temperature and of H/D Isotopic Dilution". *J. Chem. Phys.* **95** (8): 5565-5573.
- Marechal, Y. (2003). "Observing the water molecule in macromolecules and aqueous media using infrared spectrometry". *J. Mol. Struct.* **648** (1-2): 27-47.
- Martin, T. W. und Derewenda, Z. S. (1999). "The name is bond - H bond". *Nat. Struct. Biol.* **6** (5): 403-406.
- Meng, E. C. und Kollman, P. A. (1996). "Molecular Dynamics Studies of the Properties of Water around Simple Organic Solutes". *J. Phys. Chem.* **100** (27): 11460-11470.
- Mentzafos, D., Mavridis, I. M., Lebas, G. und Tsoucaris, G. (1991). "Structure of the 4-Tert-Butylbenzyl Alcohol Beta-Cyclodextrin Complex - Common Features in the Geometry of Beta-Cyclodextrin Dimeric Complexes". *Acta Crystallogr. B* **47**: 746-757.
- Nickolov, Z. S., Ohno, K. und Matsuura, H. (1999). "FTIR-ATR studies of the hydration of 15-crown-5 and 18-crown-6 in aqueous solutions". *J. Phys. Chem. A* **103** (37): 7544-7551.
- Nielsen, O. F. (1993). "Low-Frequency Spectroscopic Studies of Interactions in Liquids". *Annu. Rep. Prog. Chem., Sect. C, Phys. Chem.* **90**: 3-44.
- Nielsen, O. F. (1997). "Low-Frequency spectroscopic studies and intermolecular vibrational energy transfer in liquids". *Annu. Rep. Prog. Chem., Sect. C, Phys. Chem.* **93**: 57-99.
- Novak, A. (1974). "Hydrogen bonding in solids". *Struct. Bonding* **18**: 177-216.

- Odutola, J. A. und Dyke, T. R. (1980). "Partially deuterated water dimers: Microwave spectra and structure". *J. Chem. Phys.* **72** (9): 5062-5070.
- Pye, C. C. und Rudolph, W. W. (1998). "An ab initio and Raman investigation of magnesium(II) hydration". *J. Phys. Chem. A* **102** (48): 9933-9943.
- Reed, A. E., Curtiss, L. A. und Weinhold, F. (1988). "Intermolecular interactions from a natural bond orbital, donor-acceptor viewpoint". *Chem. Rev.* **88** (6): 899 - 926.
- Reich, R. (1993). "Thermodynamik, Grundlagen und Anwendungen in der allgemeinen Chemie". Weinheim: VCH.
- Rekharsky, M. V. und Inoue, Y. (1998). "Complexation Thermodynamics of Cyclodextrins". *Chem. Rev.* **98**: 1875-1917.
- Rice, S. A., Bergren, M. S., Belch, A. C. und Nielsen, G. (1983). "A theoretical analysis of the hydroxyl stretching spectra of ice Ih, liquid water, and amorphous solid water". *J. Phys. Chem.* **87** (21): 4295 - 4308.
- Römpp (1997). "Lexikon Chemie - Version 1.3". CD-ROM. Georg Thieme Verlag.
- Rundle, R. E. und Parasol, M. (1952). "O-H Stretching Frequencies in Very Short and Possibly Symmetrical Hydrogen Bonds". *J. Phys. Chem.* **20** (9): 1487-1488.
- Saenger, W. (1979). "Circular Hydrogens Bonds". *Nature* **279**: 343-344.
- Saenger, W. (1980a). "Cyclodextrin Inclusion-Compounds in Research and Industry". *Angew. Chem.-Int. Edit. Engl.* **19** (5): 344-362.
- Saenger, W. (1980b). "Cyclodextrin-Einschlußverbindungen in Forschung und Industrie". *Angew. Chem.* **92** (5): 343-361.
- Saenger, W. R., Jacob, J., Gessler, K., Steiner, T., Hoffmann, D., Sanbe, H., Koizumi, K., Smith, S. M. und Takaha, T. (1998). "Structures of the common cyclodextrins and their larger analogues - Beyond the doughnut". *Chem. Rev.* **98** (5): 1787-1802.
- Sammon, C., Mura, C., Yarwood, J., Everall, N., Swart, R. und Hodge, D. (1998). "FTIR-ATR studies of the structure and dynamics of water molecules in polymeric matrixes. A comparison of PET and PVC". *J. Phys. Chem. B* **102** (18): 3402-3411.
- Sceats, M. G. und Rice, S. A. (1982). "Amorphous solid water and its relationship to liquid water: a random network model for water" in: *Water: A Comprehensive Treatise*, Hg. F. Franks. **7**: 83. New York: Plenum.
- Schardinger, F. (1903). "Über thermophile Bakterien aus verschiedenen Speisen und Milch sowie über einige Umsetzungsprodukte derselben in kohlenhydrathaltigen Nährlösungen, darunter kristallisierte Polysaccharide (Dextrine) aus Stärke". *Z. Unters. Nahr. u. Genussm.* **6**: 865-880.
- Scherer, J. R. (1978). "The Vibrational Spectroscopy of Water, Chapter 3" in: *Advances in Infrared and Raman Spectroscopy*, Hg. R. J. H. Clark und R. E. Hester: 149-216. Philadelphia: Heyden.
- Scherer, J. R., Go, M. K. und Kint, S. (1973). "Raman spectra and structure of water in dimethyl sulfoxide". **77** (17): 2108-2117.

- Scherer, J. R., Go, M. K. und Kint, S. (1974). "Raman spectra and structure of water from -10 to 90.deg". *J. Phys. Chem.* **78** (13): 1304.
- Schmid, R. (2001). "Recent advances in the description of the structure of water, the hydrophobic effect, and the like-dissolves-like rule". *Mon. Chem.* **132** (11): 1295-1326.
- Schrader, B. (1995). "General survey of vibrational spectroscopy" in: *Infrared and Raman spectroscopy: methods and applications*, Hg. B. Schrader: 7-62. Weinheim: VCH.
- Sciortino, F., Geiger, A. und Stanley, H. E. (1992). "Network defects and molecular mobility in liquid water". *J. Chem. Phys.* **96** (5): 3857-3865.
- Sharp, K. A., Madan, B., Manas, E. und Vanderkooi, J. M. (2001). "Water structure changes induced by hydrophobic and polar solutes revealed by simulations and infrared spectroscopy". *J. Chem. Phys.* **114** (4): 1791-1796.
- Silverstein, K. A. T., Haymet, A. D. J. und Dill, K. A. (2000). "The strength of hydrogen bonds in liquid water and around nonpolar solutes". *J. Am. Chem. Soc.* **122** (33): 8037-8041.
- Steiner, T., Hirayama, F. und Saenger, W. (1996). "Crystal structures of hexakis-(2,6-di-O-methyl)-cyclomaltohexaose (dimethyl-alpha-cyclodextrin) crystallized from acetone, and crystallized from hot water". *Carbohydr. Res.* **296**: 69-82.
- Steiner, T. und Saenger, W. (1995). "Crystal-Structure of Anhydrous Heptakis-(2,6-Di-O-Methyl) Cyclomaltoheptaose (Dimethyl-Beta-Cyclodextrin)". *Carbohydr. Res.* **275** (1): 73-82.
- Steiner, T. und Saenger, W. (1996). "Crystal structure of anhydrous hexakis-(2,3,6-tri-O-methyl)-cyclomaltohexaose (permethyl-alpha-cyclodextrin) grown from hot water and from cold NaCl solutions". *Carbohydr. Res.* **282** (1): 53-63.
- Steiner, T. und Saenger, W. (1998a). "Closure of the cavity in permethylated cyclodextrins through glucose inversion, flipping, and kinking". *Angew. Chem.-Int. Edit.* **37** (24): 3404-3407.
- Steiner, T. und Saenger, W. (1998b). "Hohlraumverkleinerung in permethylierten Cyclodextrinen durch Inversion, Drehen und Kippen der Glucosebausteine". *Angew. Chem.* **110** (24): 3628-3632.
- Szejtli, J. (1982). "Cyclodextrins and their Inclusion Complexes". Budapest: Akademiai Kiado.
- Szejtli, J. (1988). "Cyclodextrin Technology". Dordrecht.
- Szejtli, J. (1992). "The Properties and Potential Uses of Cyclodextrin Derivatives". *J. Includ. Phenom. Mol.* **14**: 25-36.
- Takaha, T., Yanase, M., Takata, H., Okada, S. und Smith, S. M. (1996). "Potato D-Enzyme Catalyzes the Cyclation of Amylose to Produce Cycloamylose, a Novel Cyclic Glucan". *J. Biol. Chem.* **271**: 2902-2908.
- Tanford, C. (1997). "How protein chemists learned about the hydrophobic factor". *Protein Sci.* **6** (6): 1358-1366.
- Tursi, A. J. und Nixon, E. R. (1970). "Matrix-Isolation Study of the Water Dimer in Solid Nitrogen". *J. Chem. Phys.* **52**: 1521-1528.

- Udachin, K. A. und Ripmeester, J. A. (1999). "A complex clathrate hydrate structure showing bimodal guest hydration". *Nature* **397** (6718): 420-423.
- Uekama, K. und Irie, T. (1987). "Pharmaceutical Applications of Methylated Cyclodextrin Derivatives" in: *Cyclodextrins and Their Industrial Uses*, Hg. D. Duchêne: 395-439. Paris: Editions de Santé.
- Venyaminov, S. und Prendergast, F. G. (1997). "Water (H₂O and D₂O) molar absorptivity in the 1000-4000 cm⁻¹ range and quantitative infrared spectroscopy of aqueous solutions". *Anal. Biochem.* **248** (2): 234-245.
- Voet, D. und Voet, J. G. (1992). "Biochemie". Weinheim: VCH.
- Walrafen, G. E. (1968). "Raman Spectral Studies of HDO in H₂O". *J. Chem. Phys.* **48** (1): 244-251.
- Walrafen, G. E. (1974). "Structure of Water and Aqueous Solutions". Weinheim: Verlag Chemie.
- Walrafen, G. E., Hokmabadi, M. S. und Yang, W. H. (1986). "Raman isosbestic points from liquid water". *J. Chem. Phys.* **85** (12): 6964-6969.
- Walrafen, G. E., Yang, W.-H., Chu, Y. C. und Hokmabadi, M. S. (1996). "Raman OD-Stretching Overtone Spectra from Liquid D₂O between 22 and 152 C". *J. Phys. Chem.* **100** (4): 1381-1391.
- Welfle, H. (1996). "Optische Spektroskopie" in: *Methoden in der Proteinanalytik*, Hg. M. Holtzauer: 70-170. Berlin: Springer-Verlag.
- Wenz, G. (1994). "Cyclodextrins As Building-Blocks For Supramolecular Structures and Functional Units". *Angew. Chem.-Int. Edit. Engl.* **33** (8): 803-822.
- Whalley, E. (1977). "A detailed assignment of the O-H stretching bands of ice I". *Can. J. Chem.* **55** (19): 3429-3441.