

8. Literatur

- Abdalla S, Weatherall DJ, Wickramasinghe S, Hughes M. The anaemia of *Plasmodium falciparum* malaria. *Br J Haematol* 1980; 46: 171-183.
- Abu-Zeid YA, Abdulhadi NH, Hviid L, Theander TG, Saeed BO, Jepsen S, Jensen JB, Bayoumi RA. Lymphoproliferative responses to *Plasmodium falciparum* antigens in children with and without the sickle cell trait. *Scand J Immunol* 1991; 34: 237-242.
- Achidi EA, Perlmann H, Salimonu LS, Perlmann P, Walker O, Asuzu MC. A longitudinal study of seroreactivities to *Plasmodium falciparum* antigens in Nigerian infants during their first year of life. *Acta Trop* 1995; 59: 173-183.
- Afari EA, Nakano T, Binka F, Owusu-Agyei S, Asigbee J. Seasonal characteristics of malaria infection in under-five children of a rural community in southern Ghana. *W Afr J Med* 1993; 12: 39-42.
- Agarwal A, Guindo A, Cissoko Y, Taylor JG, Coulibaly D, Koné A, Kayentao K, Djimde A, Plowe CV, Doumbo O, Wellemes TE, Diallo D. Hemoglobin C associated with protection from severe malaria in the Dogon of Mali, a West African population with a low prevalence of hemoglobin S. *Blood* 2000; 96: 2358-2363.
- Aidoo M, McElroy PD, Kolczak MS, Terlouw DJ, ter Kuile FO, Nahlen B, Lal AA, Udhayakumar V. Tumor necrosis factor-alpha promoter variant 2 (TNF2) is associated with pre-term delivery, infant mortality, and malaria morbidity in western Kenya: Asembo Bay Cohort Project IX. *Genet Epidemiol* 2001; 21: 201-211.
- Aidoo M, Terlouw DJ, Kolczak MS, McElroy P, ter Kuile FO, Kariuki S, Nahlen BL, Lal AA, Udhayakumar V. Protective effects of the sickle cell gene against malaria morbidity and mortality. *Lancet* 2002; 359: 1311-1312.
- Alfrangis M, Lemnge MM, Moon R, Theisen M, Bygbjerg I, Ridley RG, Jakobsen PH. IgG reactivities against recombinant Rhoptry-Associated Protein-1 (rRAP-1) are associated with mixed *Plasmodium* infections and protection against disease in Tanzanian children. *Parasitology* 1999; 119: 337-342.
- Allen RA, Lee EM, Roberts DH, Park BK, Pirmohamed M. Polymorphisms in the TNF-alpha and TNF-receptor genes in patients with coronary artery disease. *Eur J Clin Invest* 2001; 31: 843-851.
- Allen SJ, Rowe P, Allsopp CEM, Riley EM, Jakobsen PH, Hill AVS, Greenwood BM. A prospective study of the influence of α -thalassaemia on morbidity from malaria and immune responses to defined *Plasmodium falciparum* antigens in Gambian children. *Trans R Soc Trop Med Hyg* 1993; 87: 282-285.
- Allen SJ, O'Donnell A, Alexander ND, Alpers MP, Peto TEA, Clegg JB, Weatherall JB. alpha+-Thalassemia protects children against disease caused by other infections as well as malaria. *Proc Natl Acad Sci U S A* 1997; 94: 14736-14741.
- Allison AC. The distribution of the sickle cell trait in Africa and elsewhere, and its apparent relationship to the incidence of subtertian malaria. *Trans R Soc Trop Med Hyg* 1954a; 48: 312-318.
- Allison AC. Protection afforded by the sickle cell trait against subtertian malarial infection. *Br Med J* 1954b; I: 290-294.
- Allison AC. The sickle-cell and haemoglobin C genes in some African populations. *Ann Hum Genet* 1956; 21: 67-89.
- Allison AC. Polymorphism and natural selection in human populations. *Cold Spring Harb Symp Quant Biol* 1964; 29: 137-149.
- Alonso PL, Lindsay SW, Armstrong Schellenberg JR, Gomez P, Hill AG, David PH, Fegan G, Cham K, Greenwood BM. A malaria control trial using insecticide-treated bed nets and targeted chemoprophylaxis in a rural area of The Gambia, west Africa. 2. Mortality and

- morbidity from malaria in the study area. *Trans R Soc Trop Med Hyg* 1993;87 (Suppl 2): 13-17.
- al-Yaman F, Genton B, Reeder JC, Anders RF, Smith T, Alpers MP. Reduced risk of clinical malaria in children infected with multiple clones of *Plasmodium falciparum* in a highly endemic area: a prospective community study. *Trans R Soc Trop Med Hyg* 1997; 91: 602-605.
- Angus BJ, Chotivanich K, Udomsangpetch R, White NJ. *In vivo* removal of malaria parasites from red blood cells without their destruction in acute falciparum malaria. *Blood* 1997; 90, 2037-2040.
- Anstey NM, Weinberg JB, Hassanali MY, Mwaikambo ED, Manyenga D, Misukonis MA, Arnelle DR, Hollis D, McDonald MI, Granger DL. Nitric oxide in Tanzanian children with malaria: inverse relationship between malaria severity and nitric oxide production/nitric oxide synthase type 2 expression. *J Exp Med* 1996;184: 557-567.
- Ariey F, Hommel D, Le Scanf C, Duchemin JB, Peneau C, Hulin A, Sarthou JL, Reynes JM, Fandeur T, Mercereau-Puijalon O. Association of severe malaria with a specific *Plasmodium falciparum* genotype in French Guiana. *J Infect Dis* 2001; 184: 237-241.
- Baruch DI, Pasloske BL, Singh HB, Bi X, Ma XC, Feldman M, Taraschi TF, Howard RJ. Cloning the *P. falciparum* gene encoding PfEMP1, a malarial variant antigen and adherence receptor on the surface of parasitized human erythrocytes. *Cell* 1995; 82: 77-87.
- Babiker HA, Abdel-Muhsin AMA, Ranford-Cartwright LC, Satti G, Walliker D. Characteristics of *Plasmodium falciparum* parasites that survive the lengthy dry season in Eastern Sudan where malaria transmission is markedly seasonal. *Am J Trop Med Hyg* 1998; 59: 582-590.
- Babiker HA, Ranford-Cartwright LC, Walliker D. Genetic structure and dynamics of *Plasmodium falciparum* infections in the Kilombero region of Tanzania. *Trans R Soc Trop Med Hyg* 1999; 93 (Suppl. 1): 11-14.
- Beck HP, Felger I, Huber W, Steiger S, Smith T, Weiss N., Alonso P, Tanner, M. Analysis of multiple *Plasmodium falciparum* infections in Tanzanian children during the phase III trial of the malaria vaccine SPf66. *J Infect Dis* 1997; 175: 921-926.
- Beck HP, Felger I, Vounatsou P, Hirt R, Tanner M, Alonso P, Menendez C. Effect of iron supplementation and malaria prophylaxis in infants on *Plasmodium falciparum* genotypes and multiplicity of infection. *Trans R Soc Trop Med Hyg* 1999; 93 (Suppl. 1): 41-44.
- Beeson JG, Brown GV, Molineaux ME, Mhango C, Dzinjalama F, Rogerson SJ. *Plasmodium falciparum* isolates from infected pregnant women and children are associated with distinct adhesive and antigenic properties. *J Infect Dis* 1999; 180: 464-472.
- Beeson JG, Rogerson SJ, Cooke BM, Reeder JC, Chai W, Lawson AM, Molyneux ME, Brown GV. Adhesion of *Plasmodium falciparum*-infected erythrocytes to hyaluronic acid in placental malaria. *Nature Med* 2000; 6: 86-90.
- Bendixen M, Msangeni HA, Pedersen BV, Shayo D, Bodker R. Diversity of *Plasmodium falciparum* populations and complexity of infections in relation to transmission intensity and host age: a study from the Usambara Mountains, Tanzania. *Trans R Soc Trop Med Hyg* 2001; 95: 143-148.
- Berendt AR, Turner GDH, Newbold CI. Cerebral malaria: the sequestration hypothesis. *Parasitol Today* 1994; 10: 412-414.
- Beutler E, Yeh M, Fairbanks VF. The normal human female as a mosaic of X chromosome activity. Studies using the gene for G-6-PD deficiency as a marker. *Proc Natl Acad Sci U S A* 1962; 48: 9-16.

- Bienzle U, Ayeni O, Lucas AO, Luzzatto L. Glucose-6-phosphate dehydrogenase and malaria. Greater resistance of females heterozygous for enzyme deficiency and of males with non-deficient variant. *Lancet* 1972; 1: 107-110.
- Binka FN, Morris SS, Ross DA, Arthur P, Aryeetey ME. Patterns of malaria morbidity and mortality in children in northern Ghana. *Trans R Soc Trop Med Hyg* 1994; 88: 381-385.
- Black J, Hommel M, Snounou G, Pinder M. Mixed infections with *Plasmodium falciparum* and *P. malariae* and fever in malaria. *Lancet* 1994; 343: 1095.
- Blampain-Azzibrouck, Lekoulou F, Snounou G, Ravollet JC, Ntoumi F. Short communication: *Plasmodium falciparum* and *P. malariae* infections in isolates from sickle cell gene carriers living in a hyperendemic area of Gabon. *Trop Med Int Health* 1999; 4: 872-874.
- Bloland PB, Kazembe PN, Oloo AJ, Himonga B, Barat LM, Ruebush TK. Chloroquine in Africa: critical assessment and recommendations for monitoring and evaluating chloroquine therapy efficacy in sub-Saharan Africa. *Trop Med Int Health* 1998; 3: 543-552.
- Bottius E, Guanzirolli A, Trape JF, Rogier C, Konate L, Druilhe P. Malaria: even more chronic in nature than previously thought; evidence for subpatent parasitaemia detectable by the polymerase chain reaction. *Trans R Soc Trop Med Hyg* 1996; 90: 15-19.
- Bouharoun-Tayoun H, Attanath P, Sabchareon A, Chongsuphajaisiddhi T, Druilhe P. Antibodies that protect humans against *Plasmodium falciparum* blood stages do not on their own inhibit parasite growth and invasion *in vitro*, but act in cooperation with monocytes. *J Exp Med* 1990; 172: 1633-1641.
- Brabin BJ. An analysis of malaria in pregnancy in Africa. *Bull World Health Organ* 1983; 61: 1005-1016.
- Brabin BJ, Perrin L. Sickle cell trait and *Plasmodium falciparum* parasitaemia in pregnancy in Western Province, Kenya. *Trans R Soc Trop Med Hyg* 1985; 79: 733.
- Brabin B. An analysis of malaria parasite rates in infants: 40 years after McDonald. *Trop Dis Bull* 1990; 87: R1-R21.
- Brair ME, Brabin BJ, Milligan P, Maxwell S, Hart CA. Reduced transfer of tetanus antibodies with placental malaria. *Lancet* 1994; 343: 208-209.
- Branch OH, Takala S, Kariuki S, Nahlen BL, Kolczak M, Hawley W, Lal AA. *Plasmodium falciparum* genotypes, low complexity of infection, and resistance to subsequent malaria in participants in the Asembo Bay Cohort Project. *Infect Immun* 2001; 69: 7783-7792.
- Brandling-Bennett AD, Oloo AJ, Watkins WM, Boriga DA, Kariuki DM, Collins WE. Chloroquine treatment of falciparum malaria in an area of Kenya of intermediate chloroquine resistance. *Trans R Soc Trop Med Hyg* 1988; 82: 833-837.
- Browne EN, Frimpong E, Sievertsen J, Hagen J, Hamelmann C, Dietz K, Horstmann RD, Burchard GD. Malariaometric update for the rainforest and savanna of Ashanti region, Ghana. *Ann Trop Med Parasitol* 2000; 94: 15-22.
- Bruce MC, Donelly CA, Alpers MP, Galinski MR, Barnwell JW, Walliker D, Day KP. Cross-species interactions between malaria parasites in humans. *Science* 2000; 287: 845-848.
- Bull PC, Lowe BS, Kortok M, Molyneux CS, Newbold CI, Marsh K. Parasite antigens on the infected red cell are targets for naturally acquired immunity to malaria. *Nature Med* 1998; 4: 356-360.
- Bulmer JN, Rasheed FN, Francis N, Morrison L, Greenwoods BM. Placental malaria. 1. Pathological classification. *Histopathol* 1993; 22: 211-218.
- Burchard GD, Radloff P, Philipps J, Nkeyi M, Knobloch J, Kremsner PG. Increased erythropoietin production in children with severe malarial anaemia. *Am J Trop Med Hyg* 1995; 53: 547-551.

- Burgner D, Xu W, Rockett K, Gravenor M, Charles IG, Hill AV, Kwiatkowski D. Inducible nitric oxide synthase polymorphism and fatal cerebral malaria. *Lancet* 1998; 352: 1193–1194.
- Burgner D, Rockett K, Ackermann H, Hull J, Usen S, Pinder M, Kwiatkowski DP. Haplotypic relationship between SNP and microsatellite markers of the NOS2A locus in two populations. *Genes Immun* 2003; 4: 506-514.
- Camacho LH, Gordeuk VR, Wilairatana P, Brittenham GM, Looraseesawan S. The course of anaemia after the treatment of acute falciparum malaria. *Ann Trop Med Parasitol* 1998; 5: 525-537.
- Cappadoro M, Giribaldi G, O'Brien E, Turrini F, Mannu F, Ulliers D, Simula G, Luzzatto L, Arese P. Early phagocytosis of glucose-6-phosphate dehydrogenase (G6PD)-deficient erythrocytes parasitized by *Plasmodium falciparum* may explain malaria protection in G6PD deficiency. *Blood* 1998; 92: 2527-2534.
- Carlier Y, Truyens C. Influence of maternal infection on offspring resistance towards parasites. *Parasitol Today* 1995; 5: 525-537.
- Carlson J, Nash GB, Gabutti V, Al-Yaman F, Wahlgren M. Natural protection against severe *Plasmodium falciparum* malaria due to impaired rosette formation. *Blood* 1994; 84: 3909-3914.
- Chen Q, Fernandez V, Sundstrom A, Schlichtherle M, Datta S, Hagblom P, Wahlgren M. Developmental selection of *var* gene expression in *Plasmodium falciparum*. *Nature* 1998; 394: 392-395.
- Ciuca M, Baluf L, Chelarescu-Vierum. Immunity in malaria. *Trans R Soc Trop Med Hyg* 1934; 27: 619-622.
- Clark IA, Chaudhri G. Tumor necrosis factor may contribute to the anaemia of malaria by causing dyserythropoiesis and erythrophagocytosis. *Br J Haematol* 1988; 70: 99–103.
- Clark IA, Cowden WB. The pathophysiology of falciparum malaria. *Pharmacol Therapeutics* 2003; 99: 221-160.
- Clendennen TE, Long GW, Baird JK. QBC and Giemsa-stained thick blood films: diagnostic performance of laboratory technologists. *Trans R Soc Trop Med Hyg* 1995; 89: 183-184.
- Cohen S, McGregor IA, Carrington S. Gamma globulin and acquired immunity to human malaria. *Nature* 1961; 192: 735-737.
- Collins WE, Jeffery GM. A retrospective examination of sporozoite- and trophozoite-induced infections with *Plasmodium falciparum* in patients previously infected with heterologous species of *Plasmodium*: effect on development of parasitologic and clinical immunity. *Am J Trop Med Hyg* 1999; 61S: 36-43.
- Contamin H, Fandeur T, Rogier C, Bonnefoy S, Konate L, Trape JF, Mercereau-Puijalon O. Different genetic characteristics of *Plasmodium falciparum* isolates collected during successive clinical malaria episodes in Senegalese children. *Am J Trop Med Hyg* 1996; 54: 632.
- Cox MJ, Kum DE, Tavul L, Raiko A, Baisor M, Alpers MP, Medley GF, Day KP. Dynamics of malaria parasitaemia associated with febrile illness in children from a rural area of Madang, Papua New Guinea. *Trans R Soc Trop Med Hyg* 1994; 88: 191-197.
- Cranston HA, Boylan CW, Carroll GL, Sutera SP, Williamson JR, Gluzman IY, Krogstad DJ. *Plasmodium falciparum* maturation abolishes physiologic red cell deformability. *Science* 1984; 223: 400-403.
- Curtis CF. Impregnated bednets, malaria control and child mortality in Africa. *Trop Med Int Health* 1996; 1:137-138.
- Curtis J, Duraisingh MT, Trigg JK, Mbwana H, Warhurst DC, Curtis CF. Direct evidence that asparagine at position 108 of the *Plasmodium falciparum* dihydrofolate reductase is

- involved in resistance to antifolate drugs in Tanzania. *Trans R Soc Trop Med Hyg* 1996; 90: 678-680.
- Day NPJ, Hien TT, Schollaardt T, Loc PP, Chuong LV, Chau TTH, Mai NTH, Phu NH, Sinh DX, White NJ, Ho M. The prognostic and pathophysiologic role of pro- and antiinflammatory cytokines in severe malaria. *J Infect Dis* 1999; 180: 1288-1297.
- De Caterina R, Libby P, Peng HB, Thannickal VJ, Rajavashist TB, Gimbrone MA Jr, Shin WS, Liao JK. Nitric oxide decreases cytokine-induced endothelial activation: nitric oxide selectively reduces endothelial expression of adhesion molecules and proinflammatory cytokines. *J Clin Invest* 1995; 96: 60-68.
- de Jong, Westendorp RGJ, Bakker AM, Huizinga TWJ. Polymorphisms in or near tumour necrosis factor(TNF)-gene do not determine levels of endotoxin-induced TNF production. *Gen Immun* 2002; 3: 25-29.
- Desowitz RS, Alpers MP. Placental *Plasmodium falciparum* parasitaemia in East Sepik (Papua New Guinea) women of different parity: the apparent absence of acute effects on mother and foetus. *Ann Trop Med Parasitol* 1992; 86: 95-102.
- Destro-Bisol G. Genetic resistance to malaria, oxidative stress and hemoglobin oxidation. *Parassitologia* 1999; 41: 203-204.
- Diagne N, Rogier C, Cisse B, Trape JF. Incidence of clinical malaria in pregnant women exposed to intense perenniel transmission. *Trans R Soc Trop Med Hyg* 1997; 91: 166-170.
- Diagne N, Rogier C, Sokhna C, Tall A, Fontenille D, Roussilhon C, Spiegel A, Trape JF. Increased susceptibility to malaria during the early postpartum period. *N Engl J Med* 2000; 342: 598-603.
- Diggs LM. In: Abramson H, Bertles JF, Wethers DL (Hrsg.): *Sickle Cell Disease: Diagnosis, Management, Education, and Research*, S. 189-229, CV Mosby, 1973.
- Djimde A, Doumbo OK, Cortese JF, Kayentao K, Doumbo S, Diourte Y, Dicko A, Su XZ, Nomura T, Fidock DA, Wellemes TE, Plowe CV, Coulibaly D. A molecular marker of chloroquine-resistant falciparum malaria. *N Engl J Med* 2001; 344: 257-262.
- Dondorp AM, Angus BJ, Chotivanich K, Silamut K, Ruangveerayuth R, Hardeman MR, Kager PA, Vreeken J, White NJ. Red blood cell deformability as a predictor of anemia in severe falciparum malaria. *Am J Trop Med Hyg* 1999; 60: 733-737.
- Doolan DL, Hoffman SL. IL-12 and NK cells are required for antigen-specific adaptive immunity against malaria initiated by CD8+ T cells in the *Plasmodium yoelii* model. *J Immunol* 1999; 163: 884-892.
- Dorman EK, Shulman CE, Kingdom J, Bulmer JN, Mwendwa J, Peshu N, Marsh K. Impaired uteroplacental blood flow in pregnancies complicated by falciparum malaria. *Ultrasound Obstet Gynecol* 2002; 19: 165-170.
- Duflo B, Maiga I, Pichard E, Diallo D, Diallo An, Coulibaly T, Mahamane T, Traore HA., Hemoglobin C in a hospital milieu in Bamako, Mali. *Bull Soc Pathol Exot Filiales* 1985; 78: 393-400.
- Duraisingham MT, Drakeley CJ, Müller O, Bailey R, Snounou G, Targett GA, Greenwood BM, Warhurst DC. Evidence for selection for the tyrosine-86 allele of the *pfmdr 1* gene of *Plasmodium falciparum* by chloroquine and amodiaquine. *Parasitology* 1997; 114: 205-211.
- Duraisingham MT, Curtis J, Warhurst DC. *Plasmodium falciparum*: detection of polymorphisms in the dihydrofolate reductase and dihydropteroate synthetase genes by PCR and restriction digestion. *Exp Parasitol* 1998; 89: 1-8.
- Echave P, Bille J, Audet C, Talla I, Vaudaux B, Gehri M. Percentage, bacterial etiology and antibiotic susceptibility of acute respiratory infection and pneumonia among children in rural Senegal. *J Trop Pediatr* 2003; 49: 28-32.

- Edington GM, Laing WN. Relationship between haemoglobins C and S and malaria in Ghana. *Br Med J* 1957; 2: 143-145.
- Eggelte TA. Production of monoclonal antibodies against antimalarial drugs for use in immunoassays. Navaratnam V, Payne D (Hrsg.) *The validation of chemical and immunochemical tests fro antimalarials in body fluids*. International Monograph Series. Penang, University Saint Malaysia, 1990, 35-63.
- El Hassan AM, Saeed AM, Fandrey J, Jelkmann W. Decreased erythropoietin response in *Plasmodium falciparum* malaria-associated anaemia. *Eur J Haematol* 1997; 59: 299–304.
- Engelbrecht F, Togel E, Beck HP, Enwezor F, Oettli A, Felger I. Analysis of *Plasmodium falciparum* infections in a village community in Northern Nigeria: determination of *msp2* genotypes and parasite-specific IgG responses. *Acta Trop* 2000; 74: 63-71.
- Engwerda CR, Mynott TL, Sawhney S, De Souza JB, Bickle QD, Kaye PM. Locally up-regulated lymphotoxin alpha, not systemic tumor necrosis factor alpha, is the principle mediator of murine cerebral malaria. *J Exp Med* 2002; 195: 1371–1377.
- Fairhurst RM, Fujioka H, Hayton K, Collins KF, Wellemes TE. Aberrant development of *Plasmodium falciparum* in hemoglobin CC red cells: implications for the malaria protective effect of the homozygous state. *Blood* 2003; 101: 3309-3315.
- Färnert A, Snounou G, Rooth I, Björkman A. Daily dynamics of *Plasmodium falciparum* subpopulations in asymptomatic children in a holoendemic area. *Am J Trop Med Hyg* 1997; 56: 538
- Färnert A, Rooth I, Svensson A, Snounou G, Björkman A. Complexity of *Plasmodium falciparum* infections is consistent over time and protects against clinical disease in Tanzanian children. *J Infect Dis* 1999; 179, 989-995.
- Faye FBK, Konate L, Rogier C, Trape JF. *Plasmodium ovale* in a highly malaria endemic area of Senegal. *Trans R Soc Trop Med Hyg* 1998; 92: 522-525.
- Faye FBK, Spiegel A, Tall A, Sokhna C, Fontenille D, Rogier C, Trape JF. Diagnostic criteria and risk factors for *P. ovale* malaria. *J Infect Dis* 2002; 186: 690-695.
- Felger I, Smith T, Edoh D, Kitua A, Alonso P, Tanner M, Beck HP. Multiple *Plasmodium falciparum* infections in Tanzanian infants. *Trans R Soc Trop Med Hyg* 1999; 93 (Suppl. 1): 29-34.
- Fidock AD, Nomura T, Talley KA, Cooper RA, Dzekunov SM, Ferdig MT, Ursos LMB, Sidhu ABS, Naudè B, Deitsch KW, Su XZ, Wootton JC, Roepe PD, Wellemes TE. Mutations in the *P. falciparum* digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance. *Mol Cell* 2000; 6: 861-871.
- Fitch CD. *Plasmodium falciparum* in owl monkeys: drug resistance and chloroquine binding capacity. *Science* 1970; 169: 289-290.
- Fleming AF, Storey J, Molineaux L, Iroko EA, Attai EDE. Abnormal haemoglobins in the Sudan savanna of Nigeria. I. Prevalence of haemoglobins and relationships between sickle cell trait, malaria and survival. *Ann Trop Med Parasitol* 1979; 73:161–172.
- Fleming Af, Harrison KA, Biggs ND, Attai EDE, Ghatoura GBS, Akintunde EA, Shah N. Anaemia in young Primigravidae in the Guinea savanna of Nigeria: sickle cell trait gives partial protection against malaria. *Ann Trop Med Parasitol* 1984; 78: 395-404.
- Flint J, Hill AV, Bowden DK, Oppenheimer SJ, Sill PR, Serjeantson SW, Bana-Koiri J, Bhatia K, Alpers MP, Boyce AJ, Weatherall DJ, Clegg JB. High frequencies of alpha-thalassaemia are the result of natural selection by malaria. *Nature* 1986; 321: 744-750.
- Flint J, Harding RM, Boyce AJ, Clegg JB. The population genetics of the haemoglobinopathies. *Bailliere's Clin Haematol* 1998; 11: 1-51
- Florquin S, Amraoui Z, Dubois C, Decuyper J, Yoldman M. The protective role of endogenously synthesized nitric oxide in staphylococcal enterotoxin B-induced shock in mice. *J Exp Med* 1994; 180: 1153–1158.

- Flüeck TP, Jelinek T, Kilian AH, Adagu IS, Kabagambe G, Sonnenburg F, Warhurst DC. Correlation of *in vivo*-resistance to chloroquine and allelic polymorphisms in *Plasmodium falciparum* isolates from Uganda. *Trop Med Int Health* 2000; 5: 174-178.
- Foster, S. Economic prospects for a new antimalarial drug. *Trans R Soc Trop Med Hyg* 1994; 88(Suppl.): 55-56.
- Fried M, Duffy PE. Adherence of *Plasmodium falciparum* to chondroitin sulfate A in the human placenta. *Science* 1996; 272: 1502-1504.
- Fried M, Nosten F, Brockman A, Brabin BJ, Duffy PE. Maternal antibodies block malaria. *Nature* 1998a; 395: 851-852.
- Fried M, Muga RO, Misore A, Duffy PE. Malaria elicits type 1 cytokines in the human placenta: IFN- γ and TNF- α associated with pregnancy outcomes. *J Immun* 1998b; 160: 2523-2530.
- Friedman MJ. Erythrocytic mechanism of sickle cell resistance to malaria. *Proc Natl Acad Sci USA* 1978; 75: 1994-1997.
- Friedman MJ, Roth EF, Nagel RL, Trager W. The role of hemoglobins C, S, and Nbalt in the inhibition of malaria parasite development *in vitro*. *Am J Trop Med Hyg* 1979; 28: 777-780.
- Gamain B, Smith JD, Miller LH, Baruch DI. Modification in the CD36 binding domain of the *Plasmodium falciparum* variant antigen are responsible for the inability of chondroitin sulfate A adherent parasites to bind CD36. *Blood* 2001; 97: 3268-3274.
- Garnham PCC. *Malaria Parasites and Other Haemosporidia*. Oxford: Blackwell Scientific Publications, 357-417.
- Genton B, al-Yaman F, Alpers MP, Mokela D. Indicators of fatal outcome in paediatric cerebral malaria: a study of 134 comatose Papua New Guinean children. *Int J Epidemiol* 1997; 26: 670-676.
- Gilbert SC, Plebanski M, Gupta S, Morris J, Cox M, Aidoo M, Kwiatkowski D, Greenwood BM, Whittle HC, Hill AVS. Association of malaria parasite population structure, HLA, and immunological antagonism. *Science* 1998; 279: 1173-1177.
- Gilles HM, Fletcher KA, Hendrickse RG, Lindner R, Reddy S, Allan N. Glucose-6-phosphate-dehydrogenase deficiency, sickling, and malaria in African children in South Western Nigeria. *Lancet* 1967; I: 138-140.
- Ginsburg H, Ward SA, Bray PG. An integrated model of chloroquine action. *Parasitol Today* 1999; 15: 357-360.
- Goel VK, Li X, Chen H, Liu SC, Chishti AH, Oh SS. Band 3 is a host receptor binding merozoite surface protein 1 during the *Plasmodium falciparum* invasion of erythrocytes. *Proc Natl Acad Sci U S A* 2003; 100: 5164-5169.
- Good MF, Berzofsky JA, Miller LH. The T cell response to the malaria circumsporozoite protein: an immunological approach to vaccine development. *Annu Rev Immunol* 1988; 6: 663-688.
- Good MF, Doolan DL. Immune effector mechanisms in malaria. *Curr Opin Immunol* 1999; 11: 412-419.
- Granja AC, Machungo F, Gomes A, Bergstrom S, Brabin B. Malaria-related maternal mortality in urban Mozambique. *Ann Trop Med Parasitol* 1998; 92: 257-263.
- Grau GE, Taylor TE, Molyneux ME, Wirima JJ, Vassalli P, Hommel M, Lambert PH. Tumor necrosis factor and disease severity in children with falciparum malaria. *N Engl J Med* 1989; 320: 1586-1591.
- Greenwood BM, Bradley AK, Greenwood AM, Byass P, Jammeh K, Marsh K, Tulloch S, Oldfield FS, Hayes R. Mortality and morbidity from malaria among children in a rural area of The Gambia, West Africa. *Trans R Soc Trop Med Hyg* 1987; 81: 478-486.

- Greenwood BM, Marsh K, Snow R. Why do some African children develop severe malaria? *Parasitol Today* 1991; 7: 277-281.
- Greenwood BM, David PH, Otoo-Forbes LN, Allen SJ, Alonso PL, Armstrong Schellenberg JR, Byass P, Hurwitz M, Menon A, Snow RW. Mortality and morbidity from malaria after stopping malaria chemoprophylaxis. *Trans R Soc Trop Med Hyg* 1995; 89 : 629-633.
- Guinet F, Diallo DA, Minta D, Dicko A, SissokoMS, Keita MM, Wellemes TE, Doumbo O. A comparison of the incidence of severe malaria in Malian children with normal and C-trait haemoglobin profiles. *Acta Trop* 1997; 68: 175-182.
- Gupta S, Snow RW, Donnelly CA, Marsh K, Newbold C. Immunity to non-cerebral severe malaria is acquired after one or two infections. *Nat Med* 1999; 5: 340-343.
- Haldane JBS. The rate of mutation of human genes. *Hereditas* 1949; 35: 267-272.
- Hellgren U, Ericsson O, Kihamia CM, Rombo L. Malaria parasites and chloroquine concentrations in Tanzanian schoolchildren. *Trop Med Parasitol* 1994; 45: 293-297.
- Hill AVS, Allsopp CE, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA, Bennett S, Brewster D, McMichael AJ, Greenwood BM. Common West African HLA alleles are associated with protection from severe malaria. *Nature* 1991; 353: 595-600.
- Ho M, White NJ, Looareesuwan S, Wattanagoon Y, Lee SH, Walport MJ, Bunnag D, Harinasuta T. Splenic Fc receptor function in host defense and anemia in acute *Plasmodium falciparum* malaria. *J Infect Dis* 1990; 161: 555-561.
- Ho M, Sexton MM, Tontaw P, Looaresuwan S, Suntharasamai P, Webster KJ. Interleukin-10 inhibits tumor necrosis factor production but not antigen-specific lymphoproliferation in acute *Plasmodium falciparum* malaria. *J Infect Dis* 1995; 172: 838-844.
- Hobbs MR, Udhayakumar V, Levesque MC, Booth J, Roberts JM, Tkachuk AN, Pole A, Coon H, Kariuki S, Nahlen BL, Mwaikambo ED, Lal AL, Granger DL, Anstey NM, Weinberg JB. A new NOS2 promoter polymorphism associated with increased nitric oxide production and protection from severe malaria in Tanzanian and Kenyan children. *Lancet* 2002; 360: 1468-1475.
- Hurme M, Lahdenpohja N, Santtila S. Gene polymorphisms of interleukins 1 and 10 in infectious and autoimmune diseases. *Ann Med* 1998; 30: 469-473.
- Hviid L. Clinical disease, immunity and protection against *Plasmodium falciparum* malaria in populations living in endemic areas. *Exp Rev Mol Med* 1998; 24 June, <http://www-ermm.cbcu.cam.ac.uk/lhc/tx001lhc.htm>
- Imbert P, Sartelet I, Rogier C, Ka S, Baujat G, Candito D. Severe malaria among children in a low seasonal transmission area, Dakar, Senegal: influence of age on clinical presentation. *Trans R Soc Trop Med Hyg* 1997; 91: 22-24.
- Imbert P, Gérardin P, Rogier C, Ka AS, Jouvencel P, Brousse V, Guyon P. Severe malaria in children: a comparative study of 1990 and 2000 WHO criteria for clinical presentation, prognosis and intensive care in Dakar, Senegal. *Trans R Soc Trop Med Hyg* 2002; 96: 278-281.
- Ismail MR, Ordi J, Menendez C, Ventura PJ Aponte JJ, Kahigwa E, Hirt R, Cardesa A, Alonso PL. Placental pathology in malaria: a histological, immunohistochemical, and quantitative study. *Hum Pathol* 2000; 31: 85-93.
- Itano HA, Neel JV. A new inherited abnormality of human hemoglobin. *Proc Natl Acad Sci U S A* 1950; 36: 613-617.
- Iyer JK, Milhous WK, Cortese JF, Kublin JG, Plowe CV. *Plasmodium falciparum* cross-resistance between trimethoprim and pyrimethamine. *Lancet* 2001 29;358: 1066-1067.
- Jarra W, Snounou G. Only viable parasites are detected by PCR following clearance of rodent malarial infections by drug treatment or immune responses. *Infect Immun* 1998; 66: 3783-3787.

- Jeffery GM. Epidemiological significance of repeated infections with homologous and heterologous strains and species of *Plasmodium*. *Bull World Health Organ* 1966; 35: 873-882.
- Jelinek T, Kilian AH, Curtis J, Duraisingh MT, Kabagambe G, von Sonnenburg F, Warhurst DC. *Plasmodium falciparum*: selection of serine 108 of dihydrofolate reductase during treatment of uncomplicated malaria with co-trimoxazole in Ugandan children. *Am J Trop Med Hyg* 1999; 61: 125-30.
- Kahigwa E, Schellenberg D, Sanz S, Aponte JJ, Wigayi J, Mshinda H, Alonso P, Menendez C. Risk factors for presentation to hospital with severe anaemia in Tanzanian children: a case-control study. *Trop Med Int Health* 2002; 7: 823-830.
- Kamchonwongpaisan S, Chandrangam G, Avery MA, Yuthavong Y. Resistance to artemisinin of malaria parasites (*Plasmodium falciparum*) infecting alpha-thalassemic erythrocytes in vitro. *J Clin Invest* 1994; 93: 467-473.
- Kamwendo DD, Dzinjalama FK, Snounou G, Kanjala MC, Mhango CG, Molyneux ME, Rogerson SJ. *Plasmodium falciparum*: PCR detection and genotyping of isolates from peripheral, placental, and cord blood of pregnant Malawian women and their infants. *Trans R Soc Trop Med Hyg* 2002; 96: 145-149.
- Karhukorpi J, Karttunen R. Genotyping interleukin-10 high and low producers with single-tube bidirectional allele-specific amplification. *Exp Clin Immunogenet* 2001; 18: 67-70.
- Khalil I, Ronn AM, Alifrangis M, Gabar HA, Satti GM, Bygbjerg IC. Dihydrofolate reductase and dihydropteroate synthase genotypes associated with *in vitro* resistance of *Plasmodium falciparum* to pyrimethamine, trimethoprim, sulfadoxine, and sulfamethoxazole. *Am J Trop Med Hyg* 2003; 68: 586-589.
- Kicska GA, Ting LM, Schramm VL, Kim K. Effect of dietary p-aminobenzoic acid on murine *Plasmodium yoelii* infection. *J Infect Dis* 2003; 188: 1776-1781.
- Kitua AY, Smith T, Alonso P, Masanja H, Menendez C, Urassa H, Kimario J, Tanner M. *Plasmodium falciparum* malaria in the first year of life in an area of intense perennial transmission. *Trop Med Int Health* 1996; 1: 475-484.
- Kitua AY, Smith TA, Alonso PL, Urassa H, Masanja H, Kimario J, Tanner M. The role of low level *Plasmodium falciparum* parasitaemia in anaemia among infants living in an area of intense and perennial transmission. *Trop Med Int Health* 1997; 2: 325-333.
- Knight JC, Udalova I, Hill AV, Greenwood BM, Peshu N, Marsh K, Kwiatkowski D. A polymorphism that affects OCT-1 binding to the TNF promoter region is associated with severe malaria. *Nat Genet* 1999; 22:145-150.
- Knowles R, White RS. Studies in the parasitology of malaria. *Ind Med Res Mem* 1930; 18.
- Konate L, Zwetyenga J, Rogier C, Bischoff E, Fontenille D, Tall A, Spiegel A, Trape JF, Mercereau-Puijalon O. Variation of *Plasmodium falciparum* *msp1* block 2 and *msp2* allele prevalence and of infection complexity in two neighbouring Senegalese villages with different transmission conditions. *Trans R Soc Trop Med Hyg* 1999; 93 (Suppl. 1), 21-28.
- Koram KA, Owusu-Agyei S, Fryauff DJ, Anto F, Atuguba F, Hodgson A, Hoffman SL, Nkrumah FK. Seasonal profiles of malaria infection, anaemia, and bednet use among age groups and communities in northern Ghana. *Trop Med Int Health* 2003; 8: 793-802.
- Kotea R, Kaeda JS, Yan SL, Sem Fa N, Beesoon S, Jankee S, Ramasawmy R, Vulliamy T, Bradnock RW, Batista J, Luzzatto L, Krishnamoorthy R, Mason PJ. Three major G6PD-deficient polymorphic variants identified among the Mauritian population. *Br J Haematol* 1999; 104: 849-854.
- Kremsner PG, Winkler S, Brandts C, Wildling E, Jenne L, Graninger W, Prada J, Bienzle U, Juillard P, Grau GE. Prediction of accelerated cure in *Plasmodium falciparum* malaria by the elevated capacity of tumor necrosis factor production. *Am J Trop Med Hyg* 1995; 53: 532-538.

- Kremsner PG, Winkler S, Wildling E, Prada J, Bienzle U, Graninger W, Nüssler AK. High plasma levels of nitrogen oxides are associated with severe disease and correlate with rapid parasitological and clinical cure in *Plasmodium falciparum* malaria. *Trans R Soc Trop Med Hyg* 1996; 90: 44–47.
- Krogstad DJ, Gluzman IY, Kyle DE, Oduola AM, Martin SK, Milhous WK, Schlesinger PH. Efflux of chloroquine from *Plasmodium falciparum*: mechanism of chloroquine resistance. *Science* 1987; 238: 1283–1285.
- Kublin JG, Dzinjalama FK, Kamwendo DD, Malkin EM, Cortese JF, Martino LM, Mukadam RA, Rogerson SJ, Lescano AG, Molyneux ME, Winstanley PA, Chimpeni P, Taylor TE, Plowe CV. Molecular markers for failure of sulfadoxine-pyrimethamine and chlorproguanil-dapsone treatment of *Plasmodium falciparum* malaria. *J Infect Dis* 2002; 185: 380–388.
- Kun JF, Schmidt-Ott RJ, Lehman LG, Lell B, Luckner D, Greve B, Matousek P, Kremsner PG. Merozoite surface antigen 1 and 2 genotypes and rosetting of *Plasmodium falciparum* in severe and mild malaria in Lambarene, Gabon. *Trans R Soc Trop Med Hyg* 1998a; 92: 110–114.
- Kun JF, Mordmuller B, Lell B, Lehman LG, Luckner D, Kremsner PG. Polymorphism in promoter region of inducible nitric oxide synthase gene and protection against malaria. *Lancet* 1998b; 351: 265–266.
- Kun JF, Mordmüller B, Perkins DJ, May J, Mercereau-Puijalon O, Alpers M, Weinberg JB, Kremsner PG. Nitric oxide synthase 2^{Lambaréné (G-954C)}, increased nitric oxide production, and protection against malaria. *J Infect Dis* 2001; 184: 330–336.
- Kurtzhals JA, Adabayeri V, Goka BQ, Akanmori BD, Oliver-Commeey JO, Nkrumah FK, Behr C, Hviid L. Low plasma concentrations of interleukin 10 in severe malarial anaemia compared with cerebral and uncomplicated malaria. *Lancet* 1998; 351: 1768–1772.
- Kwiatkowski D. Cytokines and anti-disease immunity to malaria. *Res Immunol* 1991; 142: 707–712.
- Kurtzhals JA, Helleberg M, Goka BQ, Akanmori BD. Severe malaria in west African children. *Lancet* 2003; 361: 1393.
- Labie D, Richin C, Pagnier J, Gentilini M, Nagel RL. Hemoglobins S and C in Upper Volta. *Hum Genet* 1984; 65: 300–302.
- Leke RF, Djokam RR, Mbu R, Leke RJ, Fogako J, Megnekou R, Metenou S, Sama G, Zhou Y, Cadigan T, Parra M, Taylor DW. Detection of the *Plasmodium falciparum* Antigen histidine-rich protein 2 in blood of pregnant women: implications for diagnosing placental malaria. *J Clin Microbiol* 1999; 2992–2996.
- Lell B, May J, Schmidt-Ott RJ, Lehman LG, Luckner D, Greve B, Matousek P, Schmid D, Herbich K, Mockenhaupt FP, Meyer CG, Bienzle U, Kremsner PG. The role of red blood cell polymorphisms in resistance and susceptibility to malaria. *Clin Infect Dis* 1999; 28: 794–799.
- Levesque MC, Hobbs MR, Anstey NM, Vaughn TN, Chancellor JA, Pole A, Perkins DJ, Misukonis MA, Chanok SJ, Granger DL, Weinberg JB. Nitric oxide synthase type 2 promoter polymorphisms, nitric oxide production, and disease severity in Tanzanian children with malaria. *J Infect Dis* 1999; 180: 1994–2002.
- Lindsay S, Ansell J, Selman C, Cox V, Hamilton K, Walraven G. Effect of pregnancy on exposure to malaria mosquitoes. *Lancet* 2000; 355: 1972.
- Liu YT, Old JM, Miles K, Fisher CA, Weatherall DJ, Clegg JB. Rapid detection of alpha-thalassaemia deletions and alpha-globin gene triplication by multiplex polymerase chain reactions. *Brit J Haematol* 2000; 108: 295–299.

- Looareesuwan S, Ho M, Wattanagoon Y, White NJ, Warrell DA, Bunnag D, Harinasuta T, Wyler DJ. Dynamic alteration in splenic function during acute falciparum malaria. *N Engl J Med* 1987;317: 675-679.
- Luty AJ, Lell B, Schmidt-Ott R, Lehman LG, Luckner D, Greve B, Matousek P, Herbich K, Schmid D, Migot-Nabias F, Deloron P, Nussenzweig RS, Kremsner PG. Interferon-gamma responses are associated with resistance to reinfection with *Plasmodium falciparum* in young African children. *J Infect Dis* 1999; 179: 980-988.
- Luzzatto L, Nwachukwu-Jarrett ES, Reddy S. Increased sickling of parasitised erythrocytes as mechanism of resistance against malaria in the sickle-cell trait. *Lancet* 1970; 1: 319-321.
- Luzzatto L, Usanga FA, Reddy S. Glucose-6-phosphate dehydrogenase deficient red cells: resistance to infection by malarial parasites. *Science* 1969; 164: 839-842.
- Luzzi GA, Merry AH, Newbold CI, Marsh K, Pasvol G, Weatherall DJ. Surface antigen expression on *Plasmodium falciparum*-infected erythrocytes is modified in α - and β -thalassemia. *J Exp Med* 1991; 173: 785-791.
- Magesa SM, Mdira KY, Babiker HA, Alifrangis M, Färnert A, Simonsen PE, Bygbjerg IC, Walliker D, Jakobsen PH. Diversity of *Plasmodium falciparum* clones infecting children living in a holoendemic area in north-eastern Tanzania. *Acta Trop* 2002; 84: 83-92.
- Maitland K, Williams TN, Newbold CI. *Plasmodium vivax* and *P. falciparum*: biological interactions and the possibility of cross-species immunity. *Parasitol Today* 1997; 13: 227
- Maitland K, Levin M, English M, Mithwani S, Peshu N, Marsh K, Newton CR. Severe *P. falciparum* malaria in Kenyan children: evidence for hypovolaemia. *QJM* 2003; 96: 427-434.
- Mankhambo L, Kanjala M, Rudman S, Lema VM, Rogerson SJ. Evaluation of the OptiMal rapid antigen test and species specific PCR to detect placental *Plasmodium falciparum* infection at delivery. *J Clin Microbiol* 2002; 40: 155-158.
- Marsh K, Howard RJ. Antigens induced on erythrocytes by *P. falciparum*: expression of diverse and conserved determinants. *Science* 1986; 231: 150-153.
- Marsh K, Forster D, Waruiru C, Mwangi I, Winstanley M, Marsh V, Newton C, Winstanley P, Warn P, Peshu N, Pasvol G, Snow R. Indicators of life-threatening malaria in African children. *N Engl J Med* 1995; 332: 1399-1404.
- Mason DP, McKenzie FE, Bossert WH. The blood-stage dynamics of mixed *Plasmodium malariae*-*Plasmodium falciparum* infections. *J Theor Biol* 1999; 198: 549
- Matteelli A, DonatoF, Shein A, Muchi JA, Leopardi O, Astori L, Carosi G. Malaria and anaemia in pregnant women in urban Zanzibar, Tanzania. *Ann Trop Med Parasitol* 1994; 88: 475-483.
- May J, Falusi Ag, Ademowo OG, Olumese PE, Grossterlinden L, Mockenhaupt, FP, Meyer CG, Luzzatto L, Bienzle U. Red cell glucose-6-phosphate dehydrogenase status and pyruvate kinase activity in a Nigerian population. *Trop Med Int Health* 2000a; 5: 119-123.
- May J, Lell B, Luty AJF, Meyer CG, Kremsner PG. Plasma interleukin-10:tumor necrosis factor (TNF)- α ratio is associated with TNF promoter variants and predicts malarial complications. *J Infect Dis* 2000b; 182: 1570-1573.
- May J, Meyer CG. Association of *Plasmodium falciparum* chloroquine resistance transporter variant T76 with age-related plasma chloroquine levels. *Am J Trop Med Hyg* 2003; 68: 143-146.
- McGregor IA, Carrington S, Cohen S. Treatment of East African *P. falciparum* malaria with East African human gamma-globulin. *Trans R Soc Trop Med Hyg* 1963; 50: 170-175.
- McGregor IA. Epidemiology, malaria and pregnancy. *Am J Trop Med Hyg* 1984; 33: 517-525.

- McGuire W, Hill AV, Allsopp CE, Greenwood BM, Kwiakowski D. Variation in the TNF- α promoter region associated with susceptibility to cerebral malaria. *Nature* 1994; 371: 508–510.
- McGuire W, Knight JC, Hill AVS, Allsopp CEM, Greenwood BM, Kwiatkowski D. Severe malarial anemia and cerebral malaria are associated with different tumor necrosis factor alleles. *J Infect Dis* 1999; 179: 287-290.
- McKenna KC, Tsuji M, Sarzotti M, Sacci JB, Witney AA, Azad AF. Gammadelta T cell are a component of early immunity against preerythrocytic malaria parasites. *Infect Immun* 2000; 68: 2224-2230
- McKenzie FE, Bossert WH. Mixed-species *Plasmodium* infections of humans. *J Parasitol* 1997; 83: 593-600.
- Mehlotra RK, Lorry K, Kastens W, Miller SM, Alpers MP, Bockarie M, Kazura JW, Zimmerman PA. Random distribution of mixed species malaria infections in Papua New Guinea. *Am J Trop Med Hyg* 2000; 62: 225-231.
- Mehta A, Mason PJ, Vulliamy TJ. Glucose-6-phosphate dehydrogenase deficiency. *Bailliere's Clin Haematol* 2000; 13: 21-38.
- Menendez C, Quinto LL, Kahigwa E, Alvarez L, Fernandez R, Gimenez N, Schellenberg D, Aponte JJ, Tanner M, Alonso PL. Effect of malaria on soluble transferrin receptor levels in Tanzanian infants. *Am J Trop Med Hyg* 2001; 65: 138-142.
- Menendez C, Todd J, Alonso PL, Francis N, Lulat S, Ceesay S, Ascaso C, Smith T, M'Boge B, Greenwood BM. The response to iron supplementation of pregnant women with the haemoglobin genotypes AA or AS. *Trans R Soc Trop Med Hyg* 1995; 89: 289-292.
- Menendez C, Kahigwa E, Hirt R, Vounatsou P, Aponte JJ, Font F, Acosta CJ, Schellenberg DM, Galindo CM, Kimario J, Urassa H, Brabin B, Smith TA, Kitua AY, Tanner M, Alonso PL. Randomised placebo-controlled trial of iron supplementation and malaria chemoprophylaxis for prevention of severe anaemia and malaria in Tanzanian infants. *Lancet* 1997; 350: 844-850.
- Menendez C, Fleming AF, Alonso PL. Malaria-related anaemia. *Parasitol Today* 2000a; 16: 469-476.
- Menendez C, Ordi J, Ismail MR, Ventura PJ, Aponte JJ, Kahigwa E, Font F, Alonso PL. The impact of placental malaria on gestational age and birth weight. *J Infect Dis* 2000b; 181: 1740-1745.
- Metaxotou-Mavromati AD, Antonopoulou HK, Laskari SS, Tsianta HK, Ladis VA, Kattamis CA. Developmental changes in hemoglobin F levels during the first two years of life in normal and heterozygous beta-thalassemia infants. *Pediatrics* 1982; 69: 734-738.
- Miller LH, Baruch DI, Marsh K, Doumbo Ok. The pathogenic basis of malaria. *Nature* 2002; 415: 673-679.
- Missinou MA, Lell B, Kremsner PG. Uncommon asymptomatic *Plasmodium falciparum* infections in Gabonese children. *Clin Infect Dis* 2003; 36: 1198-1202.
- Mockenhaupt FP, May J, Bienzle U, Meyer CG. Medikamenten-resistente Malaria. *Dtsch Med Wochenschr* 1997; 122: 1293-1297.
- Modiano D, Sirima BS, Sawadogo A, Sanou I, Konaté A, Pagnoni F. Severe malaria in Burkina Faso: influence of age and transmission level on clinical presentation. *Am J Trop Med Hyg* 1998; 59: 539-542.
- Modiano G, Morpurgo G, Terrenato L, Novelletto A, Di Rienzo A, Colombo B, Purpura M, Mariani M, Santachiara-Benerecetti S, Brega A, et al. Protection against malaria morbidity: near-fixation of the α -thalassemia gene in a Nepalese population. *Am J Hum Genet* 1991; 48: 390-397.
- Modiano G, Luoni G, Sirima BS, Simoré J, Verra F, Konaté A, Rastrellei E, Olivieri A, Calissano C, Paganotti GM, D'Urbano L, Sanou I, Sawadogo A, Modiano G, Coluzzi M.

- Haemoglobin C protects against clinical *Plasmodium falciparum* malaria. *Nature* 2001; 414: 305-308.
- Molineaux L, Storey J, Cohen JE, Thomas A. A longitudinal study of human malaria in the West African Savanna in the absence of control measures: relationships between different *Plasmodium* species, in particular *P. falciparum* and *P. malariae*. *Am J Trop Med Hyg* 1980; 29: 725-737.
- Molyneux ME, Taylor TE, Wirima JJ, Borgstein A. Clinical features and prognostic indicators in paediatric cerebral malaria: a study of 131 comatose Malawian children. *QJM* 1989; 71: 441-459.
- Moormann AM, Sullivan AD, Rochford RA, Chensue SW, Bock PJ, Nyirenda T, Meshnick SR. Malaria and pregnancy: Placental cytokine expression and its relationship to intrauterine growth retardation. *J Infect Dis* 1999; 180:1983-1993.
- Müller DA, Charlwood JD, Felger I, Ferreira C, do Rosario V, Smith T. Prospective risk of morbidity in relation to multiplicity of infection with *Plasmodium falciparum* in Sao Tome. *Acta Trop* 2001; 78: 155-162.
- Müller O, Traore C, Jahn A, Becher H. Severe anaemia in west African children: malaria or malnutrition? *Lancet* 2003; 361: 86-87.
- Murphy PM. Molecular mimicry and the generation of host defense protein diversity. *Cell* 1993; 72: 823-826.
- Mutabingwa TK, de Geus A, Meuwissen JH, Malle LN. Malaria chemosuppression during pregnancy. VI. Some epidemiological aspects of malaria in infants. *Trop Geogr Med* 1994; 46: 1-7.
- Nahlen BL, Akintunde A, Alakija T, Nguyen-Dinh P, Ogunbode O, Edungbola LD, Adetoro O, Breman JG. Lack of efficacy of pyrimethamine prophylaxis in pregnant Nigerian women. *Lancet* 1989; 2: 830-834.
- Ndyomugenyi R, Magnussen P. Chloroquine prophylaxis, iron-folic acid supplementation or case management of malaria attacks in primigravidae in western Uganda: effects on maternal parasitaemia and haemoglobin levels and on birthweight. *Trans R Soc Trop Med Hyg* 2000; 94: 413-418.
- Newbold C, Warn P, Black G, Berendt A, Graig A, Snow B, Msobo M, Peshu N, Marsh K. Receptor-specific adhesion and clinical disease in *Plasmodium falciparum*. *Am J Trop Med Hyg* 1997; 57: 389-398.
- Nevill CG, Some ES, Mung'ala VO, Mutemi W, New L, Marsh K, Lengeler C, Snow RW. Insecticide-treated bednets reduce mortality and severe morbidity from malaria among children on the Kenyan coast. *Trop Med Int Health* 1996; 1: 139-146.
- Nguyen-Dinh P, Parvin RM. Haemoglobin S and *in vitro* chloroquine susceptibility of *Plasmodium falciparum*. *Lancet* 1986; 2: 1278.
- Nnalue NA, Friedman MJ. Evidence for a neutrophil-mediated protective response in malaria. *Parasite Immunol* 1988; 10: 47-58.
- Nsimba SE, Massele AY, Eriksen J, Gustafsson LL, Tomson G, Warsame M. Case management of malaria in under-fives at primary health care facilities in a Tanzanian district. *Trop Med Int Health* 2002; 7: 201-209.
- Nsobya SL, Parikh S, Njama-Meya D, Kironde F, Lubega GW, Kamya MR, Rosenthal PJ, Dorsey G. Molecular Studies of asymptomatic malaria in Kampala, Uganda. 52nd Annual Meeting of the American Society of Tropical Medicine and Hygiene, Philadelphia, Dez. 3-7, 2003, abstract 1343
- Ntoumi F, Contamin H, Rogier C, Bonnefoy S, Trape JF, Mercereau-Puijalon O. Age-dependent carriage of multiple *Plasmodium falciparum* merozoite surface antigen-2 alleles in asymptomatic malaria infections. *Am J Trop Med Hyg* 1995; 52: 81-88.

- Ntoumi F, Mercereau-Puijalon O, Ossari S, Luty A, Reltien J, Georges A, Millet P. *Plasmodium falciparum*: sickle-cell trait is associated with higher prevalence of multiple infections in Gabonese children with asymptomatic infections. *Exp Parasitol* 1997a; 87 : 39-46.
- Ntoumi F, Rogier C, Dieye A, Trape JF, Millet P, Mercereau-Puijalon O. Imbalanced distribution of *Plasmodium falciparum* MSP-1 genotypes related to sickle-cell trait. *Mol Med* 1997b; 3: 581-592.
- Ntoumi F, Ekala MT, Makouwa M, Lekoulou F, Mercereau-Puijalon O, Deloron P. Sickle cell trait carriage: imbalanced distribution of IgG subclass antibodies reactive to *Plasmodium falciparum* family-specific MSP2 peptides in serum samples from Gabonese children. *Immun Lett* 2002; 84, 9-16.
- Nussenzweig V, Nussenzweig RS. Rationale for the development of an engineered sporozoite malaria vaccine. *Adv Immunol* 1989; 45: 283-334.
- Nzila AM, Mberu EK, Sulo J, Dayo H, Winstanley PA, Sibley CH, Watkins WM. Towards an understanding of the mechanism of pyrimethamine-sulfadoxine resistance in *Plasmodium falciparum*: genotyping of dihydrofolate reductase and dihydropteroate synthase of Kenyan parasites. *Antimicrob Agents Chemother* 2000; 44: 991-996.
- O'Neil-Dunne I, Archur RN, Agbor-Enoh ST, Valiyaveettil M, Naik RS, Ockenhouse CF, Zhou A, Megnekou R, Leke R, Taylor DW, Gowda DC. Gravidity-depending production of antibodies that inhibit binding of *Plasmodium falciparum*-infected erythrocytes to placental chondroitin sulfate proteoglycan during pregnancy. *Infect Immun* 2001; 69: 7487-7492.
- Ofosu-Okyere A, Mackinnon MJ, Sowa MP, Koram KA, Nkrumah F, Osei YD, Hill WG, Wilson MD, Arnot DE. Novel *Plasmodium falciparum* clones and rising clone multiplicities are associated with the increase in malaria morbidity in Ghanaian children during the transition into the high transmission season. *Parasitology* 2001; 123 (Pt. 2): 113-123.
- Ohashi J, Naka I, Patarapotikul J, Hananantachai H, Looareesuwan S, Tokunaga K. Significant association of longer forms of CCTTT microsatellite repeat in the inducible nitric oxide synthase promoter with severe malaria in Thailand. *J Infect Dis* 2002a; 186: 578-581.
- Ohashi J, Naka I, Patarapotikul J, Hananantachai H, Looareesuwan S, Tokunaga K. Lack of association between interleukin-10 gene promoter polymorphism, -1082G/A, and severe malaria in Thailand. *Southeast Asian J Trop Med Public Health* 2002b;33 Suppl 3:5-7.
- Olson JA, Nagel RL. Synchronized cultures of *P. falciparum* in abnormal red cells: the mechanism of the inhibition of growth in HbCC cells. *Blood* 1986; 67: 997-1001.
- O'Neill-Dunne I, Archur RN, Agbor-Enoh ST, et al. Gravidity-dependent production of antibodies that inhibit binding of *Plasmodium falciparum*-infected erythrocytes to placental chondroitin sulphate proteoglycan during pregnancy. *Infect Immun* 2001; 94: 477-483.
- Onori E. Differences in *Plasmodium malariae* prevalences in Uganda. *Bull WHO* 1967; 37: 330-331.
- Oppenheimer SJ, Hill AVS, Gibson FD, MacFarlane SB, Moody JB, Pringle J. The interaction of alpha thalassaemia and malaria. *Trans R Soc Trop Med Hyg* 1987; 81: 322-326.
- Ordi J, Ismail MR, Ventura PJ, Kehigwa E, Hirt R, Cardesa A, Alonso PL, Menendez C. Massive chronic intervillitis of the placenta associated with malaria infection. *Am J Surg Pathol* 1998; 22: 1006-1011.
- Orjih AU. Increased accumulation of chloroquine and desethylchloroquine in homozygous sickle cells. *Bull World Health Organ* 1987; 65: 51-55.

- Othoro C, Lal AA, Nahlen B, Koech D, Orago AS, Udhayakumar V. A low interleukin-10 tumor necrosis factor-alpha ratio is associated with malaria anemia in children residing in a holoendemic malaria region in western Kenya. *J Infect Dis* 1999; 179: 279-282.
- Owusu-Agyei S, Smith T, Beck HP, Amenga-Etego L, Felger I. Molecular epidemiology of *Plasmodium falciparum* infections among asymptomatic inhabitants of a holoendemic malarious area in northern Ghana. *Trop Med Int Health* 2002; 7: 421-428.
- Pasquetto V, Fidock DA, Gras H, Badell E, Eling W, Ballou WR, Belghiti J, Tartar A, Druilhe P. *Plasmodium falciparum* sporozoite invasion is inhibited by naturally acquired or experimentally induced polyclonal antibodies to the STARP antigen. *Eur J Immunol* 1997; 27: 2502-2513.
- Pasvol G, Weatherall DJ, Wilson RJ. Effects of foetal haemoglobin on susceptibility of red cells to *Plasmodium falciparum*. *Nature* 1977; 270: 171-173.
- Pasvol G, Weatherall DJ, Wilson RJM. Cellular mechanism for the protective effect of haemoglobin S against malaria. *Nature* 1978; 274: 701-703.
- Pasvol G. The interaction between sickle haemoglobin and the malarial parasite *Plasmodium falciparum*. *Trans R Soc Trop Med Hyg* 1980; 74: 701-705.
- Pasvol G, Wilson RJ. The interaction of malaria parasites with red blood cells. *Br Med Bull* 1986; 38: 133-140.
- Pattanapanyasat K, Yongvanitchit K, Tongtawe P, Tachavanich K, Wanachiwanawin W, Fucharoen S, Walsh DS. Impairment of *Plasmodium falciparum* growth in thalassemic red blood cells: further evidence by using biotin labeling and flow cytometry. *Blood* 1999; 93: 3116-3119.
- Petersen E. *In vitro* susceptibility of *Plasmodium falciparum* malaria to pyrimethamine, sulfadoxine, trimethoprim and sulfamethoxazole, singly and in combination. *Trans R Soc Trop Med Hyg* 1987; 81: 238-241.
- Plebanski M, Hill AV. The immunology of malaria infection. *Curr Opin Immunol* 2000; 12: 437-441.
- Plowe CV, Djimde A, Wellems TE, Diop S, Kouriba B, Doumbo OK. Community pyrimethamine-sulfadoxine use and prevalence of resistant *Plasmodium falciparum* genotypes in Mali: a model for deterring resistance. *Am J Trop Med Hyg* 1996; 55: 467-471.
- Postigo M, Mendoza-Leon A, Perez HA. Malaria diagnosis by the polymerase chain reaction: a field study in south-eastern Venezuela. *Trans R Soc Trop Med Hyg* 1998; 92: 509-511.
- Powers KG, Jacobs RL, Good WC, Koontz LC. *Plasmodium vinckeii*: production of chloroquine-resistant strain. *Exp Parasitol* 1969; 26: 193-202.
- Raper AB. Malaria and the sickling trait. *Br Med J* 1955; II: 1186-1189.
- Rathod PK, McErlean T, Lee PC. Variations in frequencies of drug resistance in *Plasmodium falciparum*. *Proc Natl Acad Sci U S A* 1997; 94: 9389-9393.
- Reed MB, Saliba KJ, Caruana SR, Kirk K, Cowman AF. Pgh1 modulates sensitivity and resistance to multiple antimalarials in *Plasmodium falciparum*. *Nature* 2000; 403: 906-909.
- Reeder JC, Cowman AF, Davern KM. The adhesion of *Plasmodium falciparum*-infected erythrocytes to chondroitin sulfate A is mediated by *P. falciparum* erythrocyte membrane protein 1. *Proc Natl Acad Sci U S A* 1999; 96: 5198-5202.
- Rees DC, Williams TN, Maitland K, Clegg JB, Weatherall DJ. Alpha thalassaemia is associated with increased soluble transferrin receptor levels. *Br J Haematol* 1998; 103: 365-369.
- Rhee MS, Akanmori BD, Waterfall M, Riley EM. Changes in cytokine production associated with acquired immunity to *Plasmodium falciparum* malaria. *Clin Exp Immunol* 2001; 126: 503-510.

- Rice AL, Sacco L, Hyder A, Black RE. Malnutrition as an underlying cause of childhood deaths associated with infectious diseases in developing countries. *Bull World Health Organ* 2000; 78: 1207-1221.
- Richie TL. Interactions between malaria parasites infecting the same vertebrate host. *Parasitology* 1988; 96: 607-639.
- Ricke CH, Staalsoe T, Koram K, Akanmori BD, Riley EM, Theander TG, Hviid L. Plasma antibodies from malaria-exposed pregnant women recognize variant surface antigens on *Plasmodium falciparum*-infected erythrocytes in a parity-dependent manner and block parasite adhesion to chondroitin sulfate A. *J Immunol* 2000; 165: 3309-3316.
- Ringelmann B, Hathorn MKS, Jilly P, Grant F, Parniczky G. A new look at the protection of haemoglobin AS and AC genotypes against *Plasmodium falciparum* infection: a census tract approach. *Am J Human Genet* 1976; 28: 270-279.
- Ringwald P, Eboumbou EC, Bickii J, Basco LK. *In vitro* activities of pyronaridine, alone and in combination with other antimalarial drugs, against *Plasmodium falciparum*. *Antimicrob Agents Chemother* 1999; 43: 1525-1527.
- Robert F, Ntoumi F, Angel G, Candito D, Rogier C, Fandeur T, Sarthou JL, Mercereau-Puijalon O. Extensive genetic diversity of *Plasmodium falciparum* isolates collected from patients with severe malaria in Dakar, Senegal. *Trans R Soc Trop Med Hyg* 1996; 90: 704-711.
- Rockett KA, Awburn MM, Cowden WB, Clark IA. Killing of *Plasmodium falciparum* *in vitro* by nitric oxide derivatives. *Infect Immun* 1991; 59: 3280-3283.
- Rockett KA, Awburn MM, Aggarwal BB, Cowden WB, Clark IA. *In vivo* induction of nitrite and nitrate by tumor necrosis factor, lymphotoxin, and interleukin-1: possible roles in malaria. *Infect Immun* 1992; 60: 3725-3730.
- Rogerson SJ, Brown HC, Pollina E, Abrams ET, Tadesse E, Lema VM, Molyneux E. Placental tumor necrosis factor alpha but not gamma interferon is associated with placental malaria and low birth weight in Malawian women. *Infect Immun* 2003; 71: 267-270.
- Rogier C, Trape J-F. Etude de l'aquisition de la prémunition en zone d'holo- et de méso-endémie palustre à Dielmo et à Ndiop (Senegal): Résultats préliminaires, 1990-1994. *Med Trop* 1995; 55: 71-76.
- Ross R. *The prevention of malaria*. John Murray, London, 1910.
- Roth EF Jr, Raventos-Suarez C, Rinaldi A, Nagel RL. Glucose-6-phosphate dehydrogenase deficiency inhibits *in vitro* growth of *Plasmodium falciparum* malaria. *Proc Natl Acad Sci U S A* 1983; 80: 298-299.
- Rowe JA, Kyes SA, Rogerson SJ, Babiker HA, Raza A. Identification of a conserved *Plasmodium falciparum* var gene implicated in malaria in pregnancy. *J Infect Dis* 2002; 185: 1207-1211.
- Ruffié J, Soumia JC. Die Seuchen in der Geschichte der Menschheit. Klett Cotta, Stuttgart, 1987
- Ruwende C, Khoo Sc, Snow RW, Yates SNR, Kwiatkowski D, Gupta A, Warn P, Allsopp CEM, Gilbert SC, Pesch B, Newbold CI, Greenwood BM, Marsh K, Hill AVS. Natural protection of hemi- and heterozygotes for G6PD deficiency in Africa by resistance to severe malaria. *Nature* 1995; 376: 246-249.
- Ruwende C, Hill AV. Glucose-6-phosphate dehydrogenase deficiency and malaria. *J Mol Med* 1998; 76: 581-588.
- Rzepczyk CM, Anderson K, Stamatou S, Townsend E, Allworth A, McCormack J, Whitby M. Gamma delta T cells: their immunobiology and role in malaria infections. *Int J Parasitol* 1997; 27: 191-200.

- Saiki RK, Scharf S, Faloona F, Mullis KB, Horn GT, Ehrlich HA, Arnheim N. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anaemia. *Science* 1985; 230: 1350-1354.
- Sanna E, Bruno R, Cosseddu GG, Floris G, Salis A, Silvetti M. Present-day G-6-PD deficit in Sardinia with respect to malarial morbidity and mortality in the past. *Z Morphol Anthropol* 1990; 78: 257-267.
- Saute F, Menendez C, Mayor A, Aponte J, Gomez-Olive X, Dgedge M, Alonso P. Malaria in pregnancy in rural Mozambique: the role of parity, submicroscopic and multiple *Plasmodium falciparum* infections. *Trop Med Int Health* 2002; 7: 19-28.
- Schellenberg D, Menendez C, Kahigwa E, Font F, Galindo C, Acosta C, Schellenberg JA, Aponte JJ, Kimario J, Urassa H, Mshinda H, Tanner M, Alonso P. African children with malaria in an area of intense *Plasmodium falciparum* transmission: features on admission to the hospital and risk factors for death. *Am J Trop Med Hyg* 1999; 61: 431-438.
- Schellenberg D, Menendez C, Kahigwa E, Aponte J, Vidal J, Tanner M, Mshinda H, Alonso P. Intermittent treatment for malaria and anaemia control at time of routine vaccinations in Tanzanian infants: a randomised, placebo-controlled trial. *Lancet* 2001; 357: 1471-1477.
- Schellenberg D, Schellenberg JR, Mushi A, de Savigny D, Mgalula L, Mbuya C, Victora CG. The silent burden of anaemia in Tanzanian children: a community-based study. *Bull World Health Organ* 2003; 81: 581-590.
- Schleiermacher D, Mercereau-Puijalon O. PCR-Genotypisierung von *Plasmodium falciparum* in der Schwangerschaft. *Chemother J* 2002; 11: 130-136.
- Schofield L, Novakovic S, Gerold P, Schwarz RT, McConville MJ, Tachado SD. Glycosylphosphatidylinositol toxin of *Plasmodium* up-regulates intercellular adhesion molecule-1, vascular cell adhesion molecule-1, and e-selection expression in vascular cells and increases leukocyte and parasite cytoadherence via tyrosine kinase-dependent signal transduction. *J Immunol* 1996; 145: 1886-1896.
- Schofield L, Hewitt MC, Evans K, Siomos NA, Seeberger PH. Synthetic GPI as a candidate antitoxic vaccine in a model of malaria. *Nature* 2002; 418: 785-789.
- Schrier SL, Rachmilewitz E, Mohandas N. Cellular and membrane properties of alpha and beta thalassemic erythrocytes are different: implication for differences in clinical manifestations. *Blood* 1989; 74: 2194-2202.
- Seguin MC, Klotz FW, Schneider I, Weir JP, Goodbary M, Slayter M, Raney JJ, Aniagolu JU, Green SJ. Induction of nitric oxide synthase protects against malaria in mice exposed to irradiated *Plasmodium berghei* infected mosquitoes: involvement of interferon gamma and CD8+ T cells. *J Exp Med* 1994; 180: 353-358.
- Seixas EM, Langhorne J. Gammadelta T cells contribute to control of chronic parasitemia in *Plasmodium chabaudi* infections in mice. *J Immunol* 1999; 162: 2837-2841.
- Senok AC, Li K, Nelson EA, Arumanayagam M, Li CK. Flow cytometric assessment of oxidant stress in age-fractionated thalassaemic trait erythrocytes and its relationship to *in vitro* growth of *Plasmodium falciparum*. *Parasitology* 1998; 116 (Pt 1): 1-6.
- Sargent E, Parrot L. L'immunité, la prémunition et la résistance innée. *Arch Inst Pasteur Algerie* 1945; 13: 279.
- Shear HL, Roth EF Jr, Fabry ME, Costantini FD, Pachnis A, Hood A, Nagel RL. Transgenic mice expressing human sickle hemoglobin are partially resistant to rodent malaria. *Blood* 1993; 81: 222-226.
- Shear HL, Grinberg L, Gilman J, Fabry ME, Stamatoyannopoulos G, Goldberg DE, Nagel RL. Transgenic mice expressing human fetal globin are protected from malaria by a novel mechanism. *Blood* 1998; 92: 2520-2526.
- Shretta R, Omumbo J, Rapuoda B, Snow RW. Using evidence to change antimalarial drug policy in Kenya. *Trop Med Int Health* 2000; 5: 755-764.

- Singh B, Bobogare A, Cox-Singh J, Snounou G, Abdullah MS, Rahman HA. A genus- and species-specific nested polymerase chain reaction malaria detection assay for epidemiologic studies. *Am J Trop Med Hyg* 1999; 60: 687-692.
- Smetanina NS, Huisman THJ. Detection of α -thalassemia-2 and its corresponding triplication $\alpha\alpha\alpha$ (anti-3.7 kb) by PCR: an improved technical change. *Am J Hematol* 1996; 53: 202-203.
- Smith JA. The natural history of sickle cell disease. *Ann New York Acad Sci* 1989; 565: 104-108.
- Smith T, Felger I, Tanner M, Beck HP. Premunition in *Plasmodium falciparum* infection: insights from the epidemiology of multiple infections. *Trans R Soc Trop Med Hyg* 1999a; 93 (Suppl. 1): 59-64.
- Smith T, Beck HP, Kitua A, Mwankusye S, Felger I, Fraser-Hurt N, Irion A, Alonso P, Teuscher T, Tanner M. Age dependence of the multiplicity of *Plasmodium falciparum* infections and of other malariological indices in an area of high endemicity. *Trans R Soc Trop Med Hyg* 1999b; 93 (Suppl. 1): 15-20.
- Smith T, Felger I, Kitua A, Tanner M, Beck HP. Dynamics of multiple *Plasmodium falciparum* infections in a highly endemic area of Tanzania. *Trans R Soc Trop Med Hyg* 1999c ; 93 (Suppl. 1): 35-39.
- Smith T, Genton B, Baea K, Gibson N, Narara A, Alpers MP. Prospective risk of morbidity in relation to malaria infection in an area of high endemicity of multiple species of *Plasmodium*. *Am J Trop Med Hyg* 2001a; 64: 262-267.
- Smith T, Hii JL, Genton B, Muller I, Booth M, Gibson N, Narara A, Alpers MP. Associations of peak shifts in age-prevalence for human malarias with bednet coverage. *Trans R Soc Trop Med Hyg* 2001b; 95: 1-6.
- Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, do Rosario VE, Thaitong S, Brown KN. High sensitivity of detection of malaria parasites by the use of nested polymerase chain reaction. *Mol Biochem Parasitol* 1993a; 61: 315-320.
- Snounou G, Pinheiro L, Goncalves A, Fonseca L, Dias F, Brown KN, do Rosario VE. The importance of sensitive detection of malaria parasites in the human and insect hosts in epidemiological studies, as shown by the analysis of field samples from Guinea Bissau. *Trans R Soc Trop Med Hyg* 1993b; 87: 649-653.
- Snounou G, Zhu X, Siripoon N, Jarra W, Thaithong S, Brown KN, Viriyakosol S. Biased distribution of *msp1* and *msp2* allelic variants in *Plasmodium falciparum* populations in Thailand. *Trans R Soc Trop Med Hyg* 1999; 93: 369-374.
- Snow RW, Bastos de Azevedo I, Lowe BS, Kabiru EW, Nevill CG, Mwankusye S, Kassiga G, Marsh K, Teuscher T. Severe childhood malaria in two areas of markedly different falciparum transmission in east Africa. *Acta Trop* 1994; 57: 289-300.
- Snow RW, Omumbo JA, Lowe B, Molyneux CS, Obiero JO, Palmer A, Weber MW, Pinder M, Nahalen B, Obonyo C, Newbold C, Gupta S, Marsh K. Relation between severe malaria morbidity in children and level of *Plasmodium falciparum* transmission in Africa. *Lancet* 1997; 349: 1650-1654.
- Snow RW, Nahalen B, Palmer A, Donnelly CA, Gupta S, Marsh K. Risk of severe malaria among African infants: direct evidence of clinical protection during early infancy. *J Infect Dis* 1998;177: 819-822.
- Snow RW, Craig M, Deichmann U, Marsh K. Estimating mortality, morbidity and disability due to malaria among Africa's non-pregnant population. *Bull World Health Organ* 1999; 77: 624-640.
- Sokhna CS, Rogier C, Dieye A, Trape JF. Host factors affecting the delay of reappearance of *Plasmodium falciparum* after radical treatment among a semi-immune population exposed to intense perennial transmission. *Am J Trop Med Hyg* 2000; 62: 266-270.

- Stevenson MM, Riley EM. Innate immunity to malaria. *Nature Rev* 2004; 4: 169-180.
- Sullivan AD, Nyirenda T, Cullinan T, Taylor T, Harlow SD, James SA, Meshnick S. Malaria infection during pregnancy: intrauterine growth retardation and preterm delivery in Malawi. *J Infect Dis* 1999; 179: 1580-1583.
- Sulzer AJ, Deloron P, Cantella RA, Collins WE, Ngyen-Dinh P. Antibodies to the ring-infected surface antigen of *Plasmodium falciparum* elicited by infection with *Plasmodium malariae*. *Infect Immun* 1998; 56: 729-733.
- Taylor TE, Borgstein A, Molyneux ME. Acid-base status in paediatric *Plasmodium falciparum* malaria. *QJM* 1993; 86: 99-109.
- Taylor-Robinson AW, Phillips RS, Severn A, Moncada S, Liew FY. The role of TH1 and TH2 cells in a rodent malaria infection. *Science* 1993; 260: 1931-1934.
- Terlouw DJ, Aidoo MA, Udhayakumar V, Kolczak MS, Oloo AJ, Kager PA, Lal AA, Nahlen BL, ter Kuile FO. Increased efficacy of sulfadoxine-pyrimethamine in the treatment of uncomplicated falciparum malaria among children with sickle cell trait in western Kenya. *J Infect Dis* 2002; 186: 1661-1668.
- Thompson GR. Significance of haemoglobins S and C in Ghana. *Br Med J* 1962; 1: 682-685.
- Tian LP, Nelson EA, Senok AC, Yu LM, Oppenheimer SJ, Li K. Red cell age and susceptibility to malaria during pregnancy. *Acta Obstet Gynecol Scand* 1998; 77: 717-721.
- Trape JF, Rogier C, Konate L, Diagne N, Bouganali H, Canque B, Legros F, Badji A, Ndiaye G, Ndiaye P, Brahimi K, Faye O, Druilhe P, da Silva LP. The Dielmo project: a longitudinal study on natural malaria infection and the mechanisms of protective immunity in a community living in a holoendemic area of Senegal. *Am J Trop Med Hyg* 1994; 51: 123-137.
- Trape JF, Pison G, Spiegel A, Enel C, Rogier, C. Combating malaria in Africa. *Trends Parasitol* 2002; 18: 224-230.
- Tsuji M, Romero P, Nussenzweig RS, Zavala F. CD4+ cytolytic T cell clone confers protection against murine malaria. *J Exp Med* 1990; 172: 1353-1357.
- Turner DM, William DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphism in the interleukin-10 gene promoter. *Eur J Immunogenet* 1997; 24: 1-8.
- Turner G. Cerebral malaria. *Brain Pathol* 1997; 7: 569-82.
- Turrini F, Schwarzer E, Arese P. The involvement of hemozoin toxicity in depression of cellular immunity. *Parasitol Today* 1993; 9: 297-300.
- van Dongen PW, van't Hof MA. Sickle cell trait, malaria, and anaemia in pregnant Zambian women. *Trans R Soc Trop Med Hyg* 1983; 77: 402-404.
- van der Poll T, Jansen J, Levi M, ten Cate H, ten Cate JW, van Deventer SJH. Regulation of interleukin 10 release by tumor necrosis factor in humans and chimpanzees. *J Exp Med* 1994; 180: 1985-1988.
- Verhoef H, West CE, Ndeto P, Burema J, Beguin Y, Kok FJ. Serum transferrin receptor concentration indicates increased erythropoiesis in Kenyan children with asymptomatic malaria. *Am J Clin Nutr* 2001; 74: 767-775.
- Verhoef H, West CE, Kraaijenhagen R, Nzyuko SM, King R, Mbandi MM, van Laatum S, Hogervorst R, Schep C, Kok FJ. Malarial anemia leads to adequately increased erythropoiesis in asymptomatic Kenyan children. *Blood* 2002; 100: 3489-3494.
- Vleugels MP, Brabin B, Eling WM, de Graaf R. Cortisol and *Plasmodium falciparum* infection in pregnant women in Kenya. *Trans R Soc Trop Med Hyg* 1989; 83: 173-177.
- Wagner G, Koram K, McGuiness D, Benett S, Nkrumah F, Riley E. High incidence of asymptomatic malaria infections in a birth cohort of children less than one year of age in Ghana, detected by multicopy gene polymerase chain reaction. *Am J Trop Med Hyg*, 1998; 59: 115-112.

- Waller D, Krishna S, Crawley J, Miller K, Nosten F, Chapman D, ter Kuile FO, Craddock C, Berry C, Holloway PA, Brewster D, Greenwood BM, White NJ. Clinical features and outcome of severe malaria in Gambian children. *Clin Infect Dis* 1995; 21: 577-587.
- Warrell DA, Molyneux ME, Beales PF (editors). Severe and complicated malaria, 2nd edition. *Trans R Soc Trop Med Hyg* 1990; 84, suppl. 2
- Webster HK, Brown AE, Chuenchitra C, Permanich B, Pipithkul J. Characterization of antibodies to sporozoites in *Plasmodium falciparum* malaria and correlation with protection. *J Clin Microbiol* 1988; 26: 923-927.
- Wernsdorfer WH. The development and spread of drug-resistant malaria. *Parasitol Today* 1991; 7: 297-303.
- Wetsteyn JC, de Vries PJ, Oosterhuis B, van Boxtel CJ. The pharmacokinetics of three multiple dose regimens of chloroquine: implications for malaria chemoprophylaxis. *Br J Clin Pharmacol* 1995; 39: 696-699.
- White NJ. Malaria. in: Cook, GC (Hrsg.): *Manson's Tropical Diseases*. S. 1087-1165. WB Saunders Company Ltd., London, 1996
- WHO. Expert Committee on Malaria. *World Health Organ Tech Rep Ser* 2000a; 892: i-v, 1-74.
- WHO. Severe falciparum malaria. *Trans R Soc Trop Med Hyg* 2000b; 94 (Suppl. 1): 1-74.
- WHO. The use of antimalarial drugs. WHO/CDS/RBM/2001.33. World Health Organization, Genf, 2001
- Wickramasinghe SN, Abdalla SH. Blood and bone marrow changes in malaria. *Balliere's Best Practice Res Clin Haematol* 2000; 13: 277-299.
- Wilairat P, Kittikalayawong A, Chaicharoen S. The thalassemic red cell membrane. *Southeast Asian J Trop Med Public Health* 1992; 23 Suppl 2: 74-78.
- Wilkinson RJ, Pasvol G. Host resistance to malaria runs into swampy water. *Trends Microbiol* 1997; 5: 213-215.
- Willcox M, Björkman A, Brohult J. Falciparum malaria and β-thalassaemia trait in northern Liberia. *Ann Trop Med Parasitol* 1983; 77: 173: 785-791.
- Williams TN, Maitland K, Bennett S, Ganczakowski M, Peto TEA, Newbold CI, Bowden DK, Weatherall DJ, Clegg JB. High incidence of malaria in α-thalassaemic children. *Nature* 1996; 383: 522-525.
- Williams TN, Weatherall DJ, Newbold CI. The membrane characteristics of *Plasmodium falciparum*-infected and -uninfected heterozygous alpha(0)thalassaemic erythrocytes. *Br J Haematol* 2002; 118: 663-670.
- Xu W, Liu L, Emson PC, Harrington CR, Charles IG. Evolution of a homopurine-homopyrimidine pentanucleotide repeat sequence upstream of the human inducible nitric oxide synthase gene. *Gene* 1997; 204: 165-170.
- Yuthavong Y, Butthep A, Bunyaratvej S, Fucharoen S, Khusmith S. Impaired parasite growth and increased susceptibility to phagocytosis of *Plasmodium falciparum* infected α-thalassemia or Hb Constant Spring red blood cells. *Am J Clin Pathol* 1988; 89: 521-525.
- Yuthavong Y, Butthep P, Bunyaratvej A, Fucharoen S. Decreased sensitivity to artesunate and chloroquine of *Plasmodium falciparum* infecting hemoglobin H and/or hemoglobin Constant Spring erythrocytes. *J Clin Invest* 1989; 83: 502-505.