

8 Literaturverzeichnis

- Abboud, S. & Haile, D.J. (2000)
A novel mammalian iron-regulated protein involved in intracellular iron metabolism.
J Biol Chem, **275**(26), 19906–19912.
- Ajioka, R.S., Phillips, J.D. & Kushner, J.P. (2006)
Biosynthesis of heme in mammals.
Biochimica et Biophysica Acta, **1763**, 723–736, review.
- Akompong, T., Inman, R.S. & Wessling-Resnick, M. (1995)
Phorbol Esters Stimulate Non-transferrin Iron Uptake by K562 Cells.
J Biol Chem, **270**(36), 20937–20941.
- Alitalo, R. (1990)
Induced differentiation of K562 leukemia cells: a model for studies of gene expression in early megakaryoblasts.
Leuk Res, **14**(6), 501–514.
- Althoff, K., Reddy, P., Voltz, N., Rose-John, S. & Mullberg, J. (2000)
Shedding of interleukin-6 receptor and tumor necrosis factor alpha. Contribution of the stalk sequence to the cleavage pattern of transmembrane proteins.
Eur J Biochem, **267**, 2624–2631.
- Altschul S.F., Madden T.L., Schaffer A.A., Zhang J., Zhang Z., Miller W. & Lipman D.J. (1997)
Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.
Nucleic Acids Res, **25**, 3389–3402.
- Alvarez, E., Gironès, N. & Davis, R.J. (1990)
Inhibition of the receptor-mediated endocytosis of diferric transferrin is associated with the covalent modification of the transferrin receptor with palmitic acid.
J Biol Chem, **265**(27), 16644–16655.
- Aro, A.L.A., Savikko, J., Pulkkinen, V. & von Willebrand, E. (2002)
Expression of insulin-like growth factors IGF-I and IGF-II, and their receptors during the growth and megakaryocytic differentiation of K562 cells.
Leuk Res, **26**, 831–837.
- Arosio, P., Adelman, T.G. & Drysdale, J.W. (1978)
On ferritin heterogeneity. Further evidence for heteropolymers.
J Biol Chem, **253**(12), 4451–4458.
- Ausubel, F.M., Brent, R., Kingston, R.E., Moore, D.D., Seidman, J.G., Smith, J.A. & Struhl, K. (1997)
Short Protocols in Molecular Biology.
John Wiley & Sons, Inc.; New York, Chichester, Weinheim, Brisbane, Singapore, Toronto.
- Aziz, N. & Munro, H.N. (1987)
Iron regulates ferritin mRNA translation through a segment of its 5' untranslated region.
Proc Natl Acad Sci USA, **84**, 8478–8482.
- Barr, R.D. & Koekebakker, M. (1990)
Acetylcholinesterase in the human erythron.III. Regulation of differentiation.
Am J Hematol, **34**(4), 259–261.

- Bartha, E., Olah, E., Szelenyi, J.G. & Hollan, S.R. (1987)
Characterisation of „fetal-type“ acetylcholinesterase in hemin-treated K562 cell culture.
Blood Cells, **12(3)**, 647–655.
- Bauminger, E.R., Harrison, P.M., Hechel, D., Hodson, N.W., Nowik, I., Treffry, A. & Yewdall, S. J. (1993)
Iron (II) oxidation and early intermediates of iron-core formation in recombinant human H-chain ferritin.
Biochem J, **296**, 709–719.
- Beaumont, C., Leneuve, P., Devaux, I., Scoazec, J.Y., Berthier, M., Loiseau, M.N., Grandchamp, B. & Bonneau, D. (1995)
Mutation in the iron responsive element of the L ferritin mRNA in a family with dominant hyperferritinaemia and cataract.
Nat Genet, **11(4)**, 444–446.
- Beinert, H., Kennedy M.C. & Stout, C.D. (1996)
Aconitase as iron-sulfurprotein, enzyme, and iron regulatory protein.
Chem Rev, **96**, 2335–2373.
- Belhacène, N., Maulon, L., Guérin, S., Ricci, J.E., Mari, B., Colin, Y., Cartron, J.P. & Auberger, P. (1998)
Differential expression of the Kell blood group and CD10 antigens: two related membrane metallopeptidases during differentiation of K562 cells by phorbol ester and hemin.
FASEB J, **12(7)**, 531–539.
- Beutler, E. (1997)
The significance of the 187G (H63D) mutation in hemochromatosis.
Am J Hum Genet, **61**, 762–764.
- Bhasker, C.R., Burgiel, G., Neupert, B., Emery-Goodman, A., Kuhn, L.C. & May, B.K. (1993)
The putative iron-responsive element in the human erythroid 5-aminolevulinate synthase mRNA mediates translational control.
J Biol Chem, **268**, 12699–12705.
- Borhani, D.W. & Harrison, S.C. (1991)
Crystallization and X-ray diffraction studies of a soluble form of the human transferring receptor.
J Mol Biol, **218**, 685–689.
- Burstein, S.A. & Ishibashi, T. (1989)
Erythropoietin and megakaryocytopoiesis.
Blood Cells, **15(1)**, 193–201.
- Cairo, G. & Pietrangelo, A. (1994)
Transferrin receptor gene expression during rat liver regeneration: evidence for post-transcriptional regulation by iron regulatory factor B, a second iron-responsive element binding protein.
J Biol Chem, **269**, 6405–6409.
- Cairo, G., Tacchini, L. & Pietrangelo, A. (1998)
Lack of coordinate control of ferritin and transferrin receptor expression during rat liver regeneration.
Hepatology, **28(1)**, 173–178.
- Canonne-Hergaux, F., Gruenheid, S., Ponka, P. & Gros, P. (1999)
Cellular and subcellular localization of the Nramp2 iron transporter in the intestinal brush border and regulation by dietary iron.
Blood, **93(12)**, 4406–4417.
- Canonne-Hergaux, F. & Gros, P. (2002)
Expression of the iron transporter DMT1 in kidney from normal and anemic *mk* mice.
Kidney Int, **62**, 147–156.

- Carey, J.O., Posekany, K.J., deVente, J.E., Pettit, G.R. & Ways, D.K. (1996)
Phorbol-ester-stimulated phosphorylation of PU.1: association with leukemic cell growth inhibition.
Blood, **87**, 4316–4324.
- Casado, M., Vallet, V.S., Kahn, A. & Vaulont, S. (1999)
Essential role in vivo of upstream stimulatory factors for a normal dietary response of the fatty acid synthase gene in the liver.
J Biol Chem, **274**, 2009–2013.
- Casey, J.L., Hentze, M.W., Koeller, D.M., Caughman, S.W., Rouault, T.A., Klausner, R.D. & Harford, J.B. (1988)
Iron responsive elements: Regulatory RNA sequences that control mRNA levels and translation,
Science **240**, 924–928.
- Cazzola, M., Beguin, Y., Bergamashi, G., Guardone, R., Cerani, P., Barella, S., Cao, A. & Galanello, R. (1999).
Soluble transferrin receptor as a potential determinant of iron loading in congenital anaemias due to ineffective erythropoiesis.
Br J Haematol, **106**, 752–755.
- Chan, R.Y., Seiser, C., Schulman, H.M., Kuhn, L.C. & Ponka, P. (1994)
Regulation of transferrin receptor mRNA expression: distinct regulatory features in erythroid cells.
Eur J Biochem, **220**, 683–692.
- Chang, C., Hora, N., Huberman, N., Hinderer, R., KuKuruga, M. & Hanash, S. (2001)
Onkoprotein 18 levels and phosphorylation mediate megakaryocyte polyplloidization in human erythroleukemia cells.
Proteomics, **1**, 1415–1423.
- Chasteen, N.D. & Harrison, P.M. (1999)
Mineralization in ferritin: an efficient means of iron storage.
J Struct Biol, **126**, 182–194.
- Chen, S.H., Lin, C.Y., Cho, C.S., Lo, C.Z. & Hsiung, C.A. (2003)
Primer Design Assistant (PDA): a web-based primer design tool.
Nucleic Acids Res, **31(13)**, 3751–3754.
- Cheng, T., Wang, Y. & Dai, W. (1994)
Transcription factor egr-1 is involved in phorbol 12-myristate-13-acetate-induced megakaryocytic differentiation of K562 cells.
J Biol Chem, **269**, 30848–30853.
- Cianetti, L., Segnalini, P., Calzolari, A., Morsilli, O., Felicetti, F., Ramoni, C., Gabbianni, M., Testa, U. & Sposi, N.M. (2005)
Expression of alternative transcripts of ferroportin-1 during human erythroid differentiation.
Haematologica, **90(12)**, 1595–1606.
- Cmejla, R., Petrak, J. & Cmejlova, J. (2006)
A novel iron responsive element in the 3'UTR of human MRCK alpha.
Biochem. Biophys. Res. Commun, **341**, 158–166.
- Coccia, E.M., Stellacci, E., Orsatti, R., Testa, U. & Battistini, A. (1995)
Regulation of ferritin H-chain expression in differentiating Friend leukemia cells.
Blood, **86(4)**, 1570–1579.
- Collawn, J.F., Lai, A., Domingo, D., Fitch, M., Hatton, S. & Trowbridge, I.S. (1993)
YTRF is the conserved internalisation signal of the transferrin receptor, and a second YTRF signal at position 31–34 enhances endocytosis.
J Biol Chem, **268(29)**, 21686–21692.

- Collawn, J.F., Stangel, M., Kuhn, L.A., Esekogwu, V., Jing, S.Q., Trowbridge, I.S. & Tainer, J.A. (1990) Transferrin receptor internalisation sequence YXRF implicates a tight turn as the structural recognition motif for endocytosis.
Cell, **63(5)**, 1061–1072.
- Conrad, M.E., Umbreit, J.N. & Moore, E.G. (1993) Rat duodenal iron-binding protein mobilferrin is a homologue of calreticulin.
Gastroenterology, **104**, 1700–1704.
- Conrad, M.E., Umbreit, J.N., Moore, E.G., Uzel, C. & Berry, M.R. (1994) The alternate iron pathway. Mobilferrin and integrin in K562 cells.
J Biol Chem, **269**, 7169–7173.
- Conrad, M.E., Umbreit, J.N., Moore, E.G. & Heiman, D. (1996) Mobilferrin is an intermediate in iron transport between transferrin and hemoglobin in K562 cells.
J Clin Invest, **98(6)**, 1449–1454.
- Conrad, M.E. & Umbreit, J.N. (2000) Iron absorption and transport - an update.
Am J Hematol, **64(4)**, 287–298.
- Conrad, M.E., Umbreit, J.N., Moore, E.G., Hainsworth, L.N., Porubcin, M., Simovich, M.J., Nakada, M.T. & Garrick, M.D. (2000) Separate pathways for cellular uptake of ferric and ferrous iron.
Am J Physiol Gastrointest Liver Physiol, **279**, G767–G774.
- Corsi, B., Cozzi, A., Arosio, P., Drysdale, J., Santambrogio, P., Campanella, A., Biasiotto, G., Alberini, A. & Levi, S. (2002) Human mitochondrial ferritin expressed in HeLa cells incorporates iron and affects cellular iron metabolism.
J Biol Chem, **277**, 22430–22437.
- Cozzi, A., Santambrogio, P., Levi, S. & Arosio, P. (1990) Iron detoxifying activity of ferritin. Effects of H and L human apoferritins on lipid peroxidation in vitro.
FEBS Lett, **277(1-2)**, 119–122.
- Cox, T.C., Bawden, M.J., Martin, A. & May, B.K. (1991) Human erythroid 5-aminolevulinate synthase: promoter analysis and identification of an IRE in the mRNA.
EMBO J, **10**, 1891–1902.
- Crichton, R.R. (2001) Inorganic Biochemistry of Iron Metabolism.
John Wiley Sons, New York.
- Davis, R.J., Johnson, G.L., Kelleher, D.J., Anderson, J.K., Mole, J.E. & Czech, M.P. (1986a) Identification of serine 24 as the unique site on the transferrin receptor phosphorylated by protein kinase C.
J Biol Chem, **261**, 9034–9041.
- Davis, J.Q., Dansereau, D., Johnstone, R.M. & Bennett, V. (1986b) Selective externalisation of an ATP-binding protein structurally related to the clathrin-uncoating ATPase/heat shock protein in vesicles containing terminal transferring receptors during reticulocyte maturation.
J Biol Chem, **261**, 15368–15371.
- De Domenico, I., Vaughn, M.B., Li, L., Bagley, D., Musci, G., Ward, D.M. & Kaplan, J. (2006) Ferroportin-mediated mobilization of ferritin iron precedes ferritin degradation by the proteasome.
EMBO J, **25(22)**, 5396–5404.
- De Domenico, I., Ward, D.M., Nemeth, E., Vaughn, M.B., Musci, G., Ganz, T. & Kaplan, J. (2005) The molecular basis of ferroportin-linked hemochromatosis.
Proc Natl Acad Sci U S A, **102(25)**, 8955–8960.

- De Domenico, I., Ward, D.M., Langelier, C., Vaughn, M.B., Nemeth, E., Sundquist, W.I., Ganz, T., Musci, G. & Kaplan, J. (2007)
The molecular mechanism of hepcidin mediated ferroportin down-regulation.
Mol Biol Cell, **18**(7), 2569–2578.
- Delgado-Canedo, A., Chies, J.A.B. & Nardi, N.B. (2005)
Induction of fetal haemoglobin expression in erythroid cells – A model based on iron availability signalling.
Med Hypoth, **65**, 932–936.
- Deliorman, S., Uyar, R. & Altuner, Y. (2000)
The Effect of Iron Bound Transferrin on Megakaryocytopoiesis in Bone Marrow Culture.
Turk J Med Sci, **30**, 35–38.
- Dempsey, P.J., Meise, K.S., Yoshitake, Y., Nishikawa, K. & Coffey, R.J. (1997)
Apical enrichment of human EGF precursor in Madin-Darby canine Kidney cells involves preferential basolateral ectodomain cleavage sensitive to a metalloprotease inhibitor.
J Cell Biol, **138**, 747–758.
- Denzer, K., Kleijmeer, M.J., Heijnen, H.F., Stoorvogel, W. & Geuze, H. (2000)
Exosome: From internal vesicle of the multivesicular body to intracellular signalling device.
J Cell Sci, **113**, 3365–3374.
- Deutsch, V.R. & Tomer, A. (2006)
Megakaryocyte development and platelet production.
Br J Haematol, **134**, 453–466, review.
- Donovan, A., Brownlie, A., Zhou, Y., Shepard, J., Pratt, S.J., Moynihan, J., Paw, B.H., Drejer, A., Barut, B., Zapata, A., Law, T.C., Brugnara, C., Lux, S.E., Pinkus, G.S., Pinkus, J.L., Kingsley, P.D., Palis, J., Fleming, M. D., Andrews, N.C. & Zon, L.I. (2000)
Positional cloning of zebrafish ferroportin1 identifies a conserved vertebrate iron exporter.
Nature, **403**, 776–781.
- Dorsey, J.F., Cunnick, J.M., Mane, S.M. & Wu, J. (2002)
Regulation of the Erk2-Elk1 signaling pathway and megakaryocytic differentiation of Bcr-Abl+ K562 leukemic cells by Gab2.
Blood, **99**, 1388–1397.
- Drakesmith, H., Sweetland, E., Schimanski, L., Edwards, J., Cowley, D., Ashraf, M., Bastin, J. & Townsend, A.R. (2002)
The hemochromatosis protein HFE inhibits iron export from macrophages.
Proc Natl Acad Sci U S A, **9**(24), 15602–15607.
- Drexler, H.G., Matsuo, Y. & MacLeod, R.A.F. (2004)
Malignant hematopoietic cell lines: in vitro models for the study of erythroleukemia.
Leuk Res, **28**, 1243–1251.
- Eisenstein, R.S. (2000)
Iron regulatory proteins and the molecular control of mammalian iron metabolism.
Annu Rev Nutr, **20**, 627–662.
- Ellman, G.L., Courtney, K.D., Anders, V. & Featherstone, R.M. (1961)
A new and rapid colorimetric determination of acetylcholinesterase activity.
Biochem Pharmacol, **7**, 88–95.
- Epsztejn, S., Glickstein, H., Picars, V., Slotki, I.N., Breuer, W., Beaumont, C. & Cabantchik, Z.I. (1999)
H-Ferritin Subunit Overexpression in Erythroid Cells Reduces Oxidative Stress Response and Induces Multidrug Resistance Properties.
Blood, **94**(10), 3593–3603.

- Evans, P. & Kemp, J. (1997)
Exon/intron structure of the human transferrin receptor gene.
Gene, **199**, 123–131.
- Feder, J. N., Gnarke, A., Thomas, W., Zsuehishashi, Z., Ruddy, D.A., Basava, A., Dormishian, F., Domingo, R., Ellis, M.C., Fullan, A., Hinton, L.M., Jones, N.L., Kimmel, B.E., Kronmal, G.S., Lauer, P., Lee, V.K., Loeb, D.B., Mapa, F.A., McCelland, E., Meyer, N.C., Mintier, G.A., Moeller, N., Moore, T., Morikang, E., Prass, C. E., Quintana, L., Starnes, S.M., Schatzman, R.C., Brunke, K.J., Drayna, D.T., Risch, N.J., Bacon, B.R. & Wolff, R. K. (1996)
A novel MHC class I-like gene is mutated in patients with hereditary haemochromatosis.
Nat Genet, **13**, 399–408.
- Feder, J.N., Penny, D.M., Irrinki, A., Lee, V.K., Lebro'n, J.A., Watson, N., Tsuehishashi, Z., Sigal, E., Bjorkman, P. J. & Schatzman, R.C. (1998)
The haemochromatosis gene product complexes with the transferrin receptor and lowers its affinity for ligand binding
Proc Natl Acad Sci U SA, **95**, 1472–1477.
- Feder, J.N., Tsuehishashi, Z., Irrinki, A., Lee, V.K., Mapa, F.A., Morikang, E., Prass, C.E., Starnes, S.M., Wolff, R.K., Parkkila, S., Sly, W.S. & Schatzman, R.C. (1997)
The haemochromatosis founder mutation in HLA H disrupts beta(2) microglobulin interaction and cell surface expression.
J Biol Chem, **272**, 14025–14028.
- Fish, W.W. (1988)
Rapid colorimetric Micromethod for the Quantitation of Complexed Iron in Biological Samples.
Meth Enzymol, **158**, 357–364.
- Fleming, R.E. & Britton, R.S. (2006)
Iron Imports. VI. HFE and regulation of intestinal iron absorption.
Am J Physiol Gastrointest Liver Physiol, **290**, G590–G594, review.
- Fleming, M.D., Romano, M.A., Su, M.A., Garrick, L.M., Garrick, M.D. & Andrews, N.C. (1998)
Nramp2 is mutated in the anemic belgrade (b) rat: evidence of a role for Nramp2 in endosomal iron transport.
Proc Natl Acad Sci U SA, **95**(3), 1148–1153.
- Flowers, C.H., Skikne, B.S., Covell, A.M. & Cook, J.D. (1989)
The clinical measurement of serum transferrin receptor.
J Lab Clin Med, **114**, 368–377.
- Frazer, D.M., Vulpe, C.D., McKie, A.T., Wilkins, S.J., Trinder, D., Cleghorn, G.J. & Anderson, G.J. (2001)
Cloning and gastrointestinal expression of rat hephaestin: relationship to other iron transport proteins.
Am J Physiol Gastrointest Liver Physiol, **281**, G931–G939.
- Frishman, D. & Hentze, M.W. (1996)
Conservation of aconitase residues revealed by multiple sequence analysis: implications for structure/function relationships.
Eur J Biochem, **239**, 197–200.
- Fuchs, H. & Geßner, R. (2002)
Iodination significantly influences the binding of human transferrin to the transferrin receptor.
Biochim Biophys Acta, **1570**(1), 19–26.
- Ganz, T. (2003)
Hepcidin, a key regulator of iron metabolism and mediator of anemia and inflammation.
Blood, **102**, 783–788.

- Galli, A., Bergamaschi, G., Recalde, H., Biasiotto G., Santambrogio, P., Boggi, S., Levi, S., Arossio, P. & Cazzola, M. (2004)
Ferroportin gene silencing induces iron retention and enhances ferritin synthesis in human macrophages.
Brit J Haematol, **127**, 598–603.
- Garrett, C., Ravazzolo, R., Bianchi Scarra, G.L., Sessarego, M., Baressi, R. & Ajmar, F. (1984)
Regulation of acetylcholinesterase expression in the K-562 cell line.
Cancer Res, **44**, 3749.
- Gasparini, P. & Camaschella, C. (2001)
New mutations inactivating transferrin receptor 2 in hemochromatosis type 3.
Blood, **97**, 2555–2560.
- Giannetti, A.M. & Björkman, P.J. (2004)
HFE and Transferrin Directly Compete for Transferrin Receptor in Solution and at the Cell Surface.
J Biol Chem, **279**(24), 25866–25875.
- Glockner, G., Scherer, S., Schattevoy, R., Boright, A., Weber, J., Tsui, L.C. & Rosenthal, A. (1998)
Large-scale sequencing of two regions in human chromosome 7q22: analysis of 650 kb of genomic sequence around the EPO and CUTL1 loci reveals 17 genes.
Genome Res, **8**(10), 1060–1073.
- Goldfarb, A.N., Wong, D. & Racke F.K. (2001)
Induction of Megakaryocytic Differentiation in Primary Human Erythroblasts
A Physiological Basis for Leukemic Lineage Plasticity.
American Journal of Pathology, **158**(4), 1191–1198.
- Goswami, T. & Andrews, N.C. (2006)
Hereditary Hemochromatosis Protein, HFE, Interaction with Transferrin Receptor 2 Suggests a Molecular Mechanism for Mammalian Iron Sensing.
J Biol Chem, **281**(39), 28494–28498.
- Graham, R.M., Morgan, E.H. & Baker, E. (1998)
Ferric citrate uptake by cultured rat hepatocytes is inhibited in the presence of transferrin.
Eur J Biochem, **253**, 139–145.
- Grootveld, M., Bell, J.D., Halliwell, B., Aruoma, O. I., Bomford, A. & Sadler P.J. (1989)
Non-transferrin-bound iron in plasma or serum from patients with idiopathic hemochromatosis. Characterization by high performance liquid chromatography and nuclear magnetic resonance spectroscopy.
J Biol Chem, **264**, 4417–4422.
- Gruer, M.J., Artymiuk, P.J. & Guest, J.R. (1997)
The aconitase family: three structural variations on a common theme.
Trends Biochem Sci, **22**, 3–6.
- Gunshin, H., Fujiwara, Y., Custodio, A.O., DiRenzo, C., Robine, S. & Andrews, N.C. (2005)
Slc11a2 is required for intestinal iron absorption and erythropoiesis but dispensable in placenta and liver.
J Clin Invest, **115**, 1258–1266.
- Gunshin, H., Mackenzie, B., Berger, U.V., Gunshin, Y., Romero, M.F., Boron, W.F., Nussberger, S., Gollan, J.L. & Hediger, M.A. (1997)
Cloning and characterization of a mammalian proton-coupled metal-ion transporter.
Nature, **388**, 482–488.
- Guo, B., Yu, Y. & Leibold, E.A. (1994)
Iron regulates cytoplasmic levels of a novel iron-responsive element-binding protein without aconitase activity.
J Biol Chem, **269**, 24252–24260.

- Guo, B., Phillips, J.D., Yu, Y. & Leibold, E.A. (1995)
Iron regulates the intracellular degradation of iron regulatory protein 2 by the proteasome.
J Biol Chem, **270**, 21645–21651.
- Gutierrez, J.A., Inman, R.S., Akompong, T., Yu, J. & Wessling-Resnick, M. (1998)
Metabolic depletion inhibits the uptake of nontransferrin-bound iron by K562 cells.
J Cell Physiol, **177**, 585–592.
- Han, A., Yu, C., Lu, L., Fujiwara, Y., Browne, C., Chin, G., Fleming, M., Leboulch, P., Orkin, S.H. & Chen, J.J. (2001)
HRI is required for translational regulation and survival of erythroid precursors in iron-deficiency.
EMBO J, **20(23)**, 6909–6918.
- Hanahan, D. (1983)
Studies on transformation of Escherichia coli with plasmids.
J Mol Biol, **166(4)**, 557–580.
- Hannuksela, J., Parkkila, S., Waheed, A., Britton, R.S., Fleming, R.E., Bacon, B.R. & Sly, W-S. (2003)
Human platelets express hemochromatose protein (HFE) and transferrin receptor.
Eur J Haematol, **70**, 201–206.
- Hanson, E.S., Foot, L.M. & Leibold E.A. (1999)
Hypoxia post-translationally activates iron-regulatory protein 2.
J Biol Chem, **274**, 5047–5052.
- Harding, C., Heuser, J. & Stahl, P. (1983)
Receptor-mediated endocytosis of transferrin and recycling of the transferrin receptor in rat reticulocytes.
J Cell Biol, **97**, 329–339.
- Harding, C., Heuser, J. & Stahl, P. (1984)
Endocytosis and intracellular processing of transferrin and colloidal gold-transferrin in rat reticulocytes.
J Cell Biol, **35**, 256–263.
- Harris, Z.L., Durley, A.P., Man, T.K. & Gitlin, J.D. (1999)
Targeted gene disruption reveals an essential role for ceruloplasmin in cellular iron efflux.
Proc Natl Acad Sci USA, **96**, 10812–10817.
- Harrison, P. M., Hempstead, P. D., Artymiuk, P. J. & Andrews S.C. (1998)
Structure-function relationships in the ferritins.
Met Ions Biol Syst, **35**, 435–477.
- Heijnen, H.F., Schiel, A.E., Fijnheer, R., Geuze, H.J. & Sixma, J.J. (1999)
Activated platelets release two types of membrane vesicles: microvesicles by surface shedding exosomes derived from exocytosis of multivesicular bodies and alpha granules.
Blood, **94**, 3791–3799.
- Heile, D.J., Rouault, T.A., Harford, J.B., Kennedy, M.C., Blondin, G.A., Beinert, H. & Klausner, R.D. (1992)
Cellular regulation of the iron-responsive element binding protein: disassembly of the cubane iron–sulfur cluster results in high-affinity RNA binding,
Proc Natl Acad Sci USA, **89**, 11735–11739.
- Henderson, B.R., Seiser, C. & Kuhn, L.C. (1993)
Characterization of a second RNA-binding protein in rodents with specificity for iron-responsive elements.
J Biol Chem, **268**, 27327–27334.
- Hentze, M.W., Muckenthaler, M.U. & Andrews, N.C. (2004)
Balancing acts: molecular control of mammalian iron metabolism.
Cell, **117**, 285–297.

- Herrera, R., Hubbell, S., Decker, S. & Petruzzelli, L. (1998)
A role for the MEK/MAPK pathway in PMA-induced cell cycle arrest: modulation of differentiation of K562 cells.
Exp Cell Res, **238**, 407–414.
- Horita, M., Andrei, E.J. & Benito, A. (2000)
Blockage of the Bcr-Abl kinase activity induces apoptosis of chronic myelogenous leukemia cells by suppressing signal transducer and activator of transcription 5-dependent expression of Bcl-XL.
J Exp Med, **191**, 977–984.
- Hubert, N., & Hentze, M.W. (2002)
Previously uncharacterized isoforms of divalent metal transporter (DMT)-1: implications for regulation and cellular function.
Proc Natl Acad Sci USA, **99**, 12345–12350.
- Huo, X-F., Yu, J., Peng, H., Du, Z-W., Liu, X-L., Ma, Y-Ni., Zhang, X., Zhang, Y., Zhao, H-L. & Zhang, J-W. (2006)
Differential expression changes in K562 cells during the hemin-induced erythroid differentiation and the phorbol myristate acetate PMA)-induced megakaryocytic differentiation.
Molecular and Cellular Biochemistry, **292**, 155–167.
- Iancu-Rubin, C., French, D.L. & Atweh, G.F. (2003)
Stathmin expression and megakaryocyte differentiation. A potential role in polyploidy.
Exp Hematol, **31**, 389–397.
- Iancu-Rubin, C., Nasrallah, C.A. & Atweh, G.F. (2005)
Stathmin Prevents the Transition from a Normal to an Endomitotic Cell Cycle during Megakaryocytic Differentiation.
Cell Cycle, **4(12)**, 1774–1782.
- Ishibashi, T., Kimura, H., Shikama, Y., Uchida, T., Kariyone, S. & Maruyama, Y. (1990)
Effect of recombinant granulocyte-macrophage colony-stimulating factor on murine thrombopoiesis in vitro and in vivo.
Blood, **75(7)**, 1433–1438.
- Ishida, Y., Levi, J., Baker, G., Stenberg, P.E., Yamada, Y., Sasaki, H. & Inoue, T. (1993)
Biological and biochemical characteristics of murine megakaryoblastic cell line L8057.
Exp Hematol, **21(2)**, 289–298
- Iwai, K., Klausner, R.D. & Rouault T.A. (1995)
Requirements for iron-regulated degradation of the RNA binding protein, iron regulatory protein 2.
EMBO J, **14**, 5350–5357.
- Iwasaki, K., MacKanzie, E.L., Hailemariam, K., Sakamoto, K. & Tsuji, Y. (2006)
Hemin-Mediated Regulation of an Antioxidant-Responsive-Element of the Human Ferritin H Gene and Role of Ref-1 during Erythroid Differentiation of K562 Cells.
Mol Cell Biol, **26(7)**, 2845–2856.
- Jackson, C.W. (1973)
Cholinesterase as a possible marker of early cells of the megakaryocytic series.
Blood, **42**, 413.
- Jacquel, A., Herrant, M., Defamie, V., Belhacene, N., Colosetti, P., Marchetti, S., Legros, L., Deckert, M., Mari, B., Cassuto, J.P., Hofman, P. & Auberger, P. (2006)
A survey of the signaling pathways involved in megakaryocytic differentiation of the human K562 leukemia cell line by molecular and c-DNA array analysis.
Oncogene, **25(5)**, 781–794.

- Jeong, S.Y. & David, S. (2003)
Glycosylphosphatidylinositol-anchored ceruloplasmin is required for iron efflux from cells in the central nervous system.
J Biol Chem, **278**, 27144–27148.
- Johnstone, R.M., Mathew, A., Mason, A.B. & Teng, K. (1991)
Exosome formation during maturation of mammalian and avian reticulocytes: evidence that exosome release is a major route for externalization of obsolete membrane proteins.
J Cell Physiol, **147(1)**, 27–36.
- Johnstone, R.M. (1992)
The Jeanne Manery-Fisher Memorial Lecture. Maturation of reticulocytes: formation of exosomes as a mechanism for shedding membrane proteins.
Biochem Cell Biol, **70**, 179–190.
- Kakhlon, O., Gruenbaum, Y. & Cabantchik, Z.I. (2001)
Repression of the heavy ferritin chain increases the labile iron pool of human K562 cells.
Biochem J, **356**, 311–316.
- Karlson, P., Doenecke, D. & Koolman, J. (1994)
Biochemie
Georg Thieme Verlag, Stuttgart. New York.
- Kaup, M., Dassler, K., Weise, C. & Fuchs, H. (2002a)
Shedding of the transferrinrezeptor is mediated constitutively by an integral membrane metalloprotease sensitve to tumor necrosis factor alpha protease inhibitor-2.
J Biol Chem, **277(41)**, 38498–38502.
- Kaup, M., Dassler, K., Reinecke, U., Weise, C., Tauber, R. & Fuchs, H. (2002b)
Prozessing of the human transferrin receptor at dinstinct positions within stalk region by neutrophil elastase and cathepsin G.
Biol Chem, **383(6)**, 1011–1020.
- Kawabata, H., Yang, R., Hirama, T., Vuong, P.T., Kawano, S., Gombart, A.F. & Koeffler, H.P. (1999)
Molecular cloning of transferrin receptor 2. A new member of the transferrin receptor-like family.
J Biol Chem, **274**, 20826–20832.
- Kawabata, H., Nakamaki, T., Ikonomi, P., Smith, R.D., Germani, R.S. & Koeffler, H.P. (2001)
Expression of transferrin receptor 2 in normal and neoplastic hematopoietic cells.
Blood, **98**, 2714–2719.
- Kawasaki, N., Morimoto, K., Tanimoto, T. & Hayakawa, T. (1996)
Control of hemoglobin synthesis in erythroid differentiating K562 cells. I. Role of iron in erythroid cell heme synthesis.
Arch Biochem Biophys, **328(2)**, 289–294.
- Ke, Y. & Theil, E.C. (2002)
An mRNA loop/bulge in the ferritin iron-responsive element forms in vivo and was detected by radical probing with Cu-1,10-phenanthroline and iron regulatory protein footprinting.
J Biol Chem, **277**, 2373–2376.
- Kim, H-Y., LaVaute, T., Iwai, K., Klausner R.D. & Rouault, T.A. (1996)
Identification of a conserved and functional iron-responsive element in the 5'-untranslated region of mammalian mitochondrial aconitase.
J Biol Chem, **271**, 24226–24230.
- Klausner, R.D., Harford, J. & van Renswoude, J. (1984)
Rapid internalization of the transferrin receptor in K562 cells is triggered by ligand binding or treatment with a phorbol ester.
Proc Natl Acad Sci U S A, **81(10)**, 3005–3009.

- Kohler, S.A., Henderson, B.R. & Kuhn, L.C. (1995)
Succinate dehydrogenase B mRNA of *Drosophila melanogaster* has a functional iron-responsive element in its 5'-untranslated region.
J Biol Chem, **270**, 307813–0786.
- Kohno, H. & Tokunaga, R. (1985)
Transferrin and iron uptake by rat reticulocytes.
J Biochem (Tokyo), **97**, 1181–1188.
- Konijn, A.M., Glickstein, H., Vaisman, B., Meyron-Holtz, E.G., Slotki, I.N. & Cabantchik, Z.I. (1999)
The Cellular Labile Iron Pool and Intracellular Ferritin in K562 Cells.
Blood, **94(6)**, 2128–2134.
- Krause, A., Neitz, S., Magert H.J., Schultz, A., Forssmann, W-G., Schultz-Knappe, P. & Andermann, K. (2000)
LEAP-1, a novel highly disulfide-bonded human peptide, exhibits antimicrobial activity.
FEBS Lett, **480**, 147–150.
- Laemmli, U.K. (1970)
Cleavage of structural proteins during the assembly of the head of bacteriophage T4.
Nature, **227**, 680–685.
- Lammich, S., Kojro, E., Postina, R., Gilbert, S., Pfeiffer, R., Jasionowski, M., Haass, C. & Fahrenholz, F. (1999)
Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by disintegrin metalloprotease.
Proc Natl Acad Sci U SA, **96**, 3922–3927.
- Larsson, R. & Nygren, P. (1989)
A rapid fluorometric method for semiautomated determination of cytotoxicity and cellular proliferation of human tumor cell lines in microculture.
Anticancer Res, **9**, 1111–1119.
- Lawrence, C.M., Ray, S., Babyonyshov, M., Galluser, R., Borhani, D.W. & Harrison, S.C. (1999)
Crystal structure of the ectodomain of human transferrin receptor.
Science, **286**, 779–782.
- Le, N.T. & Richardson D.R. (2002)
Ferroportin1: a new iron export molecule?
Int J Biochem Cell Biol, **34**, 103–108.
- Lee, P.L., Gelbart, T., West, C., Halloran, C. & Beutler, E. (1998)
The human Nramp2 gene: characterization of the gene structure, alternative splicing, promoter region and polymorphisms.
Blood Cells Mol Dis, **24**, 199–215.
- Lebron, J.A., Bennett, M.J., Vaughn, D.E., Chirino, A.J., Snow, P.M., Mintier, G.A., Feder, J.N. & Bjorkman, P. J. (1998)
Crystal structure of the haemochromatosis protein HFE and characterisation of its interaction with transferrin receptor.
Cell, **93**, 111–123.
- Levi, S., Girelli, D., Perrone, F., Pasti, M., Beaumont, C., Corrocher, R., Albertini, A. & Arosio, P. (1998)
Analysis of ferritins in lymphoblastoid cell lines and in the lens of subjects with hereditary hyperferritinemia-cataract syndrome.
Blood, **91**, 4180–4187.
- Lewin, B. (1991)
Gene: Lehrbuch der molekularen Genetik.
VCH, Weinheim, Basel, Cambridge, New York.

- Liu, C-C., Chou, Y-L. & Ch'ang, L-Y. (2004a)
Down-Regulation of Human NDR Gene in Megakaryocytic Differentiation of Erythroleukemia K562 Cells
J Biomed Sci, **11**, 104–116.
- Liu, Y.L., Ang, S.O., Weigent, D.A., Prchal, J.T. & Bloomer, J.R. (2004b)
Regulation of ferrochelatase gene expression by hypoxia.
Life Sci, **75(17)**, 2035–2043.
- Long, M.W., Heffner, C.H., Williams, J.L., Peters, C. & Prochownik, E.V. (1990)
Regulation of megakaryocyte potential in human erythroleukemic cells.
J Clin Invest, **85**, 1072–1084.
- Lozzio, C.B. & Lozzio, B.B. (1975)
Human chronic myelogenous leukemia cell-line with positive Philadelphia-chromosome.
Blood, **45**, 321–334.
- Löffler, G. & Petrides, P.E. (1997)
Biochemie und Pathobiochemie.
Springer-Verlag, Berlin Heidelberg New York.
- Lumelsky, N.L. & Forget, B.G. (1991)
Negative Regulation of Globin Gene Expression during Megakaryocytic Differentiation of a Human Erythroleukemic Cell Line.
Mol Cell Biol, **11(7)**, 3528–3536.
- Martin, S., Lennon, S., Bonham, A. & Cotter, T. (1990)
Induction of apoptosis (programmed cell death) in human leukemic cells by inhibitors of RNA or protein synthesis.
J Immunol, **145**, 1859–1864.
- Matsumura-Takeda, K., Sogo, S., Isakari, Y., Harada, Y., Nishioka, K., Kawakami, T., Ono, T. & Taki, T. (2007)
CD41+/CD45+ cells without acetylcholinesterase activity are immature and a major megakaryocytic population in murine bone marrow.
Stem Cells, **25(4)**, 862–870.
- May, W.S., Jacobs, S. & Cuatrecasas, P. (1984)
Association of phorbol ester-induced hyperphosphorylation and reversible regulation of transferrin membrane receptors in HL60 cells.
Proc Natl Acad Sci U S A, **81(7)**, 2016–2020.
- McClelland, A., Kuhn, L.C. & Ruddle, F.H. (1984)
The human transferrin receptor gene: genomic organization, and the complete primary structure of the receptor deduced from a cDNA sequence.
Cell, **39(2 Pt 1)**, 267–274.
- McGahon, A., Bissonnette, R., Schmitt, M., Cotter, K.M., Green, D.R. & Cotter, T.G. (1994)
BCR-ABL maintains resistance of chronic myelogenous leukaemia cells to apoptotic cell death.
Blood, **5**, 1179–1187.
- McKie, A.T., Barrow, D., Latunde-Dada, G.O., Rolfs, A., Sager, G., Mudaly, E., Mudaly, M., Richardson, C., Barlow, D., Bomford, A., Peters, T.J., Raja, K.B., Shirali S., Hediger, M.A., Farzaneh, F. & Simpson, R.J. (2001)
An iron-regulated ferric reductase associated with the absorption of dietary iron.
Science, **291**, 1755–1759.

- McKie, A.T., Marciani, P., Rolfs, A., Brennan, K., Wehr, K., Barrow, D., Miret, S., Bomford, A., Peters, T.J., Farzaneh, F., Hediger, M.A., Hentze, M.W. & Simpson, R.J. (2000)
A novel duodenal iron-regulated transporter, IREG1, implicated in the basolateral transfer of iron to the circulation.
Mol Cell, **5**, 299–309.
- McWhirter, J.R. & Wang, J.Y.J. (1991)
Activation of tyrosine kinase and microfilament-binding functions of c-abl by bcr sequences in bcr/abl fusion proteins.
Mol Cell Biol, **11**, 1553–1557.
- Melefors, O., Goossen, B., Johansson, H.E., Stripecke, R., Gray, N.K. & Hentze, M.W. (1993)
Translational control of 5-aminolevulinate synthase mRNA by iron-responsive elements in erythroid cells.
J Biol Chem, **268**, 5974–5978.
- Melemed, A.S., Ryder, J.W. & Vik, T.A. (1997)
Activation of the mitogen-activated protein kinase pathway is involved in and sufficient for megakaryocytic differentiation of CMK cells.
Blood, **90**, 3462–3470.
- Merryweather, C.A., Pointon, J.J., Shearman, J.D. & Robson, K.J. (1997)
Global prevalence of putative haemochromatosis mutations.
J Med Genet, **134**, 275–278.
- Mikulits, W., Schranzhofer, M., Bauer, A., Dolznig, H., Lobmayr, L., Infante, A.A., Beug, H. & Müllner, E.W. (1999)
Impaired ferritin mRNA translation in primary erythroid progenitors: shift to iron-dependent regulation by the v-ErbA oncogene.
Blood, **94(12)**, 4321–4332.
- Miller, L.L., Miller, S.C., Torti, S.V., Tsuji, Y. & Torti, F.M. (1991)
Iron-independent induction of ferritin H chain by tumor necrosis factor.
Proc Natl Acad Sci USA, **88**, 4946–4950.
- Morgan, E.H. & Baker, E. (1988)
Role of transferrin receptors and endocytosis in iron uptake by hepatic and erythroid cells.
Ann NY Acad Sci, **526**, 65–82.
- Mullis, K., Falloona, F., Scharf, S., Saiki, R., Horn, G. & Erlich, H. (1986)
Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction.
Cold Spring Harb Symp Quant Biol, **51(1)**, 263–273.
- Mura, C., Le Gac, G., Jacolot, S. & Ferec, C. (2004)
Transcriptional regulation of the human HFE gene indicates high liver expression and erythropoiesis coregulation.
FASEB J, **18**, 1922–1924.
- Murate, T., Saga, S., Hotta, T., Asano, H., Ito, T., Kato, K., Tsushita K., Kinoshita, T., Ichikawa, A., Yoshida, S. & Saito, H. (1993)
The close relationship between DNA replication and the selection of differentiation lineages of human erythroleukemia cell lines K562, HEL, and TF1 into either erythroid or megakaryocytic lineages.
Exp Cell Res, **208**, 35–43.
- Murphy, J.J. & Norton, J.D. (1993)
Phorbol ester induction of early response gene expression in lymphocytic leukemia and normal human B-cells.
Leuk Res, **8**, 657–662.

- Murray, N.R., Baumgardner, G.P., Burns, D.J. & Fields, A.P. (1993)
Protein kinase C isotypes in human erythroleukemia (K562) cell proliferation and differentiation. Evidence that beta II protein kinase C is required for proliferation
J Biol Chem, **268**, 15847–15853.
- Müllner, E.W. & Kühn, L.C. (1988)
A stem-loop in the 3' untranslated region mediates iron-dependent regulation of transferrin receptor mRNA stability in the cytoplasm.
Cell, **53**, 815–825.
- Neckers, L.M. (1991)
Regulation of transferrin receptor expression and control of cell growth.
Pathobiology, **59**, 11–18.
- Nemeth, E., Tuttle, M.S., Powelson, J., Vaughn, M.B., Donovan, A., Ward, D.M., Ganz, T. & Kaplan, J. (2004)
Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization.
Science, **306**, 2090–2093.
- Nemeth, E., Valore, E.V., Territo, M., Schiller, G., Lichtenstein, A. & Ganz T. (2003)
Hepcidin, a putative mediator of anemia of inflammation, is a type II acute-phase protein.
Blood, **101**, 2461–2463.
- Neshat, M.S., Raitano, A.B., Wang, H.G., Reed, J.C. & Sawyers, C.L. (2000)
The survival function of the Bcr-Abl oncogene is mediated by Bad dependent and -independent pathways: roles of phosphatidylinositol 3-kinase and Raf.
Mol Cell Biol, **20**, 1179–1186.
- Nicolas, G., Bennoun, M., Devaux, I., Beaumont, C., Grandchamp, B., Kahn, A. & Vaulont, S. (2001)
Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice.
Proc Natl Acad Sci U S A, **98**, 8780–8785.
- Nicolas, G., Bennoun, M., Porteu, A., Mativet, S., Beaumont, C., Grandchamp, B., Siriti, M., Sawadogo, M., Kahn, A. & Vaulont, S. (2002a)
Severe iron deficiency anemia in transgenic mice expressing liver hepcidin.
Proc Natl Acad Sci U S A, **99**, 4596–4601.
- Nicolas, G., Chauvet, C., Viatte, L., Danan, J.L., Bigard, X., Devaux, I., Beaumont, C., Kahn, A. & Vaulont, S. (2002b)
The gene encoding the iron regulatory peptide hepcidin is regulated by anemia, hypoxia and inflammation.
J Clin Invest, **110**(7), 1037–1044.
- Nunes, R., Beloqui, O., Potter, B.J. & Berk, P.D. (1984)
Iron uptake from transferrin by isolated hepatocytes: effect of ethanol.
Biochem Biophys Res Commun, **125**, 824–830.
- Ogawa, K., Igarashi, K., Nishitani, C., Shibahara, S. & Fujita, H. (2002)
Heme-regulated transcription factor Bach1.
J Health Sci, **48**, 1–6.
- Owen, D., & Kuhn, L.C. (1987)
Noncoding 3' sequences of the transferrin receptor gene are required for mRNA regulation by iron.
EMBO J, **6**, 1287–1293.
- Oyake, T., Itoh, K., Motohashi, H., Hayashi, N., Hoshino, H., Nishizawa, M., Yamamoto, M. & Igarashi, K. (1996)
Bach proteins belong to a novel family of BTB-basic leukine zipper transcription factors that interact with Mafk and regulate transcription through the NF-E2 site.
Mol Cell Biol, **16**(11), 6083–6095.

- Paoletti, F., Vannucchi, A.M., Mocali, A., Caporale, R. & Burstein, S.A. (1995)
Identification and Conditions for Selective Expression of Megakaryocytic Markers in Friend Erythroleukemia Cells.
Blood, **86**(7), 2624–2631.
- Pantopoulos, K. (2004)
Iron metabolism and the IRE/IRP regulatory system: an update.
Ann NY Acad Sci, **1012**, 1-13, *review*.
- Pantopoulos, K., Weiss, G. & Hentze, M.W. (1996)
Nitric oxide and oxidative stress (H_2O_2) control mammalian iron metabolism by different pathways.
Mol Cell Biol, **16**, 3781–3788.
- Papanikolaou, G. & Pantopoulos, K. (2005)
Iron metabolism and toxicity.
Toxicol and App Pharmacol, **202**, 199–211, *review*.
- Park, C.H., Valore, E.V., Waring, A.J. & Ganz, T. (2001a)
Hepcidin, a Urinary Antimicrobial Peptide Synthesized in the Liver.
J Biol Chem, **276**(11), 7806–7810.
- Park, J.I., Choi, H.S., Jeong, J.S., Han, J.Y. & Kim, I.H. (2001b)
Involvement of p38 kinase in hydroxyurea-induced differentiation of K562 cells.
Cell Growth Differ, **12**, 481–486.
- Pellicci, P.G., Testa, U., Thomopoulos, P., Tabilio, A., Vainchenker, W., Titeux, M., Gourdin, M.F. & Rochant, H. (1984)
Inhibition of transferring binding and iron uptake of hematopoietic cell lines by phorbol esters.
Leuk Res, **8**(4), 597–609.
- Peschon, J.J., Slack, J.L., Reddy, P., Stocking, K.L., Sunnarborg, S.W., Lee, D.C., Russell, W.E., Castner, B.J., Johnson, R.S., Fitzner, J.N., Boyce, R.W., Nelson, N., Kozlosky, C.J., Wolfson, M.F., Rauch, C.T., Cerretti, D.P., Paxton, R.J., March, C.J. & Black, R.A. (1998)
An essential role for ectodomain shedding in mammalian development.
Science, **282**, 1281–1284.
- Picard, V., Renaudie, F., Porcher, C., Hentze, M.W., Grandchamp, B. & Beaumont, C. (1996)
Overexpression of the ferritin H subunit in cultured erythroid cells changes the intracellular iron distribution.
Blood, **87**(5), 2057–2064.
- Pietroangelo, A. (2002)
Physiology of iron transport and the hemochromatosis gene.
Am J Physiol-Gastrointest Liver Physiol, **282**, 403–414, *review*.
- Pietroangelo, A. (2004)
The ferroportin disease.
Blood Cells Mol Dis, **32**, 131–138, *review*.
- Pigeon, C., Ilyin, G., Courselaud, B., Leroyer, P., Turlin, B., Brissot, P. & Loreal, O. (2001)
A New Mouse Liver-specific Gene, Encoding a Protein Homologous to Human Antimicrobial Peptide Hepcidin, Is Overexpressed during Iron Overload.
J Biol Chem, **276**(11), 7811–7819.
- Ponka, P., Sheftel, A.D. & Zhang, A-S. (2002)
Iron targeting to mitochondria in erythroid cells.
Biochemical Society Transactions, **30**(4), 735–738, *review*.
- Prezioso, V.R. (2007)
General Notes on Primer Design in PCR.
A service of Brinkmann Instruments, Inc.

- Quigley, J.G., Yang, Z., Worthington, M.T., Phillips, J.D., Sabo, K.M., Sabath, D.E., Berg, C.L., Sassa, S., Wood, B.L. & Abkowitz, J.L. (2004)
Identification of a human heme exporter that is essential for erythropoiesis.
Cell, **118**(6), 757–766.
- Racke, F.K., Wang, D., Zaidi, Z., Kelley, J., Visvader, J., Soh, J.W. & Goldfarb, A.N. (2001)
A potential role for protein kinase C-epsilon in regulating megakaryocytic lineage commitment.
J Biol Chem, **276**(1), 522–528.
- Rafie-Kolpin, M., Chefalo, P.J., Hussain, Z., Hahn, J., Ma, S., Matts, R.L. & Chen, J-J. (2000)
Two heme-binding domains of heme-regulated eukaryotic initiation factor-2alpha kinase. N terminus and kinase insertion.
J Biol Chem, **275**, 5171–5178.
- Richter, G.W. (1978)
The iron-loaded cell—the cytopathology of iron storage. A review.
Am J Pathol, **91**, 362–404.
- Roetto, A., Totano, A., Piperno, A., Piga, A., Longo, F., Garozzo, G., Cali, A., De Gobbi, M., Gasparini, P. & Camaschella, C. (2001)
New mutations inactivating transferrin receptor in hemochromatosis type 3.
Blood, **97**, 2555–2560.
- Rolfs, A., Bonkovsky H.L., Kohlroser, J.G., McNeal, K., Sharma, A., Berger, U.V. & Hediger, M.A. (2002)
Intestinal expression of genes involved in iron absorption in humans.
Am J Physiol, **282**, G598–G607.
- Rouault, T.A., Hentze, M.W., Caughman, S.W., Harford, J.B. & Klausner, R.D. (1988)
Binding of a cytosolic protein to the iron-responsive element of human ferritin messenger RNA.
Science, **241**, 1207–1210.
- Rowley, P., Farley, B., LaBella, S., Giuliano, R. & Leary, J. (1992)
Single K562 human leukemia cells express and are inducible for both erythroid and megakaryocytic antigens.
Int J Cell Cloning, **10**, 232–240.
- R'zik, S., Loo, M. & Beguin, Y. (2001)
Reticulocyte transferrin receptor (TfR) expression and contribution to soluble TfR levels.
Haematologica, **86**, 244–251.
- Salter-Cid, L., Brunmark, A., Li, Y., Leturcq, D., Peterson, P.A., Jackson, M.R. & Yang, Y. (1999)
Transferrin receptor is negatively modulated by the hemochromatosis protein HFE: implications for cellular iron homeostasis.
Proc Natl Acad Sci U S A, **96**(10), 5434–5439.
- Samaniego, F., Chin, J., Iwai, K., Rouault, T.A. & Klausner, R.D. (1994)
Molecular characterization of a second iron-responsive element binding protein, iron regulatory protein 2. Structure, function, and post-translational regulation.
J Biol Chem, **269**, 30904–30910.
- Sanchez, M., Galy, B., Dandekar, T., Bengert, P., Vainshtein, Y., Stollte, J., Muckenthaler, M.U. & Hentze, M.W. (2006)
Iron regulation and the cell cycle: Identification of an iron-responsive element in the 3' untranslated region of human cell division cycle 14A mRNA by a refined microarray-based screening strategy.
J Biol Chem, **281**(32), 22865–22874.
- Sanchez, M., Queralt, R., Bruguera, M., Rodes, J. & Oliva, R. (1998)
Cloning, sequencing and characterisation of the rat hereditary hemochromatosis promoter: comparison of the human, mouse and rat HFE promoter regions.
Gene, **225**, 77–87.

- Sanger, F., Nicklen, S. & Coulson, A.R. (1977)
DNA sequencing with chain-terminating inhibitors.
Proc Natl Acad Sci U S A, **74**(12), 5463–5467.
- Savina, A., Vidal, M. & Colombo, M.I. (2002)
The exosome pathway in K-562 cells is regulated by Rab 11.
Journal of Cell Science, **115**, 2505–2515.
- Scaccabarozzi, A., Arosio, P., Weiss, G., Valenti, L., Dongiovanni, P., Fracanzani, AL., Mattioli, M., Levi, S., Fiorelli, G. & Fargion, S. (2000)
Relationship between TNF-alpha and iron metabolism in differentiating human monocytic THP-1 cells.
Br J Haematol, **110**(4), 978–984.
- Schaefer, M., Albrecht, N., Hofmann, T., Gudermann, T. & Schultz, G. (2001)
Receptor-mediated translocation of protein kinase C isotypes reveals a diffusion-limited binding process.
FASEB J, **15**, 1634–1636.
- Scheiber-Mojdehkar, B., Zimmermann, I., Dresow, B. & Goldenberg, H. (1999)
Differential response of non-transferrin bound iron uptake in rat liver cells on long-term and short-term treatment with iron.
J Hepatol, **31**, 61–70.
- Schonhorn, J.E., Akompong, T. & Wessling-Resnick, M. (1995)
Mechanism of transferrin receptor down-regulation in K562 cells in response to protein kinase C activation.
J Biol Chem, **270**(8), 3698–3705.
- Schranzhofer, M., Schirrer, M., Cabrera, J.A., Kopp, S., Chiba, P., Beug, H. & Müllner, E.W. (2006)
Remodeling the regulation of iron metabolism during erythroid differentiation to ensure efficient heme biosynthesis.
Blood, **107**(10), 4159–4167.
- Schwenke, K., Peterson, H.P., Wangenheim, K.H. & Feinendegen, L.E. (1995)
Induction of differentiation in erythroleukemic K562 cells by gamma-irradiation.
Leuk Res, **19**(12), 955–961.
- Shih, Y.J., Baynes, R.D., Hudson, B.G., Flowers, C.H., Skikne, B.S. & Cook, J.D. (1990)
Serum transferrin receptor is a truncated form of tissue receptor.
J Biol Chem, **265**, 19077–19081.
- Shoden, A., Gabrio, B.W. & Finch, C.A. (1953)
The relationship between ferritin and hemosiderin in rabbits and man.
J Biol Chem, **204**, 823–830.
- Sillaber, C., Gesbert, F., Frank, D.A., Sattler, M. & Griffin J.D. (2000)
STAT5 activation contributes to growth and viability in Bcr/Abl-transformed cells.
Blood, **95**, 2118–2125.
- Simpson, R. J., Mohamad-Ghaffer, A. & Khan, R. (1998)
Glutathione and duodenal reduction and absorption of iron.
Biochem Soc Trans, **26**, S352.
- Smith, R.D., Malley, J.D. & Schechter, A.N. (2000)
Quantitative analysis of globin gene induction in single human erythroleukemic cells.
Nucleic Acids Res, **28**, 4998–5004.
- Sposi, N.M., Cianetti, L., Tritarelli, E., Pelosi, E., Militi, S., Barberi, T., Gabbianelli, M., Saulle, E., Kühn, L., Peschle, C. & Testa, U. (2000)
Mechanisms of differential transferrin receptor expression in normal hematopoiesis.
Eur J Biochem, **267**, 6762–6774.

- Srinivas, V., Zhu, X., Salceda, S., Nakamura, R. & Caro, J. (1998)
Hypoxia-inducible Factor 1 α (HIF-1 α) Is a Non-heme Iron Protein.
J Biol Chem, **273**(29), 18019–18022.
- Szalai, G., Larue, A. & Watson, D. (2006)
Molecular mechanisms of megakaryopoiesis
Cell Mol Life Sci, **63**(21), 2460–2476, review.
- Tabuchi, M., Yoshimori, T., Yamaguchi, K., Yoshida, T. & Kishi, F. (2000)
Human NRAMP2/DMT1, which mediates iron transport across endosomal membranes, is localized to late endosomes and lysosomes in HEp-2 cells.
J Biol Chem, **275**, 22220–22228.
- Tahara, T., Sun, J., Igarashi, K., & Taketani, S. (2004a)
Heme-dependent upregulation of the alpha-globin gene expression by transcriptional repressor Bach1 in erythroid cells.
Biochem Biophys Res Commun, **324**, 77–85.
- Tahara, T., Sun, J., Nakanishi, K., Yamamoto, M., Mori, H., Saito, T., Fujita, H., Igarashi, K. & Taketani, S. (2004b).
Heme positively regulates the expression of beta-globin at the locus control region via the transcriptional factor Bach1 in erythroid cells.
J Biol Chem, **279**(7), 5480–5487.
- Taketani, S., Adachi, Y. & Nakahashi, Y. (2000)
Regulation of the expression of human ferrochelatase by intracellular iron levels.
Eur J Biochem, **267**(15), 4685–4692.
- Tallkvist, J., Bowlus, C.L. & Lonnerdal, B. (2003)
Effect of iron treatment on nickel absorption and gene expression of the divalent metal transporter (DMT1) by human intestinal Caco-2 cells.
Pharmacol Toxicol **92**, 121–124.
- Tanimura, A., Nezu, A., Morita, T., Hashimoto, N. & Tojyo, Y. (2002)
Interplay between Calcium, Diacylglycerol, and Phosphorylation in the Spatial and Temporal Regulation of PKCalpha –GFP.
J Biol Chem, **277**(32), 29054–29062.
- Tatusova, T.A. & Madden, T.L. (1999)
BLAST 2 Sequences, a new tool for comparing protein and nucleotide sequences.
FEMS Microbiol Lett, **174**, 247–250.
- Testa, U., Kuhn, L., Petrini, M., Quaranta, M.T., Pelosi, E. & Peschle, C. (1991)
Differential regulation of iron regulatory element-binding protein(s) in cell extracts of activated lymphocytes versus monocytesmacrophages.
J Biol Chem, **266**, 13925–13930.
- Testa, U., Vainchenker, W., Beuzard, Y., Rouyer-Fessard, P., Guerrasio, A., Titeux, M., Lapotre, P., Bouquet, J., Breton-Gorius, J. & Rosa, J. (1982)
Hemoglobin Expression in Clones of K562 Cell Line.
Eur J Biochem, **121**, 649–655.
- Theil, E.C. (2003)
Ferritin: At the Crossroads of Iron and Oxygen Metabolism
J Nutr, **133**, 1549–1553.

- Thery, C., Regnault, A., Garin, J., Wolfers, J., Zitvogel, L., Ricciardi-Castagnoli, P., Raposo, G. & Amigorena, S. (1999) Molecular characterisation of dendritic cells derived exosomes. Selective accumulation of the heat shock protein hsc73. *J Cell Biol*, **147**, 599–610.
- Torti, F.M. & Torti, S.V. (2002) Regulation of ferritin genes and protein. *Blood*, **99**, 3505–3516.
- Toullec, D., Pianettis, P., Herve, C., Belleverguel, P., Grand-Perret, T., Ajakanee, M., Baudet, V., Boissin, P., Boursier, E., Loriolle, F., Duhamel, L., Charon, D. & Kirilovsky, J. (1991) The Bisindolylmaleimide GF 109203X Is a Potent and Selective Inhibitor of Protein Kinase C. *J Biol Chem*, **266(24)**, 15771–15781.
- Treffry, A., Zhao Z., Quail, M.A., Guest, J.R. & Harrison, P.M. (1995) Iron(II) oxidation by H chain ferritin: evidence from site-directed mutagenesis that a transient blue species is formed at the dinuclear iron center. *Biochemistry*, **34**, 15204–15213.
- Trinder, D. & Baker, E. (2003) Transferrin receptor 2: a new molecule in iron metabolism. *Int J Biochem Cell Biol*, **35**, 292–296.
- Tseng, C-P., Chang, P., Huang, C-L., Cheng, J-C. & Chang, S-S. (2005) Autocrine signaling of platelet-derived growth factor regulates disabled-2 expression during megakaryocytic differentiation of K562 cells. *FEBS Lett*, **579**, 4395–4401.
- Tsiftsoglou, A.S., Pappas, I.S. & Vizirianakis, I.S. (2003) Mechanisms involved in the induced differentiation of leukaemia cells. *Pharmacol Ther*, **100**, 257–290, review.
- Tsiftsoglou, A.S., Tsamadou, A.I. & Papadopoulou, L.C. (2006) Heme as key regulator of major mammalian cellular functions: molecular, cellular, and pharmacological aspects. *Pharmacol Ther*, **111(2)**, 327–345, review.
- Turkewitz, A.P., Amatruda, J.F., D Borhani, D., Harrison, S.C. & Schwartz, A.L. (1988) A high yield purification of the human transferrin receptor and properties of its major extracellular fragment. *J Biol Chem*, **263(17)**, 8318–8325.
- Umbreit, J.N., Conrad, M.E., Moore, E.G., Desai, M.P. & Turens, J. (1996) Paraferitin: A protein complex with ferrireductase activity is associated with iron absorption in rats. *Biochemistry*, **35**, 6460–6469.
- Van Bockxmeer, F.M. & Morgan, E.H. (1979) Transferrin receptors during rabbit reticulocyte maturation. *Biochim Biophys Acta*, **584**, 76–83.
- Van Campen, D. (1972) Effect of histidine and ascorbic acid on the absorption and retention of 59Fe by iron-depleted rats. *J Nutr*, **102**, 165–170.
- Vannucchi, A.M., Linari, S., Cellai, C., Koury, M.J. & Paletti, F. (1997) Constitutive and inducible expression of megakaryocyte-specific genes in Friend erythroleukemia cells. *Br J Haematol*, **99(3)**, 500–508.
- Wallander, M.L., Leibold, E.A. & Eisenstein, R.S. (2006) Molecular control of vertebrate iron homeostasis by iron regulatory proteins. *Biochim Biophys Acta*, **1763**, 668–689, review.

- Wandzik, K. (2003)
Untersuchungen zum Shedding des Transferrinrezeptors und anderer Membranproteine -Einfluss von TIMPs und ADAMs.
Diplomarbeit, Fachbereich Chemie, Freie Universität Berlin.
- Whalen, A.M., Galasinski, S.C., Shapiro, P.S., Nahreini, T.S. & Ahn, N.G. (1997)
Megakaryocytic Differentiation Induced by Constitutive Activation of Mitogen-Activated Protein Kinase Kinase.
Mol Cell Biol, 17(4), 1947–1958.
- White, K. & Munro, H.N. (1988)
Induction of ferritin subunit synthesis by iron is regulated at both the transcriptional and translational levels.
J Biol Chem, 263, 8938–8942.
- Witt, O., Sand, K. & Pekrun, A. (2000)
Butyrate-induced erythroid differentiation of human K562 leukemia cells involves inhibition of ERK and activation of p38 MAP kinase pathways.
Blood, 95, 2391–2396.
- Wu, L.J., Leenders, A.G., Cooperman, S., Meyron-Holtz, E., Smith, S., Land, W., Tsai, R.Y., Berger, U.V., Sheng, Z.H. & Rouault, T.A. (2004)
Expression of the iron transporter ferroportin in synaptic vesicles and the blood-brain barrier.
Brain Res, 1001, 108–117.
- Yang, F., Naylor, S.L., Lum, J.B., Cutshaw, S., McCombs, J.L., Naberhaus, H., McGill, J.R., Adrian, G.S., Moore, C.M., Barnett, D.R. & Bowman, B.H. (1986)
Characterization, mapping and expression of the human ceruloplasmin gene.
Proc Natl Acad Sci USA, 83, 3257–3261.
- Young, S.P., Bomford, A. & Williams, R. (1984)
The effect of the iron saturation of transferrin on its binding and uptake by rabbit reticulocytes.
Biochem J, 219, 505–510.
- Zhang, A.S., Sheftel, A.D. & Ponka, P. (2005)
Intracellular kinetics of iron in reticulocytes: evidence for endosome involvement in iron targeting to mitochondria.
Blood, 105, 368–375.
- Zhang, A.S., Sheftel, A.D. & Ponka, P. (2006)
The anemia of "haemoglobin-deficit" (hbd/hbd) mice is caused by a defect in transferrin cycling.
Exp Hematol, 34, 593–598.
- Zoller, H., Decristoforo, C. & Weiss, G. (2002)
Erythroid 5-aminolevulinate synthase, ferrochelatase and DMT1 expression in erythroid progenitors: differential pathways for erythropoietin and iron-dependent regulation.
Br J Haematol, 118, 619–626.