

5 Literatur

- Ahlgren, U., Jonsson, J. and Edlund, H.** (1996). The morphogenesis of the pancreatic mesenchyme is uncoupled from that of the pancreatic epithelium in IPF1/PDX1-deficient mice. *Development* **122**, 1409-16.
- Ahlgren, U., Jonsson, J., Jonsson, L., Simu, K. and Edlund, H.** (1998). beta-cell-specific inactivation of the mouse *Ipfl/Pdx1* gene results in loss of the beta-cell phenotype and maturity onset diabetes. *Genes Dev.* **12**, 1763-8.
- Ahlgren, U., Pfaff, S. L., Jessell, T. M., Edlund, T. and Edlund, H.** (1997). Independent requirement for ISL1 in formation of pancreatic mesenchyme and islet cells. *Nature* **385**, 257-60.
- Apelqvist, A., Li, H., Sommer, L., Beatus, P., Anderson, D. J., Honjo, T., Hrabe de Angelis, M., Lendahl, U. and Edlund, H.** (1999). Notch signalling controls pancreatic cell differentiation. *Nature* **400**, 877-81.
- Arden, S. D., Zahn, T., Steegers, S., Webb, S., Bergman, B., O'Brien, R. M. and Hutton, J. C.** (1999). Molecular cloning of a pancreatic islet-specific glucose-6-phosphatase catalytic subunit-related protein. *Diabetes* **48**, 531-42.
- Artavanis-Tsakonas, S., Rand, M. D. and Lake, R. J.** (1999). Notch signaling: cell fate control and signal integration in development. *Science* **284**, 770-6.
- Artner, I., Le Lay, J., Hang, Y., Elghazi, L., Schisler, J. C., Henderson, E., Sosa-Pineda, B. and Stein, R.** (2006). MafB: an activator of the glucagon gene expressed in developing islet alpha- and beta-cells. *Diabetes* **55**, 297-304.
- Asayesh, A., Alanentalo, T., Khoo, N. K. and Ahlgren, U.** (2005). Developmental expression of metalloproteases ADAM 9, 10, and 17 becomes restricted to divergent pancreatic compartments. *Dev. Dyn.* **232**, 1105-14.
- Bali, D., Svetlanov, A., Lee, H. W., Fusco-DeMane, D., Leiser, M., Li, B., Barzilai, N., Surana, M., Hou, H., Fleischer, N. et al.** (1995). Animal model for maturity-onset diabetes of the young generated by disruption of the mouse glucokinase gene. *J. Biol. Chem.* **270**, 21464-7.
- Bell, G. I. and Polonsky, K. S.** (2001). Diabetes mellitus and genetically programmed defects in beta-cell function. *Nature* **414**, 788-91.
- Birney, E., Andrews, D., Caccamo, M., Chen, Y., Clarke, L., Coates, G., Cox, T., Cunningham, F., Curwen, V., Cutts, T. et al.** (2006). Ensembl 2006. *Nucleic Acids Res.* **34**, D556-61.
- Breslin, M. B., Zhu, M. and Lan, M. S.** (2003). NeuroD1/E47 regulates the E-box element of a novel zinc finger transcription factor, IA-1, in developing nervous system. *J. Biol. Chem.* **278**, 38991-7.
- Breslin, M. B., Zhu, M., Notkins, A. L. and Lan, M. S.** (2002). Neuroendocrine differentiation factor, IA-1, is a transcriptional repressor and contains a specific DNA-binding domain: identification of consensus IA-1 binding sequence. *Nucleic Acids Res.* **30**, 1038-45.
- Brissova, M., Shiota, M., Nicholson, W. E., Gannon, M., Knobel, S. M., Piston, D. W., Wright, C. V. and Powers, A. C.** (2002). Reduction in pancreatic transcription factor PDX-1 impairs glucose-stimulated insulin secretion. *J. Biol. Chem.* **277**, 11225-32.
- Bunting, M., Bernstein, K. E., Greer, J. M., Capecchi, M. R. and Thomas, K. R.** (1999). Targeting genes for self-excision in the germ line. *Genes Dev.* **13**, 1524-8.
- Carty, M. D., Lillquist, J. S., Peshavaria, M., Stein, R. and Soeller, W. C.** (1997). Identification of cis- and trans-active factors regulating human islet amyloid polypeptide gene expression in pancreatic beta-cells. *J. Biol. Chem.* **272**, 11986-93.

- Cissell, M. A., Zhao, L., Sussel, L., Henderson, E. and Stein, R. (2003). Transcription factor occupancy of the insulin gene in vivo. Evidence for direct regulation by Nkx2.2. *J. Biol. Chem.* **278**, 751-6.
- Collombat, P., Hecksher-Sorensen, J., Broccoli, V., Krull, J., Ponte, I., Mundiger, T., Smith, J., Gruss, P., Serup, P. and Mansouri, A. (2005). The simultaneous loss of Arx and Pax4 genes promotes a somatostatin-producing cell fate specification at the expense of the alpha- and beta-cell lineages in the mouse endocrine pancreas. *Development* **132**, 2969-80.
- Collombat, P., Mansouri, A., Hecksher-Sorensen, J., Serup, P., Krull, J., Gradwohl, G. and Gruss, P. (2003). Opposing actions of Arx and Pax4 in endocrine pancreas development. *Genes Dev.* **17**, 2591-603.
- Cook, M. J. (1965). *The Anatomy of the Laboratory Mouse*. London, New York: Academic Press.
- Cooper, G. J., Day, A. J., Willis, A. C., Roberts, A. N., Reid, K. B. and Leighton, B. (1989). Amylin and the amylin gene: structure, function and relationship to islet amyloid and to diabetes mellitus. *Biochim. Biophys. Acta* **1014**, 247-58.
- Cooper, G. J., Willis, A. C., Clark, A., Turner, R. C., Sim, R. B. and Reid, K. B. (1987). Purification and characterization of a peptide from amyloid-rich pancreases of type 2 diabetic patients. *Proc. Natl. Acad. Sci. USA* **84**, 8628-32.
- Dukes, I. D., Sreenan, S., Roe, M. W., Levisetti, M., Zhou, Y. P., Ostrega, D., Bell, G. I., Pontoglio, M., Yaniv, M., Philipson, L. et al. (1998). Defective pancreatic beta-cell glycolytic signaling in hepatocyte nuclear factor-1alpha-deficient mice. *J. Biol. Chem.* **273**, 24457-64.
- Edlund, H. (2001). Developmental biology of the pancreas. *Diabetes* **50 Suppl 1**, S5-9.
- Edlund, H. (2002). Pancreatic organogenesis--developmental mechanisms and implications for therapy. *Nat. Rev. Genet.* **3**, 524-32.
- Fajans, S. S., Bell, G. I. and Polonsky, K. S. (2001). Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. *N. Engl. J. Med.* **345**, 971-80.
- Froguel, P., Vaxillaire, M., Sun, F., Velho, G., Zouali, H., Butel, M. O., Lesage, S., Vionnet, N., Clement, K., Fougere, F. et al. (1992). Close linkage of glucokinase locus on chromosome 7p to early-onset non-insulin-dependent diabetes mellitus. *Nature* **356**, 162-4.
- Fujitani, Y., Fujitani, S., Boyer, D. F., Gannon, M., Kawaguchi, Y., Ray, M., Shiota, M., Stein, R. W., Magnuson, M. A. and Wright, C. V. (2006). Targeted deletion of a cis-regulatory region reveals differential gene dosage requirements for Pdx1 in foregut organ differentiation and pancreas formation. *Genes Dev.* **20**, 253-66.
- Gebre-Medhin, S., Mulder, H., Pekny, M., Westermark, G., Tornell, J., Westermark, P., Sundler, F., Ahren, B. and Betsholtz, C. (1998). Increased insulin secretion and glucose tolerance in mice lacking islet amyloid polypeptide (amylin). *Biochem. Biophys. Res. Commun.* **250**, 271-7.
- Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J. et al. (2004). Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol.* **5**, R80.
- Gierl, M. S., Karoulias, N., Wende, H., Strehle, M. and Birchmeier, C. (2006). The zinc-finger factor Insm1 (IA-1) is essential for the development of pancreatic beta cells and intestinal endocrine cells. *Genes Dev.* **20**, 2465-78.
- Gluzman, Y. (1981). SV40-transformed simian cells support the replication of early SV40 mutants. *Cell* **23**, 175-82.

- Goto, Y., De Silva, M. G., Toscani, A., Prabhakar, B. S., Notkins, A. L. and Lan, M. S.** (1992). A novel human insulinoma-associated cDNA, IA-1, encodes a protein with "zinc-finger" DNA-binding motifs. *J. Biol. Chem.* **267**, 15252-7.
- Gradwohl, G., Dierich, A., LeMeur, M. and Guillemot, F.** (2000). neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. *Proc. Natl. Acad. Sci. U S A* **97**, 1607-11.
- Grapin-Botton, A.** (2005). Antero-posterior patterning of the vertebrate digestive tract: 40 years after Nicole Le Douarin's PhD thesis. *Int. J. Dev. Biol.* **49**, 335-47.
- Grimes, H. L., Chan, T. O., Zweidler-McKay, P. A., Tong, B. and Tsichlis, P. N.** (1996). The Gfi-1 proto-oncoprotein contains a novel transcriptional repressor domain, SNAG, and inhibits G1 arrest induced by interleukin-2 withdrawal. *Mol. Cell. Biol.* **16**, 6263-72.
- Gu, G., Brown, J. R. and Melton, D. A.** (2003). Direct lineage tracing reveals the ontogeny of pancreatic cell fates during mouse embryogenesis. *Mech. Dev.* **120**, 35-43.
- Gu, G., Dubauskaite, J. and Melton, D. A.** (2002). Direct evidence for the pancreatic lineage: NGN3+ cells are islet progenitors and are distinct from duct progenitors. *Development* **129**, 2447-57.
- Gupta, R. K., Vatamaniuk, M. Z., Lee, C. S., Flaschen, R. C., Fulmer, J. T., Matschinsky, F. M., Duncan, S. A. and Kaestner, K. H.** (2005). The MODY1 gene HNF-4alpha regulates selected genes involved in insulin secretion. *J. Clin. Invest.* **115**, 1006-15.
- Guz, Y., Montminy, M. R., Stein, R., Leonard, J., Gamer, L. W., Wright, C. V. and Teitelman, G.** (1995). Expression of murine STF-1, a putative insulin gene transcription factor, in beta cells of pancreas, duodenal epithelium and pancreatic exocrine and endocrine progenitors during ontogeny. *Development* **121**, 11-8.
- Habener, J. F., Kemp, D. M. and Thomas, M. K.** (2005). Minireview: transcriptional regulation in pancreatic development. *Endocrinology* **146**, 1025-34.
- Hagenfeldt-Johansson, K. A., Herrera, P. L., Wang, H., Gjinovci, A., Ishihara, H. and Wollheim, C. B.** (2001). Beta-cell-targeted expression of a dominant-negative hepatocyte nuclear factor-1 alpha induces a maturity-onset diabetes of the young (MODY)3-like phenotype in transgenic mice. *Endocrinology* **142**, 5311-20.
- Harrison, K. A., Thaler, J., Pfaff, S. L., Gu, H. and Kehrl, J. H.** (1999). Pancreas dorsal lobe agenesis and abnormal islets of Langerhans in Hlxb9-deficient mice. *Nat. Genet.* **23**, 71-5.
- Heller, R. S., Jenny, M., Collombat, P., Mansouri, A., Tomasetto, C., Madsen, O. D., Mellitzer, G., Gradwohl, G. and Serup, P.** (2005). Genetic determinants of pancreatic epsilon-cell development. *Dev. Biol.* **286**, 217-24.
- Heremans, Y., Van De Casteele, M., in't Veld, P., Gradwohl, G., Serup, P., Madsen, O., Pipeleers, D. and Heimberg, H.** (2002). Recapitulation of embryonic neuroendocrine differentiation in adult human pancreatic duct cells expressing neurogenin 3. *J. Cell Biol.* **159**, 303-12.
- Herrera, P. L.** (2000). Adult insulin- and glucagon-producing cells differentiate from two independent cell lineages. *Development* **127**, 2317-22.
- Holland, A. M., Hale, M. A., Kagami, H., Hammer, R. E. and MacDonald, R. J.** (2002). Experimental control of pancreatic development and maintenance. *Proc. Natl. Acad. Sci. U S A* **99**, 12236-41.

- Horikawa, Y., Iwasaki, N., Hara, M., Furuta, H., Hinokio, Y., Cockburn, B. N., Lindner, T., Yamagata, K., Ogata, M., Tomonaga, O. et al. (1997). Mutation in hepatocyte nuclear factor-1 beta gene (TCF2) associated with MODY. *Nat. Genet.* **17**, 384-5.
- Jacquemin, P., Durviaux, S. M., Jensen, J., Godfraind, C., Gradwohl, G., Guillemot, F., Madsen, O. D., Carmeliet, P., Dewerchin, M., Collen, D. et al. (2000). Transcription factor hepatocyte nuclear factor 6 regulates pancreatic endocrine cell differentiation and controls expression of the proendocrine gene *ngn3*. *Mol. Cell. Biol.* **20**, 4445-54.
- Jenny, M., Uhl, C., Roche, C., Duluc, I., Guillermin, V., Guillemot, F., Jensen, J., Kedinger, M. and Gradwohl, G. (2002). Neurogenin3 is differentially required for endocrine cell fate specification in the intestinal and gastric epithelium. *EMBO J.* **21**, 6338-47.
- Jensen, J. (2004). Gene regulatory factors in pancreatic development. *Dev. Dyn.* **229**, 176-200.
- Jensen, J., Heller, R. S., Funder-Nielsen, T., Pedersen, E. E., Lindsell, C., Weinmaster, G., Madsen, O. D. and Serup, P. (2000a). Independent development of pancreatic alpha- and beta-cells from neurogenin3-expressing precursors: a role for the notch pathway in repression of premature differentiation. *Diabetes* **49**, 163-76.
- Jensen, J., Pedersen, E. E., Galante, P., Hald, J., Heller, R. S., Ishibashi, M., Kageyama, R., Guillemot, F., Serup, P. and Madsen, O. D. (2000b). Control of endodermal endocrine development by Hes-1. *Nat. Genet.* **24**, 36-44.
- Jonsson, J., Carlsson, L., Edlund, T. and Edlund, H. (1994). Insulin-promoter-factor 1 is required for pancreas development in mice. *Nature* **371**, 606-9.
- Karlsson, E. (2001). The role of pancreatic chromogranins in islet physiology. *Curr. Mol. Med.* **1**, 727-32.
- Kataoka, K., Han, S. I., Shioda, S., Hirai, M., Nishizawa, M. and Handa, H. (2002). MafA is a glucose-regulated and pancreatic beta-cell-specific transcriptional activator for the insulin gene. *J. Biol. Chem.* **277**, 49903-10.
- Kawaguchi, Y., Cooper, B., Gannon, M., Ray, M., MacDonald, R. J. and Wright, C. V. (2002). The role of the transcriptional regulator Ptf1a in converting intestinal to pancreatic progenitors. *Nat. Genet.* **32**, 128-34.
- Kim, S. K. and Hebrok, M. (2001). Intercellular signals regulating pancreas development and function. *Genes Dev.* **15**, 111-27.
- Kojima, M., Hosoda, H., Date, Y., Nakazato, M., Matsuo, H. and Kangawa, K. (1999). Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* **402**, 656-60.
- Kozak, M. (1987). An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res.* **15**, 8125-48.
- Krapp, A., Knofler, M., Frutiger, S., Hughes, G. J., Hagenbuchle, O. and Wellauer, P. K. (1996). The p48 DNA-binding subunit of transcription factor PTF1 is a new exocrine pancreas-specific basic helix-loop-helix protein. *EMBO J.* **15**, 4317-29.
- Krapp, A., Knofler, M., Ledermann, B., Burki, K., Berney, C., Zoerkler, N., Hagenbuchle, O. and Wellauer, P. K. (1998). The bHLH protein PTF1-p48 is essential for the formation of the exocrine and the correct spatial organization of the endocrine pancreas. *Genes Dev.* **12**, 3752-63.
- Kühn, R., Rajewsky, K. and Müller, W. (1991). Generation and analysis of interleukin-4 deficient mice. *Science* **254**, 707-10.

- Kuzin, A., Brody, T., Moore, A. W. and Odenwald, W. F.** (2005). Nerfin-1 is required for early axon guidance decisions in the developing *Drosophila* CNS. *Dev. Biol.* **277**, 347-65.
- Lammert, E., Brown, J. and Melton, D. A.** (2000). Notch gene expression during pancreatic organogenesis. *Mech. Dev.* **94**, 199-203.
- Lan, M. S., Li, Q., Lu, J., Modi, W. S. and Notkins, A. L.** (1994). Genomic organization, 5'-upstream sequence, and chromosomal localization of an insulinoma-associated intronless gene, IA-1. *J. Biol. Chem.* **269**, 14170-4.
- Lan, M. S., Russell, E. K., Lu, J., Johnson, B. E. and Notkins, A. L.** (1993). IA-1, a new marker for neuroendocrine differentiation in human lung cancer cell lines. *Cancer Res.* **53**, 4169-71.
- Larsson, L. I., St-Onge, L., Hougaard, D. M., Sosa-Pineda, B. and Gruss, P.** (1998). Pax 4 and 6 regulate gastrointestinal endocrine cell development. *Mech. Dev.* **79**, 153-9.
- Le Douarin, N. M.** (1982). *The Neural Crest*. Cambridge: Cambridge University Press.
- Lee, C. S., Perreault, N., Brestelli, J. E. and Kaestner, K. H.** (2002a). Neurogenin 3 is essential for the proper specification of gastric enteroendocrine cells and the maintenance of gastric epithelial cell identity. *Genes Dev.* **16**, 1488-97.
- Lee, C. S., Sund, N. J., Behr, R., Herrera, P. L. and Kaestner, K. H.** (2005). Foxa2 is required for the differentiation of pancreatic alpha-cells. *Dev. Biol.* **278**, 484-95.
- Lee, C. S., Sund, N. J., Vatamaniuk, M. Z., Matschinsky, F. M., Stoffers, D. A. and Kaestner, K. H.** (2002b). Foxa2 controls Pdx1 gene expression in pancreatic beta-cells in vivo. *Diabetes* **51**, 2546-51.
- Lee, E. C., Yu, D., Martinez de Velasco, J., Tessarollo, L., Swing, D. A., Court, D. L., Jenkins, N. A. and Copeland, N. G.** (2001a). A highly efficient *Escherichia coli*-based chromosome engineering system adapted for recombinogenic targeting and subcloning of BAC DNA. *Genomics* **73**, 56-65.
- Lee, J. C., Smith, S. B., Watada, H., Lin, J., Scheel, D., Wang, J., Mirmira, R. G. and German, M. S.** (2001b). Regulation of the pancreatic pro-endocrine gene neurogenin3. *Diabetes* **50**, 928-36.
- Li, H., Arber, S., Jessell, T. M. and Edlund, H.** (1999). Selective agenesis of the dorsal pancreas in mice lacking homeobox gene Hlxb9. *Nat. Genet.* **23**, 67-70.
- Li, Q., Notkins, A. L. and Lan, M. S.** (1997). Molecular characterization of the promoter region of a neuroendocrine tumor marker, IA-1. *Biochem. Biophys. Res. Commun.* **236**, 776-81.
- Malecki, M. T., Jhala, U. S., Antonellis, A., Fields, L., Doria, A., Orban, T., Saad, M., Warram, J. H., Montminy, M. and Krolewski, A. S.** (1999). Mutations in NEUROD1 are associated with the development of type 2 diabetes mellitus. *Nat. Genet.* **23**, 323-8.
- Marcinkiewicz, M., Ramla, D., Seidah, N. G. and Chretien, M.** (1994). Developmental expression of the prohormone convertases PC1 and PC2 in mouse pancreatic islets. *Endocrinology* **135**, 1651-60.
- Matschinsky, F. M.** (1990). Glucokinase as glucose sensor and metabolic signal generator in pancreatic beta-cells and hepatocytes. *Diabetes* **39**, 647-52.
- Mellitzer, G., Bonne, S., Luco, R. F., Van De Casteele, M., Lenne-Samuel, N., Collombat, P., Mansouri, A., Lee, J., Lan, M., Pipeleers, D. et al.** (2006). IA1 is NGN3-dependent and essential for differentiation of the endocrine pancreas. *EMBO J.* **25**, 1344-52.

- Miura, A., Yamagata, K., Kakei, M., Hatakeyama, H., Takahashi, N., Fukui, K., Nammo, T., Yoneda, K., Inoue, Y., Sladek, F. M. et al.** (2006). Hepatocyte nuclear factor-4alpha is essential for glucose-stimulated insulin secretion by pancreatic beta-cells. *J. Biol. Chem.* **281**, 5246-57.
- Moltz, J. H. and McDonald, J. K.** (1985). Neuropeptide Y: direct and indirect action on insulin secretion in the rat. *Peptides* **6**, 1155-9.
- Murtaugh, L. C. and Melton, D. A.** (2003). Genes, signals, and lineages in pancreas development. *Annu. Rev. Cell Dev. Biol.* **19**, 71-89.
- Naya, F. J., Huang, H. P., Qiu, Y., Mutoh, H., DeMayo, F. J., Leiter, A. B. and Tsai, M. J.** (1997). Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/neuroD-deficient mice. *Genes Dev.* **11**, 2323-34.
- Naya, F. J., Stellrecht, C. M. and Tsai, M. J.** (1995). Tissue-specific regulation of the insulin gene by a novel basic helix-loop-helix transcription factor. *Genes Dev.* **9**, 1009-19.
- Nishi, M., Chan, S. J., Nagamatsu, S., Bell, G. I. and Steiner, D. F.** (1989). Conservation of the sequence of islet amyloid polypeptide in five mammals is consistent with its putative role as an islet hormone. *Proc. Natl. Acad. Sci. U S A* **86**, 5738-42.
- Offield, M. F., Jetton, T. L., Labosky, P. A., Ray, M., Stein, R. W., Magnuson, M. A., Hogan, B. L. and Wright, C. V.** (1996). PDX-1 is required for pancreatic outgrowth and differentiation of the rostral duodenum. *Development* **122**, 983-95.
- Ohlsson, H., Karlsson, K. and Edlund, T.** (1993). IPF1, a homeodomain-containing transactivator of the insulin gene. *EMBO J.* **12**, 4251-9.
- Olbrot, M., Rud, J., Moss, L. G. and Sharma, A.** (2002). Identification of beta-cell-specific insulin gene transcription factor RIPE3b1 as mammalian MafA. *Proc. Natl. Acad. Sci. U S A* **99**, 6737-42.
- Opara, E. C., Burch, W. M., Taylor, I. L. and Akwari, O. E.** (1991). Pancreatic hormone response to neuropeptide Y (NPY) perfusion in vitro. *Regul. Pept.* **34**, 225-33.
- Orci, L., Thorens, B., Ravazzola, M. and Lodish, H. F.** (1989). Localization of the pancreatic beta cell glucose transporter to specific plasma membrane domains. *Science* **245**, 295-7.
- Osoegawa, K., Tateno, M., Woon, P. Y., Frengen, E., Mammoser, A. G., Catanese, J. J., Hayashizaki, Y. and de Jong, P. J.** (2000). Bacterial artificial chromosome libraries for mouse sequencing and functional analysis. *Genome Res.* **10**, 116-28.
- Pettersson, M., Ahren, B., Lundquist, I., Bottcher, G. and Sundler, F.** (1987). Neuropeptide Y: intrapancreatic neuronal localization and effects on insulin secretion in the mouse. *Cell Tissue. Res.* **248**, 43-8.
- Pictet, R. and Rutter, W. J.** (1972). Development of the embryonic endocrine pancreas. In *Handbook of Physiology, section 7, vol. 1*, (ed. D. F. Steiner and N. Frenkel), pp. 25-66. Washington DC: American Physiological Society.
- Pontoglio, M., Sreenan, S., Roe, M., Pugh, W., Ostrega, D., Doyen, A., Pick, A. J., Baldwin, A., Velho, G., Froguel, P. et al.** (1998). Defective insulin secretion in hepatocyte nuclear factor 1alpha-deficient mice. *J. Clin. Invest.* **101**, 2215-22.
- Prado, C. L., Pugh-Bernard, A. E., Elghazi, L., Sosa-Pineda, B. and Sussel, L.** (2004). Ghrelin cells replace insulin-producing beta cells in two mouse models of pancreas development. *Proc. Natl. Acad. Sci. U S A* **101**, 2924-9.
- R Development Core Team.** (2005). R: A language and environment for statistical computing. Vienna, Austria.

- Rindi, G., Leiter, A. B., Kopin, A. S., Bordi, C. and Solcia, E. (2004). The "normal" endocrine cell of the gut: changing concepts and new evidences. *Ann. N. Y. Acad. Sci.* **1014**, 1-12.
- Sambrook, J. and Russell, D. W. (2001). *Molecular Cloning: A Laboratory Manual*: Cold Spring Harbor Laboratory Press.
- Sander, M., Neubuser, A., Kalamaras, J., Ee, H. C., Martin, G. R. and German, M. S. (1997). Genetic analysis reveals that PAX6 is required for normal transcription of pancreatic hormone genes and islet development. *Genes Dev.* **11**, 1662-73.
- Sander, M., Sussel, L., Connors, J., Scheel, D., Kalamaras, J., Dela Cruz, F., Schwitzgebel, V., Hayes-Jordan, A. and German, M. (2000). Homeobox gene Nkx6.1 lies downstream of Nkx2.2 in the major pathway of beta-cell formation in the pancreas. *Development* **127**, 5533-40.
- Schwitzgebel, V. M., Scheel, D. W., Connors, J. R., Kalamaras, J., Lee, J. E., Anderson, D. J., Sussel, L., Johnson, J. D. and German, M. S. (2000). Expression of neurogenin3 reveals an islet cell precursor population in the pancreas. *Development* **127**, 3533-42.
- Shih, D. Q., Screenan, S., Munoz, K. N., Philipson, L., Pontoglio, M., Yaniv, M., Polonsky, K. S. and Stoffel, M. (2001). Loss of HNF-1alpha function in mice leads to abnormal expression of genes involved in pancreatic islet development and metabolism. *Diabetes* **50**, 2472-80.
- Slack, J. M. (1995). Developmental biology of the pancreas. *Development* **121**, 1569-80.
- Smyth, G. K. (2005). Limma: linear models for microarray data. In *Bioinformatics and Computational Biology Solutions using R and Bioconductor*, (ed. R. Gentleman V. Carey S. Dudoit R. Irizarry and W. Huber), pp. 397-420. New York: Springer.
- Sonnenberg-Riethmacher, E., Walter, B., Riethmacher, D., Godecke, S. and Birchmeier, C. (1996). The c-ros tyrosine kinase receptor controls regionalization and differentiation of epithelial cells in the epididymis. *Genes Dev.* **10**, 1184-93.
- Sosa-Pineda, B., Chowdhury, K., Torres, M., Oliver, G. and Gruss, P. (1997). The Pax4 gene is essential for differentiation of insulin-producing beta cells in the mammalian pancreas. *Nature* **386**, 399-402.
- Stivers, C., Brody, T., Kuzin, A. and Odenwald, W. F. (2000). *Nerfin-1* and *-2*, novel *Drosophila* Zn-finger transcription factor genes expressed in the developing nervous system. *Mech. Dev.* **97**, 205-10.
- Stoffers, D. A., Ferrer, J., Clarke, W. L. and Habener, J. F. (1997). Early-onset type-II diabetes mellitus (MODY4) linked to IPF1. *Nat. Genet.* **17**, 138-9.
- St-Onge, L., Sosa-Pineda, B., Chowdhury, K., Mansouri, A. and Gruss, P. (1997). Pax6 is required for differentiation of glucagon-producing alpha-cells in mouse pancreas. *Nature* **387**, 406-9.
- Su, A. I., Wiltshire, T., Batalov, S., Lapp, H., Ching, K. A., Block, D., Zhang, J., Soden, R., Hayakawa, M., Kreiman, G. et al. (2004). A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc. Natl. Acad. Sci. U S A* **101**, 6062-7.
- Sund, N. J., Vatamaniuk, M. Z., Casey, M., Ang, S. L., Magnuson, M. A., Stoffers, D. A., Matschinsky, F. M. and Kaestner, K. H. (2001). Tissue-specific deletion of Foxa2 in pancreatic beta cells results in hyperinsulinemic hypoglycemia. *Genes Dev.* **15**, 1706-15.
- Sussel, L., Kalamaras, J., Hartigan-O'Connor, D. J., Meneses, J. J., Pedersen, R. A., Rubenstein, J. L. and German, M. S. (1998). Mice lacking the homeodomain transcription factor Nkx2.2 have diabetes due to arrested differentiation of pancreatic beta cells. *Development* **125**, 2213-21.

- Taupenot, L., Harper, K. L. and O'Connor, D. T.** (2003). The chromogranin-secretogranin family. *N. Engl. J. Med.* **348**, 1134-49.
- Teitelman, G., Alpert, S., Polak, J. M., Martinez, A. and Hanahan, D.** (1993). Precursor cells of mouse endocrine pancreas coexpress insulin, glucagon and the neuronal proteins tyrosine hydroxylase and neuropeptide Y, but not pancreatic polypeptide. *Development* **118**, 1031-9.
- Terauchi, Y., Sakura, H., Yasuda, K., Iwamoto, K., Takahashi, N., Ito, K., Kasai, H., Suzuki, H., Ueda, O., Kamada, N. et al.** (1995). Pancreatic beta-cell-specific targeted disruption of glucokinase gene. Diabetes mellitus due to defective insulin secretion to glucose. *J. Biol. Chem.* **270**, 30253-6.
- Thomas, S. A., Matsumoto, A. M. and Palmiter, R. D.** (1995). Noradrenaline is essential for mouse fetal development. *Nature* **374**, 643-6.
- Thorens, B., Sarkar, H. K., Kaback, H. R. and Lodish, H. F.** (1988). Cloning and functional expression in bacteria of a novel glucose transporter present in liver, intestine, kidney, and beta-pancreatic islet cells. *Cell* **55**, 281-90.
- Tomasetto, C., Karam, S. M., Ribieras, S., Masson, R., Lefebvre, O., Staub, A., Alexander, G., Chenard, M. P. and Rio, M. C.** (2000). Identification and characterization of a novel gastric peptide hormone: the motilin-related peptide. *Gastroenterology* **119**, 395-405.
- Turque, N., Plaza, S., Radvanyi, F., Carriere, C. and Saule, S.** (1994). Pax-QNR/Pax-6, a paired box- and homeobox-containing gene expressed in neurons, is also expressed in pancreatic endocrine cells. *Mol. Endocrinol.* **8**, 929-38.
- Vilen, H., Eerikainen, S., Tornberg, J., Airaksinen, M. S. and Savilahti, H.** (2001). Construction of gene-targeting vectors: a rapid Mu in vitro DNA transposition-based strategy generating null, potentially hypomorphic, and conditional alleles. *Transgenic Res.* **10**, 69-80.
- Vionnet, N., Stoffel, M., Takeda, J., Yasuda, K., Bell, G. I., Zouali, H., Lesage, S., Velho, G., Iris, F., Passa, P. et al.** (1992). Nonsense mutation in the glucokinase gene causes early-onset non-insulin-dependent diabetes mellitus. *Nature* **356**, 721-2.
- Waeber, G., Thompson, N., Nicod, P. and Bonny, C.** (1996). Transcriptional activation of the GLUT2 gene by the IPF-1/STF-1/IDX-1 homeobox factor. *Mol. Endocrinol.* **10**, 1327-34.
- Wang, H., Brun, T., Kataoka, K., Sharma, A. J. and Wollheim, C. B.** (2007). MAFA controls genes implicated in insulin biosynthesis and secretion. *Diabetologia* **50**, 348-58.
- Wang, J., Elghazi, L., Parker, S. E., Kizilocak, H., Asano, M., Sussel, L. and Sosa-Pineda, B.** (2004a). The concerted activities of Pax4 and Nkx2.2 are essential to initiate pancreatic beta-cell differentiation. *Dev. Biol.* **266**, 178-89.
- Wang, L., Coffinier, C., Thomas, M. K., Gresh, L., Eddu, G., Manor, T., Levitsky, L. L., Yaniv, M. and Rhoads, D. B.** (2004b). Selective deletion of the Hnf1beta (MODY5) gene in beta-cells leads to altered gene expression and defective insulin release. *Endocrinology* **145**, 3941-9.
- Wells, J. M. and Melton, D. A.** (1999). Vertebrate endoderm development. *Annu. Rev. Cell Dev. Biol.* **15**, 393-410.
- Wilson, M. E., Scheel, D. and German, M. S.** (2003). Gene expression cascades in pancreatic development. *Mech. Dev.* **120**, 65-80.
- Wu, J., Duggan, A. and Chalfie, M.** (2001). Inhibition of touch cell fate by egl-44 and egl-46 in *C. elegans*. *Genes Dev.* **15**, 789-802.

- Xie, J., Cai, T., Zhang, H., Lan, M. S. and Notkins, A. L.** (2002). The Zinc-Finger Transcription Factor INSM1 Is Expressed during Embryo Development and Interacts with the Cbl-Associated Protein. *Genomics* **80**, 54-61.
- Yagi, T., Nada, S., Watanabe, N., Tamemoto, H., Kohmura, N., Ikawa, Y. and Aizawa, S.** (1993). A novel negative selection for homologous recombinants using diphtheria toxin A fragment gene. *Anal. Biochem.* **214**, 77-86.
- Yamagata, K., Furuta, H., Oda, N., Kaisaki, P. J., Menzel, S., Cox, N. J., Fajans, S. S., Signorini, S., Stoffel, M. and Bell, G. I.** (1996a). Mutations in the hepatocyte nuclear factor-4alpha gene in maturity-onset diabetes of the young (MODY1). *Nature* **384**, 458-60.
- Yamagata, K., Oda, N., Kaisaki, P. J., Menzel, S., Furuta, H., Vaxillaire, M., Southam, L., Cox, R. D., Lathrop, G. M., Boriraj, V. V. et al.** (1996b). Mutations in the hepatocyte nuclear factor-1alpha gene in maturity-onset diabetes of the young (MODY3). *Nature* **384**, 455-8.
- Yu, D., Ellis, H. M., Lee, E. C., Jenkins, N. A., Copeland, N. G. and Court, D. L.** (2000). An efficient recombination system for chromosome engineering in *Escherichia coli*. *Proc. Natl. Acad. Sci. U S A* **97**, 5978-83.
- Yu, H., Pretot, R. F., Burglin, T. R. and Sternberg, P. W.** (2003). Distinct roles of transcription factors EGL-46 and DAF-19 in specifying the functionality of a polycystin-expressing sensory neuron necessary for *C. elegans* male vulva location behavior. *Development* **130**, 5217-27.
- Zhang, C., Moriguchi, T., Kajihara, M., Esaki, R., Harada, A., Shimohata, H., Oishi, H., Hamada, M., Morito, N., Hasegawa, K. et al.** (2005). MafA is a key regulator of glucose-stimulated insulin secretion. *Mol. Cell. Biol.* **25**, 4969-76.
- Zhijin, W., Irizarry, R. A., Gentleman, R., Martinez-Murillo, F. and Spencer, F.** (2004). A Model Based Background Adjustment for Oligonucleotide Expression Arrays. *J. Am. Stat. Assoc.* **99**, 909-917.
- Zhou, Q. Y., Quaife, C. J. and Palmiter, R. D.** (1995). Targeted disruption of the tyrosine hydroxylase gene reveals that catecholamines are required for mouse fetal development. *Nature* **374**, 640-3.
- Zhu, M., Breslin, M. B. and Lan, M. S.** (2002). Expression of a novel zinc-finger cDNA, IA-1, is associated with rat AR42J cells differentiation into insulin-positive cells. *Pancreas* **24**, 139-45.