

6 Literatur

- Ahmad M, Rees RC, Ali SE. 2004. Escape from immunotherapy: possible mechanisms that influence tumor regression/progression. *Cancer Immunol. Immunother.* 53: 844-854.
- Allan GG, Peyron M. 1997. Depolymerisation of chitosan by means of nitrous acid. In: Muzzarelli RAA, Peter MG (eds). *Chitin Handbook*. Grottammare: Atec Edizioni. p. 175-180.
- Alonso MJ, Gupta RK, Min C, Siber GR, Langer R. 1994. Biodegradable microspheres as controlled-release tetanus toxoid delivery systems. *Vaccine* 12: 299-306.
- Anderson JM, Shive MS. 1997. Biodegradation and biocompatibility of PLA and PLGA microspheres. *Adv. Drug Del. Rev.* 28: 5-24.
- Anton D, Dabadghao S, Palucka K, Holm G, Yi Q. 1998. Generation of dendritic cells from peripheral blood adherent cells in medium with human serum. *Scand. J. Immunol.* 47: 116-121.
- van Apeldoorn AA, van Manen H-J, Bezemer JM, de Bruijn JD, van Blitterswijk AC, Otto C. 2004. Raman imaging of PLGA microsphere degradation inside macrophages. *J. Am. Chem. Soc.* 126: 13226-13227.
- Banchereau J, Briere F, Caux C, Davoust J, Lebecque S, Liu Y-J, Pulendran B, Palucka K. 2000. Immunobiology of dendritic cells. *Annu. Rev. Immunol.* 18: 767-811.
- Banchereau J, Palucka AK. 2005. Dendritic cells as therapeutic vaccines against cancer. *Nat. Rev. Immunol.* 5: 296-306.
- Barrera DA, Zylstra E, Lansbury PT, Langer R. 1993. Synthesis and RGD peptide modification of a new biodegradable copolymer: Poly(lactic acid-co-lysine). *J. Am. Chem. Soc.* 115: 11010-11011.
- Bendix D. 1998. Chemical synthesis of polylactide and its copolymers for medical application. *Polym. Degrad. Stab.* 59: 129-135.
- Berger TG, Feuerstein B, Strasser E, Hirsch U, Schreiner D, Schuler G, Schuler-Turner B. 2002. Large-scale generation of mature monocyte-derived dendritic cells for clinical application in cell factoriesTM. *J. Immunol. Meth.* 268: 131-140.
- Berger TG, Strasser E, Smith E, Carste C, Schuler-Turner B, Kaempgen E, Schuler G. 2005. Efficient elutriation of monocytes within a closed system (ElutraTM) for clinical-scale generation of dendritic cells. *J. Immunol. Meth.* 298: 61-72.
- Berger J, Reist M, Mayer JM, Felt O, Peppas NA, Gurny R. 2004. Structure and interactions in covalently and ionically crosslinked chitosan hydrogels for biomedical applications. *Eur. J. Pharm. Biopharm.* 57: 19-34.

- Bernhard H, Disis ML, Heimfeld S, Hand S, Gralow JR, Cheever MA. 1995. Generation of immunostimulatory dendritic cells from human CD34⁺ hematopoietic progenitor cells of the bone marrow and peripheral blood. *Cancer Res.* 55: 1099-1104.
- Bernkop-Schnürch A, Hornof M, Guggi D. 2004. Thiolated chitosans. *Eur. J. Pharm. Biopharm.* 57: 9-17.
- Bilati U, Allémann E, Doelker E. 2005a. Strategic approaches for overcoming peptide and protein instability within biodegradable nano- and microparticles. *Eur. J. Pharm. Biopharm.* 59: 375-388.
- Bilati U, Allémann E, Doelker E. 2005b. Poly(D,L-lactide-co-glycolide) protein-loaded nanoparticles prepared by a double emulsion method – processing and formulation issues for enhanced entrapment efficiency. *J. Microencapsulation* 22: 205-214.
- Bingaman A, Huxley VH, Rumbaut RE. 2003. Fluorescent dyes modify properties of proteins used in microvascular research. *Microcirculation* 10: 221-231.
- Bittner B, Morlock M, Koll H, Winter G, Kissel T. 1998. Recombinant human erythropoietin (rhEPO) loaded poly(lactide-co-glycolide) microspheres: Influence of the encapsulation technique and polymer purity on microsphere characteristics. *Eur. J. Pharm. Biopharm.* 45: 295-305.
- Bittner B, Kissel T. 1999. Ultrasonic atomization for spray-drying: a versatile technique for the preparation of protein loaded biodegradable microspheres. *J. Microencapsulation* 16: 325-341.
- Blanco D, Alonso MJ. 1997. Development and characterization of protein-loaded poly(lactide-co-glycolide) nanospheres. *Eur. J. Pharm. Biopharm.* 43: 287-294.
- Blanco D, Alonso MJ. 1998. Protein encapsulation and release from poly(lactide-co-glycolide) microspheres: effect of the polymer and protein properties and of the co-encapsulation of surfactants. *Eur. J. Pharm. Biopharm.* 45: 285-294.
- Blasi P, D'Souza SS, Selmin F, DeLuca PP. 2005. Plasticizing effect of water on poly(lactide-co-glycolide). *J. Controlled Release* 108 : 1-9.
- Boon T, Cerottini JC, van den Eynde B, van der Bruggen P, van Pel A. 1994. Tumor antigens recognized by T lymphocytes. *Annu. Rev. Immunol.* 12: 337-365.
- Boratynski J. 1985. Colorimetric micromethod for the determination of protein in solution with silver ions and dithizone. *Anal. Biochem.* 148: 213-219.
- Borchard G. 2001. Chitosans for gene delivery. *Adv. Drug Del. Rev.* 52: 145-150.
- Bos OJM, Labro FA, Fischer MJE, Wilting J, Janssen LHM. 1989. The molecular mechanism of the neutral-to-base transition of human serum albumin. *J. Biol. Chem.* 264: 953-959.
- Boury F, Ivanova T, Panaiotov I, Proust IE, Bois A, Richou J. 1995. Dynamic properties of poly(D,L-lactide) and polyvinyl alcohol monolayers at the air/water and dichlormethane/water interfaces. *J. Colloid Interface Sci.* 169: 380-392.

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of proteins utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248-254.

Brady JM, Cutright DE, Miller RA, Battistone GC. 1973. Resorption rate, route of elimination, and ultrastructure of implant site of polylactic acid in the abdominal wall of the rat. *J. Biomed. Mater. Res.* 7: 155-166.

Bright F. 1988. Bioanalytical application of fluorescence spectroscopy. *Anal. Chem.* 60: 1031-1039.

Briones M, Singh M, Ugozzoli M, Kazzaz J, Klakamp S, Ott G, O'Hagan D. 2001. The preparation, characterization, and evaluation of cationic microparticles for DNA vaccine delivery. *Pharm. Res.* 18: 709-712.

Brode S, Macary PA. 2004. Cross-presentation: dendritic cells and macrophages bite off more than they can chew! *Immunology* 112: 345-351.

Brunner A, Mäder K, Göpferich A. 1999. pH and osmotic pressure inside biodegradable microspheres during erosion. *Pharm. Res.* 16: 847-853.

Bummer PM, Koppenol S. 2000. Chemical and physical considerations in protein stability. In: McNally EJ (ed). *Protein formulation and delivery*. New York: Marcel Dekker. p. 5-69.

von Burkersroda F, Schedl L, Göpferich A. 2002. Why degradable polymers undergo surface erosion or bulk erosion. *Biomaterials* 23: 4221-4231.

Cai Q, Shi G, Bei J, Wang S. 2003. Enzymatic degradation behavior and mechanism of poly(lactide-co-glycolide) foams by trypsin. *Biomaterials* 24: 629-638.

Calvör A, Müller B. 1998. Production of microparticles by high-pressure homogenization. *Pharm. Dev. Technol.* 3: 297-305.

Caponetti G, Hrkach JS, Kriwet B, Poh M, Lotan N, Colombo P, Langer R. 1999. Microparticles of novel branched copolymers of lactic acid and amino acid: Preparation and characterization. *J. Pharm. Sci.* 88: 136-141.

Carbone FR, Kurts C, Bennett SR, Miller JF, Heath WR. 1998. Cross-presentation: a general mechanism for CTL immunity and tolerance. *Immunol. Today* 19: 368-373.

Carrasquillo KG, Aponte Carro JC, Alejandro A, Toro DD, Griebenow K. 2001a. Reduction of structural perturbations in bovine serum albumine by a non-aqueous microencapsulation. *J. Pharm. Pharmacol.* 53: 115-120.

Carrasquillo KG, Stanley AM, Aponte Carro JC, De Jesús P, Costantino HR, Bosques CJ, Griebenow K. 2001b. Non-aqueous encapsulation of excipient-stabilized spray-freeze dried BSA into poly(lactide-co-glycolide) microspheres results in release of native protein. *J. Controlled Release* 76: 199-208.

- Carrio A, Schwach G, Coudane J, Vert M. 1995. Preparation and degradation of surfactant-free PLGA microspheres. *J. Controlled Release* 37: 113-121.
- Carter DC, He X-M, Munson SH, Twiggs PD, Gernert KM, Broom MB, Miller TY. 1989. Three dimensional structure of human serum albumin. *Science* 244: 1195-1198.
- Carter DC, He X-M. 1990. Structure of human serum albumin. *Science* 249: 302-303.
- Carter DC, Ho JX. 1994. Structure of serum albumin. *Adv. Protein Chem.* 45: 153-203.
- Castellanos IJ, Cuadrado WL, Griebenow K. 2001. Prevention of structural perturbations and aggregation upon encapsulation of bovine serum albumin into poly(lactide-co-glycolide) microspheres using the solid-in-oil-in-water technique. *J. Pharm. Pharmacol.* 53: 1099-1107.
- Castellanos IJ, Cruz G, Crespo R, Griebenow K. 2002. Encapsulation-induced aggregation and loss in activity of γ -Chymotrypsin and their prevention. *J. Controlled Release* 81: 307-319.
- Castellanos IJ, Griebenow K. 2003. Improved α -Chymotrypsin stability upon encapsulation on PLGA microspheres by solvent replacement. *Pharm. Res.* 20: 1873-1880.
- Castellanos IJ, Crespo R, Griebenow K. 2003. Poly(ethylen glycol) as a stabilizer and emulsifying agent: a novel stabilization approach preventing aggregation and inactivation of proteins upon encapsulation in bioerodible polyester microspheres. *J. Controlled Release* 88: 135-145.
- Caux C, Dezutter-Dambuyant C, Schmitt D, Banchereau J. 1992. GM-CSF and TNF α cooperate in the generation of dendritic Langerhans cells. *Nature* 360: 258-261.
- Chen RF, Knutson JR. 1988. Mechanism of fluorescence concentration quenching of carboxyfluorescein in liposomes: energy transfer to nonfluorescent dimers. *Anal. Biochem.* 172: 61-77.
- Chen L, Apte RN, Cohen S. 1997. Characterisation of PLGA microspheres for controlled delivery of IL-1 α for tumor immunotherapy. *J. Controlled Release* 43: 261-272.
- Cherry WB. 1970. Fluorescence emission with special reference to standardization in immunofluorescence. In: Holborow EJ (ed). *Standardization in Immunofluorescence*. Oxford: Blackwell Scientific Publications. p. 127-153.
- Chu CC. 1981. The in-vitro degradation of poly(glycolic acid) sutures – effect of pH. *J. Biomed. Mat. Res.* 15: 795-804.
- Cleland JL, Johnson OF, Putney S, Jones AJS. 1997. Recombinant human growth hormone poly(lactic-co-glycolic acid) microsphere formulation development. *Adv. Drug Del. Rev.* 28: 71-84.
- Cleland JL, Duenas ET, Park A, Daugherty A, Kahn J, Kowalski J, Cuthbertson A. 2001. Development of poly-(D,L-lactide-co-glycolide) microsphere formulations containing recombinant human vascular endothelial growth factor to promote local angiogenesis. *J. Controlled Release* 72: 13-24.

Cohen S, Yoshioka T, Lucarelli M, Hwang LH, Langer R. 1991. Controlled delivery systems for proteins based on poly(lactic/glycolic acid) microspheres. *Pharm. Res.* 8: 713-720.

Conway BR, Alpar HO. 1996. Double emulsion microencapsulation of proteins as model antigens using polylactide polymers: Effect of emulsifier on the microsphere characteristics and release kinetics. *Eur. J. Pharm. Biopharm.* 42: 42-48.

Cortese JD, Voglino AL, Hackenbrock CR. 1991. Ionic strength of the intermembran space of intact mitochondria as estimated with Fluorescein-BSA delivered by low pH fusion. *J. Cell Biol.* 113: 1331-1340.

Costantino HR, Firouzabadian L, Hogeland K, Wu C, Beganski C, Carrasquillo KG, Córdova M, Giebenow K, Zale SE, Tracy MA. 2000. Protein spray freeze drying. Effect of atomization conditions on particle size and stability. *Pharm. Res.* 17: 1374-1383.

Costantino HR, Firouzabadian L, Wu C, Carrasquillo KG, Giebenow K, Zale SE, Tracy MA. 2002. Protein spray freeze drying. 2. Effect of formulation variables on size and stability. *J. Pharm. Sci.* 91: 388-395.

Costantino HR, Johnson OFL, Zale SE. 2004. Relationship between encapsulated drug particle size and initial release of recombinant human growth hormone from biodegradable microspheres. *J. Pharm. Sci.* 93: 2624-2634.

Couvreur P, Blanco-Prieto MJ, Puisieux F, Roques B, Tattal E. 1997. Multiple emulsion technology for the design of microspheres containing peptides and oligopeptides. *Adv. Drug Del. Rev.* 28: 85-96.

Crotts R, Park TG. 1998. Protein delivery from poly(lactic-co-glycolic acid) biodegradable microspheres: release kinetics and stability issues. *J. Microencapsulation* 15: 699-713.

Cui C, Schwendeman SP. 2001. Surface entrapment of polylysine in biodegradable poly(D,L-lactide-co-glycolide) microparticles. *Macromolecules* 34: 8426-8433.

Cutright DE, Beasley JD, Perez B. 1971. Histologic comparison of polylactic and polyglycolic acid sutures. *Oral Surg.* 32: 165-173.

Dailey LA, Kleemann E, Wittmar M, Gessler T, Schmehl T, Roberts C, Seeger W, Kissel T. 2003. Surfactant-free, biodegradable nanoparticles for aerosol therapy based on the branched polyesters, DEAPA-PVAL-g-PLGA. *Pharm. Res.* 20: 2011-2020.

Debenedetti PG, Tom JW, Yeo S-D, Lim B-B. 1993. Application of supercritical fluids for the production of sustained delivery devices. *J. Controlled Release* 24: 27-44.

Denis-Mize KS, Dupuis M, Singh M, Woo C, Ugozzoli M, O'Hagan D, Donnelly III JJ, Ott G, McDonald DM. 2003. Mechanisms of increased immunogenicity for DNA-based vaccines adsorbed onto cationic microparticles. *Cell. Immunol.* 225: 12-20.

Der-Balian GP, Kameda N, Rowly GL. 1988. Fluorescein labeling of F_{ab} while preserving single thiol. *Anal. Biochem.* 173: 59-63.

Diehl H, Markuszewski R. 1989. Studies on Fluorescein VII. The fluorescence of fluorescein as a function of pH. *Talanta* 36: 416-418.

Ding AG, Schwendeman SP. 2004. Determination of water-soluble acid distribution in poly(lactide-co-glycolide). *J. Pharm. Sci.* 93: 322-331.

Diwan M, Elamanchili P, Lane H, Gainer A, Samuel J. 2003. Biodegradable nanoparticle mediated antigen delivery to human cord blood derived dendritic cells for induction of primary T cell responses. *J. Drug Targeting* 11: 495-507.

Domb AJ. 1994. Implantable biodegradable polymers for site-specific drug delivery. In: Domb AJ (ed.) *Polymeric site-specific pharmacotherapy*. Chichester: Wiley. p. 1-26.

Dong Y, Qui W, Ruan Y, Wu Y, Wang M, Xu C. 2001. Influence of molecular weight on critical concentration of chitosan/formic acid liquid crystalline solution. *Polym. J.* 33: 387-389.

Doolittle AK. 1954. Properties of individual solvents. In: *The technology of solvents and plasticizers*. Chpt. 12. New York: Wiley. p. 492-742.

Dunne M, Corrigan OI, Ramtoola Z. 2000. Influence of particle size and dissolution conditions on the degradation properties of polylactide-co-glycolide particles. *Biomaterials* 21: 1659-1668.

Ebeling W, Hennrich N, Klockow M, Metz H, Orth HD, Lang H. 1974. Proteinase K from *Tritirachium album* Limber. *Eur. J. Biochem.* 47: 91-97.

Endl E, Steinbach P, Scharfe J, Fickweiler S, Worle K, Hofstadter F. 1996. Cell-type-specific response to shock waves of suspended or pelleted cells as analysed by flow cytometry or electrical cell volume determination. *Ultrasound Med. Biol.* 22: 515-525.

Fahr A, Kissel T. 1998. Mikropartikel und Implantate: Arzneiformen zur parenteralen Applikation. In: Müller RH, Hildebrand GE (ed). *Pharmazeutische Technologie: Moderne Arzneiformen*. 2. Aufl. Stuttgart: Wissenschaftliche Verlagsgesellschaft. p. 243-258.

Fanger NA, Wardwell K, Shen L, Tedder TF, Guyre PM. 1996. Type I (CD64) and type II (CD32) Fc gamma receptor-mediated phagocytosis by human blood dendritic cells. *J. Immunol.* 157: 541-548.

Faraasen S, Vörös J, Csúcs G, Textor M, Merkle HP, Walter E. 2003. Ligand-specific targeting of microspheres to phagocytes by surface modification with poly(L-lysine)-grafted poly(ethylene glycol) conjugate. *Pharm. Res.* 20: 237-246.

Farkas A, Conrad C, Tonel G, Borbenyi Z, Kemeny L, Dobozy A, Nestle FO. 2006. Current state and perspectives of dendritic cell vaccination in cancer immunotherapy. *Skin Pharmacol. Physiol.* 19: 124-131.

FDA (Food and Drug Administration). 1997. International Conference on Harmonisation; Draft Guideline on Impurities, Residual Solvents, Availability. *Fed. Regist.* 62: 24302-24309.

- Felt O, Buri P, Gurny R. 1998. Chitosan: a unique polysaccharide for drug delivery. *Drug Dev. Ind. Pharm.* 24: 979-993.
- Ferrer ML, Duchowicz R, Carrasco B, de la Torre JG, Acuña AL. 2001. The conformation of serum albumin in solution: A combined phosphorescence depolarisation-hydrodynamic modelling study. *Biophys. J.* 80: 2422-2430.
- Finley JH. 1961. Spectrophotometric determination of polyvinyl alcohol in paper coatings. *Anal. Chem.* 33: 1925-1927.
- Fischer S, Foerg C, Ellenberger S, Merkle HP, Gander B. 2006. One-step preparation of polyelectrolyte-coated PLGA microparticles and their functionalisation with model antigens. *J. Controlled Release* in press.
- Flamand V, Sornasse T, Thielemans K, Demanet C, Bakkus M, Bazin H, Tielemans F, Leo O, Urbain J, Moser M. 1994. Murine dendritic cells pulsed in vitro with tumor antigen induce tumor resistance in vivo. *Eur. J. Immunol.* 24: 605-610.
- Folta-Stogniew E, Williams KR. 1999. Determination of molecular masses of proteins in solution: Implementation of an HPLC size exclusion chromatography and laser scattering service in a core laboratory. *J. Biomol. Techn.* 10: 51-63.
- Förster T, König E. 1957. Absorptionsspektren und Fluoreszenzeigenschaften konzentrierter Lösungen organischer Stoffe. *Zeitschrift für Elektrochemie* 61: 344-348.
- Förster T (ed). 1982. Fluoreszenz organischer Verbindungen. Nachdruck der 1. Auflage. Göttingen: Vandenhoeck & Ruprecht.
- Foster JF. 1960. Plasma albumin. In: Putnam FW (ed). *The plasma proteins*. Vol. 1. New York: Academic Press. p. 179-239.
- Frangione-Beebe M, Rose RT, Kaumaya PTP, Schwendeman SP. 2001. Microencapsulation of a synthetic peptide epitope for HTLV-1 in biodegradable poly(D,L-lactide-co-glycolide) microspheres using a novel encapsulation technique. *J. Microencapsulation* 18: 663-677.
- Freitas S, Walz A, Merkle HP, Gander B. 2003. Solvent extraction employing a static micromixer: a simple, robust and versatile technology for the microencapsulation of proteins. *J. Microencapsulation* 20: 67-85.
- Freitas S, Merkle HP, Gander B. 2004. Ultrasonic atomisation into reduced pressure atmosphere – envisaging aseptic spray-drying for microencapsulation. *J. Controlled Release* 95: 185-195.
- Freitas S, Merkle HP, Gander B. 2005. Microencapsulation by solvent extraction/evaporation: reviewing the state of the art of microsphere preparation process technology. *J. Controlled Release* 102: 313-332.
- Fu K, Griebenow K, Hsieh L, Klibanov AM, Langer R. 1999. FTIR characterisation of the secondary structure of proteins encapsulated within PLGA microspheres. *J. Controlled Release* 58: 357-366.

- Fu K, Klibanov AM, Langer R. 2000. Protein stability in controlled-release systems. *Nat. Biotechnol.* 18: 24-25.
- Fu K, Harrel R, Zinski K, Jaklenec A, Frazier J, Lotan N, Burke P, Klibanov AM, Langer R. 2003. A potential approach for decreasing the burst effect of protein from PLGA microspheres. *J. Pharm. Sci.* 92: 1583-1591.
- Gatza E, Okada CY. 2006. Adjuvant IL-15 does not enhance the efficacy of tumor cell lysate-pulsed dendritic cell vaccines for active immunotherapy of T cell lymphoma. *Cancer Immunol. Immunother.* 55: 420-432.
- Geissler M, Weth R. 2002. Immunotherapie: Neue Entwicklungen. *Praxis* 91: 2236-2246.
- Giancola C, Sena C, Fessa D, Graziano G, Barone G. 1997. DCS studies on bovine serum albumin denaturation: Effects of ionic strength and SDS concentration. *Int. J. Biol. Macromol.* 20: 193-204.
- Göpferich A, Langer R. 1993. Modeling of polymer erosion. *Macromol.* 26: 4105-4112.
- Göpferich A. 1996a. Polymer degradation and erosion: Mechanisms and applications. *Eur. J. Pharm. Biopharm.* 42: 1-11.
- Göpferich A. 1996b. Mechanisms of polymer degradation and erosion. *Biomaterials* 17: 103-104.
- Graf L, Craik CS, Patthy A, Rocznik S, Fletterick RJ, Rutter WJ. 1987. Selective alteration of substrate specificity by replacement of aspartic acid-189 with lysine in the binding pocket of trypsin. *Biochemistry* 26: 2616-2623.
- Grayson ACR, Cima MJ, Langer R. 2005. Size and temperature effects on poly(lactic-co-glycolic acid) degradation and microreservoir device performance. *Biomaterials* 26: 2137-2145.
- Grizzi I, Garreau H, Li S, Vert M. 1995. Hydrolytic degradation of devices based on poly(D,L-lactic acid) size-dependence. *Biomaterials* 16: 305-311.
- Guermonprez P, Valladeau J, Zitvogel L, Théry C, Amigorena S. 2002. Antigen presentation and T cell stimulation by dendritic cells. *Annu. Rev. Immunol.* 20: 621-667.
- Hackham DJ, Rotstein OD, Zhang W-J, Demaurex N, Woodside M, Tsai O, Grinstein S. 1997. Regulation of phagosomal acidification. Differential targeting on Na⁺/H⁺-exchangers, Na⁺/K⁺-ATPases, and vacuolar-type ATPases. *J. Biol. Chem.* 272: 29810-29820.
- Hakkarainen M. 2002. Aliphatic polyesters: Abiotic and biotic degradation and degradation products. *Adv. Polym. Sci.* 157: 113-138.
- Hamilton JA. 2004. Fatty acid interaction with proteins: what X-ray crystal and NMR solution structures tell us. *Prog. Lipid Res.* 43: 177-199.
- Haugland RP. 1996. Handbook of fluorescent probes and research products, 6th ed., Molecular Probes, Eugene, USA

Haugland RP. 2002. Handbook of fluorescent probes and research products, 9th ed., Molecular Probes, Eugene, USA

Haverkamp V, Ehrenfeld M, Gebauer K, Hessel V, Löwe H, Richter T, Wille C. 1999. The potential of micromixers for contacting of disperse liquid phases. *Fresenius J. Anal. Chem.* 364: 617-624.

Hed J, Hallden G, Johansson SGO, Larsson P. 1987. The use of fluorescence quenching in flow cytometry to measure the attachment and ingestion phase in phagocytosis in peripheral blood without prior cell separation. *J. Immunol. Meth.* 101: 119-125.

Herbert P, Murphy K, Johnson OF, Dong N, Jaworowicz W, Tracy MA, Cleland JL, Putney SD. 1998. A large-scale process to produce microencapsulated proteins. *Pharm. Res.* 15: 357-361.

Herrmann WO, Haehnel W. 1927. Über den Polyvinylalkohol. *Berichte der Deutschen Chemischen Gesellschaft* 60: 1658-1663.

Herrmann J, Bodmeier R. 1995. Somatostatin containing biodegradable microspheres prepared by a modified solvent evaporation method based on w/o/w multiple emulsions. *Int. J. Pharm.* 126: 129-138.

Herrmann J, Bodmeier R. 1998. Biodegradable, somatostatin acetate containing microspheres prepared by various aqueous and non-aqueous solvent evaporation methods. *Eur. J. Pharm. Biopharm.* 45: 72-85.

Herschel JFW. 1845. On a case of superficial colour presented by a homogeneous liquid internally colourless. *Phil. Trans. R. Soc.* 135: 143-145. abgedruckt bei Lakowicz 1999.

Hessel V, Löwe H, Schönfeld F. 2005. Micromixers - a review on passive and active mixing principles. *Chem. Eng. Sci.* 60: 2479-2501.

Heukeshoven J, Dernick R. 1985. Simplified method for silver staining of proteins in polyacrylamide gels and the mechanism of silver staining. *Electrophoresis* 6: 103-112.

Hirayama K, Akashi S, Furuya M, Fukuhara K-I. 1990. Rapid confirmation and revision of the primary structure of bovine serum albumin by ESIMS and FRIT-FAB LC/MS. *Biochem. Biophys. Res. Comm.* 173: 639-646.

Holtzhauer M (ed). 1996. *Methoden in der Proteinanalytik*. Berlin: Springer Verlag.

Hornung V, Rothenfusser S, Britsch S, Krug A, Jahrsdorfer B, Giese T, Endres S, Hartmann G. 2002. Quantitative expression of toll-like receptor 1-10 mRNA in cellular subsets of human peripheral blood mononuclear cells and sensitivity to CpG oligodeoxynucleotides. *J. Immunol.* 168: 4531-4537.

Howell PB, Mott DR, Golden JP, Ligler FS. 2004. Design and evaluation of a Dean vortex based micromixer. *Lab. Chip.* 4: 663-669.

Hrkach JS, Ou J, Lotan N, Langer R. 1995. Synthesis of poly(L-lactic acid-co-L-lysine). *Macromolecules* 28: 4736-4739.

Hsu FJ, Benike C, Fagnoni F, Liles TM, Czerwinski D, Taidi B, Engleman EG, Levy R. 1996. Vaccination of patients with B-cell lymphoma using autologous antigen-pulsed dendritic cells. *Nat. Med.* 2: 52-58.

Hunter AK, Carta G. 2001. Effects of bovine serum albumin heterogeneity on frontal analysis with anion-exchange media. *J. Chromatogr. A* 937: 13-19.

Jacques YV, Bainton DF. 1978. Changes in pH within phagolytic vacuoles of human neutrophils and monocytes. *Laboratory Investigation* 39: 179-185.

Jain RA, Rhodes CT, Railkar AM, Malick AW, Shah NH. 2000. Controlled release of drug from injectable in situ formed biodegradable PLGA microspheres: effect of various formulative variables. *Eur. J. Pharm. Biopharm.* 50: 257-262.

Janatova J, Fuller JK, Hunter MJ. 1968. The heterogeneity of bovine albumin with respect to sulfhydryl and dimer content. *J. Biol. Chem.* 243: 3612-3622.

Janeway CA, Travers P, Walport M, Shlomchik M. 2002. *Immunologie*. 5. Auflage. Heidelberg: Spektrum Akademischer Verlag.

Jeon Y-I, Jeong Y-I, Jang M-K, Park Y-H, Nah J-W. 2000. Effect of solvent on the preparation of surfactant-free poly(D,L-lactide-co-glycolide) nanoparticles and norfloxacin release characteristics. *Int. J. Pharm.* 207: 99-108.

Jeong Y-I, Shim Y-H, Choi C, Jang M-K, Shin GM, Nah J-W. 2003. Surfactant-free nanoparticles of poly(D,L-lactide-co-glycolide) prepared with poly(L-lactide)/poly(ethylene glycol). *J. Appl. Polym. Sci.* 89: 1116-1123.

Jiang G, Woo BH, Kang F, Singh J, DeLuca PP. 2002. Assessment of protein release kinetics, stability and protein polymer interaction of lysozyme encapsulated poly(D,L-lactide-co-glycolide) microspheres. *J. Controlled Release* 79: 137-145.

Jiang G, Qui W, DeLuca PP. 2003. Preparation and in vitro/in vivo evaluation of Insulin-loaded poly(acryloyl-hydroxyethyl starch)-PLGA composite microspheres. *Pharm. Res.* 20: 452-459.

Jilek S, Ulrich M, Merkle HP, Walter E. 2004. Composition and surface charge of DNA-loaded microparticles determine maturation and cytokine secretion in human dendritic cells. *Pharm. Res.* 21: 1240-1247.

Jobmann M, Rafler G. 1998. Controlled release systems of biodegradable polymers. 6th Communication: Control of particle size and size distribution of microparticles produced by the solvent evaporation technique. *Pharm. Ind.* 60: 979-982.

Johanson P, Men Y, Audran R, Corradin G, Merkle HP, Gander B. 1998. Improved stability and release kinetics of microencapsulated tetanus toxoid by co-encapsulation of additives. *Pharm. Res.* 15: 1103-1110.

Johanson P, Merkle HP, Gander B. 2000a. Technological considerations related to the up-scaling of protein microencapsulation by spray-drying. *Eur. J. Pharm. Biopharm.* 50: 413-417.

Johanson P, Men Y., Merkle HP, Gander B. 2000b. Revisiting PLA/PLGA microspheres: an analysis of their potential in parenteral vaccination. *Eur. J. Pharm. Biopharm.* 50: 129-146.

Johnson OF, Jaworowicz W, Cleland JL, Bailey L, Charnis M, Duenas E, Wu C, Shepard D, Magil S, Last T, Jones AJS, Putney SD. 1997. The stabilization and encapsulation of human growth hormone into biodegradable microspheres. *Pharm. Res.* 14: 730-735.

Jones E. 1996. Methylene chloride – An overview of human and environmental effects. *Pharm. Tech. Eur.* 8: 30-32.

Jonuleit H, Kuhn U, Muller G, Steinbrink K, Paragnik L, Schmitt E, Knop J, Enk AH. 1997. Pro-inflammatory cytokines and prostaglandins induce maturation of potent immunostimulatory dendritic cells under fetal calf serum-free conditions. *Eur. J. Immunol.* 27: 3135-3142.

Joshi DP, Lan-Chun-Fung YL, Pritchard JG. 1979. Determination of poly(vinyl alcohol) via its complex with boric acid and iodine. *Anal. Chim. Acta* 104: 153-160.

Kalinski P, Hilkens CM, Wierenga EA, Kapsenberg ML. 1999. T-cell priming by type-1 and type-2 polarized dendritic cells: the concept of a third signal. *Immunol. Today* 20: 561-567.

Kang F, Singh J. 2003. Conformational stability of a model protein (bovine serum albumin) during primary emulsification process of PLGA microsphere synthesis. *Int. J. Pharm.* 260: 149-156.

Kao JY, Zhang M, Chen CM, Chen JJ. 2005. Superior efficacy of dendritic cell-tumor fusion vaccine compared with tumor lysate-pulsed dendritic cell vaccine in colon cancer. *Immunol. Lett.* 101: 154-159.

Karpyuk AD, Kolyada NS, Pshenichnikova EY. 1993. Determination of polyacrylamid and polyvinyl alcohol by a turbidimetric method using potassium permanganate in technological solutions. *Zavod. Lab.* 59: 4-6; übersetzt in *Industrial Laboratory* 59: 357-360.

Kasturi SP, Sachaphibulkij K, Roy K. 2005. Covalent conjugation of polyethyleneimine on biodegradable microparticles for delivery of plasmid DNA vaccines. *Biomaterials* 26: 6375-6385.

Kawakatsu T, Komori H, Nakajima M, Kikuchi Y, Yonemoto T. 1999. Production of a monodisperse oil-in-water emulsion using a crossflow-type silicon microchannel plate. *J. Chem. Eng. Japan* 32: 241-244.

Kawamura A (ed). 1977. *Fluorescent antibody techniques and their applications*. Second Edition. Tokyo: University of Tokyo Press.

Kawaski A. 1983. Excitation energy transfer and its manifestation isotropic media. *Photochem. Photobiol.* 38: 487-508.

- Kim DS, Lee SH, Kwon TH, Ahn CH. 2005. A serpentine laminating micromixer combining splitting/recombination and advection. *Lab. Chip.* 5: 739-747.
- Kissel T, Li XY, Volland C, Görich S, Koneberg R. 1996. Parenteral protein delivery systems using biodegradable polyesters of ABA block structure, containing hydrophobic poly(lactide-co-glycolide) A blocks and hydrophilic poly(ethylen oxide) B blocks. *J. Controlled Release* 39: 315-326.
- Kissel T, Li X, Unger F. 2002. ABA-triblock copolymers from biodegradable polyester A blocks and hydrophilic poly(ethylen oxide) B blocks as a candidate for in situ forming hydrogel delivery systems for proteins. *Adv. Drug Del. Rev.* 54: 99-134.
- Kloetzel P-M, Sijts A. 2002. Processing and presentation of tumor-associated antigens. In: Stuhler G, Walden P (eds). *Cancer immune therapy*. Weinheim: Wiley-VCH. p. 30-39.
- Kosa T, Maruyama T, Otagiri M. 1998. Species difference of serum albumins: II Chemical and thermal stability. *Pharm. Res.* 15: 449-454.
- Kranz H, Brazeau GB, Napaporn J, Martin RL, Millard W, Bodmeier R. 2001. Myotoxicity study of biodegradable in-situ forming drug delivery devices. *Int. J. Pharm.* 212: 11-18.
- Krystal G, MacDonald C, Munt B, Ashwell S. 1985. A method for quantitating nanogram amounts of soluble protein using the principle of silver binding. *Anal. Biochem.* 148: 451-460.
- Kumar G, Smith PJ, Payne GF. 1999. Enzymatic grafting of a natural product onto chitosan to confer water solubility under basic conditions. *Biotechnol. Bioeng.* 63: 154-165.
- Kumar NMVR, Bakowsky U, Lehr CM. 2004. Preparation and characterization of cationic PLGA nanospheres as DNA carriers. *Biomaterials* 25: 1771-1777.
- Kuznetsov AN, Ebert B, Lassmann G, Shapiro AB. 1975. Adsorption of small molecules to bovine serum albumin studied by the spin-probe method. *Biochem. Biophys. Acta* 379: 139-146.
- Lakowicz JR (ed). 1999. *Principles of fluorescence spectroscopy*. 2. Edition. New York: Kluwer Academic/Plenum Publishers.
- Lam XM, Duenas ET, Cleland JL. 2001. Encapsulation and stabilisation of nerve growth factor into poly(lactide-co-glycolide) acid microspheres. *J. Pharm. Sci.* 90: 1356-1365.
- Laurencin CT, Elgendy H. 1994. The biocompatibility and toxicity of degradable polymeric materials: Implications for drug delivery. In: Domb AJ (ed). *Polymeric site-specific pharmacotherapy*. Chichester: Wiley. p. 27-46.
- Leach WT, Simpson DT, Val TN, Anuta EC, Yu Z, Williams RO, Johnston KP. 2005. Uniform encapsulation of stable protein nanoparticles produced by spray freezing for the reduction of burst release. *J. Pharm. Sci.* 94: 56-69.

- Lee AC, Oh JT, Jang MH, Chung SI. 1999. Quantitative analysis of polyvinyl alcohol on the surface of poly(D,L-lactide-co-glycolide) microparticles prepared by solvent evaporation method: effect of particle size and PVA concentration. *J. Controlled Release* 59: 123-132.
- Lee NY, Yamada M, Seki M. 2005. Development of a passive micromixer based on repeated fluid twisting and flattening, and its application to DNA purification. *Anal. Bioanal. Chem.* 383: 776-782.
- Leong KW, Mao H-Q, Truong-Le VL, Roy K, Walsh SM, August JT. 1998. DNA-polycation nanospheres as non-viral gene delivery vehicles. *J. Controlled Release* 53: 183-193.
- Li SM, Garreau H, Vert M. 1990a. Structure-property relationships in the case of the degradation of massive aliphatic poly(α -hydroxy acids) in aqueous media. Part 1: Degradation of lactide-glycolide copolymers: PLA37.5GA25 and PLA75GA25. *J. Mater. Sci., Mater. Med.* 1: 131-139.
- Li SM, Garreau H, Vert M. 1990b. Structure-property relationships in the case of the degradation of massive poly(α -hydroxy acids) in aqueous media. Part 2: Poly(DL-lactic acid). *J. Mater. Sci., Mater. Med.* 1: 123-130.
- Li X, Kissel T. 1993. Synthesis and properties of a biodegradable ABA triblock copolymers consisting of poly(L-lactic acid) or poly(L-lactic-co-glycolic acid) A-blocks attached to central poly(oxyethylene) B-blocks. *J. Controlled Release* 27: 247-257.
- Li X, Volland C, Kissel T. 1994. In-vitro degradation and bovine serum albumin release of the ABA triblock copolymers consisting of poly(L(+))lactic acid, or poly(L(+))lactic acid-co-glycolic acid) A-blocks attached to central poly(oxyethylene) B-blocks. *J. Controlled Release* 32: 121-128.
- Li S. 1999. Hydrolytic degradation characteristics of aliphatic polyesters derived from lactic and glycolic acids. *J. Biomed. Mater. Res. (Appl. Biomater.)* 48: 342-353.
- Li X, Zhang Y, Yan R, Jia W, Yuan M, Deng X, Huang Z. 2000. Influence of process parameters on the protein stability encapsulated in poly-DL-lactide-poly(ethylene glycol) microspheres. *J. Controlled Release* 68: 41-52.
- Li L, Schwendeman SP. 2005. Mapping neutral microclimate pH in PLGA microspheres. *J. Controlled Release* 101: 163-173.
- Litvinova LS, Gankina ES, Belen'kii BG. 1990. Determination of polyvinyl alcohol in the presence of triethanolamine by thin-layer chromatography. *Zavod. Lab.* 59: 7-9; übersetzt in *Industrial Laboratory* 56: 254-257.
- Lin J-L, Lee K-H, Lee G-B. 2005. Active mixing inside microchannels utilizing dynamic variation of gradient zeta potentials. *Electrophoresis* 26: 4605-4615.
- Lindahl PE. 1948. Principle of a counter-streaming centrifuge for the separation of particles of different sizes. *Nature* 161: 648-649.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. 1951. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193: 265-275.

- Lottspeich F, Zorbas H (eds). 1998. Bioanalytik. Heidelberg: Spektrum Akademischer Verlag.
- Lotze MT, Farhood H, Wilson CC, Storkus WJ. 1999. Dendritic cell therapy of cancer and HIV infection. In: Lotze MT, Thomson AW (eds). Dendritic cells, biology and clinical application. San Diego: Academic Press. p. 459-486.
- Lu L, Garcia CA, Mikos AG. 1999. In vitro degradation of thin poly(D,L-lactic-glycolic acid) films. *J. Biomed. Mater. Res.* 46: 236-244.
- van der Lubben IM, Verhoef JC, Borchard G, Junginger HE. 2001a. Chitosan for mucosal vaccination. *Adv. Drug Del. Rev.* 52: 139-144.
- van der Lubben IM, Verhoef JC, van Aelst AC, Borchard G, Junginger HE. 2001b. Chitosan microparticles for oral vaccination: preparation, characterisation and preliminary in vivo uptake studies in murine Peyer's patches. *Biomaterials* 22: 687-694.
- van der Lubben IM, Kersten G, Fretz MM, Beuvery C, Verhoef JC, Junginger HE. 2003. Chitosan microparticles for mucosal vaccination against diphtheria: oral and nasal efficacy studies in mice. *Vaccine* 21: 1400-1408.
- Lucke A, Kiermaier J, Göpferich A. 2002. Peptide acylation by poly(α -hydroxy esters). *Pharm. Res.* 19: 175-181.
- Ludewig B, McCoy K, Pericin M, Ochsenbein AF, Dumrese T, Odermatt B, Toes REM, Melief CJM, Hengartner H, Zinkernagel RM. 2001. Rapid peptide turnover and inefficient presentation of exogenous antigen critically limit the activation of self-reactive CTL by dendritic cells. *J. Immunol.* 166: 3678-3687.
- Ma W, Tang C, Lau L. 2005. Specificity of trypsin and chymotrypsin: Loop-motion-controlled dynamic correlation as a determinant. *Biophys. J.* 89: 1183-1193.
- Mäder K, Bittner B, Li Y, Wohlauf W, Kissel T. 1998. Monitoring microviscosity and microacidity of the albumin environment inside degrading microparticles from poly(lactide-co-glycolide) (PLG) or ABA-triblock polymers containing hydrophobic poly(lactide-co-glycolide) A blocks and hydrophilic poly(ethyleneoxide) B blocks. *Pharm. Res.* 15: 787-793.
- Mahnke K, Schmitt E, Bonifaz L, Henk A, Jonulei H. 2002. Immature, but not inactive: the tolerogenic function of immature dendritic cells. *Immunol. Cell Biol.* 80: 477-483.
- Malawista SE, Gee JBL, Bensch KG. 1971. Cytochalasin B reversibly inhibits phagocytosis: Functional, metabolic, and ultrastructural effects in human blood leukocytes and rabbit alveolar macrophages. *Yale J. Biol. Med.* 44: 286-300.
- Mansouri S, Lavigne P, Corsi K, Benderdour M, Beaumont E, Fernandes JC. 2004. Chitosan-DNA nanoparticles as non-viral vectors in gene therapy: strategies to improve transfection efficacy. *Eur. J. Pharm. Biopharm.* 57: 1-8.
- Manuel WS, Zheng JI, Hornsby PJ. 2001. Transfection by polyethyleneimine-coated microspheres. *J. Drug Target.* 9: 15-22.

- Maraskovsky E, Daro E, Roux E, Teepe M, Maliszewski CR, Hoek J, Caron D, Lebsack ME, McKenna HJ. 2000. In vivo generation of human dendritic cell subsets by Flt3 ligand. *Blood* 96: 878-884.
- Martinou A, Tsigos I, Bouriotis V. 1997. Preparation of chitosan by enzymatic deacetylation. In: Muzzarelli RAA, Peter MG (eds). *Chitin Handbook*. Grottammare: Atec Edizioni. p. 501-505.
- Mathiowitz E, Jacob J, Pekarek K, Chickering III D. 1993. Morphological characterisation of bioerodible polymers. 3. Characterisation of the erosion and intact zones in polyanhydrides using scanning electron microscopy. *Macromol.* 26: 6756-6765.
- Matsuno K, Ezaki T, Kudo S, Uehara Y. 1996. A life stage of particle-loaden rat dendritic cells in vivo: their terminal division, active phagocytosis, and translocation from the liver to the draining lymph. *J. Exp. Med.* 183: 1865-1878.
- Mayordomo JI, Zorina T, Storkus WJ, Zitvogel L, Celluzzi C, Falo LD, Melief CJ, Ildstad ST, Kast WM, Deleo AB, et al. 1995. Bone marrow-derived dendritic cells pulsed with synthetic tumour peptides elicit protective and therapeutic antitumour immunity. *Nat. Med.* 1: 1297-1302.
- Mazzoni A, Segal DM. 2004. Controlling the Toll road to dendritic cell polarization. *J Leukoc. Biol.* 75: 721-730.
- McEwen CR, Stallard RW, Juhos ET. 1968. Separation of biological particles by centrifugal elutriation. *Anal. Biochem.* 23: 369-377.
- Medzhitov R, Preston-Hurlburt P, Janeway CA Jr. 1997. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 388: 394-397.
- Medzhitov R, Janeway CA Jr. 2002. Decoding the patterns of self and nonself by the innate immune system. *Science* 296: 298-300.
- Meinel L, Illi OE, Zapf J, Malfanti M, Merkle HP, Gander B. 2001. Stabilizing insulin-like growth factor-I in poly(D,L-lactide-co-glycolide) microspheres. *J. Controlled Release* 70: 193-202.
- Melin J, Giménez G, Roxhed N, van der Wijngaart W, Stemme G. 2004. A fast passive and planar liquid sample micromixer. *Lab. Chip.* 4: 214-219.
- Müller RH (ed). 1996. *Zetapotential und Partikelladung in der Laborpraxis*. Stuttgart: Wissenschaftliche Verlagsgesellschaft.
- Müller RH, Schuhmann R (ed). 1996. *Teilchengrößenmessung in der Laborpraxis*. Stuttgart: Wissenschaftliche Verlagsgesellschaft.
- Murata J, Ohya Y, Ouchi T. 1996. Possibility of application of quaternary chitosan having pendant galactose residues as gene delivery tool. *Carbohydr. Polym.* 29: 69-74.

Murata J, Ohya Y, Ouchi T. 1997. Design of quarternary chitosan conjugate having antennary residues as a gene delivery tool. *Carbohydr. Polym.* 32: 105-109.

Muzzarelli RAA, Tubertini O. 1969. Chitin and Chitosan as chromatographic supports and adsorbents for collection of metall ions from organic and aqueous solutions and sea-water. *Talanta* 16: 1571-1577.

Muzzarelli RAA. 1998. Colorimetric determination of chitosan. *Anal. Biochem.* 260: 255-257.

Muzzarelli C, Tosi G, Francescangeli O, Muzzarelli RAA. 2003. Alkaline chitosan solutions. *Carbohydr. Res.* 338: 2247-2255.

Na DH, Youn YS, Lee SD, Son M-W, Kim W-B, DeLuca PP, Lee KC. 2003. Monitoring of peptide acylation inside degrading PLGA microspheres by capillary electrophoresis and MALDI-TOF mass spectrometry. *J. Controlled Release* 92: 291-299.

Najar HM, Ruhl S, Bru-Capdeville AC, Peters JH. 1990. Adenosine and its derivatives control human monocyte differentiation into highly accessory cells versus macrophages. *J. Leukoc. Biol.* 47: 429-439.

Nau P, Peters JH. 1986. Human peripheral blood accessory cell: isolation by hypotonic density gradient, functional, and phenotypical characterization. *Immunobiology* 173: 82-97.

Nelson D, Cox M (eds). 2001. *Lehninger Biochemie*. 3. Auflage. Berlin: Springer Verlag.

Nestle FO, Alijagic S, Gilliet M, Sun Y, Grabbe S, Dummer R, Burg G, Schadendorf D. 1998. Vaccination of melanoma patients with peptide- or tumor lysate-pulsed dendritic cells. *Nature Med.* 4: 328-332.

Neuhoff V, Stamm R, Eibl H. 1985. Clear background and high sensitive protein staining with Coomassie Blue dyes in polyacrylamide gels: A systematic analysis. *Electrophoresis* 6: 427-448.

Neuhoff V, Arold N, Taube D, Ehrhardt W. 1988. Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis* 9: 255-262.

Niu C-H, Chiu H-H. 1998. FDA perspective on peptide formulation and stability issues. *J. Pharm. Sci.* 87: 1331-1334.

No HK, Park NY, Lee SH, Meyers SP. 2002. Antibacterial activity of chitosans and chitosan oligomers with different molecular weights. *Int. J. Food Microbiology* 74: 65-72.

Oakes J. 1976. Thermally denaturated proteins. *J. Chem. Soc. Faraday I* 72: 228-237.

Ogawa Y, Yamamoto M, Okada H, Yashiki T, Shimamoto. 1988. A new technique to efficiently entrap leuprolide acetate into microcapsules of polylactic acid or copoly(lactic/glycolic) acid. *Chem. Pharm. Bull.* 36: 1095-1103.

- O'Hagan DT, Singh M. 2004. Microparticles as vaccine adjuvants and delivery systems. In: Kaufmann SHE (ed). Novel vaccination strategies. Weinheim: Wiley-VCH Verlag. p. 147-172.
- O'Hagan DT, Singh M, Dong C, Ugozzoli M, Berger K, Glazer E, Selby M, Wining M, Ng P, Crawford P, Paliard X, Coates S, Houghton M. 2004. Cationic microparticles are a potent delivery system for a HCV DNA vaccine. *Vaccine* 23: 672-680.
- Oster CG, Kim N, Grode L, Barbu-Tudoran L, Schaper AK, Kaufmann SHE, Kissel T. 2005. Cationic microparticles consisting of poly(lactide-co-glycolide) and polyethylenimine as carrier systems for parenteral DNA vaccination. *J. Controlled Release* 104: 359-377.
- Otto M (ed). 1995. Analytische Chemie. Weinheim: VCH Verlagsgesellschaft..
- Pack JW, Kim SH, Park SY, Lee Y-W, Kim YH. 2004. Ring opening polymerisation of L-lactide and preparation of its microsphere in supercritical fluids. *Macromol. Biosci.* 4: 340-345.
- Packhaeuser CB, Schnieders J, Oster CG, Kissel T. 2004. In situ forming parenteral drug delivery systems: an overview. *Eur. J. Pharm. Biopharm.* 58: 445-455.
- Paredes S, Tribout M. 1984. Enthalpies of micellization of quarternary tetradecyl- and cetyltrimethylammonium. *J. Phys. Chem.* 88: 1871-1875.
- Park TG. 1995. Degradation of poly(lactic-co-glycolic acid) microspheres: effect of copolymer composition. *Biomaterials* 16: 1123-1130.
- Park TG, Lu W, Crofts G. 1995. Importance of in vitro experimental conditions on protein release kinetics, stability and polymer degradation in protein encapsulated poly(D,L-lactic acid-co-glycolic acid) microspheres. *J. Controlled Release* 33: 211-222.
- Park TG, Lee HY, Nam YS. 1998. A new preparation method for protein loaded poly(D,L-lactic-co-glycolic acid) microspheres and protein release mechanism study. *J. Controlled Release* 55: 181-191.
- Péan J-M, Boury F, Venier-Julienne M-C, Menei P, Proust J-E, Benoit J-P. 1999. Why does PEG 400 co-encapsulation improve NGF stability and release from PLGA biodegradable microspheres. *Pharm. Res.* 16: 1294-1299.
- Perez C, Griebenow K. 2001. Improved activity and stability of lysozyme at the water/CH₂Cl₂ interface: enzyme unfolding and aggregation and its prevention by polyols. *J. Pharm. Pharmacol.* 53: 1217-1226.
- Peters T. 1985. Serum Albumin. *Adv. Protein Chem.* 37: 161-245.
- Peters JH, Ruhl S, Friedrich D. 1987. Veiled accessory cells deduced from monocytes. *Immunobiology* 176: 154-166.
- Peters JH, Ruppert J, Gieseler RK, Najjar HM, Xu H. 1991. Differentiation of human monocytes into CD14 negative accessory cells: Do dendritic cells derive from the monocytic lineage? *Pathobiology* 59: 122-126.

Poehling H-M, Neuhoff V. 1981. Visualization of proteins with a silver stain: A critical analysis. *Electrophoresis* 2: 141-147.

Poltorak A, He X, Smirnova I, Liu M-Y, van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, Beutler B. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: Mutations in TLR 4 gene. *Science* 282: 2085-2088.

Poole S, West SI, Fry JC. 1987. Effects of basic proteins on the denaturation and heat-gelation of acidic proteins. *Food Hydrocolloids* 1: 301-316.

Pritchard JG, Akintola DA. 1972. Complexation of polyvinyl alcohol with iodine. Analytical precision and mechanism. *Talanta* 19: 877-888.

Raffler G, Müller B. 1993. Verfahren zur Herstellung biologisch abbaubarer Polyester. Europäische Patentanmeldung, EP 0573024.

Rafler G, Jobmann M. 1994. Controlled release systems of biodegradable polymers. 3rd Communication: Degradation and release properties of poly(glycolide (50)-co-lactide (50)) with random and non-random monomer distribution. *Pharm. Ind.* 56: 753-755.

Rafler G, Jobmann M. 1997. Controlled release systems of biodegradable polymers. 5th Communication: Microparticle preparation with a salting-out process. *Pharm. Ind.* 59: 620-624.

Randolph DA, Fathman CG. 2006. CD4(+)CD25(+) regulatory T cells and their therapeutic potential. *Annu. Rev. Med.* 57: 381-402.

Rauen HM (ed). 1964. *Biochemisches Taschenbuch*. 2. Auflage in 2 Bänden. Berlin: Springer Verlag. p. 91-105, 2. Band.

Rege PR, Block LH. 1999. Chitosan processing: influence of process parameters during acidic and alkaline hydrolysis and effect of the processing sequence on the resultant chitosan's properties. *Carbohydr. Res.* 321: 235-245.

Reich G. 1998. Ultrasound-induced degradation of PLA and PLGA during microsphere processing: influence of formulation variables. *Eur. J. Pharm. Biopharm.* 45: 165-171.

Rescigno M, Granucci F, Ricciardi-Castagnoli P. 1999. Dendritic cells at the end of the millennium. *Immunol. Cell Biol.* 77: 404-410.

Rietschel ET, Cavaillon J-M. 2003. Richard Pfeiffer and Alexander Besredka: creators of the concept of endotoxin and anti-endotoxin. *Microbes Infect.* 5: 1407-1414.

Roberts EJ. 1985. A quantitative method for dextran analysis. *Int. Sugar Jnl.* 85: 10-13.

Roberts G. 1997. Determination of the degree of N-acetylation of chitin und chitosan. In: Muzzarelli RAA, Peter MG (eds). *Chitin Handbook*. Grottammare: Atec Edizioni. p. 127-132.

- Roldo M, Hornof M, Caliceti P, Bernkop-Schnürch A. 2004. Mucoadhesive thiolated chitosans as platforms for oral controlled drug delivery: synthesis and in vitro evaluation. *Eur. J. Pharm. Biopharm.* 57: 115-121.
- Romani N, Gruner S, Brang D, Kämpgen E, Lenz A, Trockenbacher B, Konwalinka G, Fritsch PO, Steinmann RM, Schuler G. 1994. Proliferation of dendritic cells progenitors in human blood. *J. Exp. Med.* 180: 83-92.
- Romani N, Holzmann S, Tripp CH, Koch F, Stoitzner P 2003. Langerhans cells - dendritic cells of the epidermis. *Apmis* 111: 725-740.
- Rosenblatt J, Kufe D, Avigan D. 2005. Dendritic cell fusion vaccines for cancer immunotherapy. *Expert Opin. Biol. Ther.* 5: 703-715.
- Rüegg M, Moor U, Blanc B. 1977. A calorimetric study on the thermal denaturation of whey proteins in simulated milk ultrafiltrate. *J. Dairy Res.* 44: 509-520.
- Ruiz JM, Benoit JP. 1991. In vivo peptide release from poly(D,L lactic acid-co-glycolic acid) copolymer 50/50 microspheres. *J. Controlled Release* 16: 177-186.
- Ruppert J, Schutt C, Ostermeier D, Peters JH. 1993. Down-regulation and release of CD14 on human monocytes by IL-4 depends on the presence of serum or GM-CSF. *Adv. Exp. Med. Biol.* 329: 281-286.
- Sabnis S, Block LH. 2000. Chitosan as an enabling excipient for drug delivery systems. I. Molecular modifications. *Int. J. Biol. Macromol.* 27: 181-186.
- Sah H, Smith MS, Chern RT. 1996. A novel method of preparing PLGA microcapsules utilizing methylethyl ketone. *Pharm. Res.* 13: 360-367.
- Sah H. 1997. Microencapsulation techniques using acetate as a dispersed solvent: effects of its extraction rate on the characteristics of PLGA microspheres. *J. Controlled Release* 47: 233-245.
- Sah H. 1999. Stabilization of proteins against methylene chloride/water interface-induced denaturation and aggregation. *J. Controlled Release* 58: 143-151.
- Sah H. 2000. Ethyl formate – alternative dispersed solvent useful in preparing PLGA microspheres. *Int. J. Pharm.* 195: 103-113.
- Sahlin S, Hed J, Rundquist I. 1983. Differentiation between attached and ingested immune complexes by a fluorescence quenching cytofluorometric assay. *J. Immunol. Meth.* 60: 115-124.
- Sahoo SK, Panyam J, Prabha S, Labhaestwar V. 2002. Residual polyvinyl alcohol associated with poly (D,L-lactide-co-glycolide) nanoparticles affects their physical properties and cellular uptake. *J. Controlled Release* 82: 105-114.
- Sallusto F, Lanzavecchia A. 1994. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by granulocyte/macrophage colony-stimulating factor

plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J. Exp. Med.* 179: 1109-1118.

Sallusto F, Cella M, Danieli C, Lanzavecchia A. 1995. Dendritic cells use macropinocytosis and the mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: downregulation by cytokines and bacterial products. *J. Exp. Med.* 182: 389-400.

Salthouse TN, Matlaga BF. 1976. Polyglactin 910 suture absorption and the role of cellular enzymes. *Surg. Gynecol. Obstet.* 142: 544-550.

Samanen JM. 1991. Physical biochemistry of peptide drugs: Structure, properties, and stabilities of peptide compared with proteins. In: Lee VHL (ed). *Peptide and protein drug delivery*. New York: Marcel Dekker. p. 137-166.

Sánchez A, Villamayor B, Guo Y, McIver J, Alonso MJ. 1999. Formulation strategies for the stabilisation of tetanus toxoid in poly(lactide-co-glycolide) microspheres. *Int J. Pharm.* 185: 255-266.

Sargent MG. 1987. Fiftyfold amplification of the Lowry protein assay. *Anal. Biochem.* 163: 476-481.

Sato T, Kanke M, Schroeder HG, DeLuca PP. 1988. Porous biodegradable microspheres for controlled drug delivery. I. Assessment of processing conditions and solvent removal techniques. *Pharm. Res.* 5: 21-30.

Schalper K. 2002. Herstellung hochdispenser Systeme mit kontinuierlichen Mikromischern. Dissertation. Freie Universität Berlin.

Schalper K, Harnisch S, Müller RH, Hildebrand GE. 2005. Preparation of microparticles by micromixers – Characterisation of o/w process and prediction of particle size. *Pharm. Res.* 22: 276-284.

Schenk R, Hessel V, Hofmann C, Kiss J, Löwe H, Ziogas A. 2004a. Numbering-up of micro devices: a first liquid-flow splitting unit. *Chem. Eng. J.* 101: 421-429.

Schenk R, Hessel V, Hofmann C, Kiss J, Löwe H, Schönfeld F, Ziogas A. 2004b. Numbering up von Mikroreaktoren: Ein neues Flüssigkeitsverteilsystem. *Chem. Ing. Tech.* 76: 584-597.

Schmut O, Faulborn J, Trummer G. 1999. Quantifizierung der Schädigung von Bindehaut und Hornhautzellkulturen durch UV-Licht mit Hilfe des CASY (Zellanalyse)-Systems. *Ophthalmologie* 96: 375-381.

Schnurr M, Galambos P, Scholz C, Dauer M, Krug A, Hartmann G, Eigler A, Endres S. 2002. Dendritische Zellen - Träger tumorgerichteter Immuntherapie. *Deutsches Ärzteblatt* 99: C1929-1934.

Scholes PD, Coombes AGA, Illum L, Davis SS, Watts JF, Ustariz C, Vert M, Davies MC. 1999. Detection and determination of surface levels of poloxamer and PVA surfactant on biodegradable nanospheres using SSIMS and XPS. *J. Controlled Release* 59: 261-278.

Schwendeman SP, Cardamone M, Klivanov A, Langer R, Brandon MR. 1996. Stability of proteins and their delivery from biodegradable polymer microspheres. In: Cohen S, Bernstein H (eds). *Microparticulate systems for the delivery of proteins and vaccines*. New York: Marcel Decker. p 1-49.

Shakesheff KM, Evora C, Soriano I, Langer R. 1997. The adsorption of poly(vinyl alcohol) to biodegradable microparticles studied by X-ray photoelectron spectroscopy (XPS). *J. Colloid Interface Sci.* 185: 538-547.

Shakweh M, Besnard M, Nicolas V, Fattal E. 2005. Poly(lactide-co-glycolide) particles of different physicochemical properties and their uptake by peyer's patches in mice. *Eur. J. Pharm. Biopharm.* 61: 1-13.

Shortman K, Liu Y-J. 2002. Mouse and human dendritic cell subtypes. *Nature Rev. Immunol.* 2: 151-161.

Siepmann J, Elkharraz K, Siepmann F, Klose D. 2005. How autocatalysis accelerates drug release from PLGA-based microparticles: A quantitative treatment. *Biomacromolecules* 6: 2312-2319.

Sieval AB, Thanou M, Kotze AF, Verhoef JC, Brussee J, Junginger HE. 1998. Preparation and NMR characterisation of highly substituted N-trimethyl chitosan chloride. *Carbohydr. Polym.* 36: 157-165.

Simard JR, Zunszain PA, Ha C-E, Yang JS, Bhagavan NV, Petitpas I, Curry S, Hamilton JA. 2005. Locating high-affinity fatty acid-binding sites on albumin by x-ray crystallography and NMR spectroscopy. *Proc. Natl. Acad. Sci. USA* 102: 17958-17963.

Singh M, Briones M, Ott G, O'Hagan D. 2000. Cationic microparticles: A potent delivery system for DNA vaccines. *Proc. Natl. Acad. Sci. USA* 97: 811-816.

Singh M, Ott G, Kazzaz J, Ugozzoli M, Briones M, Donnelly J, O'Hagan D. 2001. Cationic microparticles are an effective delivery system for immune stimulatory CpG DNA. *Pharm. Res.* 18: 1476-1479.

Smith DS, Al-Hakiem MHH, Landon J. 1981. A review of fluoroimmunoassay and immunofluorometric assay. *Ann. Clin. Biochem.* 18: 253-274.

Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goeke NM, Olson BJ, Klenk DC. 1985. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* 150: 76-85.

Smith SA, Pretorius WA. 2002. Spectrophotometric determination of pK_a values for fluorescein using activity coefficient corrections. *Water SA* 28: 395-402. Verfügbar unter <http://www.wrc.org.za>.

Solier P, Denuziere A, Viton C, Domaed A. 2001. Relation between the degree of acetylation and the electrostatic properties of chitin and chitosan. *Biomacromolecules* 2: 765-772.

Song L, Varma C, Verhoeven JW, Tanke HJ. 1996. Influence of the triplet excited state on the photobleaching kinetics of fluorescein in microscopy. *Biophys. J.* 70: 2959-2968.

- Song M, Ma H-M, Huang Y-X, Liang S-C. 1997. Determination of poly(vinyl alcohol) by adsorptive stripping voltammetry based on its copper(II) complex. *Anal. Chim. Acta* 338: 103-108.
- Sprang S, Standing T, Fletterick RJ, Stroud RM, Finer-Moore J, Xuong N-H, Hamlin R, Rutter WJ, Craik CS. 1987. The three-dimensional structure of Asn 102 mutant of trypsin: Role of Asp 102 in serin protease catalysis. *Science* 237: 905-909.
- Steinmann RM, Cohn ZA. 1973: Identification of a novel cell type in peripheral lymphoid organs of mice. I. Morphology, quantitation, tissue distribution. *J. Exp. Med.* 137: 1142-1162.
- Stokes GG. 1852. On the change of refrangibility of light. *Phil. Trans. R. Soc.* 142: 463-562. abgedruckt bei Lakowicz 1999.
- Storch WB. 1997. *Immunfluoreszenzfibel*. 2. Auflage. Berlin: Blackwell Wissenschaftsverlag.
- Stoschek CM. 1990. Quantitation of proteins. *Methods Enzymol.* 182: 50-68.
- Stridsberg KM, Ryner M, Albertsson A-M. 2002. Controlled ring-opening polymerisation: Polymers with designed macromolecular architectures. *Adv. Polym. Sci.* 157: 41-65.
- Stroud RM, Kay LM, Dickerson RE. 1972. The crystal and molecular structure of DIP-inhibited bovine trypsin at 2,7 Å resolution. *Cold Spring Harbor Symp. Quant. Biol.* 36: 125-140.
- Stuart LM, Ezekowitz RAB. 2005. Phagocytosis: Elegant Complexity. *Immunity* 22: 539-550.
- Sturesson C, Carlfors J. 2000. Incorporation of protein in PLG-microspheres with retention of bioactivity. *J. Controlled Release* 67: 171-178.
- Sugiura S, Nakajima M, Tong J, Nabetani H, Seki M. 2000. Preparation of monodispersed solid lipid microspheres using a microchannel emulsification procedure. *J. Coll. Interface Sci.* 227: 95-103.
- Tamada JA, Langer R. 1993. Erosion kinetics of hydrolytically degradable polymers. *Proc. Natl. Acad. Sci. USA* 90: 552-556.
- Tamulis A, Tamuliene J, Balevicius ML, Rinkevicius Z, Tamulis V. 2003. Quantum mechanical studies of the intensity in electronic spectra of fluorescein dianion and monoanion forms. *Struct. Chem.* 14: 643-648.
- Terreux R, Domard M, Viton C, Domard A. 2006. Interactions study between the copper II ion and constitutive elements of chitosan structure by DFT calculation. *Biomacromolecules* 7: 31-37.
- Thiele L, Rothen-Rutishauser B, Jilek S, Wunderli-Allenspach H, Merkle HP, Walter E. 2001. Evaluation of particle uptake in human blood monocyte-derived cells in vitro. *Does*

phagocytosis activity of dendritic cells measure up with macrophages? *J. Controlled Release* 76: 59-71.

Thies C. 1992. Formation of biodegradable drug-loaded microparticles by in-liquid drying processes. In: Donbrow M (ed). *Microcapsules and nanoparticles in medicine and pharmacy*. Boca Raton: CRC Press. p. 47-71.

Thumann P, Moc I, Humrich J, Berger TG, Schultz ES, Schuler G, Jenne L. 2003. Antigen loading of dendritic cells with whole tumor cell preparations. *J. Immunol. Methods* 277: 1-16.

Tian ZM, Wan MX, Wang B, Wang SP, Wu XM, Ruan YS. 2003. Effects of ultrasound on the structure and function of tumor necrosis factor- α . *Ultrasound Med. Biol.* 29: 1331-1339.

Tian ZM, Wan MX, Wang SP, Kang JQ. 2004. Effects of ultrasound and additives on the function and structure of trypsin. *Ultrason. Sonochem.* 11: 399-404.

Tong J, Nakajima M, Nabetani H, Kikuchi Y, Maruta Y. 2001. Production of oil-in-water microspheres using a stainless steel microchannel. *J. Coll. Interface Sci.* 237: 239-248.

Trakatelli M, Toungouz M, Blocklet D, Dodoo Y, Gordower L, Vereecken MLP, Sales F, Mortier L, Mazouz N, Lambermont M, Goldman S, Coulie P, Goldman M, Velu T. 2006. A new dendritic cell vaccine generated with interleukin-3 and interferon- β induces CD8⁺ T cell responses against NA17-A2 tumor peptide in melanoma patients. *Cancer Immunol. Immunother.* 55: 469-474.

Tsuji K, Lewis AR. 1978a. Dry-heat destruction of lipopolysaccharide: A mathematical approach process evaluation. *Appl. Environ. Microbiol.* 36: 715-719.

Tsuji K, Lewis AR. 1978b. Dry-heat destruction of lipopolysaccharide: Dry-heat destruction kinetics. *Appl. Environ. Microbiol.* 36: 710-714.

Uchida T, Hosaka S, Miura K. 1985. Direct measurement of phagolysosomal esterase activity. *Biochem. Biophys. Res. Comm.* 127: 584-589.

Verdijk RM, Mutis T, Esendam B, Kamp J, Melief CJ, Brand A, Goulmy E. 1999. Polyriboinosinic polyribocytidylic acid (poly(I:C)) induces stable maturation of functionally active human dendritic cells. *J. Immunol.* 163: 57-61.

van Voorhis WC, Hair LS, Steinman RM, Kaplan G. 1982. Human dendritic cells. Enrichment and characterization from peripheral blood. *J. Exp. Med.* 155: 1172-1187.

Waeckerle-Men Y, Scandella E, Uetz-von Allmen E, Ludewig B, Gillesen S, Merkle HP, Gander B, Groettrup M. 2004. Phenotype and functional analysis of human monocyte-derived dendritic cells loaded with biodegradable poly(lactide-co-glycolide) microspheres for immunotherapy. *J. Immunol. Methods* 287: 109-124.

Wackerle-Men Y, Groettrup M. 2005. PLGA microspheres for improved antigen delivery to dendritic cells as cellular vaccines. *Adv. Drug Del. Rev.* 57: 475-482.

Waeckerle-Men Y, Uetz-von Allmen E, Gander B, Scandella E, Schlosser E, Schmidtke G, Merkle HP, Groettrup M. 2006. Encapsulation of proteins and peptides into biodegradable poly(D,L-lactide-co-glycolide) microspheres prolongs and enhances antigen presentation by human dendritic cells. *Vaccine* 24: 1847-1857.

Walter B. 1888. Die Änderung des Fluoreszenzvermögens mit der Konzentration. *Z. Physik. Chem.* 2: 551.

Walter B. 1889. Über den Nachweis des Zerfalls von Molekulargruppen in Lösungen durch Fluoreszenz- und Absorptionserscheinungen. *Z. Physik. Chem.* 3: 234.

Wang W. 1999. Instability, stabilisation, and formulation of liquid protein pharmaceuticals. *Int. J. Pharm.* 185: 129-188.

Wang XH, Li DP, Wang WJ, Feng QL, Cui FZ, Xu YX, Song XH. 2003. Covalent immobilisation of chitosan and heparin on PLGA surface. *Int. J. Biol. Macromol.* 33: 95-100.

Waterborg JH, Matthews HR. 1994. The Lowry method for protein quantitation. In: Walker JM (ed). *Methods in molecular biology*, Vol. 32: Basic protein and peptide protocols. Totowa: Humana Press. p. 1-4.

van der Weert M, Hoehstetter J, Hennink WE, Crommelin DJA. 2000a. The effect of a water organic solvent interface on the structural stability of lysozyme. *J. Controlled Release* 68: 351-359.

van de Weert M, Hennink WE, Jiskoot W. 2000b. Protein instability in poly(lactic-co-glycolic acid) microparticles. *Pharm. Res.* 17: 1159-1167.

Wetzel R, Becker M, Behlke J, Billwitz H, Böhm S, Ebert B, Hamann H, Krumbiegel J, Lassmann G. 1980. Temperature behaviour of human serum albumin. *Eur. J. Biochem.* 104: 469-478.

Wichert B, Rohdewald P. 1990. A new method for the preparation of drug containing polylactic acid microparticles without using organic solvents. *J. Controlled Release* 14: 269-283.

Williams RT, Bridges JW. 1964. Fluorescence of solutions: A review. *J. Clin. Path.* 71: 371-394.

Wilson WD, Foster JF. 1971. Conformation-dependent limited proteolysis of bovine plasma albumin by an enzyme present in commercial albumin preparations. *Biochemistry* 10: 1772-1780.

Wischke C, Zimmermann J, Lorenzen DR, Borchert H-H. 2004. Impact of microparticle phagocytosis on dendritic cell maturation. *Immunobiol.* 209: 371.

Wischke C, Lorenzen DR, Zimmermann J, Borchert H-H. 2005. Physical stability of CTAB-PLGA microparticles for DNA vaccination. *Immunobiol.* 210: 484.

Wischke C, Borchert H-H. 2005. Fluorescent-labeled proteins used to characterise microparticulate vehicles for the uptake into dendritic cells. In: Annual Meeting of the German Pharmaceutical Society. ISBN 3-00-016844-3.

Wischke C, Borchert H-H. 2006a. Influence of the primary emulsification procedure on the characteristics of small protein loaded PLGA microparticles for antigen delivery. *J. Microencapsulation* 23: 435-448.

Wischke C, Borchert H-H. 2006b. FITC-BSA as a model protein drug: opportunities and drawbacks. *Die Pharmazie* in press.

Wischke C, Lorenzen D, Zimmermann J, Borchert H-H. 2006a. Preparation of protein loaded poly(D,L-lactide-co-glycolide) microparticles for the antigen delivery to dendritic cells using a static micromixer. *Eur. J. Pharm. Biopharm.* 62: 247-253.

Wischke C, Borchert H-H, Zimmermann J, Siebenbrodt I, Lorenzen DR. 2006b. Stable cationic microparticles for enhanced model antigen delivery to dendritic cells. *J. Controlled Release* in press.

Witschi C, Doelker E. 1998a. Influence of the microencapsulation method and peptide loading on poly(lactic acid) and poly(lactic-co-glycolic acid) degradation during in vitro testing. *J. Controlled Release* 51: 327-341.

Witschi C, Doelker E. 1998b. Peptide degradation during preparation and in vitro release testing of poly(L-lactic acid) and poly(D,L-lactic-co-glycolic acid) microparticles. *Int. J. Pharm.* 171: 1-18.

Xia D, Chan T, Xiang J. 2005. Dendritic cell/myeloma hybrid vaccine. *Methods Mol. Med.* 113: 225-233.

Xing DKL, Crane DT, Bolgiano B, Corbel MJ, Jones C, Sesardic D. 1996. Physicochemical and immunological studies on the stability of free and microsphere-encapsulated tetanus toxoid in vitro. *Vaccine* 14: 1205-1213.

Yang J, Cleland JL. 1997. Factors affecting the in vitro release of recombinant human interferon- γ (rhIFN- γ) from PLGA microspheres. *J. Pharm. Sci.* 86: 908-914.

Yamamoto H, Kuno Y, Sugimoto S, Takeuchi H, Kawashima Y. 2005. Surface-modified PLGA nanosphere with chitosan improved pulmonary delivery of calcitonin by mucoadhesion and opening of the intercellular tight junctions. *J. Controlled Release* 102: 373-381.

Yeh M-K, Davis SS, Coombes AGA. 1996. Improving protein delivery from microparticles using blends of poly(D,L-lactide-co-glycolide) and poly(ethylene oxide)-poly(propylene oxide) copolymers. *Pharm. Res.* 13: 1693-1698.

Yeh M-K. 2000. The stability of insulin in biodegradable microparticles based on blends of lactide polymers and polyethylene glycol. *J. Microencapsulation* 17: 743-756.

Yeo Y, Chen AU, Basaran OA, Park K. 2004. Solvent exchange method: A novel microencapsulation technique using dual microdispensers. *Pharm. Res.* 21: 1419-1427.

Yi H, Wu L-Q, Sumner JJ, Gillespie JB, Payne GF, Bentley WE. 2003. Chitosan scaffolds for biomolecular assembly: Coupling nucleic acid probes for detecting hybridization. *Biotechnol. Bioeng.* 83: 646-652.

Yu Z, Johnston KP, Williams III RO. 2006. Spray freezing into liquid versus spray-freeze drying: Influence of atomization on protein aggregation and biological activity. *Eur. J. Pharm. Sci.* 27: 9-18.

Zanker V, Peter W. 1958. Die prototropen Formen des Fluoresceins. *Chem. Ber.* 91: 572-580.

Zeng J, Hornsby PJ. 1999. Production of microspheres with surface amino groups from blends of poly(lactide-co-glycolide) and poly(ϵ -CBZ-L-lysine) and use for encapsulation. *Biotechnol. Prog.* 15: 763-767.

Zhou LJ, Tedder TF. 1996. CD14⁺ blood monocytes can differentiate into functionally mature CD83⁺ dendritic cells. *Proc. Natl. Acad. Sci. USA* 93: 2588-2592.

Zimmermann J, Peters JH, Wischke C, Borchert H-H, Lorenzen DR. 2003. Delivery of PLGA-microsphere encapsulated antigens to monocyte-derived dendritic cells prolongs their antigen presentation and promotes Th1-type cytokine production. *Immunobiol.* 208: 270.