

7 Literaturverzeichnis

- 1 Anonymous: Consensus development conference: diagnosis, prophylaxis and treatment of osteoporosis. Am J Med 1993; 94:646-650.
- 2 Abendschein W, Hyatt GW: Ultrasonic and selected physical properties of bone. Clin Orthop 1970; 69:294-301
- 3 Agren M, Karella A, Leahey D, Marks S, Baran D: Ultrasound attenuation of the calcaneus: a sensitive and specific discriminator of osteopenia in postmenopausal women. Calcif Tissue Int 1991; 48:240-4.
- 4 Ahmed AIH, Blake GM, Rymer JM, et al.: Screening for osteopenia and osteoporosis: do the accepted normal ranges lead to overdiagnosis? Osteoporosis Int 1997; 7:432-8.
- 5 Alenfeld FE, Wuster C, Funck C, et al.: Ultrasound measurements at the proximal phalanges in healthy women and patients with hip fractures. Osteoporosis Int 1998; 8(5): 393 – 8
- 6 Ashman RB, Cowin SC, van Buskirk WC, Rice JC: Acontinuous wave technique for the measurement of elastic properties of cortical bone. J Biomechanics 1984; 17:349-361.
- 7 Ashman RB, Rho JY, Turner CH: Anatomical variation of orthotropic elastic moduli of the proximal human tibia. J Biomechan 1989; 22:859 – 900.
- 8 Baran DT, Kelley AM, Karella A, et al.: Ultrasound attenuation of the os calcis in women with osteoporosis and hip fractures. Calcif Tissue Int 1988; 43:138-42.
- 9 Baran DT, McCarthy CK, Leahey D, Lew R: Broadband ultrasound attenuation of the calcaneus predicts lumbar and femoral density in Caucasian women: a preliminary study. Osteoporosis Int 1991; 1:110-3.
- 10 Bauer DC, Browner WS, Cauley JA, Orwoll ES, Scott JC, Black DM, Tao JL, Cummings SR: Factors associated with appendicular bone mass in older women. The study of Osteoporotic Fractures Research Group. Ann Intern Med 1993 may1; 118(9):741-2.
- 11 Bauer DC, Glüer CC, Cauley JA, Vogt TM, Ensrud KE, Genant HK, Black DM: Broadband ultrasound attenuation predicts fractures strongly and independently of densitometry in older women. A prospective study. Study of osteoporotic Fractures Research Group. Arch Intern Med 1997; 157(6):629-34.

- 12 Bauer DC, Gluer CC, Pressman AR, Vogt TM, Ensrud KE, Cauley JA, et al.: BUA and the risk of fracture: a prospective study. *J Bone Miner Res (suppl)* 1995; 10:175.
- 13 Bonfield W, Tully AE: Ultrasonic analysis of the Young's modulus of cortical bone. *J Biomed Eng* 1982; 4:23 – 27.
- 14 Bouxsein ML, Courtney AC, Hayes WC: Ultrasound and densitometry of the calcaneus correlate with the failure loads of cadaveric femurs. *Calcif Tissue Int* 1995; 56:99 - 103.
- 15 Bouxsein ML, Radloff SE: Quantitative ultrasound of the calcaneus reflects the mechanical properties of calcaneal trabecular bone. *J Bone Miner Res* 1977; 12: 839 - 846.
- 16 Cann CE: Low dose CT. *Radiology* 1987; 164: 419 – 423.
- 17 Cepollaro C, Zachei F, Borracelli D; et al.: Precision of new ultrasound bone densitometers: correlation with absorptiometry methods. In: Current research in osteoporosis and bone mineral measurement I London: British Institute of Radiology 1992.
- 18 Cheng S, Njeh CF, Fan B, Cheng X, Hans D, Wang L, Fuerst T, Genant HK: Influence of region of interest and bone size on calcaneal BMD: implications for the accuracy of quantitative ultrasound assessments at the calcaneus. *Br J Radiol* 2002; 75: 59 - 68.
- 19 Ciarelli MJ, Goldstein SA, Kuhn JL, Cody DD, Brown MB: Evaluation of orthogonal mechanical properties and density of human trabecular bone from major metaphyseal regions with materials testing and computed tomography. *J Orthop Res* 1991; 9: 674 - 82.
- 20 Consensus Development Diagnosis: Diagnosis, prophylaxis and treatment of osteoporosis. Hong-Kong; april 1993. Osteologie Sonderdruck
- 21 Cummings SR, Kelsey JL, Nevitt MC, O'Dowd KJ: Epidemiology of osteoporosis and osteoporotic fractures. *Epidemiol Rev* 1985; 7: 178 - 208.
- 22 Cummings SR, Nevitt MC, Browner WS, et al.: Risk factors for hip fracture in white women. *New England Journal of Medicine* 1995; 332: 767 - 73.
- 23 Damilakis JE, Dretakis E, Gourtsoyiannis NC: Ultrasound attenuation of the calcaneus in the female population: normative data. *Calcif Tissue Int* 1992; 155:825-8
- 24 Dargent-Molina P, Favier F, Grandjean H, Baudois C, Schott AM, Hausherr E, Meunier PJ, Breart G: Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet* 1996; 348:145-49.

- 25 Dargent-Molina P, Piault S, Breart G: A comparison of different screening strategies to identify elderly women at high risk of hip fracture: results from the EPIDOS prospective study. *Osteoporosis Int* 2003;14(12):969-77.
- 26 Duquette J, Lin J, Hoffmann A, Houde J, Ahmadi S, Baran D: Correlation among bone mineral density, broadband ultrasound attenuation, mechanical identification testing, and bone orientation in bovine femoral neck samples. *Calcif Tissue Int* 1997; 60:181-6.
- 27 Eastell R: Bone density of the calcaneus measured by ultrasound and DXA. In: Private symposium on ultrasound for bone measurement, Florence april 1992.
- 28 Ekman A, Michaelson K, Petren-Mallmin M, Ljunghall S, Mallmin H: DXA of the hip and heel ultrasound but not densitometry of the fingers can discriminate female hip fracture patients from controls: A comparison between four different methods. *Osteoporosis Int* 2000; 12:185-91.
- 29 Evans JA, Tavakoli MB: Ultrasonic attenuation and velocity in bone. *Phys Med Biol* 1990; 35:1387 – 1396.
- 30 Faulkner KG, Roberts LA, McClung MR: Discrepancies in normative data between Lunar and Hologic DXA systems. *Osteoporosis Int* 1996; 6:432-6.
- 31 Felsenberg D, Fobbe F, Wolf KJ: Ultraschallmessungen zur Bestimmung der Knochenmineralisation. *Jahrbuch der Radiologie* 1993
- 32 Felsenberg D, Kalender W, Rüegsegger P: Osteodensitometrische Untersuchungsverfahren. *Osteologie* 1993; Band 2. Sonderdruck
- 33 Felsenberg D, Kalender WA, Trinkwalter W et al: CT-Untersuchungen mit reduzierter Strahlendosis. *Fortschr Röntgenstr* 1990; 153: 516 - 21
- 34 Felson DT, Kiel DP, Anderson JJ, Kannel WB: Alcohol consumption and hip fractures: the Framingham study. *Am J Epidemiol* 1988; 128:1102-10.
- 35 Fortinsky RH, Iannuzzi-Sucich M, Baker DI, Gottschalk M, King MB, Brown CJ, Tinetti ME: Fall-risk assessment and management in clinical practice: Views from healthcare providers. *J Am Geriatr Soc* 2004; 52:1522-26.
- 36 Frost ML, Blake GM, Fogelman I: Can the WHO criteria for diagnosing osteoporosis be applied to calcaneal quantitativ ultrasound? *Osteoporosis Int* 2000; 11:321-30.

- 37 Genant HK, Grampp S, Glüer CC, Faulkner KG, Jergas M, Engelke K, et al.: Universal standardization for dual x-ray absorptiometry: patient and phantom cross-calibration results. *J Bone Miner Res* 1994; 9:1503-14.
- 38 Gerdhem P, Magnusson H, Karlsson MK, Kaesson K: Ultrasound of the phalanges is not related to a previous fracture. *J Clin Densitom* 2002; 5:159-66:
- 39 Glüer CC, Blake G, Lu Y, Blunt BA, Jergas M, Genant HK: Accurate Assessment of precision errors: How to measure the reproducibility of bone densitometry techniques. *Osteoporosis Int* 1995; 5:262-70.
- 40 Glüer CC, Cummings SR, Pressmann A: Prediction of hip fractures from pelvic radiographs: the study of osteoporotic fractures. *J Bone Miner Res* 1994; 9:671-7
- 41 Glüer CC, Hans D: How to use ultrasound for risk assessment: a need for defining strategies (editorial). *Osteoporosis Int* 1999; 9(3):193 – 5.
- 42 Glüer CC, Wu CY, Jergas M, Goldstein SA, Genant HK: Three quantitative ultrasound parameters reflect bone strukture. *Calcif Tissue Int* 1994; 55:46-52.
- 43 Glüer CC, Wu CY, Engelke K, et al.: Assessment of bone structure by quantitativ ultrasound. *J Bone Miner Res* 1993; 8:357.
- 44 Glüer CC, Wu CY, Genant HK: Broadband ultrasound attenuation signals depend on trabecular orientation: an in-vitro study. *Osteoporosis Int* 1993; 3:185-91.
- 45 Glüer CC: Quantitative ultrasound techniques for the assessment of osteoporosis: expert agreement on current status. *J Bone Miner Res* 1997; 12:1280-8.
- 46 Goldstein SA, Goulet R, McCubbrey D: Measurement and significance of three-dimensional architecture in the mechanical integrety of trabecular bone. *Calcif Tissue Int* 1993; 53:127-33.
- 47 Graafmans WC, Bouter LM, Lips P: The influence of physical activity and fractures on ultrasound parameters in elderly people. *Osteoporosis Int* 1998; 8(5):449-54.
- 48 Grampp S, Genant HK, Mathur A, Lang P, Jergas M, Takada M, et al.: Comparisons on noninvasive bone mineral measurements in assessing age-related loss fracture discrimination, and diagnostic classification. *J Bone Miner Res* 1997; 12:697-711.

- 49 Grimm MJ, Chung HW, Wehrli FW, Williams JL: Dependence of ultrasound attenuation on plate separation in trabekular bone. In: Proceedings of 40th annual meeting, Orthopaedic research Society, New Orleans, Louisiana, USA; february 1994:442.
- 50 Hans D, Dargent-Molina P, Schott AM, Sebert JL, Cormier C, Kotzki PO, Delmas PD, Pouilles JM, Breart G, Meunier PJ: Ultrasonographic heel measurements to predict hip fractures in elderly women: the EPIDOS prospective study. Lancet 1996; 348(9026):511-4.
- 51 Hans d, Hartl F, Krieg MA: Device-specific weighted T-score for two quantitative ultrasounds: operational propositions for the management of osteoporosis for 65 years and older women in Switzerland. Osteoporosis Int 2003;14 (3):251 - 8.
- 52 Hans D, Schott AM, Arlot ME, Sorrey E, Delmas PD, Meunier PJ: Influence of anthropometric parameters on ultrasound measurements of os calcis. Osteoporosis Int 1995; 5(5):371-6.
- 53 Hans D, Schott AM, Duboeuf F, Durosier C, Meunier PJ: Does follow-up duration influence the ultrasound and Dxa prediction of hip fracture? The EPIDOS prospective study. Bone 2004; 35:357-63.
- 54 Hans D, Schott AM, Meunier PJ: Ultrasonic assessment of bone: a review. EJM 1993; 2: 157 – 163.
- 55 Hayes WC, Piazza SJ, Zysset PK: Biomechanics of fracture risk prediction of the hip and spine by quantitative computer tomography. Radiol Clin North Am 1991; Am 29:1-18.
- 56 Hemenway D, Colditz GA, Willett WC, Stampfer MJ, Speizer ME: Fractures and lifestyle: effect of cigarette smoking, alcohol intake, and related weight on the risk of hip and forearm fractures in middle-aged women. Am J Public Health 1988; 78:1554-8.
- 57 Hernandez-Avila M, Stampfer MJ, Ravnikar VA, et al.: Caffeine and other predictors of bone density among pre- and perimenopausal women. Epidemiology 1993; 4:128-34.
- 58 Hologic Clinical user' guide Sahara
- 59 Ingle BM, Sherwood KE, Eastell R: Comparison of two methods for measuring ultrasound properties of the heel in postmenopausal women. Osteoporosis Int 2001; 12:500-5.
- 60 Jergas M, Glüer CC: Assessment of fracture risk by bone density measurements. Semin Nucl Med 1997; 27(3):261-75.
- 61 KalenderWA: Effektivedose values in bone mineral measurements by photon absorptiometryand computed tomography. Osteoporosis Int 1992; 2: 82 - 87.

- 62 Kang C, Speller R: Comparison of ultrasound and dual energy X-ray absorptiometry measurements in the calcaneus. *Br J Radiol* 1998; 71(848):861-7.
- 63 Kanis JA, McCloskey EV: Epidemiology of vertebral osteoporosis. *Bone* 1992; 13:1-10.
- 64 Kanis JA: Diagnosis of osteoporosis and assessment of fracture risk. *Lancet* 2002; 359:1929-36.
- 65 Kiel DP, Felson DT, Hannan MT, Anderson JJ, Wilson PWF: Caffeine and the risk of hip fracture: the Framingham study. *Am J Epidemiol* 1990; 132:675-84.
- 66 Kinsler LE, Frey AR, Coppens AB, Sanders JV: Fundamentals of acoustics. New York, NY: Wiley 1982.
- 67 Kotz S, Johnson NL: Encyclopedia of statistical sciences. Vol.8, New York: Wiley 1982
- 68 Krieg MA, Thiebaud D, Burckhardt P: Quantitative ultrasound of bone in institutionalized elderly women: a cross- sectional and longitudinal study. *Osteoporosis Int* 1996; 6(3):189-95.
- 69 Kung AWC, Tang GWK, Luk KDK, Chu LW: Evaluation of a new calcaneal quantitative ultrasound system and determination of normative ultrasound values in southern Chinese women. *Osteoporosis Int* 1999; 9:312-7.
- 70 Lang SM, Moyle DD, Berg EW, Detorie N, Gilpin AT, Pappan Jr NJ, Reynolds JC, Tkacik M, Waldron RL: Correlation of mechanical properties of vertebral trabecular bone with equivalent mineral density as measured by computed tomography. *J Bone Joint Surg* 1988; 70-A:1531-8.
- 71 Langton CM, Njeh CF, Hodgkinson R, Currey JD: Prediction of the human calcaneus by broadband ultrasound attenuation. *Bone* 1996; 18:495-503.
- 72 Langton CM, Palmer SB, Porter RW: The measurements of broadband ultrasonic attenuation in cancellous bone. *Engineering in Medicine* 1984; 13: 89 – 91.
- 73 Laugier P, Giat P, Berger G: Broadband ultrasound attenuation imaging: a new imaging technique of the os calcis. *Calcif Tissue Int* 1994; 54:83-6.
- 74 Laugier P, Giat P, Berger G: New ultrasonic methods of quantitative assessment of bone status. *Eur J Ultrasound* 1994; 1: 23 – 38.
- 75 Lees B, Stevenson J: Precision and sensitivity of a new ultrasound bone densitometer. In: Current research in osteoporosis and bone mineral measurement I London: British Institute of Radiology. 1992

- 76 Lees B, Stevenson JC: Preliminary evaluation of new ultrasound bone densitometer. *Calcif Tissue Int* 1993; 53:149-52.
- 77 Lewis MK, Blake GM, Fogelman I: Patient dose in Dual X-ray absorptiometry. *Osteoporosis Int* 1994; 4: 11 – 15.
- 78 Lotz JC, Gerhart TN, Hayes WC: Mechanical properties of trabecular bone from the proximal femur: a quantitative CT study. *J Comput Assist Tomogr* 1990; 14:107-14.
- 79 Lunar GmbH: ACHILLES+ Ultraschall-Densitometer. Bedienungsanleitung version 12/95
- 80 Madison, WI: Lunar Corporation: Lunar News. june 1994
- 81 Mazess RB, Wiener SA, Hanson JA, Bonnick SL: Ultrasound measurements of the os calcis. *Osteoporosis Int* 1991; 1: 194.
- 82 McCloskey EV, Murray SA, Miller C, et al.: Broadband ultrasound attenuation in the os calcis: relationship to bone mineral at other skeletal sites. *Clin Sci* 1990; 78:227-33.
- 83 Melton LJI: Epidemiology of hip fractures: implications of the exponential increase with age. *Bone* 1996;18:121S-5S.
- 84 Mosekilde L, Bentzen SM, Roft G, Jørgensen J: The predictive value of quantitative computed tomography for vertebral body compressive strength and as density. *Bone* 1989; 10:465-70.
- 85 Nicholson PHF, Haddaway MJ, Davie MWJ: The dependence of ultrasonic properties on orientation in human vertebral bone. *Phys Med Biol* 1994; 39:1013-1024.
- 86 Njeh CF, Boivin CM, Langton CM: The role of ultrasound in the assessment of osteoporosis: a review. *Osteoporosis Int* 1997; 7:7-22.
- 87 Njeh CF, Hans D, Li J, Fan B, Fuerst T, He YQ, et al.: Comparison of six calcaneal quantitative ultrasound devices:precision and hip fracture discrimination. *Osteoporosis Int* 2000; 11:1051-62.
- 88 Patel R, Blake GM, Jefferies A, et al.: A comparison of a peripheral DXA system with conventional densitometry of the spine and femur. *Clin Densitometry* 1998; 1:235-44.
- 89 Pfeifer M, Pollaehne W, Minne HW: Ultrasound analyses of the calcaneus predict relative risk of the presence of at least one vertebral fracture and reflect different physical qualities of bone in different regions of the skeleton. *Horm Metab Res* 1997; 29(2):76-9.
- 90 Pluijm SMF, Graafmans WC, Bouter LM, Lips P: Ultrasound measurements for prediction of osteoporotic fractures in elderly people. *Osteoporosis Int* 1999;9(6):550-6.

- 91 Pocock NA, Babichev A, Culton N, et al.: Temperatur dependency of quantitative ultrasound. *Osteoporosis Int* 2000; 11:316-20.
- 92 Pocock NA, Sambrook PN, Nguyen T, Kelly P, Freund J, Eisman J: Assessment of spinal and femoral bone density by dual x-ray absorptiometry: comparison of Lunar and Hologic instruments. *J Bone Miner Res* 1992; 7:1081-4.
- 93 Porter RW, Miller CG, Grainger D, Palmer SB: Prediction of hip fracture in elderly women: a prospective study. *BMJ* 1990; 301:638-41.
- 94 Ramalingham T, Lees B, Blake GM, Stevenson J, Miller CG, Fogelman I: Ultrasonic studies of the calcaneus: a comparison of measurement precision in three commercial scanners; Ringe EFJ, editors. *Current research in osteoporosis and bone mineral measurements*: London: British Institute of Radiology 1992.
- 95 Reeve J, Lunt M: Assessing the rate of bone loss:a technique revisited. *J Bone Miner Res* 1999; 14:1990.
- 96 Richardson RB: Past and revised risk for cancer induced by irradiation and their influence on dose limits. *Br J Radiol* 1990; 63:235-245
- 97 Riggs BL, Hodgson SF; O'Fallon WM; Chao EYS, Wahner HW, Muhs JM, Cedel SL, Melton LJ: Effect of fluoride treatment on the fracture rate in postmenopausal women with osteoporosis. *N Engl J Med* 1990; 322:802-9.
- 98 Schott AM, Hans D, Sorrey-Rendu E, Delmas PD, Meunier PJ: Ultrasound measurements on os calcis: precision and age- related changes in normal female population. *Osteoporosis Int* 1993; 3:249-54.
- 99 Schott AM, Weill-Engerer S, Hans D, et al.: Ultrasound discriminates patients with hip fracture equally well as dual energy x-ray absorptiometry and indepently of bone mineral density. *J Bone Miner Res* 1995; 10:243-9
- 100 Sosa M, Saavedra P, Munos-Torres M: Quantitative ultrasound calcaneus measurements: Normative data and precision in the spanish population. *Osteoporosis Int* 2002; 13:487-92.
- 101 Steiger P, Cummings SR, Black DM, et al.: Age-related degrements in bone mineral density in women over 65. *J Bone Miner Res* 1992; 7:697-711.
- 102 Strelitzki R, Clarke AJ, Truscott JG, Evans JA: Ultrasonic measurements: an evaluation of three heel bone scanners compared with a bench-top system. *Osteoporosis Int* 1996; 6(6):471-9.
- 103 Strelitzki R, Evans JA: Diffraction and interface losses in broadband ultrasound attenuation measurementsof the calcaneum. *Physiol Meas* 1998; 19(2):197-204.

- 104 Tavakoli MB, Evans JA: Dependence of velocity and attenuation of ultrasound in bone on the mineral content. *Phys Med Biol* 1991; 36(11): 1529-37.
- 105 Thompson PW, Taylor J, Oliver R, Fisher A: QUS of the heel predicts wrist and osteoporosis related fractures in women age 45-75 years. *J Clin Densitom* 1998; 1(3):219-26.
- 106 Trempe J, Genske T, Wiener S, Mazess RB: Ultrasound measurements of the os calcis. Current research in osteoporosis and bone mineral measurements: London: British Institute of Radiology 1992
- 107 Turner CH, Eich M: Ultrasonic velocity as a predictor of strength in bovine cancellous bone. *Calcif Tissue Int* 1991; 49:116-9.
- 108 Van Daele PLA, Burger H, Algra D, Hofman A, Grobbee DE, Birkenhäger JC, Pols HAP: Age-associated changes in ultrasound measurements of the calcaneus in men and women: The Rotterdam Study. *J Bone Miner Res* 1994; 9(11):1751-7.
- 109 Vogel JM, Wasnich RD, Ross PD: The clinical relevance of calcaneal bone mineral measurements: a review. *Bone Miner* 1988; 5:35-38.
- 110 Waud CE, Lew R, Baran DT: The relationship between ultrasound and densitometric measurements of bone mass at the calcaneus in women. *Calcif Tissue Int* 1992; 51:415-8.
- 111 Wuster C: Knochendichthymessung mittels Ultraschall – reif für die Praxis? *Zentralbl. Gynäkol* 1998;120 (6):269-74.
- 112 Yamada M, Ito M, Hayashi K, et al.: Dual energy x-ray absorptiometry of the calcaneus: comparison with other techniques to assess bone density and value in predicting risk of spine fracture. *AJR* 1994; 163:1435-40.
- 113 Yamazaki K, Kushida K, Ohmura A, Sano M, Inoue T: Ultrasound bone densitometry of os calcis in Japanese women. *Osteoporosis Int* 1994; 4:220-225.