

## 8 Literatur

Andre P, Nannizzi-Alaimo L, Prasad SK, Phillips DR. Platelet-derived CD40L: the switch-hitting player of cardiovascular disease. *Circulation*. 2002 Aug;106(8):896-9.

Apple FS, Wu AH, Mair J, Ravkilde J, Panteghini M, Tate J, Pagani F, Christenson RH, Mockel M, Danne O, Jaffe AS. Future biomarkers for detection of ischemia and risk stratification in acute coronary syndrome. *Clin Chem*. 2005 May;51(5):810-24.

Baldus S, Heeschen C, Meinertz T, Zeiher AM, Eiserich JP, Munzel T, Simoons ML, Hamm CW; CAPTURE Investigators. Myeloperoxidase serum levels predict risk in patients with acute coronary syndromes. *Circulation*. 2003 Sep;108(12):1440-5.

Barger SW, Wolchok SR, Van Eldik LJ. Disulfide-linked S100 beta dimers and signal transduction. *Biochim Biophys Acta*. 1992 Nov;1160(1):105-12.

Baudier J, Gerard D. Ions binding to S100 proteins: Conformational studies and calcium-induced conformational changes in S100 alpha alpha protein: the effect of acidic pH and calcium incubation on subunit exchange in S100a (alpha beta) protein. *J Biol Chem*. 1986 Jun;261(18):8204-12.

de Beer FC, Hind CR, Fox KM. Measurement of serum C-reactive protein concentration in myocardial ischaemia and infarction. *Br Heart J*. 1982 Mar;47(3):239-43.

Biasucci LM, Liuzzo G, Fantuzzi G, Caligiuri G, Rebuffi AG, Ginnetti F, Dinarello CA, Maseri A. Increasing levels of interleukin (IL)-1Ra and IL-6 during the first 2 days of hospitalization in unstable angina are associated with increased risk of in-hospital coronary events. *Circulation*. 1999 Apr;99(16):2079-84.

Blankenberg S, Rupprecht HJ, Poirier O, Bickel C, Smieja M, Hafner G, Meyer J, Cambien F, Tiret L, AtheroGene Investigators. Plasma concentrations and genetic variation of matrix metalloproteinase 9 and prognosis of patients with cardiovascular disease. *Circulation*. 2003 Apr;107(12):1579-85.

Bouma G, Lam-Tse WK, Wierenga-Wolf AF, Drexhage HA, Versnel MA. Increased serum levels of MRP-8/14 in type 1 diabetes induce an increased expression of CD11b and an enhanced adhesion of circulating monocytes to fibronectin. *Diabetes*. 2004 Aug;53(8):1979-86.

Borderie D, Allanore Y, Meune C, Devaux JY, Ekindjian OG, Kahan A. High ischemia-modified albumin concentration reflects oxidative stress but not myocardial involvement in systemic sclerosis. *Clin Chem*. 2004 Nov;50(11):2190-3.

Brown DL, Desai KK, Vakili BA, Nouneh C, Lee HM, Golub LM. Clinical and biochemical results of the metalloproteinase inhibition with subantimicrobial doses of doxycycline to prevent acute coronary syndromes (MIDAS) pilot trial. *Arterioscler Thromb Vasc Biol*. 2004 Apr;24(4):733-8.

Buffon A, Biasucci LM, Liuzzo G, D'Onofrio G, Crea F, Maseri A. Widespread coronary inflammation in unstable angina. *N Engl J Med*. 2002 Jul;347(1):5-12.

Burke AP, Kolodgie FD, Zieske A, Fowler DR, Weber DK, Varghese PJ, Farb A, Virmani R. Morphologic findings of coronary atherosclerotic plaques in diabetics: a postmortem study. *Arterioscler Thromb Vasc Biol.* 2004 Jul;24(7):1266-71.

Burkhardt K, Radespiel-Troger M, Rupprecht HD, Goppelt-Struebe M, Riess R, Renders L, Hauser IA, Kunzendorf U. An increase in myeloid-related protein serum levels precedes acute renal allograft rejection. *J Am Soc Nephrol.* 2001 Sep;12(9):1947-57.

Cipollone F, Marini M, Fazia M, Pini B, Iezzi A, Reale M, Paloscia L, Materazzo G, D'Annunzio E, Conti P, Chiarelli F, Cuccurullo F, Mezzetti A. Elevated circulating levels of monocyte chemoattractant protein-1 in patients with restenosis after coronary angioplasty. *Arterioscler Thromb Vasc Biol.* 2001 Mar;21(3):327-34.

Cornish CJ, Devery JM, Poronnik P, Lackmann M, Cook DI, Geczy CL. S100 protein CP-10 stimulates myeloid cell chemotaxis without activation. *J Cell Physiol.* 1996 Feb;166(2):427-37.

DeWood MA, Spores J, Notske R, Mouser LT, Burroughs R, Golden MS, Lang HT. Prevalence of total coronary occlusion during the early hours of transmural myocardial infarction. *N Engl J Med.* 1980 Oct;303(16):897-902.

Donato R. Intracellular and extracellular roles of S100 proteins. *Microsc Res Tech.* 2003 Apr 15;60(6):540-51.

Eggers KM, Oldgren J, Nordenskjold A, Lindahl B. Diagnostic value of serial measurement of cardiac markers in patients with chest pain: limited value of adding myoglobin to troponin I for exclusion of myocardial infarction. *Am Heart J* 2004;148: 574-581.

Eue I, Pietz B, Storck J, Klempt M, Sorg C. Transendothelial migration of 27E10+ human monocytes. *Int Immunol.* 2000 Nov;12(11):1593-604.

Eue I, Langer C, Eckardstein A, Sorg C. Myeloid related protein (MRP) 14 expressing monocytes infiltrate atherosclerotic lesions of ApoE null mice. *Atherosclerosis.* 2000 Aug;151(2):593-7.

Every NR, Larson EB, Litwin PE, Maynard C, Fihn SD, Eisenberg MS, Hallstrom AP, Martin JS, Weaver WD. The association between on-site cardiac catheterization facilities and the use of coronary angiography after acute myocardial infarction. Myocardial Infarction Triage and Intervention Project Investigators. *N Engl J Med.* 1993 Aug;329(8):546-51.

Frosch M, Strey A, Vogl T, Wulffraat NM, Kuis W, Sunderkötter C, Harms E, Sorg C, Roth J. Myeloid-related proteins 8 and 14 are specifically secreted during interaction of phagocytes and activated endothelium and are useful markers for monitoring disease activity in pauciarticular-onset juvenile rheumatoid arthritis. *Arthritis Rheum.* 2000 Mar;43(3):628-37.

Furukawa Y, Matsumori A, Ohashi N, Shioi T, Ono K, Harada A, Matsushima K, Sasayama S. Anti-monocyte chemoattractant protein-1/monocyte chemotactic and activating factor antibody inhibits neointimal hyperplasia in injured rat carotid arteries. *Circ Res.* 1999 Feb;84(3):306-14.

Garlichs CD, Eskafi S, Raaz D. Patients with acute coronary syndromes express enhanced CD40 ligand/CD154 on platelets. *Heart.* 2001 Dec;86(6):649-55.

Gibler WB, Hoekstra JW, Weaver WD. A randomized trial of the effects of early cardiac serum marker availability on reperfusion therapy in patients with acute myocardial infarction: the serial markers, acute myocardial infarction and rapid treatment trial (SMARTT). *J Am Coll Cardiol* 2000;36:1500-1506.

Heeschen C, Dimmeler S, Fichtlscherer S. Prognostic value of placental growth factor in patients with acute chest pain. *JAMA* 2004;291:435-441.

Heeschen C, Dimmeler S, Hamm CW. Soluble CD40 ligand in acute coronary syndromes. *N Engl J Med* 2003;348:1104-1111.

Hessian PA, Fisher L. The heterodimeric complex of MRP-8 (S100A8) and MRP-14 (S100A9). Antibody recognition, epitope definition and the implications for structure. *Eur J Biochem.* 2001 Jan;268(2):353-63.

Hessian PA, Edgeworth J, Hogg N. MRP-8 and MRP-14, two abundant Ca(2+)-binding proteins of neutrophils and monocytes. *J Leukoc Biol.* 1993 Feb;53(2):197-204.

Hofmann MA, Drury S, Hudson BI, Gleason MR, Qu W, Lu Y, Lalla E, Chitnis S, Monteiro J, Stickland MH, Bucciarelli LG, Moser B, Moxley G, Itescu S, Grant PJ, Gregersen PK, Stern DM, Schmidt AM. RAGE and arthritis: the G82S polymorphism amplifies the inflammatory response. *Genes Immun.* 2002 May;3(3):123-35.

Hollander JE, Muttreja MR, Dalesandro MR, Shofer FS. Risk stratification of emergency department patients with acute coronary syndromes using P-selectin. *J Am Coll Cardiol.* 1999 Jul;34(1):95-105.

Hunter MJ, Chazin WJ. High level expression and dimer characterization of the S100 EF-hand proteins, migration inhibitory factor-related proteins 8 and 14. *J Biol Chem.* 1998 May 15;273(20):12427-35.

Jaresova M, Striz I, Cermakova J, Lacha J, Sedlacek J, Mudra K, Hana I, Vitko S. Serum procalcitonin concentrations in transplant patients with acute rejection and bacterial infections. *Immunol Lett.* 1999 Sep;69(3):355-8.

Kai H, Ikeda H, Yasukawa H. Peripheral blood levels of matrix metalloproteinases-2 and -9 are elevated in patients with acute coronary syndromes. *J Am Coll Cardiol.* 1998 Aug;32(2):368-72.

Katus HA, Remppis A, Looser S, Hallermayer K, Scheffold T, Kübler W. Enzyme Linked Immuno Assay of Cardiac Troponin T for the Detection of an acute Myocardial Infarction in Patients. *J.Mol.Cell Cardiol* 1989;21:1349–53.

Kerkhoff C, Eue I, Sorg C. The regulatory role of MRP8 (S100A8) and MRP14 (S100A9) in the transendothelial migration of human leukocytes. *Pathobiology*. 1999;67(5-6):230-2.

Kerkhoff C, Klempt M, Kaever V, Sorg C. The two calcium-binding proteins, S100A8 and S100A9, are involved in the metabolism of arachidonic acid in human neutrophils. *J Biol Chem*. 1999 Nov;274(46):32672-9.

Kerkhoff C, Klempt M, Sorg C. Novel insights into structure and function of MRP8 (S100A8) and MRP14 (S100A9). *Biochim Biophys Acta*. 1998 Dec;1448(2):200-11.

Kislinger T, Tanji N, Wendt T, Qu W, Lu Y, Ferran LJ Jr, Taguchi A, Olson K, Bucciarelli L, Goova M, Hofmann MA, Cataldegirmen G, D'Agati V, Pischetsrieder M, Stern DM, Schmidt AM. Receptor for advanced glycation end products mediates inflammation and enhanced expression of tissue factor in vasculature of diabetic apolipoprotein E-null mice. *Arterioscler Thromb Vasc Biol*. 2001 Jun;21(6):905-10.

Klempt M, Melkonyan H, Nacken W, Wiesmann D, Holtkemper U, Sorg C. The heterodimer of the Ca<sup>2+</sup>-binding proteins MRP8 and MRP14 binds to arachidonic acid. *FEBS Lett*. 1997 May;408(1):81-4.

Lackmann M, Rajasekariah P, Iismaa SE, Jones G, Cornish CJ, Hu S, Simpson RJ, Moritz RL, Geczy CL. Identification of a chemotactic domain of the pro-inflammatory S100 protein CP-10. *J Immunol*. 1993 Apr;150(7):2981-91.

Lau W, Devery JM and Geczy CL. A chemotactic S100 peptide enhances scavenger receptor and Mac-1 expression and cholesteryl ester accumulation in murine peritoneal macrophages in vivo. *J Clin Invest* 1995(5):1957-65.

Lawrence JB, Oxvig C, Overgaard MT, Sottrup-Jensen L, Gleich GJ, Hays LG, Yates JR 3rd, Conover CA. The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A. *Proc Natl Acad Sci U S A*. 1999 Mar;96(6):3149-53.

Loomans HJ, Hahn BL, Li QQ, Phadnis SH, Sohnle PG. Histidine-based zinc-binding sequences and the antimicrobial activity of calprotectin. *J Infect Dis*. 1998 Mar;177(3):812-4.

Lugering N, Stoll R, Kucharzik T, Schmid KW, Rohlmann G, Burmeister G, Sorg C, Domschke W. Immunohistochemical distribution and serum levels of the Ca(2+)-binding proteins MRP8, MRP14 and their heterodimeric form MRP8/14 in Crohn's disease. *Digestion*. 1995;56(5):406-14.

Luttun A, Tjwa M, Carmeliet P. Placental growth factor (PlGF) and its receptor Flt-1 (VEGFR-1): novel therapeutic targets for angiogenic disorders. *Ann NY Acad Sci* 2002;979:80-93.

Luttun A, Tjwa M, Moons L. Revascularization of ischemic tissues by PlGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nat Med* 2002;8:831-840.

- Maier W, Altwegg LA, Corti R. Inflammatory Markers at the Site of Ruptured Plaque in Acute Myocardial Infarction. *Circulation* 2005;111:1355-1361.
- Manten A, de Winter RJ, Minnema MC. Procoagulant and proinflammatory activity in acute coronary syndromes. *Cardiovasc Res.* 1998 Nov;40(2):389-95.
- Matsumori A, Furukawa Y, Hashimoto T. Plasma levels of the monocyte chemotactic and activating factor/monocyte chemoattractant protein-1 are elevated in patients with acute myocardial infarction. *J Mol Cell Cardiol.* 1997 Jan;29(1):419-23.
- McCord J, Nowak RM, McCullough PA. Ninety-minute exclusion of acute myocardial infarction by use of quantitative point-of-care testing of myoglobin and troponin I. *Circulation* 2001;104:1483-1488.
- McCormick MM, Rahimi F, Bobryshev YV, Gaus K, Zreiqat H, Cai H, Lord RS, Geczy CL. S100A8 and S100A9 in human arterial wall. Implications for atherogenesis. *J Biol Chem.* 2005 Dec;280(50):41521-9.
- Moreno PR, Falk E, Palacios IF, Newell JB, Fuster V and Fallon JT. Macrophage infiltration in acute coronary syndromes. Implications for plaque rupture. *Circulation* 1994 Aug;90(2):775-8.
- Moroz OV, Dodson GG, Wilson KS, Lukanidin E, Bronstein IB. Multiple structural states of S100A12: A key to its functional diversity. *Microsc Res Tech.* 2003 Apr;60(6):581-92.
- Murono S, Yoshizaki T, Sato H, Takeshita H, Furukawa M, Pagano JS. Aspirin inhibits tumor cell invasiveness induced by Epstein-Barr virus latent membrane protein 1 through suppression of matrix metalloproteinase-9 expression. *Cancer Res.* 2000 May 1;60(9):2555-61.
- Nisapakultorn K, Ross KF, Herzberg MC. Calprotectin expression inhibits bacterial binding to mucosal epithelial cells. *Infect Immun.* 2001 Jun;69(6):3692-6.
- Oakley BR, Kirsch DR, Morris NR. A simplified ultrasensitive silver stain for detecting proteins in polyacrylamide gels. *Annal Biochem.* 1980 Jul;105(2):361-3.
- Ono K, Matsumori A, Furukawa Y, Igata H, Shioi T, Matsushima K, Sasayama S. Prevention of myocardial reperfusion injury in rats by an antibody against monocyte chemotactic and activating factor/monocyte chemoattractant protein-1. *Lab Invest.* 1999 Feb;79(2):195-203.
- Passey RJ, Williams E, Lichanska AM, Wells C, Hu S, Geczy CL, Little MH, Hume DA. A null mutation in the inflammation-associated S100 protein S100A8 causes early resorption of the mouse embryo. *J Immunol.* 1999 Aug;163(4):2209-16.
- Pietas A, Schluns K, Marenholz I, Schafer BW, Heizmann CW, Petersen I. Molecular cloning and characterization of the human S100A14 gene encoding a novel member of the S100 family. *Genomics.* 2002 Apr;79(4):513-22.

Quiles J, Roy D, Gaze D, Garrido IP, Avanzas P, Sinha M, Kaski JC. Relation of ischemia-modified albumin (IMA) levels following elective angioplasty for stable angina pectoris to duration of balloon-induced myocardial ischemia. *Am J Cardiol.* 2003 Aug;92(3):322-4.

Rammes A, Roth J, Goebeler M, Klempt M, Hartmann M, Sorg C. Myeloid-related protein (MRP) 8 and MRP14, calcium-binding proteins of the S100 family, are secreted by activated monocytes via a novel, tubulin-dependent pathway. *J Biol Chem.* 1997 Apr;272(14):9496-502.

Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med.* 1997 Apr;336(14):973-9.

Ridker PM, Rifai N, Stampfer MJ, Hennekens CH. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation.* 2000 Apr 18;101(15):1767-72.

Robinson MJ, Tessier P, Poulsom R, Hogg N. The S100 family heterodimer, MRP-8/14, binds with high affinity to heparin and heparan sulfate glycosaminoglycans on endothelial cells. *J Biol Chem.* 2002 Feb;277(5):3658-65.

Rouleau JL, Moye LA, Pfeffer MA, Arnold JM, Bernstein V, Cuddy TE, Dagenais GR, Geltman EM, Goldman S, Gordon D. A comparison of management patterns after acute myocardial infarction in Canada and the United States. The SAVE investigators. *N Engl J Med.* 1993 Mar;328(11):779-84.

Ross, R. Atherosclerosis – an inflammatory disease. *N Engl J Med.* 1999 Jan;340:115-126.

Schafer BW, Heizmann CW. The S100 family of EF-hand calcium-binding proteins: functions and pathology. *Trends Biochem Sci.* 1996 Apr;21(4):134-40.

Schluesener HJ, Kreamsner PG and Meyermann R. Widespread expression of MRP8 and MRP14 in human cerebral malaria by microglial cells. *Acta Neuropathol (Berl).* 1998 Dec;96(6):575-80.

Schmidt AM, Yan SD, Yan SF, Stern DM. The biology of the receptor for advanced glycation end products and its ligands. *Biochim Biophys Acta.* 2000 Dec;1498(2-3):99-111.

Sinha MK, Gaze DC, Tippins JR, Collinson PO, Kaski JC. Ischemia modified albumin is a sensitive marker of myocardial ischemia after percutaneous coronary intervention. *Circulation.* 2003 May;107(19):2403-5.

Sinha MK, Roy D, Gaze DC, Collinson PO, Kaski JC. Role of "Ischemia modified albumin", a new biochemical marker of myocardial ischaemia, in the early diagnosis of acute coronary syndromes. *Emerg Med J.* 2004 Jan;21(1):29-34.

Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goeke NM, Olson BJ, Klenk DC. Measurement of protein using bicinchoninic acid. *Anal Biochem.* 1985 Oct;150(1):76-85.

Soejima H, Ogawa H, Yasue H, Kaikita K, Takazoe K, Nishiyama K, Misumi K, Miyamoto S, Yoshimura M, Kugiyama K, Nakamura S, Tsuji I. Angiotensin-converting enzyme inhibition reduces monocyte chemoattractant protein-1 and tissue factor levels in patients with myocardial infarction. *J Am Coll Cardiol.* 1999 Oct;34(4):983-8.

Stary HC, Chandler AB, Glagov S, Guyton JR, Insull W Jr, Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation.* 1994 May;89(5):2462-78.

Steinberg D, Parthasarathy S, Carew TE, Khoo JC, Witztum JL. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med.* 1989 Apr;320(14):915-24.

Stockley RA, Dale I, Hill SL, Fagerhol MK. Relationship of neutrophil cytoplasmic protein (L1) to acute and chronic lung disease. *Scand J Clin Lab Invest.* 1984 Nov;44(7):629-34.

Strandberg TE, Vanhanen H, Tikkanen MJ. Effect of statins on C-reactive protein in patients with coronary artery disease. *Lancet.* 1999 Jan;353(9147):118-9.

Striz I, Trebichavsky I. Calprotectin - a pleiotropic molecule in acute and chronic inflammation. *Physiol Res.* 2004;53(3):245-53.

Suefuji H, Ogawa H, Yasue H, Sakamoto T, Miyao Y, Kaikita K, Soejima H, Misumi K, Miyamoto S, Kataoka K. Increased plasma level of soluble E-selectin in acute myocardial infarction. *Am Heart J.* 2000 Aug;140(2):243-8.

Szmitko PE, Wang CH, Weisel RD, de Almeida JR, Anderson TJ, Verma S. New markers of inflammation and endothelial cell activation: Part I. *Circulation.* 2003 Oct;108(16):1917-23.

The Joint European Society of Cardiology/American College of Cardiology Committee.. Myocardial infarction redefined - A consensus document of the Joint European Society of Cardiology/American College of Cardiology Committee for the Redefinition of Myocardial Infarction. *Eur Heart J* 2000;21:1502-1513.

Tjwa M, Luttun A, Autiero M, Carmeliet P. VEGF and PlGF: two pleiotropic growth factors with distinct roles in development and homeostasis. *Cell Tissue Res.* 2003 Oct;314(1):5-14.

Toss H, Lindahl B, Siegbahn A, Wallentin L. Prognostic influence of increased fibrinogen and C-reactive protein levels in unstable coronary artery disease. FRISC Study Group. Fragmin during Instability in Coronary Artery Disease. *Circulation.* 1997 Dec;96(12):4204-10.

Tracy RP, Arnold AM, Ettinger W, Fried L, Meilahn E, Savage P. The relationship of fibrinogen and factors VII and VIII to incident cardiovascular disease and death in the elderly: results from the cardiovascular health study. *Arterioscler Thromb Vasc Biol.* 1999 Jul;19(7):1776-83.

Varo N, de Lemos JA, Libby P. Soluble CD40L: risk prediction after acute coronary syndromes. *Circulation* 2003;108:1049-1052.

Viemann D, Strey A, Janning A, Jurk K, Klimmek K, Vogl T, Hirono K, Ichida F, Foell D, Kehrel B, Gerke V, Sorg C, Roth J. Myeloid-related proteins 8 and 14 induce a specific inflammatory response in human microvascular endothelial cells. *Blood.* 2005 Apr;105(7):2955-62.

Xu Z, Zhao S, Zhou H, Ye H, Li J. Atorvastatin lowers plasma matrix metalloproteinase-9 in patients with acute coronary syndrome. *Clin Chem.* 2004 Apr;50(4):750-3.

Yusuf S, Flather M, Pogue J, Hunt D, Varigos J, Piegas L, Avezum A, Anderson J, Keltai M, Budaj A, Fox K, Ceremuzynski L. Variations between countries in invasive cardiac procedures and outcomes in patients with suspected unstable angina or myocardial infarction without initial ST elevation. OASIS (Organisation to Assess Strategies for Ischaemic Syndromes) Registry Investigators. *Lancet.* 1998 Aug;352(9127):507-14.

Zapico-Muniz E, Santalo-Bel M, Merce-Muntanola J, Montiel JA, Martinez-Rubio A, Ordonez-Llanos J. Ischemia-modified albumin during skeletal muscle ischemia. *Clin Chem.* 2004 Jun;50(6):1063-5.

Zhang R, Brennan ML, Fu X, Aviles RJ, Pearce GL, Penn MS, Topol EJ, Sprecher DL, Hazen SL. Association between myeloperoxidase levels and risk of coronary artery disease. *JAMA.* 2001 Nov;286(17):2136-42.

Zheng L, Teschler H, Guzman J, Hubner K, Striz I, Costabel U. Alveolar macrophage TNF-alpha release and BAL cell phenotypes in sarcoidosis. *Am J Respir Crit Care Med.* 1995 Sep;152(3):1061-6.