

7 LITERATURVERZEICHNIS

- Abremski K.**, Hoess R. Bacteriophage P1 site-specific recombination. Purification and properties of the Cre recombinase protein. *J Biol Chem.* 1984; 259(3): 1509-1514.
- Achermann J.C.**, Ozisik G., Ito M., Orun U.A. Harmanci K., Gurakan B., Jameson J.L. Gonadal determination and adrenal development are regulated by the orphan nuclear receptor steroidogenic factor-1, in a dose-dependent manner. *J. Clin. Endocrinol. Metab.* 2002; 87: 1829-1833
- Aebi M.**, Hornig H., Weissman C. 5' cleavage site in eukaryotic pre-mRNA splicing is determined by overall 5' splice region, not by the conserved 5' GU. *Cell* 1987; 50: 237-246
- Aigueperse C.**, Martinez A., Lefrancois-Martinez A.M., Veyssiere G., Jean Cl. Cyclic AMP regulates expression of the gene coding for a mouse vas deferens protein related to the aldose reductase superfamily in human and murine adrenalcortical cells. *J. Endocrinol.* 1999; 160: 147-154
- Aigueperse C.**, Val P., Pacot C., Darne C., Lalli E., Sassone-Corsi P., Veyssiere G., Jean Cl., Martinez A. SF-1 (Steroidogenic Factor-1), C/EBP β (CCAAT/Enhancer Binding Protein), Ubiquitous Transcription Factors NF1 (Nuclear Factor 1) and Sp1 (Selective Promoter Factor 1) Are Required for Regulation of the Mouse Aldose Reduktase-Like Gene (AKR1B7) Expression in Adrenocortical Cells. *Mol. Endocrinol.* 2001; 15: 93-111
- Aitken R.J.** The role of free oxygen radicals and sperm function. *Int. J. Androl.* 1989; 12(2): 95-97.
- Aitken R.J.** A free radical theory of male infertility. *Reprod. Fertil. Dev.* 1994; 6: 19-23
- Aitken R.J.**, Fisher H. Reactive oxygen species generation and human spermatozoa: the balance of benefit and risk. *Bioessays* 1994; 16: 259-267
- Alary J.**, Fernandez Y., Debrauwer L., Perdu E., Gueraud F. Identification of intermediate pathways of 4-hydroxynonenal metabolism in the rat. *Chem. Res. Toxicol.* 2003a; 16: 320-327
- Alary J.**, Gueraud F., Cravedi J. Fate of 4-hydroxynonenal in vivo: disposition and metabolic pathways. *Mol. Aspects of Med.* 2003b; 24: 177-187
- Angele M.K.**, Schwacha M.G., Ayala A. Effect of gender and sex hormones on immune responses following shock. *Shock* 2000; 14: 81-90
- Aschoff J.** Laws of biological diurnal periodicity. *Dtsch. Med. Wochenschr.* 1963; 88: 1930-1937.
- Aveldano M.I.**, Rotstein N.P., Vermouth N.T. Occurrence of long and very long polyenoic fatty acids of the n-9 series in rat spermatozoa. *Lipids* 1992; 27(9): 676-680.
- Babu P.S.**, Bavers D.L., Shah S., Hammer G.D. Role of phosphorylation, gene dosage and Dax-1 in SF-1 mediated steroidogenesis. *Endocr. Res.* 2000; 26(4): 985-994
- Babu PS**, Bavers DL, Beuschlein F, Shah S, Jeffs B, Jameson JL, Hammer GD. Interaction between Dax-1 and steroidogenic factor-1 in vivo: increased adrenal responsiveness to ACTH in the absence of Dax-1. *Endocrinology.* 2002;143(2): 665-673.
- Bagnasco S.M.**, Uchida S., Balaban P.F., Kador P.F., Burg M.B. Induction of aldose reductase and sorbitol in renal inner medullary cells by elevated extracellular NaCl. *Proc. Natl. Acad. Sci. USA* 1987; 84: 1718-1720
- Balhorn R.**, Reed S., Tanphaichitr N. Aberrant protamine 1/protamine 2 ratios in sperm of infertile human males. *Experientia* 1988; 44: 52-55

- Barazzone-Argiroffo C.**, Pagano A., Juge C., Metrailler I., Rochchat A., Vesin C., Donati Y. Glucocorticoids aggravate hyperoxia-induced lung injury through decreased nuclear factor-kappa B activity. *Am. J. Physiol. Lung. Cell. Mol. Physiol.* 2003; 284(1): 197-204
- Barnes W.M.** PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates. *Proc. Natl. Acad. Sci. USA.* 1994; 91(6): 2216-2220
- Baxter J.D.**, Tyrell J.B. The adrenal cortex. In: Felig P., Baxter J.D., Frohman L.A. eds., S. 385 *Endocrinology and metabolism.* McGraw-Hill, New York (1981)
- Beuschlein F.**, Mutch C., Bavers D.L., Ulrich-Lai Y.M., Engeland W.C., Keegan C., Hammer G.D. Steroidogenic Factor-1 Is Essential for Compensatory Adrenal Growth Following Unilateral Adrenalectomy. *Endocrinology* 2002; 143(8): 3122-3135
- Bianchi P.G.**, Manicardi G.C., Urner F., Campana A., Sakkas D. Chromatin packaging and morphology in ejaculated human spermatozoa: evidence of hidden anomalies in normal spermatozoa. *Mol. Hum. Reprod.* 1996; 2: 139-144
- Bjerrum O.J.**, Schafer-Nielsen C. *Analytical Electrophoresis.* M.J. Dunn ed., S. 315; Verlag Chemie, Weinheim (1986)
- Bohren K.**, Bullock B., Wermuth B., Gabbay K., The aldo-keto reductase superfamily. *J. Biol. Chem.* 1989; 264: 9547-9551
- Bradford M.M.** A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 1976; 72: 248-254.
- Breathnach R.**, Chambon P. Organization and expression of split genes coding for proteins. *A. Rev. Biochem.* 1981; 50: 349-383
- Brockstedt E.**, Peters-Kottig M., Badock V., Hegele-Hartung C., Lessl M. Luteinizing Hormone Induces Mouse Vas Deferns Protein Expression in the Murine Ovary. *Endocrinology* 2000; 141: 2574-2581
- Burstein S.**, Gut M. Biosynthesis of pregnenolone. *Recent Prog. Horm. Res.* 1971; 27: 303-349
- Cammass F.M.**, Pullinger G.D., Barker S., Clark A.J. The mouse adrenocorticotropin receptor gene: cloning and characterization of its promoter and evidence for a role of orphan nuclear receptor steroidogenic factor-1. *Mol. Endocrinol.* 1997; 11: 867-876
- Chau Y.M.**, Crawford P.A., Woodson K.G., Polish J.A., Olson L.M., Sadovsky Y. Role of steroidogenic-factor 1 in basal and 3',5'-cyclic adenosine monophosphate-mediated regulation of cytochrome P450 side-chain cleavage enzyme in the mouse. *Biol. Reprod.* 1997; 57: 765-771
- Christian J.J.** Population density and reproductive efficiency. *Biol. Reprod.* 1971; 4(3): 248-94.
- Chung S.M.**, Ho E.C.M., Lam K.S.L., Chung S.K. Contribution of polyol pathway to diabetes-induced oxidative stress. *J. Am. Soc. Nephrol.* 2003; 14: S233-236
- Cohen J.J.** Glucocorticoids. In Roitt I.M., Delves P.J. eds., S. 616 *Encyclopedia of Immunology,* Academic Press, London (1992)
- Constantopoulos G.**, Carpenter A., Satoh P., Tchen T.T. Formation of isocaproaldehyd in the enzymatic cleavage of cholesterol side chain by adrenal extract. *Biochemistry* 1966; 5: 1650-1652
- Crawford P.A.**, Dorn C., Sadovsky Y., Milbrandt J. Nuclear receptor DAX-1 recruits nuclear receptor corepressor N-CoR to steroidogenic factor 1. *Mol. Cell. Biol.* 1998; 18: 2949-2956

- Cross J.C.**, Werb Z., Fisher S.J. Implantation and the placenta: key pieces of the development puzzle. *Science* 1994; 266: 1508-1518
- Daoudal S.**, Berger M., Pailhoux E., Tournaire C., Veysiere G., Jean C. Cloning and expression of mouse aldose reductase. Comparison with the androgen-dependent protein from mouse vas deferens. *Life Sci. Adv. (Steroid Biochem.)* 1995; 14: 45-58
- Davis D.W.**, Hamilton R.F. Jr., Holian A. 4-Hydroxynonenal inhibits interleukin-1 beta converting enzyme. *J. Interferon Cytokine. Res.* 1997; 17(4): 205-210.
- Devenport L.**, Knehans A., Sundstrom A., Thomas T. Corticosterone's dual metabolic actions. *Life Sci.* 1989; 45: 1389-1396
- de Vos P.**, Lefebvre A., Shribo I., Fruchart J., Auwerx J. Glucocorticoids induce the expression of the leptin gene through a non-classical mechanism of transcriptional activation. *Eur. J. Biochem.* 1998; 253: 619-626
- Dubuc PU**, Carlisle HJ. Food restriction normalizes somatic growth and diabetes in adrenalectomized ob/ob mice. *Am. J. Physiol.* 1988; 255: R787-793.
- Duval D.L.** Nelson S.E., Clay C.M. A binding site for steroidogenic factor-1 is part of a complex enhancer that mediates expression of the murine gonadotropin-releasing hormone receptor gene. *Biol. Reprod.* 1997; 56: 160-168
- Elkenov I.J.**, Chrousos G.P. Stress hormones, Th1/Th2 patterns, pro/anti-inflammatory cytokines and susceptibility to disease. *Trends Endocrinol. Metab.* 1999; 10: 359-386
- Esterbauer H.**, Schaur R.J., Zollner H. Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free. Radic. Biol. Med.* 1991; 11: 81-128
- Evenson D.P.**, Jost L.K., Corzett M., Balhorn R. Characteristics of human sperm chromatin structure following an episode of influenza and high fever: a case study. *J. Androl.* 2000; 21: 739-746
- Fabre S**, Darne C, Veysiere G, Jean C. Characterization of the promoter of the gene for a mouse vas deferens protein related to the aldo-keto reductase superfamily: effect of steroid hormones and phorbol esters. *J. Steroid Biochem. Mol. Biol.* 1995; 5(3-4): 315-325.
- Facchinetti F.**, Dawson V.L., Dawson T.M. Free radicals as mediators of neuronal injury. *Cell. Mol. Neurobiol.* 1998; 18: 667-682
- Fedorova L**, Fedorov A. Introns in gene evolution. *Genetica* 2003; 118(2-3): 123-231.
- Fevold H.L.** Synergism of follicle stimulating and luteinizing hormones in producing estrogen secretion. *Endocrinology* 1941; 28: 33-36
- Frederic F.**, Oliver C., Wollman E., Delhaye-Bouchaud N., Mariani J. IL-1 and LPS induce a sexually dimorphic response of the hypothalamo-pituitary-adrenal axis in several mouse strains. *Eur. Cytokine Netw.* 1993; 4(5): 321-329
- Freeman, M. E.** The ovarian cycle of the rat. in *The Physiology of Reproduction*. E. Knobil and J. D. Neill, eds., S. 1893-1928 Raven Press, New York (1988)
- Gabbay KH.** The sorbitol pathway and the complications of diabetes. *N. Engl. J. Med.* 1973 Apr 19; 288(16): 831-836.
- Gonzalez A.M.**, Sochor M., McLean P. The effect of an aldose reductase inhibitor (Sorbitol) on the level of metabolites in lenses of diabetic rats. *Diabetes* 1983; 32: 482-485
- Grisham M.B.**, Granger D.N. Metabolic sources of reactive oxygen metabolites during oxidant stress and ischemia with reperfusion. *Clin. Chest. Med.* 1989; 10(1): 71-81
- Gui T.**, Tanimoto T., Kokai Y., Nishimura C. Presence of a closely related subgroup in the aldo-ketoreductase family of the mouse. *Eur. J. Biochem.* 1995; 227(1-2): 448-453
- Hall P.F.** In: *The Testis* (Johnson A.D., Gomes W.R., Vandemark N.L. eds., S. 1-72, Academic Press, New York (1970)

- Hamilton** D.W. Structure and function of the epithelium lining the ductuli efferentes, ductus epididymis and ductus deferens of the rat. In: Hamilton D.W., Greep R.O., eds. Handbook of Physiology. S. 259-301, Am. Physiological Society Washington, DC (1975)
- Hawkins** E.F., Young P.N., Hawkins A.M., Bern H.A. Adrenocortical function: corticosterone levels in female BALB/c and C3H mice under various conditions. *J. Exp. Zool.* 1975; 194(3): 479-484
- Hemachand** T., Shaha C. Functional role of sperm surface glutathione S-transferases and extracellular glutathione in the haploid spermatozoa under oxidative stress. *FEBS letters* 2003; 538: 14-18
- Herbst** U., Toborek M., Kaiser S., Mattson M.P., Hennig B. 4-Hydroxynonenal induces dysfunction and apoptosis of cultured endothelial cells. *J. Cell Physiol.* 1999; 181(2): 295-303
- Hers** H.G. The mechanism of transformation of glucose to fructose by seminal vesicles. *Biochem. Biophys. Acta* 1956; 22: 202-203
- Ho** H.T.B.; Jenkins N.A., Copeland N.G., Golbert D.J., Winkles J.A., Louie H.W.Y., Lee F.K.L., Chung S.M., Chung S.K. Comparisons of genomic structures and chromosomal locations of the mouse aldose reductase and aldose reductase-like genes. *Eur. J. Biochem.* 1999; 259: 726-730
- Hoess** R.H., Abremski K. Mechanism of strand cleavage and exchange in the Cre-lox site-specific recombination system. *J. Mol. Biol.* 1985; 181(3): 351-362.
- Hogan** B., Constantini F., Lacy E. Manipulating the Mouse Embryo. A Laboratory Manual. Cold Spring Harbor Laboratory (1986)
- Hyndman** D., Bauman D.R., Heredia V.V., Penning T.M. The aldo-keto reductase superfamily homepage. *Chem. Biol. Interact.* 2003; 143-144:621-631
- Ikeda** Y., Shen W.H., Ingraham H.A., Parker K.L. Developmental expression of mouse steroidogenic factor-1, an essential regulator of steroid hydroxylases. *Mol. Endocrinol.* 1994; 8:654-662
- Ito** M., Yu R., Jameson J.L. DAX-1 inhibits mediated transactivation via a carboxy-terminal domain that is deleted in adrenal hypoplasia congenita. *Mol. Cell. Biol.* 1997; 17: 1476-1483
- James** V.H.T., Honour J.W., Fraser R., Makin H.L.J., Gower D.B., Kirk D.N. (Eds.) S. 229 *Steroid Analysis.* Blackie, Glasgow (1995)
- Jez** J, Flynn T., Penning T. 1997. A new nomenclature for the aldo-keto reductase superfamily. *Biochem. Pharmacol.* 1997; 54: 639-647
- Jez**, J. M., and T. M. Penning. 2001. The aldo-keto reductase (AKR) superfamily: an update. *Chem. Biol. Interact.* 130–132: 499–525.
- Jez** J.M., Bennett J.M., Schlegel B.P., Lewis M. Pennig T.M. Comparative anatomy of the aldo–keto reductase superfamily. *Biochem. J.* 1997; 326: 625-636
- Joyce** C.L., Nuzzo N.A., Wilson Jr. L., Zaneveld L.J. Evidence for a role of cyclooxygenase (prostaglandin synthetase) and prostaglandins in the sperm acrosome reaction and fertilization. *J. Androl.* 1987; 8: 74-82
- Kalinich** J.F., Ramakrishnan R., McClain D.E., Ramakrishnan N. 4-Hydroxynonenal, an end-product of lipid peroxidation, induces apoptosis in human leukemic T- and B-cell lines. *Free Radic. Res.* 2000; 33(4): 349-358
- Kawabe** K., Shikayama T., Tsuboi H., Oka S., Oba K., Yanase T., Nawata H., Morohashi K. DAX-1 is one of the target genes of Ad4BP/SF-1. *Mol. Endocrinol.* 1999; 13: 1267-1284

- Keri R.A.**, Nilson J.H. A steroidogenic factor-1 binding site is required for activity of the luteinizing hormone b subunit promoter in gonadotropes of transgenic mice. *J. Biol. Chem.* 1996; 271: 10782-10785
- Kinoshita JH**, Nishimura C. The involvement of aldose reductase in diabetic complications. *Diabetes Metab. Rev.* 1988; 4(4): 323-337.
- Klatt P.**, Esterbazer H. Oxidative hypothesis of atherogenesis. *J Cardivasc Risk* 1996; 3: 346-351
- Klebe RJ**, Harriss JV, Sharp ZD, Douglas MG. A general method for polyethylene-glycol-induced genetic transformation of bacteria and yeast. *Gene.* 1983; 25(2-3): 333-341.
- Klein S.L.** The effects of hormones on sex differences in infection: From genes to behavior. *Neurosci. Biobehav. Rev.* 2000; 24: 627-638
- Kruman I.**, Bruce-Keller A.J., Bredesen D., Waeg G., Mattson M.P. Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. *J. Neurosci.* 1997; 17(13): 5089-5100
- Lalli E.**, Melner M.H., Stocco D.M., Sassone-Corsi P. DAX-1 blocks steroid production at multiple levels. *Endocrinology* 1998; 139: 4237-4243
- Lalli E.**, Ohe K., Himdelang C., Sassone-Corsi P. Orphan receptor DAX-1 is a shuttling RNA binding protein associated with polyribosomes via mRNA. *Mol. Cell. Biol.* 2000; 20: 4910-4921
- Laemmli U.K.** Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 1970; 227(259): 680-685
- Lau E. T.**, Cao D., Lin C., Chung S.K., Chung S.S. Tissue-specific expression of two aldose reductase-like genes in mice: abundant expression of mouse vas deferens protein and fibroblast growth factor-regulated protein in the adrenal gland. *Biochem. J.* 1995; 312: 609-615
- Leal-Cerro A.**, Soto A., Martinez M.A., Dieguez C., Casanueva F.F. influence of cortisol on leptin secretion. *Pituitary* 2002; 4(1): 111-116
- Lee A.Y.**, Chung S.K., Chung S.S. Demonstration that polyol accumulation is responsible for diabetic cataract by use of transgenic mice expression the aldose reductase gene in the lens. *Proc. Natl. Acad. Sci. USA* 1995; 92: 2784
- Lee A.Y.**, Chung S.S. Contribution of polyol pathway to oxidative stress in diabetic cataract. *FASEB J.* 1999; 13: 23-30,
- Lefer A.M.** Leukotrienes as mediators of ischemia and shock. *Biochem. Pharmacol.* 1986; 35: 123-127
- Lefrancois-Martinez A.M.**, Tournaire C., Martinez A., Berger M., Daoudal S., Tritsch D., Veyssiere G., Jean Cl. Product of side-chain cleavage of cholesterol, isocaproaldehyde, is an endogenous specific substrate of mouse vas deferens protein, an aldose reductase-like protein in adrenocortical cells. *JBiol Chem* 1999; 274: 32875-32880
- Licino J.**, Mantzoros C., Negrao A.B., Cizza G., Wong M.L., Bongiorno P.B., Chrousos G.P., Karp B., Allen C., Flier J.S., Gold P.W. Human leptin levels are pulsatile and inversely related to pituitary-adrenal function. *Nat. Med.* 1997; 3: 575-579
- Majdic G.**, Young M., Gomez-Sanchez E., Anderson P., Szczepaniak L.S., Dobbins R.L., McGarry J.D., Parker K.L. Knockout mice lacking steroidogenic factor 1 are a novel genetic model of hypothalamic obesity. *Endocrinology* 2002; 143(2): 607-614
- Manin M.**, Martinez A., Van der Schueren B., Reynaert I., Jean C. Acquisition of androgen-mediated expression of mouse vas deferens protein (MVDP) gene in cultured epithelial cells and in the vas deferens during postnatal development. *J Androl.* 2000; 21: 641-650
- Martinez A.**, Berger M., Jean C.L. Androgen Regulation of Major mRNAs for Proteins of the Immature and Adult Mouse Vas Deferens. *J Endocrinol.* 1989; 120: 67-74

- Martinez A.**, Pailhoux E., Berger M., Jean Cl. Androgen Regulation of the mRNA Encoding a Major Protein of the Mouse Vas Deferens. *Mol. Cell. Endocrinol.* 1990; 72: 201-211
- Martinez A.**, Berger M., Jean-Faucher Ch., Veyssière G., Jean Cl. Radioimmunoassay of mouse vas deferens protein: developmental profile and hormonal regulation. *J. Endocrinol.* 1993; 138: 99-105
- Martinez A.**, Lefrancois-Martinez A., Manin M, Guyot S., Jean-Faucher Ch., Veyssiere G., Kahn A., Jean C. 5'-Flanking and intragenic sequences confer androgenic and developmental regulation of mouse aldose reductase-like gene in vas deferens and adrenal in transgenic mice. *Endocrinology* 1999; 140: 1338-1348
- Martinez A.**, Aigueperse C., Val C., Dussault M., Tournaire C., Berger M., Veyssiere G., Jean C., Lefrancois-Martinez A. Physiological functions and hormonal regulation of mouse vas deferens protein (AKR1B7) in steroidogenic tissues. *Chem. Biol. Interact.* 2001; 130-132: 903-917
- Martinez A.**, Val P., Sahut-Barnola I., Aigueperse C., Veyssiere G., Lefrancois-Martinez A. Steroidogenic factor-1 controls the aldose reductase *akr1b7* gene promoter in transgenic mice through an atypical binding site. *Endocrinology* 2003; 144: 2111-2120
- Medzhitov R.** Toll-like receptors and innate immunity. *Nat. Rev. Immunol.* 2001; 1: 135-145
- Melov S.** Mitochondrial oxidative stress. Physiologic consequences and potential for a role in aging. *Ann. N. Y. Acad. Sci.* 2000; 908: 219-225.
- Morohashi K.I.**, Zanger U.M., Honda S.I., Hara M., Waterman M.R., Omura T. Activation of CYP11A and CYP11B gene promoters by steroidogenic cell-specific transcription factor, Ad5BP. *Mol. Endocrinol.* 1993; 7: 1196-1204
- Moustafa M.H.**, Sharma R.K., Thornton J., Mascha E., Abdel-Hafez M.A., Thomas A.J., Agarwal A. Relationship between ROS production, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. *Hum. Reprod.* 2004; 1: 129-138
- Mukai T.**, Kusaka M., Kawabe K., Goto K., Nawata H., Fujieda K., Morohashi K. Sexually dimorphic expression of Dax-1 in the adrenal cortex. *Genes to Cells* 2002; 7: 717-729
- Nichols DJ**, Chevins PF. Plasma corticosterone fluctuations during the oestrous cycle of the house mouse. *Experientia.* 1981; 37(3): 319-320.
- Oates PJ**, Mylari BL. Aldose reductase inhibitors: therapeutic implications for diabetic complications. *Expert Opin. Investig. Drugs.* 1999; 8(12): 2095-2119.
- Pailhoux E.A.**, Martinez A., Veyssiere G.M., Jean C.G. Androgen-dependent protein from mouse vas deferens. cDNA cloning and protein homology with the aldo-keto reductase superfamily. *J. Biol. Chem.* 1990; 265(32): 19932-19936.
- Pailhoux E.**, Martinez A, Veyssiere G, Tournaire C, Berger M, Jean C. Characterization of cDNA and of the gene corresponding to androgen-dependent protein of the vas deferens in mice. *Ann. Endocrinol. (Paris).* 1991; 52(6): 437-440
- Pailhoux E.**, Veyssiere G., Fabre S., Tournaire C., Jean C. The genomic organization and DNA sequence of the mouse vas deferens androgen-regulated protein gene. *J. Steroid. Biochem. Mol. Biol.* 1992; 42(6): 561-568.
- Parker K.L.**, Rice D.A., Lala D.S., Ikeda Y., Luo X., Wong M., Bakke M., Zhao L., Frigeri C., Hanley N.A., Stallings N., Schimmer B.P. Steroidogenic factor 1: an essential mediator of endocrine development. *Recent Prog. Horm. Res.* 2002; 57: 19-36
- Pedersen T**, Peters H. Proposal for a classification of oocytes and follicles in the mouse ovary. *J. Reprod. Fertil.* 1968; 17(3): 555-557.
- Peng X.R.**, Hsueh A.J., LaPolt P.S., Bjersing L., Ny T. Localization of luteinizing hormone receptor messenger ribonucleic acid expression in ovarian cell types during follicle development and ovulation. *Endocrinology* 1991; 129: 3200-3207

- Peters-Kottig M.** Funktionale Charakterisierung des Maus Vas Deferens Proteins. Dissertation eingereicht am Fachbereich Biologie, Chemie, Pharmazie der Freien Universität Berlin (2001)
- Petrides P.E.** Endokrine Gewebe III : Hypothalamisch-hypophysäres System und Zielgewebe. In Biochemie und Pathobiochemie. Löffler G. und Petrides P.E. eds. Kapitel 29; Springer Verlag Berlin, Heidelberg, New York (1998)
- Plante M., Lamirande E., Gagnon C.** Reactive oxygen species released by activated neutrophils, but not by deficient spermatozoa, are sufficient to affect normal sperm motility. *Fertil. Steril.* 1994; 62: 387-393
- Poli G., et al.** Separation and characterization of the aldehydic products of lipid peroxidation stimulated by carbon tetrachloride or ADP-iron in isolated rat hepatocytes and rat liver microsomal suspensions. *Biochem.* 1985; 227: 629-638
- Quinn S.J., Williams G.H.** Regulation of aldosterone secretion. *Annu. Rev. Physiol.* 1988; 50: 409-426
- Radulovic M., Dautzenberg F.M., Sydow S., Radulovic J., Spiess J.** Corticotropin-releasing factor receptor 1 in mouse spleen: expression after immun stimulation and identification of receptor-bearing cells. *J. Immunol.* 1999; 162: 3013-3021
- Rindgen D, Nakajima M, Wehrli S, Xu K, Blair IA.** Covalent modifications to 2'-deoxyguanosine by 4-oxo-2-nonenal, a novel product of lipid peroxidation. *Chem. Res. Toxicol.* 1999; 12(12): 1195-1204.
- Rittner H.L., Hafner V., Klimiuk P.A., Szweda L.I., Goronzy J.J., Weyand C.M.** Aldose reductase functions as a detoxification system for lipid peroxidation products in vasculitis. *J. Clin. Invest.* 1999; 103(7): 1007-1013
- Rugh R.** The mouse, its reproduction and development. Burgess Puv. Co Minneapolis (1968)
- Sadovsky Y., Crawford P.A., Woodson K.G., Polish J., Clements M.A., Tourtellotte L.M., Simburger K., Milbrandt J.** Mice deficient in the orphan receptor steroidogenic factor 1 lack adrenal glands and gonads but express P450 side-chain-cleavage enzyme in the placenta and have normal embryonic serum levels of corticoids. *Proc. Natl. Acad. Sci.* 1995; 92: 10939-10943
- Sahut-Barnola I., Lefranciose-Martinez A.M., Jean C., Veyssiere G., Martinez A.** Adrenal tumorigenesis targeted by the corticotropin-regulated promoter of the aldo-keto reductase AKR1B7 gene in transgenic mice. *Endocr. Res.* 2000; 26: 885-898
- Saito M, Bray GA.** Adrenalectomy and food restriction in the genetically obese (ob/ob) mouse. *Am. J. Physiol.* 1984; 246: R20-25
- Schade U.F.** The effect of endotoxin on the lipoxygenase-mediated conversion of exogenous and endogenous arachidonic acid in mouse peritoneal macrophages. *Prostaglandins.* 1987; 34(3): 385-400.
- Schmid W., Cole T.J., Blendy J.A., Schutz G.** Molecular genetic analysis of glucocorticoid signalling in development. *J. Steroid. Biochem. Mol. Biol.* 1995; 53: 33-35
- Schroder J., Kahlke V., Staubach K.H.** Gender differences in sepsis: Genetically determined? *Shock* 2000; 14: 307-310
- Seery L.T., Nestor P.V., FitzGerald G.A.** Molecular Evolution of the aldo-keto reductase gene superfamily. *J. Mol. Evol.* 1998; 46: 139-146
- Serhan C.N., Haeggstrom J.Z., Leslie C.C.** Lipid mediator networks in cell signalling: update and impact of cytokines. *FASEB J.* 1996; 10: 1147-1158
- Sharma R.K., Agarwal A.** Role of reactive oxygen species in male infertility. *Urology* 1996; 48(6): 835-850

- Silver** L.M. Mouse genetics. Concepts and applications. Kapitel 4.1, Oxford University Press (1995)
- Simpson** E.R., Waterman M.R. Regulation of the synthesis of steroidogenic enzymes in adrenocortical cells by ACTH. *Annu. Rev. Physiol.* 1988; 50: 427-440
- Smith** A.G. Culture and differentiation of embryonic stem cells. *J. Tiss. Cult. Meth.* 1991; 3: 89-94.
- Spinedi** E., Chisari A., Pralong F., Gaillard R.C. Sexual dimorphism in the mouse hypothalamic-pituitary-adrenal axis function after endotoxin and insulin stresses during development. *Neuroimmunomodulation* 1997; 4(2): 77-83
- Srivastava** S., Dixit B.L., Cai J., Sharma S., Hurst H.E., Bhatnagar A., Srivastava S.K. Metabolism of lipid peroxidation product, 4-hydroxynonenal (HNE) in rat erythrocytes: role of aldose reductase. *Free. Radic. Biol. Med.* 2000; 29: 642-651
- Srivastava** S., Harter T.M., Chandra A., Bhatnagar A., Srivastava S.K., Petrash J.M. Kinetic studies of FR-1, a growth factor-inducible aldo-keto reductase. *Biochemistry* 1998; 37: 12909-12917
- Stocco** D.M., Clark B.J. Regulation of the acute production of steroids in steroidogenic cells. *Endocr. Rev.* 1996; 17: 221-244
- Suescun** M., Chisari A.N., Carino M., Hadid R., Gaillard R.C., Spinedi E. Sex steroid regulation of the hypothalamo-pituitary-adrenal axis activity in middle-aged mice during endotoxic shock. *Neuroimmunomodulation* 1994; 1(5): 315-320
- Sugawara** T, Kiriakidou M., Mc Allister J.M. Holt J.A., Arakane F., Strauss J.F. Regulation of expression of the steroidogenic acute regulatory protein (StAR) gene: a central role for steroidogenic factor 1. *Steroids* 1997; 62: 5-9
- Sun** A.Y., Chen Y.M. Oxidativ stress and neurodegenerative disorders. *J. Biomed. Sci.* 1998; 5: 401-414
- Suresh** C.S. Oxidative Stress and role of antioxidants in normal and abnormal sperm function. *Front. Biosc.* 1996; 1: 78-86
- Taniguchi** M., Sanbo M., Watanabe S., Naruse I., Mishina M., Yagi T. Efficient production of Cre-mediated site-directed recombinants through the utilization of the puromycin resistance gene, *pac*: a transient gene-integration marker for ES cells. *Nucleic. Acids. Res.* 1998; 26(2): 679-680.
- Taragnat** C., Berger M., Jean C.L. Identification and androgen dependence of proteins in the mouse vas deferens. *Int. J. Androl.* 1986; 9: 299-311
- Taragnat** C., Berger M., Jean C.L. Preliminary characterization of androgen-dependence and ontogeny of an abundant protein from mouse vas deferens. *J. Reprod. Fertil.* 1988; 83: 835-842
- Taragnat** C., Berger M., Jean C.L. Tissue and species specificity of mouse vas deferens protein. *J. Androl.* 1990; 11: 279-286
- Threadgill** D.W., Dlugosz A.A., Hansen L.A., Tennenbaum T., Lichti U., Yee D., LaMantia C., Mourtou T., Herrup K., Harris R.C. Targeted disruption of mouse EGF receptor: effect of genetic background on mutant phenotype. *Science.* 1995; 269(5221): 230-234
- Tracey** K.J., Beutler B., Lowry S.F., Merryweather J., Wolpe S., Milsark I.W., Hariri R.J., Fahey T.J. 3rd, Zentella A., Albert J.D., *et al.* Shock and tissue injury induced by recombinant human cachectin. *Science* 1986; 234: 470-474
- Tou** J.C., Wade C.E. Determinants affecting physical activity levels in animal models. *Exp. Biol. Med.* (Maywood). 2002; 227(8): 587-600

- Ulich** T.R., Guo K.Z., Remick D., del Castillo J., Yin S.M. Endotoxin-induced cytokine gene expression in vivo. III. IL-6 mRNA and serum protein expression and the in vivo hematologic effects of IL-6. *J. Immunol.* 1991; 146(7): 2316-2323
- Underhill** D.M., Ozinsky A. Toll-like receptors: key mediators of microbe detection. *Curr. Opin. Immunol.* 2002; 14: 103-110
- Val** P., Aigueperse C., Lefrancois-Martinez A.M., Jean Cl., Veyssiere G., Martinez A. Role of three SF-1 binding sites in the expression of the MVDP/AKR1-B7 isocaproaldehyde reductase in Y1 cells. *Endocrine Res.* 2002a; 28: 527-533
- Val** P., Martinez A., Sahut-Barnola I., Jean C., Veyssiere G., Lefrancois-Martinez A. A 77-Base pair LINE-like sequence elicits androgen-dependent mvdp/akr1-b7 expression in mouse vas deferens, but is dispensible for adrenal expression in rats. *Endocrinology* 2002b; 143(9); 3435-3448
- Vilain** E., Guo W., Zhang Y.H., Mc Gabe E.R.B. DAX-1 gene expression upregulated by steroidogenic factor-1 in an adrenocortical carcinoma cell line. *Biochem. Mol. Med.* 1997; 61: 1-8
- Wermuth** B. Aldo-keto reductases. *Prog Clin Biol Res.* 1985;174: 209-230.
- Yagihashi** S., Yamagishi S., Wada R., Baba M., Hohman T.C., Yabe-Nishimura C., Kokai Y. Neuropathy in diabetic mice overexpressing human aldose reductase and effects of aldose reductase inhibitor. *Brain* 2001; 124: 2448-2458
- Yamaoka** T., Nishimura C., Yamashita K., Itakura M., Yamada T., Fujimoto J., Kokai Y. Acute onset of diabetic pathological changes in transgenic mice with human aldose reductase cDNA. *Diabetologia.* 1995; 38(3): 255-261
- Yi** P., Zhan D., Samokyszyn V.M., Doerge D.R., Fu P.P. Synthesis and ³²P-postlabeling/high-performance liquid chromatography separation of diastereomeric 1N2-(1,3-propano)-2'-deoxyguanosine 3'-phosphate adducts formed from 4-hydroxy-2-nonenal. *Chem. Res. Toxicol.* 1997; 10: 1259-1265
- Yu** N. R., Ito M., Saunders T.L., Camper S.A., Jameson J.L. Role of Ahch in gonadal development and gametogenesis. *Nat. Genetics* 1998; 20: 353-357
- Young** E.A. Normal glucocorticoid fast feedback following chronic corticosterone pellet treatment. *Psychoneuroendocrinology* 1995; 20: 771-784
- Zanaria** E., Muscatelli F., Bardoni B. An unusual member of the nuclear hormone receptor superfamily responsible for X-linked adrenal hypoplasia congenita. *Nature* 1994; 372: 635-641
- Zazopoulos** E., Lalli E., Stocco D.M., Sassone-Corsi P. DNA-binding and transcriptional repression by DAX-1 blocks steroidogenesis. *Nature* 1997; 390: 311-315
- Zellweger** R., Wichmann M.W., Ayala A. Females in proestrus state maintain splenic immun functions and tolerate sepsis better than males. *Crit. Care Med.* 1997; 25: 106-110
- Zhang** H., Thomsen J.S., Gustafsson J.A., Eckardt T. DAX-1 functions as an LXXLL-containing corepressor for activated estrogen receptors. *J. Biol. Chem.* 2000; 275: 39855-39859
- Zhao** J., Karalis K.P. Regulation of nuclear factor-kappaB by corticotropin-releasing hormone in mouse thymocytes. *Mol. Endocrinol.* 2002; 16(11): 2561-2570.