

7 Literaturverzeichnis

1. Bubien JK, Zhou LJ, Bell PD, et al. Transfection of the CD20 cell surface molecule into ectopic cell types generates a Ca²⁺ conductance found constitutively in B lymphocytes. *J Cell Biol* 1993;121 (5): 1121-32.
2. Bhan AK, Nadler LM, Stashenko P, et al. Stages of B cell differentiation in human lymphoid tissue. *J Exp Med* 1981;154 (3): 737-49.
3. Zubler RH and Kanagawa O Requirement for three signals in B cell responses. II. Analysis of antigen- and Ia-restricted T helper cell-B cell interaction. *J Exp Med* 1982;156 (2): 415-29.
4. Janeway CA, Jr. The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. *Annu Rev Immunol* 1992;10: 645-74.
5. Kabelitz D, Marischen L, Oberg HH, et al. Epithelial defence by gamma delta T cells. *Int Arch Allergy Immunol* 2005;137 (1): 73-81.
6. Kabelitz D and Wesch D Features and functions of gamma delta T lymphocytes: focus on chemokines and their receptors. *Crit Rev Immunol* 2003;23 (5-6): 339-70.
7. Hayday AC [gamma][delta] cells: a right time and a right place for a conserved third way of protection. *Annu Rev Immunol* 2000;18: 975-1026.
8. Kaufmann SH and Kabelitz D Gamma/delta T lymphocytes and heat shock proteins. *Curr Top Microbiol Immunol* 1991;167: 191-207.
9. Malizia G, Calabrese A, Cottone M, et al. Expression of leukocyte adhesion molecules by mucosal mononuclear phagocytes in inflammatory bowel disease. *Gastroenterology* 1991;100 (1): 150-9.
10. Mahida YR, Patel S, Gionchetti P, et al. Macrophage subpopulations in lamina propria of normal and inflamed colon and terminal ileum. *Gut* 1989;30 (6): 826-34.
11. Rogler G, Hausmann M, Vogl D, et al. Isolation and phenotypic characterization of colonic macrophages. *Clin Exp Immunol* 1998;112 (2): 205-15.
12. Germain RN MHC-dependent antigen processing and peptide presentation: providing ligands for T lymphocyte activation. *Cell* 1994;76 (2): 287-99.
13. Brigi M and Brenner MB CD1: antigen presentation and T cell function. *Annu Rev Immunol* 2004;22: 817-90.
14. Mowat AM Anatomical basis of tolerance and immunity to intestinal antigens. *Nat Rev Immunol* 2003;3 (4): 331-41.

15. Wittig BM and Zeitz M The gut as an organ of immunology. *Int J Colorectal Dis* 2003;18 (3): 181-7.
16. Cornes JS Peyer's patches in the human gut. *Proc R Soc Med* 1965;58 (9): 716.
17. O'Leary AD and Sweeney EC Lymphoglandular complexes of the colon: structure and distribution. *Histopathology* 1986;10 (3): 267-83.
18. MacDonald TT The mucosal immune system. *Parasite Immunol* 2003;25 (5): 235-46.
19. Neutra MR, Frey A, and Kraehenbuhl JP Epithelial M cells: gateways for mucosal infection and immunization. *Cell* 1996;86 (3): 345-8.
20. Ohtsuka A, Piazza AJ, Ermak TH, et al. Correlation of extracellular matrix components with the cytoarchitecture of mouse Peyer's patches. *Cell Tissue Res* 1992;269 (3): 403-10.
21. Brandtzaeg P and Bjerke K Immunomorphological characteristics of human Peyer's patches. *Digestion* 1990;46 Suppl 2: 262-73.
22. Tseng J Expression of immunoglobulin isotypes by lymphoid cells of mouse intestinal lamina propria. *Cell Immunol* 1982;73 (2): 324-36.
23. MacLennan IC, Gulbranson-Judge A, Toellner KM, et al. The changing preference of T and B cells for partners as T-dependent antibody responses develop. *Immunol Rev* 1997;156: 53-66.
24. Hamann A, Andrew DP, Jablonski-Westrich D, et al. Role of alpha 4-integrins in lymphocyte homing to mucosal tissues in vivo. *J Immunol* 1994;152 (7): 3282-93.
25. Hamann A, Jablonski-Westrich D, Jonas P, et al. Homing receptors reexamined: mouse LECAM-1 (MEL-14 antigen) is involved in lymphocyte migration into gut-associated lymphoid tissue. *Eur J Immunol* 1991;21 (12): 2925-9.
26. Bargatze RF, Jutila MA, and Butcher EC Distinct roles of L-selectin and integrins alpha 4 beta 7 and LFA-1 in lymphocyte homing to Peyer's patch-HEV in situ: the multistep model confirmed and refined. *Immunity* 1995;3 (1): 99-108.
27. Schon MP, Arya A, Murphy EA, et al. Mucosal T lymphocyte numbers are selectively reduced in integrin alpha E (CD103)-deficient mice. *J Immunol* 1999;162 (11): 6641-9.
28. Wysocka J, Hassmann E, Lipska A, et al. Naive and memory T cells in hypertrophied adenoids in children according to age. *Int J Pediatr Otorhinolaryngol* 2003;67 (3): 237-41.

29. Akbar AN, Terry L, Timms A, et al. Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. *J Immunol* 1988;140 (7): 2171-8.
30. Testi R, Phillips JH, and Lanier LL T cell activation via Leu-23 (CD69). *J Immunol* 1989;143 (4): 1123-8.
31. Zeitz M, Schieferdecker HL, Ullrich R, et al. Phenotype and function of lamina propria T lymphocytes. *Immunol Res* 1991;10 (3-4): 199-206.
32. Weiner HL Oral tolerance: immune mechanisms and the generation of Th3-type TGF-beta-secreting regulatory cells. *Microbes Infect* 2001;3 (11): 947-54.
33. Neurath MF, Finotto S, and Glimcher LH The role of Th1/Th2 polarization in mucosal immunity. *Nat Med* 2002;8 (6): 567-73.
34. Moser M and Murphy KM Dendritic cell regulation of TH1-TH2 development. *Nat Immunol* 2000;1 (3): 199-205.
35. Croft M and Swain SL B cell response to fresh and effector T helper cells. Role of cognate T-B interaction and the cytokines IL-2, IL-4, and IL-6. *J Immunol* 1991;146 (12): 4055-64.
36. Stout RD and Bottomly K Antigen-specific activation of effector macrophages by IFN-gamma producing (TH1) T cell clones. Failure of IL-4-producing (TH2) T cell clones to activate effector function in macrophages. *J Immunol* 1989;142 (3): 760-5.
37. Henkart PA Lymphocyte-mediated cytotoxicity: two pathways and multiple effector molecules. *Immunity* 1994;1 (5): 343-6.
38. Chen YH and Weiner HL Dose-dependent activation and deletion of antigen-specific T cells following oral tolerance. *Ann N Y Acad Sci* 1996;778 111-21.
39. Friedman A Induction of anergy in Th1 lymphocytes by oral tolerance. Importance of antigen dosage and frequency of feeding. *Ann N Y Acad Sci* 1996;778 103-10.
40. Shevach EM Certified professionals: CD4(+)CD25(+) suppressor T cells. *J Exp Med* 2001;193 (11): F41-6.
41. Nakamura K, Kitani A, Fuss I, et al. TGF-beta 1 plays an important role in the mechanism of CD4+CD25+ regulatory T cell activity in both humans and mice. *J Immunol* 2004;172 (2): 834-42.
42. Makita S, Kanai T, Oshima S, et al. CD4+CD25bright T cells in human intestinal lamina propria as regulatory cells. *J Immunol* 2004;173 (5): 3119-30.
43. Maul J, Loddenkemper C, Mundt P, et al. Peripheral and intestinal regulatory CD4+ CD25(high) T cells in inflammatory bowel disease. *Gastroenterology* 2005;128 (7): 1868-78.

44. Tsuji NM, Mizumachi K, and Kurisaki J Interleukin-10-secreting Peyer's patch cells are responsible for active suppression in low-dose oral tolerance. *Immunology* 2001.103 (4): 458-64.
45. Spahn TW, Fontana A, Faria AM, et al. Induction of oral tolerance to cellular immune responses in the absence of Peyer's patches. *Eur J Immunol* 2001.31 (4): 1278-87.
46. Mowat AM and Weiner HL Oral Tolerance: physiological basis and clinical applications. In: Bienenstock J. *Handbook of Mucosal Immunology*. 2 ed London, England: Academic Press Inc. 1999:587-617.
47. Newberry RD and Lorenz RG Organizing a mucosal defense. *Immunol Rev* 2005.206 6-21.
48. Makala LH, Suzuki N, and Nagasawa H Peyer's patches: organized lymphoid structures for the induction of mucosal immune responses in the intestine. *Pathobiology* 2002.70 (2): 55-68.
49. McCracken VJ and Lorenz RG The gastrointestinal ecosystem: a precarious alliance among epithelium, immunity and microbiota. *Cell Microbiol* 2001.3 (1): 1-11.
50. Kanamori Y, Ishimaru K, Nanno M, et al. Identification of novel lymphoid tissues in murine intestinal mucosa where clusters of c-kit+ IL-7R+ Thy1+ lympho-hemopoietic progenitors develop. *J Exp Med* 1996.184 (4): 1449-59.
51. MacDonald TT, Spencer J, Viney JL, et al. Selective biopsy of human Peyer's patches during ileal endoscopy. *Gastroenterology* 1987.93 (6): 1356-62.
52. Nagata S, McKenzie C, Pender SL, et al. Human Peyer's patch T cells are sensitized to dietary antigen and display a Th cell type 1 cytokine profile. *J Immunol* 2000.165 (9): 5315-21.
53. Monteleone G, Holloway J, Salvati VM, et al. Activated STAT4 and a functional role for IL-12 in human Peyer's patches. *J Immunol* 2003.170 (1): 300-7.
54. Farstad IN, Halstensen TS, Lien B, et al. Distribution of beta 7 integrins in human intestinal mucosa and organized gut-associated lymphoid tissue. *Immunology* 1996.89 (2): 227-37.
55. Farstad IN, Halstensen TS, Fausa O, et al. Do human Peyer's patches contribute to the intestinal intraepithelial gamma/delta T-cell population? *Scand J Immunol* 1993.38 (5): 451-8.
56. Resendiz-Albor AA, Esquivel R, Lopez-Revilla R, et al. Striking phenotypic and functional differences in lamina propria lymphocytes from the large and small intestine of mice. *Life Sci* 2005.76 (24): 2783-803.
57. McNeill L, Cassady RL, Sarkardei S, et al. CD45 isoforms in T cell signalling and development. *Immunol Lett* 2004.92 (1-2): 125-34.

58. Janossy G, Tidman N, Papageorgiou ES, et al. Distribution of t lymphocyte subsets in the human bone marrow and thymus: an analysis with monoclonal antibodies. *J Immunol* 1981;126 (4): 1608-13.
59. Johnson GD, Ling NR, Nathan PD, et al. Use of monoclonal antibodies reactive with secretory epithelial cells for the immunocytochemical identification of plasma cells. *Immunol Lett* 1986;12 (5-6): 295-300.
60. MacDonald TT and Monteleone G IL-12 and Th1 immune responses in human Peyer's patches. *Trends Immunol* 2001;22 (5): 244-7.
61. Fujihashi K and McGhee JR Mucosal immunity and tolerance in the elderly. *Mech Ageing Dev* 2004;125 (12): 889-98.
62. Smith PD, Janoff EN, Mosteller-Barnum M, et al. Isolation and purification of CD14-negative mucosal macrophages from normal human small intestine. *J Immunol Methods* 1997;202 (1): 1-11.
63. Zhou LJ and Tedder TF Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. *J Immunol* 1995;154 (8): 3821-35.
64. McWilliams M, Lamm ME, and Phillips-Quagliata JM Surface and intracellular markers of mouse mesenteric and peripheral lymph node and Peyer's patch cells. *J Immunol* 1974;113 (4): 1326-33.
65. Lyscom N and Brueton MJ Intraepithelial, lamina propria and Peyer's patch lymphocytes of the rat small intestine: isolation and characterization in terms of immunoglobulin markers and receptors for monoclonal antibodies. *Immunology* 1982;45 (4): 775-83.
66. Garside P, Ingulli E, Merica RR, et al. Visualization of specific B and T lymphocyte interactions in the lymph node. *Science* 1998;281 (5373): 96-9.
67. Tseng J Induction of indigenous immunoglobulin-containing cells by gamma-irradiation and lymphoid cell transfer in mice. *J Immunol* 1981;127 (5): 2140-6.
68. Stashenko P, Nadler LM, Hardy R, et al. Expression of cell surface markers after human B lymphocyte activation. *Proc Natl Acad Sci U S A* 1981;78 (6): 3848-52.
69. Banchereau J and Rousset F Human B lymphocytes: phenotype, proliferation, and differentiation. *Adv Immunol* 1992;52: 125-262.
70. Laichalk LL and Thorley-Lawson DA Terminal differentiation into plasma cells initiates the replicative cycle of Epstein-Barr virus in vivo. *J Virol* 2005;79 (2): 1296-307.
71. Schieferdecker HL, Ullrich R, Hirselund H, et al. T cell differentiation antigens on lymphocytes in the human intestinal lamina propria. *J Immunol* 1992;149 (8): 2816-22.

72. Kelsall BL and Strober W The role of dendritic cells in antigen processing in the Peyer's patch. *Ann N Y Acad Sci* 1996;778:47-54.
73. Salmi M, Andrew DP, Butcher EC, et al. Dual binding capacity of mucosal immunoblasts to mucosal and synovial endothelium in humans: dissection of the molecular mechanisms. *J Exp Med* 1995;181(1):137-49.
74. Tedder TF, Penta AC, Levine HB, et al. Expression of the human leukocyte adhesion molecule, LAM1. Identity with the TQ1 and Leu-8 differentiation antigens. *J Immunol* 1990;144(2):532-40.
75. Munro JM, Briscoe DM, and Tedder TF Differential regulation of leucocyte L-selectin (CD62L) expression in normal lymphoid and inflamed extralymphoid tissues. *J Clin Pathol* 1996;49(9):721-7.
76. Seibold F, Seibold-Schmid B, Cong Y, et al. Regional differences in L-selectin expression in murine intestinal lymphocytes. *Gastroenterology* 1998;114(5):965-74.
77. Gallatin WM, Weissman IL, and Butcher EC A cell-surface molecule involved in organ-specific homing of lymphocytes. *Nature* 1983;304(5921):30-4.
78. Arbones ML, Ord DC, Ley K, et al. Lymphocyte homing and leukocyte rolling and migration are impaired in L-selectin-deficient mice. *Immunity* 1994;1(4):247-60.
79. Testi R, Phillips JH, and Lanier LL Leu 23 induction as an early marker of functional CD3/T cell antigen receptor triggering. Requirement for receptor cross-linking, prolonged elevation of intracellular [Ca⁺⁺] and stimulation of protein kinase C. *J Immunol* 1989;142(6):1854-60.
80. Hara T, Jung LK, Bjorndahl JM, et al. Human T cell activation. III. Rapid induction of a phosphorylated 28 kD/32 kD disulfide-linked early activation antigen (EA 1) by 12-o-tetradecanoyl phorbol-13-acetate, mitogens, and antigens. *J Exp Med* 1986;164(6):1988-2005.
81. Lanier LL, Buck DW, Rhodes L, et al. Interleukin 2 activation of natural killer cells rapidly induces the expression and phosphorylation of the Leu-23 activation antigen. *J Exp Med* 1988;167(5):1572-85.
82. Cebrian M, Yague E, Rincon M, et al. Triggering of T cell proliferation through AIM, an activation inducer molecule expressed on activated human lymphocytes. *J Exp Med* 1988;168(5):1621-37.
83. Takahata Y, Nomura A, Takada H, et al. CD25+CD4+ T cells in human cord blood: an immunoregulatory subset with naive phenotype and specific expression of forkhead box p3 (Foxp3) gene. *Exp Hematol* 2004;32(7):622-9.
84. Choy MY, Walker-Smith JA, Williams CB, et al. Differential expression of CD25 (interleukin-2 receptor) on lamina propria T cells and macrophages in the intestinal lesions in Crohn's disease and ulcerative colitis. *Gut* 1990;31(12):1365-70.

85. Springer TA Traffic signals on endothelium for lymphocyte recirculation and leukocyte emigration. *Annu Rev Physiol* 1995.57 827-72.
86. Springer TA Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* 1994.76 (2): 301-14.
87. Wagner N, Lohler J, Kunkel EJ, et al. Critical role for beta7 integrins in formation of the gut-associated lymphoid tissue. *Nature* 1996.382 (6589): 366-70.
88. Salmi M and Jalkanen S Lymphocyte homing to the gut: attraction, adhesion, and commitment. *Immunol Rev* 2005.206 100-13.
89. Ceppek KL, Shaw SK, Parker CM, et al. Adhesion between epithelial cells and T lymphocytes mediated by E-cadherin and the alpha E beta 7 integrin. *Nature* 1994.372 (6502): 190-3.
90. Lehmann J, Huehn J, de la Rosa M, et al. Expression of the integrin alpha Ebeta 7 identifies unique subsets of CD25+ as well as CD25- regulatory T cells. *Proc Natl Acad Sci U S A* 2002.99 (20): 13031-6.
91. Cerf-Bensussan N, Jarry A, Brousse N, et al. A monoclonal antibody (HML-1) defining a novel membrane molecule present on human intestinal lymphocytes. *Eur J Immunol* 1987.17 (9): 1279-85.
92. Kilshaw PJ and Murant SJ Expression and regulation of beta 7(beta p) integrins on mouse lymphocytes: relevance to the mucosal immune system. *Eur J Immunol* 1991.21 (10): 2591-7.
93. Hadley GA, Bartlett ST, Via CS, et al. The epithelial cell-specific integrin, CD103 (alpha E integrin), defines a novel subset of alloreactive CD8+ CTL. *J Immunol* 1997.159 (8): 3748-56.
94. Stassen M, Fondel S, Bopp T, et al. Human CD25+ regulatory T cells: two subsets defined by the integrins alpha 4 beta 7 or alpha 4 beta 1 confer distinct suppressive properties upon CD4+ T helper cells. *Eur J Immunol* 2004.34 (5): 1303-11.
95. Beagley KW, Fujihashi K, Lagoo AS, et al. Differences in intraepithelial lymphocyte T cell subsets isolated from murine small versus large intestine. *J Immunol* 1995.154 (11): 5611-9.
96. Boll G and Reimann J Lamina propria T cell subsets in the small and large intestine of euthymic and athymic mice. *Scand J Immunol* 1995.42 (2): 191-201.
97. Boll G, Rudolphi A, Spiess S, et al. Regional specialization of intraepithelial T cells in the murine small and large intestine. *Scand J Immunol* 1995.41 (2): 103-13.

98. Ernst PB, Clark DA, Rosenthal KL, et al. Detection and characterization of cytotoxic T lymphocyte precursors in the murine intestinal intraepithelial leukocyte population. *J Immunol* 1986;136 (6): 2121-6.
99. Barrett TA, Gajewski TF, Danielpour D, et al. Differential function of intestinal intraepithelial lymphocyte subsets. *J Immunol* 1992;149 (4): 1124-30.
100. Hayday A, Theodoridis E, Ramsburg E, et al. Intraepithelial lymphocytes: exploring the Third Way in immunology. *Nat Immunol* 2001;2 (11): 997-1003.
101. Arstila T, Arstila TP, Calbo S, et al. Identical T cell clones are located within the mouse gut epithelium and lamina propria and circulate in the thoracic duct lymph. *J Exp Med* 2000;191 (5): 823-34.