

7. Literatur

- Akdis CA, Akdis M, Trautmann A et al.: Immune regulation in atopic dermatitis. *Curr Opin Immunol* 2000; 12(6): 641-46.
- Akdis M, Trautmann A, Klunker S et al.: T helper 2 predominance in atopic diseases is due to preferential apoptosis of circulating memory/effector Th1 cells. *FASEB* 2003; 17: 1026-35.
- Antonelli A, Bianchi M, Crinelli R et al.: Modulation of ICAM-1 expression in ECV304 cells by macrophage-released cytokines. *J Immunol* 2001; 27(6): 978-91.
- Apostolou I, Sarukhan A, Klein L: Origin of regulatory T cells with known specificity for antigen. *Nat Immunol* 2002; 3: 756-63.
- Azuma M, Ito D, Yagita H et al.: The B70 antigen is a second ligand for CTLA-4 and CD28. *Nature* 1993; 366: 76-9.
- Babu KS, Belgı G: Management of cutaneous drug reactions. *Curr Allergy Asthma Rep* 2002 Jan; 2(1): 26-33.
- Bancherau J, Steinman R: Dendritic cells and the control of immunity. *Nature* 1998; 329(19): 245-252.
- Barbaud AM, Bene MC, Reichert-Penetrat S et al.: Immunocompetent cells and adhesion molecules in 14 cases of cutaneous drug reactions induced with the use of antibiotics. *Arch Dermatol* 1998; 134: 1040-1.
- Barbaud AM, Bene MC, Schmutz JL et al.: Role of delayed cellular hypersensitivity and adhesion molecules in amoxicillin-induced morbilliform rashes. *Arch Dermatol* 1997; 133: 481-6.
- Barbaud AM, Reichert-Penetrat S, Trechot P et al.: The use of skin testing in the investigation of cutaneous adverse drug reactions. *Br J Dermatol* 1998; 139:49-58.
- Baskar S, Glimcher L, Nabavi N et al.: MHC class II+ B7-1+ tumor cells are potent vaccines for stimulation tumor rejection in tumor-bearing mice. *J Exp Med* 1995; 181(2): 619-29.
- Becker S, Daniel E: Antagonistic and additive effects of IL-4 and IFN- γ on human monocytes and macrophages. *Cell Immunol* 1990; 129, 351-62.
- Beissert S, Hosoi J, Grabbe S et al.: IL-10 inhibits tumor antigen presentation by epidermal antigenpresenting cells. *J Immunol* 1995; 154: 2668-77.

- Berg E, Robinson M: The human peripheral lymph node vascular addressin is a ligand for LECAM-1, the peripheral lymph node homing receptor. *J Cell Biol* 1991; 108: 2435-47.
- Bevilacqua MP, Nelson RM, Warnock RA: Selectins. *J Clin Invest* 1993; 91: 379-87.
- Birk RW, Gratchev A, Hakki N et al.: Alternative activation of antigen-presenting cells: concepts and clinical relevance. *Hautarzt* 2001; 52(3): 193-200.
- Bordet R, Gautier S, Le Lovet M et al.: Analysis of the direct cost of adverse drug reactions in hospitalized patients. *Eur J Clin Pharmacol* 2001; 56: 935-41.
- Bork K: Unerwünschte Arzneimittelreaktionen an der Haut. *Arzneimitteltherapie* 2000; 4: 109.
- Brasch J, Burgard J, Sterry W: Common pathogenetic pathways in allergic and irritant dermatitis. *J Invest Dermatol* 1992; 98(2): 166-70.
- Breathnach S.: Mechanisms of drug eruptions. *Austral J Dermatol* 1995; 36:121-7.
- Büdinger L, Hertl M: The skin as target for IgE-mediated allergic reactions. *Allergy* 2000; 55(2): 108.
- Chen C, Nabavi N: In vitro induction of T cell anergy by blocking B7 and early T cell costimulatory molecule ETC-1/b7-2. *Immunity* 1994; 1: 147-54.
- Coffman RL, Chatelain R, Leal LM et al.: *Leishmania major* infection in mice: a model system for study of CD4+ T cell subset differentiation. *Res Immunol* 1991; 142: 36-9.
- Dascalau DI, Klatter Y, Baretz M et al.: Langerhans' cell distribution in drug eruption. *Acta Derm Venerol* 1992; 72: 175-7.
- Del Prete GF, De Carli M, Ricci M et al.: Helper activity for immun-globulin synthesis of T helper type 1 (Th1) and (Th2) human T cell clones: the help of Th1 clones is limited by their cytolytic capacity. *J Exp Med* 1991; 174: 809-13.
- Del Prete GF: Human Th1 and Th2 lymphocytes: their role in the pathophysiology of atopy. *Allergy* 1992; 47: 450-5.
- Denis M.: Involvement of cytokines in determining resistance and acquired immunity in murine tuberculosis. *J Leu Biol* 1991; 50: 495- 501.
- Detmar M: Mechanismen der Interaktion von Leukozyten und dermalen Endothelzellen in der kutanen Entzündung. *Hautarzt* 1992; 43:679-86.
- Dupperay A, Mantovani A: Endothelial cell regulation of leukocyte infiltration in inflammatory tissues. *Med Inflam* 1995; 4: 322-30.

- Enk A, Saloga J: Induction of hapten-specific tolerance by IL-10 in vivo. *J Exp Med* 1994; 179: 1397-1402.
- Fitch FW, Stack R, Fields P et al: Regulation of T lymphocyte subsets. *Ciba Found Symp* 1995; 195: 68-85.
- Friedmann PS, Strickland I, Memon AA et al.: Early time course of recruitment of immune surveillance in human skin. *Clin Exp Immunol* 1993; 91(3): 351-6.
- Fries JW, Williams AJ: Expression of VCAM-1 and E-Selectin in an in vivo model of endothelial activation. *Am J Pathol* 1993; 143: 725-737.
- Fuccao T, Satoshi M, Shigeo K: Synergistic effects of IL-4 and IL-8 on IL-12-dependent IFN- γ production by dendritic cells. *J Immunol* 2000; 164: 64-71.
- Fuhlbridge RC, Kieffer JD, Armerding D et al.: Cutaneous lymphocyte antigen is a specialised form of PSGL-1 expressed on skin homing T cells. *Nature* 1997; 389: 978-981.
- Gajewski TF, Lancki DW, Stack R et al.: ‘Anergy’ of Th0 helper T lymphocytes induces downregulation of Th1 characteristics and a transition to a Th2-like phenotype. *J Exp Med* 1994; 179: 481-91.
- Gawkrodger DJ, McVittie E, Carr MM et al.: Phenotypic characterisation of the early cellular responses in allergic and irritant contact dermatitis. *Clin Exp Immunol* 1986; 66: 590-598.
- Gawkrodger DJ, McVittie E, Hunter JA et al.: Immunophenotyping of the eczematous flare-up reaction in a nickel-sensitive subject. *Deramtologica* 1987; 175: 171-177.
- Goebeler M, Meinardus-Hager G, Roth J et al.: Nickel chloride and cobalt chloride, two common contact sensitizers, directly induce expression of intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leucocyte adhesion molecule (ELAM-1) by endothelial cells. *J Invest Dermatol* 1993; 100: 759-65.
- Goerdt S, Walsh L, Murphy GF et al.: Identification of a novel high molecular weight protein prefentially expressed by sinusoidal endothelial cells in normal human tissues. *J Cell Biol* 1991; 113(6): 1425-37.
- Goerdt S, Kodelja V, Schnuth M et al.: The mononuclear phagocyte-dendritic cell dichotomy: myths, facts and a revised concept. *Clin Exp Immunol* 1996; 105: 1-9.

- Goerdt S, Orfanos C: Other Functions, other Gens: Alternative activation of antigen-presenting cells. *Immunity* 1999; 10: 137-42.
- Gonzalez F, Leyva L, Posadas S et al.: Participation of T lymphocytes in cutaneous allergic reactions to drugs. *Clin Exp Allergy* 1998; 28(4): 3-6.
- Grabbe S, Varga G, Beissert S et al.: β 1-Integrins are required for skin homing of primed T cells but not for priming naïve T cells. *J Clin Invest* 2002; 109: 183-192.
- Gratschev A, Schledzewski K, Guillot P et al.: Alternatively activated antigen-presenting cells. *Skin Pharmacol Appl Skin Physiol* 2001; 14: 272-79.
- Griffiths CE: The immunological basis of psoriasis. *J Invest Dermatol* 1999; 7: 354-9.
- Groves RW, Allen MH, Barker JN et al.: Endothelial leucocyte adhesion molecule-1 (ELAM-1) expression in cutaneous inflammation. *Br J Dermatol* 1991; 124: 117-23.
- Hallas J, Gram LF, Grodun E et al.: Drug related admissions to medical wards. *Brit J Clin Pharmacol* 1992; 33: 61-8.
- Hartmann K, Kuhn J, Gatmann J et al.: Spontaneous reporting system for the assessment of new, unknown and rare undesirable effects of drugs. *Schweiz Med Wschr* 1992; 122: 1409-13.
- Hinden M, Christensen OB: Delayed hypersensitivity reactions following allergic and irritant inflammation. *Acta Derm Venereol* 1992; 72(3): 220-1.
- Hogg N.: Roll, roll, roll your leucocyte gently down the vein. *Immunol today* 1992; 13: 113-5.
- Horn TD, Burke PJ, Karp JE et al.: Intravenous administration of recombinant human granulocyte-macrophage colony-stimulating factor causes a cutaneous eruption. *Arch Dermatol* 1991; 127: 49-52.
- Hunyadi J, Simon M, Kenderessy AS et al.: Expression of monocyte/macrophage markers (CD13, CD14, CD68) on human keratinocytes in healthy and diseased skin. *J Dermatol* 1993; 20(6): 341-5.
- Hynes OH: Integrins: bidirectional, allosteric signaling machines. *Cell* 2002; 110: 673-87.
- Inaba K., Inaba M. et al.: Granulocytes macrophages and dendritic cells arise from a common major histocompatibility complex class II-negative progenitor in mouse bone marrow. *J Immunol* 1993; 90: 3039-42.

- Ives TJ, Bentz EJ, Gwyther RE et al.: Dermatologic adverse drug reactions in a family medicine setting. *Arch Fam Med* 1992; 1: 241-45.
- Kaburagi Y, Hasegawa M, Nagaoka T: The cutaneous reverse arthus reaction requires intercellular adhesion molecule 1 and L-Selectin expression. *J Immunol* 2002; 168: 2970-8.
- Kalinski P, Schuitemaker JH, Hilkens CM et al.: Final maturation of dendritic cells is associated with impaired responsiveness to IFN- γ and to bacterial IL-12 inducers. *J Immunol* 1999; 162(6): 3231-6.
- Kalish RS: Recent developments in the pathogenesis of allergic contact dermatitis. *Arch Dermatol* 1991; 127: 1558-63.
- Kapsenberg ML, Hilkens C, Wierenga E et al.: The paradigm of type 1 and type 2 antigen-presenting cells. *Clin Exp Allergy* 1999; 29(2): 33-36.
- Kapsenberg ML, Wierenga E et al.: Functional subsets of allergen-reactive human CD4+ T-cells. *Immunol Today* 1991; 12: 392-5.
- Kapsenberg ML, Hilkens CM, van Der Pouw et al.: Atopic allergy: a failure of antigen-presenting cells to properly polarize helper T cells? *Am J Respir Crit Care Med*. 2000 Sep; 162: 76-80.
- Kaufmann R, Weber L, Klein CE et al.: Integrine - neue Rezeptormoleküle: ihre Bedeutung für die Differenzierung und Immunantwort der Haut. *Hautarzt* 1990; 41: 265-61.
- Klein CE: Adhäsionsrezeptoren. *Hautarzt* 1994; 45:263-84.
- Knowles S, Shapiro L, Shear NH: Serious dermatologic reactions in children. *Curr Opin Pediatr* 1997; 9:388-395.
- Kodelja V, Goerdt S: Dissection of macrophage differentiation pathways in cutaneous macrophage disorders and *in vitro*. *Exp Dermatol* 1994; 3: 257-68.
- Kodelja V, Müller C, Tenorio S et al.: Differences in angiogenetic potential of classically vs alternatively activated macrophages. *Immunobiology* 1997; 197: 478-93.
- Kuchroo VK, Das MP, Brown JA et al.: B7-1 and B7-2 costimulatory molecules activate differentially the Th1/Th2 developmental pathways: application to autoimmune disease therapy. *Cell* 1995; 80:707-18.

- Kumasaka T, Doyle NA, Quinlan WM et al.: Role of CD11/CD18 in neutrophil emigration during acute and recurrent *Pseudomonas aeruginosa*-induced pneumonia in rabbits. Am J Pathol 1996; 148: 1297-1305.
- Lasky LA: Selectins: Interpreters of cell specific carbohydrate information during inflammation. Science 1992, 258: 964-69.
- Lawley TJ, Caughman SW, Swerlick RA et al.: The role of adhesions molecules in cutaneous inflammation. J Dermatol 1994; 21: 790- 4.
- Lazarou J, Pomeranz BH, Correy PN: Incidence of adverse drug reactions in hospitalized patients. JAMA 1998; 279:1200-1205.
- Leung DY, Bhan AK, Schneeberger EE et al.: Characterization of mononuclear cell infiltrate in atopic dermatitis using monoclonal antibodies. J Allergy Clin Immunol 1983; 71: 47-55.
- Lewis RE, Buchsbaum M: Intercellular adhesion molecule expression in the evolving human cutaneous delayed hypersensitivity reaction. J Invest Dermatol 1989; 93: 672-7.
- Lingnau K, Hoehn P, Kerdine S et al.: IL-4 in combination with TGF- favors an alternative pathway Th1 development independent of IL-12. J Exp Med 1998; 188: 2175.
- Little MC, Metcalfe RA, Haycock JW et al.: The participation of proliferative keratinocytes in the preimmune response to sensitising agents. Br J Dermatol 1998; 138: 45-56.
- Lowitt MH, Shear NH: Pharmacogenomics and dermatological therapeutics. Arch Dermatol 2001; 137: 1512-4.
- Martin KH, Slack JK, Scott AB et al.: Integrin connections map: to infinity and beyond. Science 2002; 296: 1652-3.
- Mitchell AA, Lacouture P, Sheehan JE et al.: Adverse drug reactions in children leading to hospital admission. Pediatrics 1988; 82: 24-9.
- Mizgerd JP, Horwitz BH, Quillen HC et al.: Effects of CD18 deficiency on the emigration of murine neutrophils during pneumonia. J Immunol 1999; 163: 995-999.
- Mockenhaupt M, Schopf E: Epidemiology of drug-induced severe skin reactions. Semin Cutan Med Surg 1996; 15(4): 236-43.

- Modolell M, Corraliza I, Link F et al.: Reciprocal regulation of the nitric oxide synthase/arginase balance in mouse bone marrow macrophages by Th1 and Th2 cytokines. *Eur J Immunol* 1995; 25: 1101-04.
- Moser M: Regulation of Th1/Th2 development by antigen-presenting cells in vivo. *Immunobiology* 2001; 204: 551-557.
- Mosmann TR, Moore KW: The role of IL-10 in cross-regulation of Th1 and Th2 responses. *Immunol today* 1991; 12: A49-53.
- Mosser D, Handman E: Treatment of murine macrophages with interferon- γ inhibits their ability to bind leishmania promastigotes. *J Leukoc Biol* 1992; 52: 369-76.
- Narvaez D, Kanitakis J, Faure D et al.: Immunohistochemical study of CD34-positive dendritic cells of human dermis. *Am J Dermatopathol* 1996; 18(3): 283-8.
- Novak N, Kruse S, Kraft S et al.: Dichotomic nature of atopic dermatitis. *J Invest Dermatol* 2002; 119: 870-875.
- Ohki O, Yokozeki H, Katayama I et al.: Functional CD86 (B7-2/B70) is predominantly expressed on Langerhans cells in atopic dermatitis. *Br J Dermatol* 1997; 136: 838-845.
- Orlikowsky T, Wang ZQ, Duhane A et al.: The cell surface marker phenotype of macrophages. *J Interferon Cytokine Res* 1996; 16(11): 957-62.
- Osborn L: Leukocyte adhesion to endothelium in inflammation. *Cell* 1990; 62: 3-6.
- Paul W, Seder R: Lymphocyte responses and cytokines. *Cell* 1994; 76: 241-51.
- Pichler WJ: Medikamentenallergien. *Wien Klin Wochenschr* 1993; 105/22:
- Poppema S, Bhan AK, Reinherz EL et al.: Distribution of T-cell subsets in human lymph nodes. *J Exp Med* 1981; 153: 30-41.
- Probst P, Küntzlin D, Fleischer B: Th2-type infiltrating T cells in nickel-induced contact dermatitis. *Cell Immunol* 1995; 165: 134-40.
- Prussick R: Adverse cutaneous reactions to chemotherapeutic agents and cytokine therapy. *Semin Cutan Med Surg* 1996; 15(4): 267-76.
- Radi Z, Kehrli M, Ackermann MR: Cell adhesion molecules, leucocyte trafficking and strategies to reduce leucocyte infiltration. *J Vet Intern Med* 2001, 15: 516-529.
- Ranki A, Kanerva L, Forstrom L et al.: T and B lymphocytes, macrophages and langerhans cells during the course of contact allergic and irritant skin reactions in man. *Acta Derm Venereol* 1983; 63: 376-383.

- Reid CD, Stackpoole A, Meager A et al.: Interactions of tumor necrosis factor with granulocyte-macrophage colony-stimulating factor and other cytokines in the regulation of dendritic cell growth in vitro from early bipotent CD34+ progenitors in human bone marrow. *J Immunol* 1992; 149: 2681-8.
- Rocha C, de Maubeuge J, Sarfati M et al.: Characterization of cellular infiltrates in skin lesions of atopic eczema by means of monoclonal antibodies. *Dermatologica* 1984; 169: 330-338.
- Rudikoff D, Lebwohl M: Atopic dermatitis. *Lancet* 1998; 351: 1715-1721.
- Rutherford MS, Schook LB: Differential immunocompetence of macrophages derived using macrophage or granulocyte-macrophage colony-stimulating factor. *J Leukoc Biol* 1992; 51: 69-76
- Rutherford MS, Witsell A, Schook LB: Mechanisms generating functionally heterogeneous macrophages: chaos revisited. *J Leukoc Biol* 1993; 53: 602-18.
- Sacerdoti G, Vozza A, Ruocco V et al.: Identifying drug reactions to drugs. *Int J Dermatol* 1993; 32(7): 469-79.
- Sallusto F, Lanzavecchia A: Efficient presentation of soluble antigen by cultured human dendritic cells in maintained by granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor- α . *J Exp Med* 1994; 179: 1109-18.
- Santiago-Schwarz F, Belilos E, Diamond B et al.: THF in combination with GM-CSF enhances the differentiation of neonatal cord blood stem cells into dendritic cells and macrophages. *J Leukoc Biol* 1992; 52: 274-81.
- Sasaki M, Namioka Y, Ito T et al.: Role of ICAM-1 in the aggregation and adhesion of HMA in response to TNF- α and IFN- γ . *Mediators Inflamm* 2001; 10, 309-313.
- Schönemann KW: Adverse drug reaction on the skin. *J Immunol* 1998; 25: 576.
- Shimizu Y, Newman W, Gopal TV et al.: Four molecular pathways of T cell adhesion to endothelial cells: roles of LFA-1, VCAM-1 and ELAM-1 and changes in pathway hierarchy under different activation conditions. *J Cell Biol* 1991; 113(5): 1203-12.
- Shimizu Y, Newman W, Tamaka Y et al.: Lymphocyte interactions with endothelial cells. *Immunol today* 1992; 13: 106-12.
- Simon Jr. M, Hunyadi J, Gomez RS: Macrophage markers 25F9 and 27E10 on human keratinocytes in normal and diseased skin. *J Dermatol* 1993; 20: 618-22.

- Smith JW, Seidl LG, Cluff LE: Studies on the epidemiology of adverse drug reactions. *Annals Intern Med* 1966; 65(4): 629.
- Smits HH, de Jong EC, Schnitemaker JH: Intercellular adhesion molecule-1/LFA-1 ligation favors human Th1 Development. *J Immunol* 2002; 168(4): 1710-16.
- Spertini O, Kansas GS, Munro JM et al.: Regulation of leucocyte migration by activation of the leucocyte adhesion molecule-1 selectin. *Nature* 1991; 349(6311): 691-4.
- Stein M, Keshav S, Harris N et al.: Interleukin-4 potently enhances murine makrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. *J Exp Med* 1992; 176: 287-92.
- Szepietowski JC, McKenzie RC, Keohane SG et al.: Atopic and nonatopic individuals react to nickel challenge in a similar way. *Dermatol* 1997; 137: 195-200.
- Tedder TF, Steeber DA, Chen A et al.: The selectins: Vascular adhesions molecules. *FASEB J* 1995; 9: 866-73.
- Thompson C: Distinct roles for the costimulatory ligands B7-1 and B7-2 in T helper cell differentiation? *Cell* 1995; 81: 979-82.
- Trautmann A, Akdis M, Bröcker E: New insights into the role of T-cells in atopic dermatitis and allergic contact dermatitis. *Trends Immunol* 2001; 22(10): 530-32.
- Valks R, Vargas E Munoz E et al.: Dermal infiltrate of enlarged macrophages in patients receiving chemotherapy. *J Cutan Pathol* 1998; 25: 259-64.
- Van de Stolpe A, van der Saag P: Intercellular adhesions molecule-1. *J Mol Med* 1996; 74: 13-33.
- Vervloet D, Durham S: Adverse reactions to drugs. *BMJ* 1998; 316: 1511- 14.
- Vestweber D, Blanks JE: Mechanisms that regulate the function of the selectins and their ligands. *Physiol Rev* 1999; 56: 181-213.
- Villaneuva PO, Harris TS, Ricklan PE et al.: Macrophages from schistosomal egg granulomas induce unresponsiveness in specific cloned TH-1 lymphocytes in vitro and down-regulate schistosomal granulomatous disease in vivo. *J Immunol* 1994; 152: 1847-55.
- Weck de A: Pharmacologic and immunochemical mechanisms of drug hypersensitivity. *Immunol Allergol Clin N Am* 1991; 11: 461-74.

- Weyl A, Vanscheidt W, Weiss JM et al.: Expression of the adhesion molecules ICAM-1, VCAM-1, and E-Selectin and their ligands VLA-4 and LFA-1 in chronic venous leg ulcers. *J Am Acad Dermatol* 1996; 34: 418-23.
- Wiktor-Jedrzejczak, Ansari A, Szperl M et al.: Distinct in vivo functions of two macrophage subpopulations as evidenced by studies using macrophage-deficient op/op mouse. *Eur J Immunol* 1992; 22: 1951-4.
- Williams ME, Chang TL, Burke SK et al.: Activation of functionally distinct subsets of CD4+ T lymphocytes. *Res Immunol* 1991; 142: 23-7.
- Yawalkar N, Pichler WJ: Pathogenesis of drug-induces exanthema. *Int Arch Allergy Immunol* 2001; 124: 336-8.