
Quellen

- [1] Addington J, Addington D (1998)
Visual attention and symptoms in schizophrenia: A 1 year follow-up.
Schizophr Res 34: 95-99
- [2] Aleman A, Hijman R, De Haan E, Kahn R (1999)
Memory impairment in schizophrenia: A meta-analysis.
Am J Psychiatry 156: 1358-1366
- [3] Beblo T, Herrmann M (2000)
Neuropsychologische Defizite bei depressiven Störungen.
Fortschr Neurol Psychiatrie 68: 1-11
- [4] Benkert O, Hippius H (2001)
Kompendium der psychiatrischen Pharmakotherapie.
Springer, Berlin – Heidelberg – New York, 3. Auflage, S 127-137
- [5] Brébion G, Amadar X, Smith MJ, Malaspia D, Sharif Z, Gorman J (1999)
Opposite links of positive and negative symptomatology with memory errors in schizophrenia.
Psychiatry Res 88: 15-24
- [6] Bryson G, Pill M, Kaplan E, Greig T (1998)
The functional consequences of memory impairment on initial work performance in people with schizophrenia.
J Nerv Ment Dis 186: 610-615
- [7] Campbell MJ (1990)
Medical statistics. A common-sense approach.
J Wiley and Sons, Chichester

-
- [8] Clementz B, Geyer M, Braff D (1998)
Poor P 50 suppression among schizophrenia patients and their first-degree biological relatives.
Am J Psychiatry 155: 1691-1694
- [9] Curtis C, Calhins M, Grove W, Feil K, Lacono W (2001)
Saccadic disinhibition in patients with acute and remitted schizophrenia and their first-degree biological relatives.
Am J Psychiatry 158: 100-106
- [10] Dolan R, Fletcher P, McKenna P, Friston K, Frith C (1999)
Abnormal neural integration related to cognition in schizophrenia.
Acta Psychiatr Scand Suppl 395: 58-67
- [11] Erlenmeyer-Kimling L, Rock D, Roberts S, Janal M, Kestenbaum C, Cornblatt B, Hilldöf Adamo U, Gottesman II (2000)
Attention, memory and motor skills as childhood predictors of schizophrenia-related psychoses: The New York high-risk project.
Am J Psychiatry 157: 1416-1422
- [12] Franke P, Maier W, Hardt J, Hain C, Comblatt B (1994)
Attentional abilities and measures of schizotypy: Their variation and covariation in schizophrenic patients, their siblings, and normal control subjects.
Psychiatry Res 54: 259-272
- [13] Friedl-Francesconi H, Binder H
Leistungsinseln bei Demenzpatienten: Diagnostische und therapeutische Möglichkeiten der Neuropsychologie.
In: Hinterhuber H (Hrsg): Dementielle Syndrome.
Innsbruck, Integrative Psychiatrie VIP, S 86-91

-
- [14] Goldberg T, Gold J (1995)
Neurocognitive functioning in patients with schizophrenia.
In: Bloom F, Kupfer D: Psychopharmacology – the fourth generation of progress.
Raven Press, New York, S 1245-1247
- [15] Goldman-Rahic P (1999)
The relevance of the dopamine-D1 receptor in the cognitive symptoms of schizophrenia.
Neuropsychopharmacology 21: 170-180
- [16] Green M (1998)
Schizophrenia from a neurocognitive perspective: Probing the impenetrable darkness.
Allyn & Bacon, Boston
- [17] Heinrichs RW, Zakzanis K (1998)
Neurocognitive deficit in schizophrenia: A quantitative review of the evidence.
Neuropsychology 12: 426-445
- [18] Hofer A, Weiss E (2002)
Advances in the neuroimaging of cognitive functions in schizophrenia.
Curr Opin Psychiatry 15: 3-7
- [19] Holzmann P, Solomon C, Levin S, Waternaux C (1984)
Pursuit eye movements dysfunctions in schizophrenia. Family evidence for specificity.
Arch Gen Psychiatrie 41: 136-139
- [20] Höschel K, Uhlendorff V, Biegel K, Kunert H, Weniger G, Irle E (1997)
Effektivität eines ambulanten neuropsychologischen Aufmerksamkeits- und Gedächtnistrainings in der Spätphase nach Schädelhirntrauma.
Zeitschrift für Neuropsychologie 7 Heft 2: 69-82

-
- [21] Iacono WG, Beise M (1992)
Are males more likely than females to develop schizophrenia?
Am J Psychiatry 149: 1070-1074
- [22] Kay SR, Opler LA, Fiszbein A (1986)
The Positive and Negative Syndrome Scale (PANSS).
Bewertungsleitfaden
- [23] Kay SR, Fiszbein A, Opler LA (1987)
The positive and negative syndrome scale (PANSS) for schizophrenia.
Schizophr Bull 13: 261-276
- [24] Kern R, Green M, Goldstein M
Modification of performance on the Span of Apprehension, a putative marker of vulnerability to schizophrenia.
J Abnorm Psychol 104: 386-389
- [25] Kurtz M, Rayland J, Bilker W, Ger R (2001)
Comparison of the continuous performance test with and without working memory demands in healthy controls and patients with schizophrenia.
Schizophr Res 48: 307-316
- [26] Lautenbacher S (1999)
Zur Notwendigkeit, die Schizophrenie aus einer neuropsychologischen Perspektive zu verstehen.
Verhaltensther Verhaltensmed 20: 223-248
- [27] Liebermann JA, Murray RM (2001)
Comprehensive care of schizophrenia.
London

-
- [28] Lussier I, Slip E (2001)
Memory and attention deficits in drug naïve patients with schizophrenia.
Schizophr Res 48: 45-55
- [29] Medalia A, Aluma M, Tryon W, Merriam A (1998)
Effectiveness of attention training in schizophrenia.
Schizophr Bull 24: 147-152
- [30] Meltzer H, McGurk S (1999)
The effects of clozapine, risperidone and olanzapine on cognitive functioning in schizophrenia.
Schizophr Bull 25: 233-255
- [31] Möller HJ, Laux G, Deister A (2001)
Psychiatrie und Psychotherapie.
Thieme, Stuttgart, 2. Auflage, S 134
- [32] Moritz S, Andresen B, Perro C (2002)
Neurocognitive performance in first-episode and chronic schizophrenic patients.
Eur Arch Psychiatry Clin Neurosci 252: 33-37
- [33] Mortimer AM (1997)
Cognitive function in schizophrenia – do neuroleptics make a difference?
Pharmacol Biochem Behav 56: 789-795
- [34] Nüchterlein K (1983)
Signal detection in vigilance tasks and behavioral attributes among offspring of schizophrenic mothers and among hyperactive children.
J Abnorm Psychol 92: 4-28

-
- [35] Olbrich R (2001)
Computergestütztes Training in der Psychiatrie: Trainingsprogramm, Implementierung in klinischen Einrichtungen, erste Ergebnisse, Forschungsperspektiven.
Nervenheilkunde 20: 90-95
- [36] Olbrich R, Mussgay L (1990)
Reduction of schizophrenic deficits by cognitive training.
Psychiatry Clin Neurosci 239: 366-369
- [37] Paus T, Zatarre R, Hofle N (1997)
Time-changes in neural systems underlying attention and arousal during the performance of an auditory vigilance task.
Journal of Cognitive Neuroscience 9: 392-408
- [38] Pfleger U (1996)
Computergestütztes kognitives Trainingsprogramm mit schizophrenen Patienten.
Münster – New York, Waxmann – Internationale Hochschulschriften Bd 204
- [39] Preetz N (1992)
Untersuchung zur Validierung eines computergestützten neuropsychologischen Gedächtnis- und Konzentrations-Trainingsprogramms für zerebralgeschädigte Patienten an einer Klinik für neurologische und orthopädische Rehabilitation.
Dissertation an der Medizinischen Akademie Magdeburg
- [40] Ramm B, Hofmann G (1987)
Biomathematik.
Stuttgart, Enke, S 157-195
- [41] Regel H, Fritsch A (1997)
Evaluationsstudie zum computergestützten Training psychischer Basisfunktionen.
Abschlussbericht zum geförderten Forschungsprojekt.
Bonn, Kuratorium ZNS

-
- [42] Rund B, Berg N (1999)
Cognitive deficits and cognitive training in schizophrenic patients: A review.
Acta Psychiatr Scad 100: 85-95
- [43] Snitz BE, Daum J (2001)
The neuropsychology of schizophrenia: A selective review.
Neuropsychology 12: 3-7
- [44] Sobizack N, Albus M, Hubmann W, Mohr F (1999)
Neuropsychologische Defizite bei ersterkrankten schizophrenen Patienten.
Nervenarzt 70: 408-415
- [45] Spaulding W, Fleming S, Reed D, Sullivan M, Storzbach D, Lam M (1999)
Cognitive functioning in schizophrenia for psychiatric rehabilitation.
Schizophr Bull 25: 257-289
- [46] Statistisches Bundesamt
Mikrozensus
Personen im Alter von 15 und mehr Jahren.
Statistisches Bundesamt, März 2004
- [47] Sturm W (2005)
Aufmerksamkeitsstörungen.
Hogrefe Verlag, Göttingen, S 2-5
- [48] Suslow T, Arolt V (1998)
Effektivität eines computergestützten Aufmerksamkeitstrainings bei schizophrenen Patienten.
Psychiatr Prax 25: 105-110

-
- [49] Van Zomeren AW, Brouwer WH (1994)
Clinical neuropsychology of attention.
Oxford Univ Press, New York
- [50] Volz H (2000)
Kognitionsstörungen sind schizophrene Kernsymptome.
In: Volz H, Kasper S, Möller HJ, Sach G, Höse A: Die Rolle der Kognition in der Therapie schizophrener Störungen.
Deutscher Universitätsverlag, Wiesbaden, S 50-73
- [51] Weisbrod M, Hill H, Neithammer R, Sauer H (1999)
Genetic influence on auditory information processing in schizophrenia: P 300 in monozygotic twins.
Biol Psychiatry 46: 721-725
- [52] Wexler B, Hawkins K, Rounsville B, Anderson M, Sernyak M, Green M (1997)
Normal neurocognitive performance after extended practice in patients with schizophrenia.
Schizophr Res 26: 173-180