

LITERATURVERZEICHNIS

ACRI, J.B., Brown K.J., Saah, M.I., Grunberg, N.E.:

Strain and age differences in acoustic startle responses and effects of nicotine in rats.

Pharmacol. Biochem. Behav. 50: 191-198, 1995

AMERICAN PSYCHIATRIC ASSOCIATION:

Diagnostic and Statistical Manual of Mental Disorders, 4. Auflage, American Psychiatric Association (Hrsg.), Washington, D.C., 1994

AMIN, A.H., Crawford, T.B.B., Gaddum, J.H.:

The distribution of substance P and 5-HT in the central nervous system of the dog.

J. Physiol. 126: 596-618, 1954

ANDREWS, N., File, S.E.:

Handling history of rats modifies behavioural effects of drugs in the elevated plus-maze test of anxiety.

Eur. J. Pharmacol. 235: 109-112, 1993

ARBORELIUS, L., Chergui, K., Murase S., Nomikos, G.G., Höök, B.B., Chouvet, G., Hacksell, U., Svensson, T.H.:

The 5-HT_{1A} receptor selective ligands, (R)-8-OH-DPAT and (S)-UH-301, differentially affect the activity of midbrain dopamin neurons.

Naunyn-Schmiedberg's Arch. Pharmacol. 347: 353-362, 1993

AZZAM, R.M., Notarianni, L.J., Ali, H.M.:

Rapid and simple chromatographic method for the determination of diazepam and its major metabolites in human plasma and urine.

J. Chromatogr. 708: 304-309, 1998

BALFOUR, D.J.K.:

Effects of GABA and diazepam on ³H-serotonin release from hippocampal synaptosomes.

Eur. J. Pharmacol. 68: 11-16, 1980

BARNARD, E.A., Skolnick, P., Olsen, R.W., Möhler, H., Sieghart, W., Biggio, G., Braestrup, C., Bateson, A.N., Langer, S.Z.:

International union of pharmacology. XV. Subtypes of γ -aminobutyric acid_A receptors: classification on the basis of subunit structure and receptor function.

Pharmacol. Rev. 50: 291-313, 1998

BARNES, N.M., Sharp, T.:

A review of central 5-HT receptors and their function.

Neuropharmacol. 38: 1083-1152, 1999

BIRMELIN, I.:

Behaviour of pet animals.

Dtsch. Tierärztl. Wochenschr. 97: 243-247, 1990

BLANCHARD, D.C., Blanchard, R.J., Rodgers R.J.:

Risk assessment and animal models of anxiety.

In: *Animal models in Psychopharmacology*, B. Olivier, J. Mos, J.L. Slanger (Hrsg.), Birkhäuser Verlag, Basel, S. 117-134, 1991

BLANCHARD, R.J., Yudko, E.B., Rodgers, R.J., Blanchard D.C.:

Defense system psychopharmacology: an ethological approach to the pharmacology of fear and anxiety.

Behav. Brain Res. 58: 155-165, 1993

BOADLE-BIBER, M.C.:

Regulation of serotonin synthesis.

Prog. Biophys. molec. Biol. 60: 1-15, 1993

BOIX, F., Teruel, A.F., Escorihuela, R.M., Tobena, A.:

Handling-habituation prevents the effects of diazepam and alprazolam on brain serotonin levels in rats.

Behav. Brain Res. 36: 209-215, 1990

BODNOFF, S.R., Suranyi-Cadotte, B., Quirion, R., Meany, M.J.:

A comparison of the effects of diazepam versus several typical and atypical anti-depressant drugs in an animal model of anxiety.

Psychopharmacol. 97: 277-279, 1989

BORMANN, J.:

Electrophysiology of GABA_A and GABA_B subtypes.

Trends Neurosci. 11: 112-116, 1988

BORMANN, J.:

The 'ABC' of GABA receptors.

Trends Pharmacol. Sci. 21: 16-19, 2000

BOWERY, N.G.:

GABA_B receptors and their significance in mammalian pharmacology.

Trends Pharmacol. Sci. 10: 401-407, 1989

BRADLEY, P.B., Engel, G., Fenuik, W., Fozard, J.R., Humphrey, P.P., Middlemiss, D.N.,

Mylecharane, E.J., Richardson B.P., Saxena P.R.:

Proposal for the classification and nomenclature of functional receptors for 5-hydroxytryptamine.

Neuropharmacol. 25: 563-576, 1986

BRETT, R.R., Pratt, J.A.:

Chronic handling modifies the anxiolytic effect of diazepam in the elevated plus-maze.

Eur. J. Pharmacol. 178: 135-138, 1990

BRITTON, D.R., Britton, K.T.:

A sensitive open field measure of anxiolytic drug activity.

Pharmacol. Biochem. Behav. 15: 577-582, 1981

BRITTON, K.T., Page, M., Baldwin, H.A., Koob, G.F.:

Anxiolytic activity of steroid anesthetic alphaxolone.

J. Pharmacol. Exp. Ther. 258: 124-129, 1991

BROADHURST, P.L.:

Determinants of emotionality in the rat: III. Strain differences.

J. comp. physiol. Psychol. 51: 55-59, 1958

BROADHURST, P.L.:

Experiments in psychogenetics.

In: Experiments in personality, Psychogenetics and Psychopharmacology, Vol. I, H.J. Eysenck (Hrsg.), Routledge and Kegan Paul, London, S. 729-763, 1960

BRUCE, J.H., Ramirez, A.M., Lin, L., Oracion, A., Agarwal, R.P., Norenberg, M.D.:

Peripheral-type benzodiazepines inhibit proliferation of astrocytes in culture.

Brain Res. 564: 167-170, 1991

BURT, D.R., Kamatchi, G.L.:

GABA_A receptor subtypes: from pharmacology to molecular biology.

FASEB J. 5: 2916-2923, 1991

CHAOULOFF, F., Kulikov, A., Sarieau, A., Castanon, N., Mormède, P.:

Male Fischer 344 and Lewis rats display differences in locomotor reactivity, but not in anxiety-related behaviours: relationship with the hippocampal serotonergic system.

Brain Res. 693: 169-178, 1995

CHOJNACKA-WÓJCIK, E.:

5-Hydroxytryptamine in the central nervous system.

Pol. J. Pharmacol. 47: 219-235, 1995

COLE, J.C., Rodgers, R.J.:

An ethological analysis of the effects of chlordiazepoxide and bretazenil (Ro16-6028) in the murine elevated plus-maze.

Behav. Pharmacol. 4: 573-580, 1993

COLLINGE, J., Pycock, C.:

Differential actions of diazepam on the release of [³H]5-hydroxytryptamine from cortical and midbrain raphe slices in the rat.

Eur. J. Pharmacol. 85: 9-14, 1982

COLPAERT, F.C., Meert, T.F., Niemegeers, C.J.E., Janssen, P.A.J.:

Behavioral and 5-HT antagonist effects of ritanserin: a pure and selective antagonist of LSD discrimination in rat.

Psychopharmacol. 86: 45-54, 1985

COMMISSARIS, R.L., Harrington G.M., Altman, H.J.:

Benzodiazepine anticonflict effects in Maudsley reactive (MR/HAR) and non-reactive (MNRA/HAR) rats.

Psychopharmacol. 100: 287-292, 1990

COSTA, E., Guidotti, A., Mao, C.C.:

Evidence for the involvement of GABA in the action of benzodiazepine: studies on rat cerebellum.

In: Mechanism of Action of Benzodiazepines, E. Costa, P. Greengard (Hrsg.), Raven Press, N.Y., S. 113-130, 1975

COSTALL, B., Jones, B.J., Kelly, M.E., Naylor, R.J., Tomkins, D.M.:

Exploration of mice in a black and white test box: validation as a model of anxiety.

Pharmacol. Biochem. Behav. 32: 777-785, 1989

CRABBE, J.C., Wahlsten, D., Dudek, B.C.:

Genetics of mouse behavior: interactions with laboratory environment.

Science 284: 1670-1672, 1999

CRAWLEY, J.N., Davis, L.G.:

Baseline exploratory activity predicts anxiolytic responsiveness to diazepam in five mouse strains.

Brain. Res. Bull. 8: 609-612, 1982

CRAWLEY, J.N., Goodwin, F.K.:

Preliminary report of a simple animal behavior model for the anxiolytic effects of benzodiazepines.

Pharmacol. Biochem. Behav. 13: 167-170, 1980

CUNHA, J.M., Masur, J.:

Evaluation of psychotropic drugs with a modified open field test.

Pharmacology 16: 259-267, 1978

CURTIS, D.R., Johnston, G.A.R.:

Amino acid transmitters in the mammalian central nervous system.

Ergebn. Physiol. Biol. Chem. Exp. Pharmacol. 69: 97-188, 1974

DAHLSTRÖM, A., Fuxe, K.:

Evidence for the existence of monoamine-containing neurons in the central nervous system.

I. Demonstration of monoamines in the cell bodies of brainstem neurons.

Acta Physiol. Scand. 62: 1-55, 1964

DAWSON, G.R., Tricklebank, M.D.:

Use of the elevated plus maze in the search for novel anxiolytic agents.

Trends Pharmacol. Sci. 16: 33-36, 1995

DE VRY, J., Schreiber, R.:

Comparison of acute and repeated treatment with the 5-HT_{1A} receptor ligands 8-OH-DPAT and ipsapirone in animal models of anxiety and depression.

Drug Dev. Res. 30: 91-103, 1993

DOMINIC, J.A., Sinha, A.K., Barchas, J.D.:

Effect of benzodiazepine compounds on brain amine metabolism.

Eur. J. Pharmacol. 32: 124-127, 1975

DREW, C.A., Johnston, G.A.R., Weatherby, R.P.:

Bicuculline-insensitive GABA receptors: studies on the binding of (-)-baclofen to rat cerebellar membranes.

Neurosci. Lett. 52: 317-321, 1984

FALCK, A., Hillarp, N.A., Thieme, G., Thorp, A.:

Fluorescence of catecholamines and related compounds with formaldehyde.

J. Histochem. Cytochem. 10: 348-354, 1962

FERNANDEZ-GUASTI, A., Hong, E., Lopez-Rubalcava, C.:

Species differences in the mechanism through which the serotonergic agonists indoreneate and ipsapirone produce their anxiolytic action.

Psychopharmacol. 107: 61-68, 1992

FILE, S:E.:

Models of anxiety.

Br. J. Clin. Pract. Symp. Suppl. 39: 15-19, 1985

FILE, S.E.:

The contribution of behavioural studies to the neuropharmacology of anxiety.
Neuropharmacol. 26: 877-886, 1987

FILE, S.E.:

One-trial tolerance to the anxiolytic effects of chlordiazepoxide in the elevated plus-maze.
Psychopharmacol. 100: 281-282, 1990a

FILE, S.E.:

Age and anxiety: increased anxiety, decreased anxiolytic, but enhanced sedative, response to chlordiazepoxide in old rats.
Hum. Psychopharmacol. 5: 169-173, 1990b

FREUD, S., Breuer, J.:

Studien über Hysterie.
Fischer Taschenbuch-Verlag, Frankfurt a.M., 1993, 1. Auflage

FRUSSA-FILHO, R., Otoboni, J.R., Uema, F.T., Sa-Rocha, L.C.:

Evaluation of memory and anxiety in rats observed in the elevated plus-maze: effects of age and isolation.
Braz. J. Med. Biol. Res. 24: 725-728, 1991

GADDUM, J.H., Picarelli, Z.P.:

Two kinds of tryptamine receptors.
Br. J. Pharmacol. 12: 323-328, 1957

GALLAGER, D.W.:

Benzodiazepine potentiation of a GABA inhibitory response in the dorsal raphe nucleus.
Eur. J. Pharmacol. 49: 133-143, 1978

GALLAHER, E.J., Hollister, L.E., Gionet, S.E., Crabbe, J.C.:

Mouse lines selected for genetic difference in diazepam sensitivity.
Psychopharmacol. 93: 25-30, 1987

GARDNER, C.R.:

Recent developments in 5HT-related pharmacology of animal models of anxiety.
Pharmacol. Biochem. Behav. 24: 1479-1485, 1986

GELLER, I., Seifter, J.:

The effects of meprobamate, barbiturates, d-amphetamine and promazine on experimentally induced conflict in the rat.

Psychopharmacologia 1: 482-495, 1960

GENTSCH, C., Lichtsteiner, M., Feer, H.:

Genetic and environmental influences on behavioral and neurochemical aspects of emotionality in rats.

Experientia 44: 482-490, 1988

GILAD, G.M., Shiller, I.:

Differences in open-field behavior and in learning tasks between two rat strains differing in their reactivity to stressors.

Behav. Brain Res. 32: 89-93, 1989

GONZALEZ, L.E., File, S.E., Overstreet, D.H.:

Selectively bred lines of rats differ in social interaction and hippocampal 5-HT_{1A} receptor function: a link between anxiety and depression?

Pharmacol. Biochem. Behav. 59: 787-792, 1998

GRIEBEL, G.:

5-Hydroxytryptamine-interacting drugs in animal models of anxiety disorders: more than 30 years of research.

Pharmacol. Biochem. Behav. 65: 319-395, 1995

GRIEBEL, G., Moreau, J.-L., Jenck, F., Martin, J.R., Misslin, J.R.:

Some critical determinants of the behaviour of rats in the elevated plus-maze.

Behav. Process. 29: 37-48, 1993

HAEFELY, W., Kulcsar, A., Möhler, H., Polc, P., Schaffner, R.:

Possible involvement of GABA in the central actions of benzodiazepines.

In: *Mechanism of Action of Benzodiazepines*, E. Costa, P. Greengard (Hrsg.), Raven Press, N.Y., S. 131-151, 1975

HALL, C.S.:

Emotional behavior in the rat. I. Defecation and urination as measures of individual differences in emotionality.

J. Comp. Psychol. 18: 385-403, 1934

HALLIDAY, G., Harding, A., Paxinos, G.:

Serotonin and tachykinin systems.

In: The rat nervous system. G. Paxinos (Hrsg.), Academic Press, San Diego, S. 929-974, 1995

HANDLEY, S.L., McBlane, J.W.:

5HT drugs in animal models of anxiety.

Psychopharmacol. 112: 13-20, 1993

HANDLEY, S.L., Mithani, S.:

Effects of alpha-adrenoceptor agonists and antagonists in a maze-exploration model of "fear"-motivated behaviour.

Naunyn Schmiedberg's Arch. Pharmacol. 327: 1-5, 1984

HERTZ, L., Tamir, H.:

Some properties of an astrocytic protein fraction that binds serotonin.

J. Neurochem. 37: 1331-1334, 1981

HILL, D.R., Bowery, N.G.:

³H-baclofen und ³H-GABA bind to bicuculline-insensitive GABA_B sites in the rat brain.

Nature 290: 149-152, 1981

HIRONAKA, T., Fuchino, K., Fujii, T.:

Absorption of diazepam and its transfer through the blood-brain barrier after intraperitoneal administration in the rat.

J. Pharmacol. Exp. Ther. 229: 809-815, 1984

HJORTH, S., Sharp, T., Hacksell, U.:

Partial postsynaptic 5-HT_{1A} agonist properties of the novel stereoselective 8-OH-DPAT analogue (+)cis-8-hydroxy-1-methyl-2-(di-n-propylamino)tetralin, (+)ALK-3.

Eur. J. Pharmacol. 170: 269-264, 1989

HOGG, S.:

A review of the validity and variability of the elevated plus-maze as an animal model of anxiety.

Pharmacol. Biochem. Behav. 54: 21-30, 1996

HOLLISTER, L.E.:

Treatment of depression with drugs.

Ann. Intern. Med. 89: 78-84, 1978

HOYER, D., Clarke, D.E., Fozard, J.R.:

International union of pharmacological classification of receptors for 5-hydroxytryptamine (serotonin).

Pharmacol. Rev. 46: 157-193, 1994

HOYER, D., Martin, G.:

5-HT receptor-classification and nomenclature: towards harmonisation with the human genome.

Neuropharmacol. 36: 419-428, 1997

IKEGWUONU, F.I., Ganem, L.G., Larson, M.C., Shen, X., Jefcoate, C.R.:

The regulation by gender, strain, dose, and feeding status of the induction of multiple form of cytochrome P450 isozymes in rat hepatic microsomes by 2,4,5,2',4',5'-hexachlorobiphenyl.

Toxicol. Appl. Pharmacol. 139: 33-41, 1996

IMHOF, J.T., Coelho, Z.M., Schmitt, M.L., Morato, G.S., Carobrez, A.P.:

Influence of gender and age on performance of rats in the elevated plus maze apparatus.

Behav. Brain Res. 56: 177-180, 1993

IVERSEN, S.D.:

5-HT and anxiety.

Neuropharmacol. 23: 1553-1560, 1984

IVERSEN, L.L., Blohm, F.E.:

Studies on the uptake of ^3H -GABA und ^3H -glycine in slices and homogenisates of rat brain and spinal cord by electron microscopic autoradiography.

Brain Res. 41: 131-143, 1972

JANOWSKA, E., Pucilowski, O., Kostowski, W.:

Chronic oral treatment with diltiazem or verapamil decreases isolation-induced activity impairment in elevated plus-maze.

Behav. Brain Res. 43: 155-158, 1991

JOHNSTON, A.L., File, S.E.:

Sex differences in animal tests of anxiety.

Physiol. Behav. 49: 245-250, 1991

KLOCKOWSKI, P.M., Levy, G.:

Kinetics of drug action in disease states. XXIV. Pharmacodynamics of diazepam and its active metabolites in rats.

J. Pharmacol. Exp. Ther. 244: 912-918, 1988

KLOTZ, U.:

Die Bedeutung der Pharmakokinetik von Benzodiazepinen.

In: *Benzodiazepine: Standortbestimmung und Perspektiven*, H. Coper, H. Rommelspacher (Hrsg.), Urban & Schwarzenberg, München, S. 1-13, 1987

KLOTZ, U., Antonin, K.-H., Biek, P.R.:

Pharmacokinetics and plasma binding of diazepam in man, dog, rabbit, guinea pig and rat.

J. Pharmacol. Exp. Ther. 199: 67-73, 1976

KOSTER, A.S., Nieuwenhuis, L., Frankhuijzen-Sierevogel, A.C.:

Comparison of microsomal drug-metabolizing enzymes in 14 rat inbred strains.

Biochem. Pharmacol. 38: 759-765, 1989

KULIKOV, A., Augerre, S., Berton, O., Ramos, A., Mormède, P., Chaouloff F.:

Central serotonergic systems in the spontaneously hypertensive and Lewis rat strains that differ in the elevatd plus-maze test of anxiety.

J. Pharmacol. Experiment. Behav. 281: 775-784, 1997

LAL, H., Shearman, G.T.:

Attenuation of chemically-induced anxiogenic stimuli as a novel method for evaluating anxiolytic drugs: a comparison of clobazam with other benzodiazepines.

Drug Dev. Res. 1: 127-134, 1982

LARSEN, M.C., Brake, P.B., Parmar, D., Jefcoate, C.R.:

The induction of five rat hepatic P450 cytochromes by phenobarbital and similarly acting compounds is regulated by a sexually dimorphic, dietary-dependent endocrine factor that is highly strain specific.

Arch. Biochem. Biophys. 315: 24-34, 1994

LISTER, R.G.:

The use of a plus-maze to measure anxiety in the mouse.

Psychopharmacol. 92: 180-185, 1987

LISTER, R.G.:

Ethologically-based animal models of anxiety disorders.

Pharmacol. Ther. 46: 321-340, 1990

LÓPEZ-RUBALCAVA, C., Saldívar, A., Fernández-Guasti, A.:

Interaction of GABA and serotonin in the anxiolytic action of Diazepam and serotonergic anxiolytics.

Pharmacol. Biochem. Behav. 43: 433-440, 1992

LORENZ, K.:

Studies in Animal and Human behavior.

Harvard University Press, Cambridge, 1971

LUCKI, I.:

The spectrum of behaviors influenced by serotonin.

Biol. Psych. 44: 151-162, 1998

MARCUCCI, F., Guaitani, A., Kvetina, J., Mussini, E., Garattini, S.:

Species differences in diazepam metabolism and anticonvulsant effect.

Eur. J. Pharmacol. 4: 467-470, 1968

MARCUCCI, F., Mussini, E., Fanelli, R., Garattini, S.:

Species differences in diazepam metabolism – I. Metabolism of diazepam metabolites.

Biochem. Pharmacol. 19: 1847-1851, 1970

MARSDEN, C.A.:

The neuropharmacology of serotonin in the central nervous system.

In: Selective serotonin re-uptake inhibitors. J.P. Feighner, W.F. Boyer (Hrsg.), Wiley Verlag, Chichester, S. 1-33, 1996

MCBLANE, J.W., Critchley, M.A.E., Handley, S.L.:

Light intensity influences response to 8-OH-DPAT in the elevated X-maze model of anxiety.

Br. J. Pharmacol. 105 (Suppl.): 221, 1992

MEERT, T.F., Janssen, P.A.J.:

Psychopharmacology of ritanserin: comparison with chlordiazepoxide.

Drug Dev. Res. 18: 119-144, 1989

MENARD, J., Treit, D.:

Effects of centrally administered anxiolytic compounds in animal models of anxiety.

Neurosci. Biobehav. Rev. 23: 591-613, 1999

MICZEK, K.A.:

Effects of scopolamine, amphetamine and benzodiazepines on conditioned suppression.

Pharmacol. Biochem. Behav. 1: 401-411, 1973

MILLAN, M.J., Brocco, M.:

Serotonin and anxiety: mixed 5-HT_{1A} agonists – 5-HT_{1C/2} antagonists as potential anxiolytic agents.

In: Anxiety: Neurobiology, clinic and therapeutic perspectives. M. Hamon, H. Ollat, M.-H. Thiébot (Hrsg.), Colloque INSERM/ John Libbey Eurotext Verlag, 232: 153-165, 1993

MÖHLER, H., Okada, T.:

Benzodiazepine receptors: demonstration in the central nervous system.

Science 198: 849-851, 1977

MOLEWIJK, H.E., van der Poel, A.M., Olivier, B.:

The ambivalent behaviour “stretched approach posture“ in the rat as a paradigm to characterize anxiolytic drugs.

Psychopharmacol. 121: 81-90, 1995

MONTGOMERY, S.A.:

The relation between fear induced by novel stimulation and exploratory behaviour.

J. comp. physiol. Psychol. 48: 254-260, 1955

ONAIVI, E.S., Maguire, P.A., Tsai, N.F., Davies, M.F., Loew, G.H.:

Comparison of behavioral and central BDZ binding profile in three rat lines.

Pharmacol. Biochem. Behav. 43 : 825-831, 1992

OROSCO, M., Gourch, A., Cohen, Y., Jacquot, C.:

Cholecystokinin-induced variations in hypothalamic serotonergic system of the "cafeteria" rat.

Brain Res. Bull. 25: 949-952, 1990

OVERSTREET, D.H., Rezvani, A.H.:

Behavioral differences between two inbred strains of Fawn-Hooded rat: a model of serotonin dysfunction.

Psychopharmacol. 128: 328-330, 1996

PALKOVITS, M.:

Isolated removal of hypothalamic or other brain nuclei of the rat.

Brain Res. 59: 449-450, 1973

PAXINOS, G., Watson, C.:

The rat brain stereotaxic coordinates.

Academic Press, Orlando, 1997

PELLOW, S., Chopin, P., File, S.E., Briley, M.:

Validation of open:closed arm entries in an elevated plus-maze as a measure of anxiety in the rat.

J. Neurosci. Methods 14: 149-167, 1985

PEROUTKA, S.J., Snyder, S.H.:

Multiple serotonin receptors: differential binding of [³H]5-hydroxytryptamine, [³H]lysergic acid diethylamide and [³H]spiroperidol.

Mol. Pharmacol. 16: 687-699, 1979

PLASSAT, J.-L., Amlaiky, N., Hen, R.:

Molecular cloning of a mammalian serotonin receptor that activates adenylate cyclase.
Mol. Pharmacol. 44: 229-236, 1993

PORSOLT, R.D., Bertin, A., Jalfre, M.:

“Behavioral despair” in rats and mice: strain differences and the effects of imipramine.
Eur. J. Pharmacol. 51: 291-294, 1978

RAMOS, A., Berton, O., Mormède, P., Chaouloff, F.:

A multiple-test study of anxiety-related behaviours in six inbred rat strains.
Behav. Brain Res. 85: 57-69, 1997

RAMPE, D., Triggle, D.J.:

Benzodiazepines and calcium channel function.
Trends Pharmacol. Sci. 7: 461-464, 1986

RAPPORT, M.M., Green, A.A., Page, I.H.:

Crystalline serotonin.
Science 108: 329, 1948

REX, A., Marsden, C.A., Fink, H.:

Effect of diazepam on cortical 5-HT release and behavior in the guinea-pig on exposure to the elevated plus maze.
Psychopharmacol. 110: 490-496, 1993

REX, A., Sondern, U., Voigt, J.P., Franck, S., Fink, H.:

Strain differences in fear-motivated behavior of rats.
Pharmacol. Biochem. Behav. 54: 107-111, 1996

REX, A., Voigt, J.-P., Fink, H.:

Behavioral and neurochemical differences between identically-raised Fischer 344 and Harlan-Wistar rats.
Behav. Genet. 29: 187-192, 1999

REX, A., Voigt, J.P., Voits, M., Fink, H.:

Pharmacological evaluation of a modified open-field test sensitive to anxiolytic drugs.
Pharmacol. Biochem. Behav. 59: 677-683, 1998

RICHARDS, J.G., Möhler, H.:

Benzodiazepine receptors.

Neuropharmacol. 23: 233-242, 1984

ROBERTSON, H.A., Martin, O.L., Candy, J.M.:

Differences in benzodiazepine binding in Maudsley reactive and Maudsley non-reactive rats.

Eur. J. Pharmacol. 50: 455-457, 1978

RODGERS, R.J.:

Animal models of ‘anxiety’: where next?

Behav. Pharmacol. 8: 477-496, 1997

RODGERS, R.J., Cole, J.C.:

The elevated plus-maze: pharmacology, methodology and ethology.

In: *Ethology and Psychopharmacology*, S.J. Cooper, C.A. Hendrie (Hrsg.), John Wiley & Sons Ltd., S. 9-44, 1994

RODGERS, R.J., Cole, J.C., Dobson, M.J., Eells, J.R.:

Effects of fluprazine and eltoprazine in the mouse elevated plus-maze and light/dark exploration test of anxiety.

J. Psychopharmacol. (Suppl. Abst. Book) 6: A 52, 1992

ROMERO, L., Jerenej, B., Bel, N., Cicin-Sain, L., Cortés, R., Artigas, F.:

Basal and stimulated extracellular serotonin concentration in the brain of rats with altered serotonin uptake.

Synapse 28: 313-321, 1998

ROSS, G.D.:

Structure and function of membrane complement receptors.

Fed. Proc. 41: 3089-3093, 1982

SANER, A., Pletcher, A.:

Effect of diazepam on cerebral 5-hydroxytryptamine synthesis.

Eur. J. Pharmacol. 55: 315-318, 1979

SANGER, D.J., Benavides, J., Perrault, G., Morel E., Cohen, C., Joly, D., Zivikovic, B.:

Recent developments in the behavioral pharmacology of benzodiazepine (ω) receptors:
evidence for the functional significance of receptor subtypes.

Neurosc. Biobehav. Rev. 18: 355-372, 1994

SCHMITT, U., Hiemke, C.:

Strain differences in open-field and elevated plus-maze behavior of rats without and with
pretest handling.

Pharmacol. Biochem. Behav. 59: 807-811, 1998

SCHWARTING, R.K.W., Thiel, C.M., Müller, C.P., Huston, J.P.:

Relationship between anxiety and serotonin in the ventral striatum.

NeuroReport 9: 1025-1029, 1998

SCHWARTZ, M.A., Koechlin, B.A., Postma, E., Palmer, S., Krol, G.:

Metabolism of diazepam in rat, dog, and man.

J. Pharmacol. Exp. Ther. 149: 423-435, 1965

SETEM, J., Pinheiro A.P., Motta, V.A., Morato, S., Cruz, A.P.:

Ethopharmacological analysis of 5-HT ligands on the rat elevated plus-maze.

Pharmacol. Biochem. Behav. 62: 515-521, 1999

SHEPHARD, R.A., Nielsen, E.A., Broadhurst, P.L.:

Sex and strain differences in benzodiazepine receptor binding in Roman rat strains.

Eur. J. Pharmacol. 77: 327-330, 1982

SHEPHERD, J.K., Grewal, S.S., Fletcher, A., Bill, D.J., Dourish, C.T.:

Behavioural and pharmacological characterisation of the elevated “zero-maze” as an animal
model of anxiety.

Psychopharmacol. 116: 56-64, 1994

SIGEL, E., Stephenson, F.A., Mamalaki, C., Barnard, E.A.:

γ -Aminobutyric acid/benzodiazepine receptor complex of bovine cerebral cortex – purification
and partial characterisation.

J. Biol. Chem. 258: 6965-6971, 1983

SNYDER, S.H.:

Angstlösende Medikamente.

In: *Chemie der Psyche. Drogenwirkung im Gehirn*, Spektrum der Wissenschaft, Verlagsgesellschaft mbH, Heidelberg, S. 155-180, 1990

SOUBRIE, P., Blas, C., Ferron, A., Glowinski, J.:

Chlordiazepoxide reduces *in vivo* serotonin release in the basal ganglia of encéphale isolé but not anesthetized cats: evidence for a dorsal raphe site action.

J. Pharmacol. Exp. Ther. 226: 526-532, 1983

SQUIRES, R.F., Braestrup, C.:

Benzodiazepine receptors in rat brain.

Nature 266: 732-734, 1977

STEIN, L., Wise, D., Berger, B.D.:

Anti-anxiety actions of benzodiazepines: decrease in activity of serotonin neurons in the punishment system.

In: *The benzodiazepines*, S. Garattini, E. Massini, L.O. Randall (Hrsg.), S. 299-326, Raven Press, New York, 1973

STEINBUSCH, H.W.M.:

Distribution of serotonin-immunoreactivity in the central nervous system of the rat-cell bodies and terminals.

Neuroscience 6: 557-618, 1981

STERNBACH, L.H.:

The benzodiazepine story.

J. Med. Chem. 22: 1-7, 1979

STEPHENS, D.N.:

Animal models of anxiety; ground for depression? Commentary on Rodgers, 'Animal models of anxiety: where next?'

Behav. Pharmacol. 8: 497-501, 1997

STÖHR, T., Schulte-Wermeling, D., Weiner, I., Feldon, J.:

Rat strain differences in open-field behavior and the locomotor stimulating and rewarding effects of amphetamine.

Pharmacol. Biochem. Behav. 59: 813-818, 1998

STUDY, R.E., Barker, J.L.:

Diazepam and (--)pentobarbital: Fluctuation analysis reveals different mechanisms for potentiation of γ -aminobutyric acid response in cultured central neurons.

Proc. Natl. Acad. Sci. 78: 7180-7184, 1981

TRABER, J., Glaser, T.:

5-HT_{1A} receptor-related anxiolytics.

Trends Pharmacol. Sci. 8: 432-437, 1987

TREIT, D.:

Animal models for the study of anti-anxiety agents: a review.

Neurosci. Biobehav. Reviews 9: 203-222, 1985

TREIT, D.:

Anxiolytic effects of benzodiazepines, and 5-HT_{1A} agonists: animal models.

In: 5-HT_{1A} Agonists, 5-HT₃ Antagonists and Benzodiazepines: Their Comparative Behavioural Pharmacology, R.J. Rodgers, S.J. Cooper (Hrsg.) Whiley, Chichester, S. 107-131, 1991

TREIT, D., Menard, J.:

Animal models of anxiety and depression.

In: Neuromethods (34): In Vivo Neuromethods, A.A. Boulton, G.B. Baker, A.N. Bateson, (Hrsg.) Humana Press Inc., Totowa, New Jersey, S. 89-148, 1998

TRULLAS, R., Skolnick, P.H.:

Differences in fear motivated behaviors among inbred mouse strains.

Psychopharmacol. 111: 323-331, 1993

TWAROG, B.M., Page, I.H.:

Serotonin content of some mammalian tissues and urine and a method of its determination.

Am. J. Physiol. 174: 157-161, 1953

VAN DER LAAN, J.W., De Boer, S.F., Van der Gutgen, J., De Groot, G.:

Differences in the duration of sedative and anxiolytic effects of desmethyldiazepam in two outbred Wistar strains.

Pharmacol. Biochem. Behav. 39: 149-153, 1991

VAN DER LAAN, J.W., De Groot, G., Wortelboer, H., Noordhoek, J.:

Pharmacokinetic differences of desmethyldiazepam in three outbred Wistar strains related to differences in liver enzyme activities.

Pharmacol. Toxicol. 73: 229-232, 1993

VAN DER STAAY, F.J.:

Shift in the performance of 24-month-old Wistar rats in the Morris water escape task: a comparison across 36 experiments.

Behav. Brain Res. 87: 213-222, 1997

VAN DER STAAY, F.J., Blokland, A.:

Behavioral differences between outbred Wistar, inbred Fischer 344, Brown Norway, and hybrid Fischer 344 x Brown Norway rats.

Physiol. & Behav. 60: 97-109; 1996

VALZELLI, L.:

The exploratory behaviour in normal and aggressive mice.

Psychopharmacologia 15: 232-235, 1969

VOGEL, J.R., Beer, B., Clody, D.E.:

A simple and reliable conflict procedure for testing anti-anxiety agents.

Psychopharmacologia 21: 1-7, 1971

WALSH, R.N., Cummins, R.A.:

The open-field test: a critical review.

Psychol. Bulletin 83: 482-504, 1976

WHEATLEY, D.:

Buspirone: multicenter efficacy study.

J. clin. Psychiat. 43: 92-94, 1982

WILCOCK, J., Broadhurst, P.L.:

Strain differences in emotionality: open-field and conditioned avoidance behavior in the rat.

J. Comp. Physiol. Psychol. 63: 335-338, 1967

WISE C.D., Berger, B.D., Stein, L.:

Benzodiazepines: anxiety-reducing activity by reduction of serotonin turnover in the rat brain.

Science, 177: 180-183, 1972

WOOLEY, S., Shaw, E.:

Some neurophysiological aspects of serotonin.

Br. Med. J. 2: 122, 1954

WORLD HEALTH ORGANISATION:

The ICD-10 Classification of Mental and Behavioural Disorders

World Health Organisation (Hrsg.), Genf, 1992