

## 5 LITERATURVERZEICHNIS

- Ahmad,M. and Cashmore,A.R. (1993). HY4 gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature* 366, 162-166.
- Ahmad,M., Jarillo,J.A., and Cashmore,A.R. (1998a). Chimeric proteins between cry1 and cry2 Arabidopsis blue light photoreceptors indicate overlapping functions and varying protein stability. *Plant Cell* 10, 197-207.
- Ahmad,M., Jarillo,J.A., Smirnova,O., and Cashmore,A.R. (1998b). Cryptochrome blue-light photoreceptors of Arabidopsis implicated in phototropism. *Nature* 392, 720-723.
- Ahmad,M., Jarillo,J.A., Smirnova,O., and Cashmore,A.R. (1998c). The CRY1 blue light photoreceptor of Arabidopsis interacts with phytochrome A in vitro. *Mol. Cell* 1, 939-948.
- Andel,F.I., Murphy,J.T., Haas,J.A., McDowell,M.T., van,d.H., I, Lugtenburg,J., Lagarias,J.C., and Mathies,R.A. (2000). Probing the photoreaction mechanism of phytochrome through analysis of resonance Raman vibrational spectra of recombinant analogues. *Biochemistry* 14, 2667-2676.
- Aravind,L. and Ponting,C.P. (1997). The GAF domain: an evolutionary link between diverse phototransducing proteins. *Trends. Biochem. Sci.* 22, 458-459.
- Arciero,D.M., Bryant,D.A., and Glazer,A.N. (1988). In vitro attachment of bilins to apophycocyanin. I Specific covalent adduct formation at cysteinyl residues involved in phycocyanobilin binding in C-phycocyanin. *J. Biol. Chem.* 263, 18343-18349.
- Assmann,S.M., Simoncini,L., and Schroeder,J.I. (1985). Blue light activates electrogenic ion pumping in guard cell protoplasts of *Vicia faba*. *Nature* 318, 285-287.
- Aukerman,M.J., Hirschfeld,M., Wester,L., Weaver,M., Clack,T., Amasino,R.M., and Sharrock,R.A. (1997). A deletion in the PHYD gene of the arabidopsis Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *Plant Cell* 9, 1317-1326.
- Bagnall,D.J., King,R.W., and Hangarter,R.P. (1996). Blue-light promotion of flowering is absent in hy4 mutants of Arabidopsis. *Planta* 200, 278-280.
- Berkelman,T.R. and Lagarias,J.C. (1986). Visualization of bilin-linked peptides and proteins in polyacrylamide gels. *Anal. Biochem.* 156, 194-201.
- Bessette,P.H., Aslund,F., Beckwith,J., and Georgiou,G. (1999). Efficient folding of proteins with multiple disulfide bonds in the Escherichia coli Cytoplasm. *Proc Nat Acad Sci USA* 96, 13703-13708.
- Bhoo,S.H., Davis,S.J., Walker,J., Karniol,B., and Vierstra,R.D. (2001). Bacteriophytochromes are photochromic histidine kinases using a biliverdin chromophore. *Nature* 414, 776-779.
- Bhoo,S.H., Hirano,T., Jeong,H.Y., Lee,J.G., Furuya,M., and Song,P.-S. (1997). Phytochrome photochromism probed by site-directed mutations and chromophore esterification. *J. Am. Chem. Soc.* 119, 11717-11718.

- Blaauw, O.H., Laauw-Jansen, G., and van Leeuwen, W.J. (1968). An irreversible red-light-induced growth response in *Avena*. *Planta* 82, 87-104.
- Borgstahl, G.E., Williams, D.R., and Getzoff, E.D. (1995). 1.4 A structure of photoactive yellow protein, a cytosolic photoreceptor: unusual fold, active site, and chromophore. *Biochemistry* 34, 6278-6287.
- Borthwick, H.A., Hendricks, S.B., and Parker M W (1952a). The reaction controlling floral initiation. *Proc. Natl. Acad. Sci. U. S. A.* 38, 929-934.
- Borthwick, H.A., Hendricks, S.B., Parker, M.W., Toole, E.H., and Toole, V.K. (1952b). A reversible photoreaction controlling seed germination. *Proc. Natl. Acad. Sci. U. S. A.* 38, 662-666.
- Borucki, B., Otto, H., Rottwinkel, G., Hughes, J., Heyn, M.P., and Lamparter, T. (2003). Mechanism of Cph1 phytochrome assembly from stopped-flow kinetics and circular dichroism. *Biochemistry* 42, 13684-13697.
- Bowler, C., Neuhaus, G., Yamagata, H., and Chua, N.H. (1994). Cyclic GMP and calcium mediate phytochrome phototransduction. *Cell* 77, 73-81.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248-254.
- Briggs, W.R., Beck, C.F., Cashmore, A.R., Christie, J.M., Hughes, J., Jarillo, J.A., Kagawa, T., Kanegae, H., Liscum, E., Nagatani, A., Okada, K., Salomon, M., Rudiger, W., Sakai, T., Takano, M., Wada, M., and Watson, J.C. (2001). The phototropin family of photoreceptors. *Plant Cell* 13, 993-997.
- Briggs, W.R. and Huala, E. (1999). Blue-light photoreceptors in higher plants. *Annu Rev Cell Dev Biol* 15, 33-62.
- Butler, W.L., Hendricks S.B., and Siegelman, H.A. (1960). *Plant Physiol Suppl* 35, xxxii.
- Butler, W.L., Lane, H.C., and Siegelman, H.W. (1963). Nonphotochemical transformations of phytochrome *in vivo*. *Plant Physiol.* 38, 514-519.
- Butler, W.L., Norris, K.H., Siegelman, H.W., and Hendricks, S.B. (1959). Detection, assay, and preliminary purification of the pigment controlling photoresponsive development of plants. *Proc. Natl. Acad. Sci. U. S. A.* 45, 1703-1708.
- Cabral, J.H.M. and et al (1998). Crystal structure and functional analysis of the HERG potassium channel N terminus: a eukaryotic PAS domain. *Cell* 95, 649-655.
- Carrascal, M., Carujo, S., Bachs, O., and Abián, J. (2002). Identification of p21<sup>Cip1</sup> binding proteins by gel electrophoresis and capillary liquid chromatography microelectrospray tandem mass spectrometry. *Proteomics* 2, 455-468.
- Casal, J.J. and Boccalandro, H. (1995). Co-action between phytochrome B and HY4 in *Arabidopsis thaliana*. *Planta* 197, 213-218.
- Casal, J.J. and Mazzella, M.A. (1998). Conditional synergism between cryptochrome 1 and phytochrome B is shown by the analysis of phyA, phyB, and hy4 simple, double, and triple mutants in *Arabidopsis*. *Plant Physiol.* 118, 19-25.
- Cashmore, A. R., Jarillo, J. A., Wu, Y. J., and Liu, D. M. Cryptochromes - Blue-Light Receptors for Plants and Animals. *Science* Vol 284[Iss 5415], 760-765. 1999.

- Cerdan,P.D., Staneloni,R.J., Casal,J.J., and Sanchez,R.A. (1997). A 146 bp fragment of the tobacco Lhcb1\*2 promoter confers very- low- fluence, low-fluence and high-irradiance responses of phytochrome to a minimal CaMV 35S promoter. *Plant Mol. Biol.* 33, 245-255.
- Cerdan,P.D., Yanovsky,M.J., Reymundo,F.C., Nagatani,A., Staneloni,R.J., Whitelam,G.C., and Casal,J.J. (1999). Regulation of Phytochrome-B Signaling by Phytochrome-A and Fhy1 in Arabidopsis-Thaliana. *Plant J Vol 18*, 499-507.
- Chang, C. und Stewart, R. C. The two-component system - regulation of diverse signaling pathways in prokaryotes and eukaryotes. *Plant Physiol.* 117[3], 723-731. 1998.
- Chapman,D.J., Cole,W.J., and Siegelman,H.W. (1967). *J. Am. Chem. Soc.* 89, 3643-3645.
- Chen,E., Lapko,V.N., Lewis,J.W., Song,P.S., and Kliger,D.S. (1996). Mechanism of native oat phytochrome photoreversion: a time- resolved absorption investigation. *Biochemistry* 35, 843-850.
- Chentao,L. (2002). Blue Light Receptors and Signal Transduction. *The Plant Cell Supplement 2002*, 207-225.
- Cherry,J.R., Hondred,D., Walker,J.M., Keller,J.M., Hershey,H.P., and Vierstra,R.D. (1993). Carboxy-terminal deletion analysis of oat phytochrome A reveals the presence of separate domains required for structure and biological activity. *Plant Cell* 5, 565-575.
- Choi,G., Yi,H., Lee,J., Kwon,Y.K., Soh,M.S., Shin,B., Luka,Z., and Song,P.S. (1999). Phytochrome signalling is mediated through nucleoside diphosphate kinase 2. *Nature* 401, 610-613.
- Christie,J.M. and Briggs,W.R. (2001). Blue light sensing in higher plants. *J Biol Chem.* 276, 11457-11460.
- Christie,J.M., Salomon,M., Nozue,K., Wada,M., and Briggs,W.R. (1999). LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide. *Proc. Natl. Acad. Sci. U. S. A.* 96, 8779-8783.
- Clack,T., Mathews,S., and Sharrock,R.A. (1994). The phytochrome apoprotein family in Arabidopsis is encoded by five genes: the sequences and expression of PHYD and PHYE. *Plant Mol. Biol.* 25, 413-427.
- Clough,R.C., Jordan-Beebe,E.T., Lohman,K.N., Marita,J.M., Walker,J.M., Gatz,C., and Vierstra,R.D. (1999). Sequences within both the N- and C-terminal domains of phytochrome A are required for PFR ubiquitination and degradation. *Plant J* 17, 155-167.
- Cole,W.J., Chapman,D.J., and Siegelman,H.W. (1967). The structure of phycocyanobilin. *J. Am. Chem. Soc.* 89, 3642-3645.
- Cosgrove,D.J. (1994). Photomodulation of growth. In *Photomorphogenesis in plants*, 2nd edition, Kluwer Academic Publishers Netherlands, pp. 631-658.
- Crosson,S. and Moffat,K. (2001). Structure of a flavin-binding plant photoreceptor domain: insights into light-mediated signal transduction. *Proc. Natl. Acad. Sci. U. S. A.* 98, 2995-3000.

- Cutler,S.R., Ehrhardt,D.W., Griffiths,J.S., and Somerville,C.R. (2000). Random GFP::cDNA fusions enable visualization of subcellular structures in cells of Arabidopsis at a high frequency. *Proc. Natl. Acad. Sci. U. S. A.* 97, 3718-3723.
- D'Agostino,I.B. and Kieber,J.J. (1999). Phosphorelay signal transduction: the emerging family of plant response regulators. *TIBS* 24, 452-456.
- Davis,S.J., Vener,A.V., and Vierstra,R.D. (1999). Bacteriophytochromes: Phytochrome-like photoreceptors from nonphotosynthetic eubacteria. *Science* 286, 2517-2520.
- Deforce,L., Tokutomi,S., and Song,P.S. (1994). Phototransformation of pea phytochrome A induces an increase in alpha- helical folding of the apoprotein: comparison with a monocot phytochrome A and CD analysis by different methods. *Biochemistry* 33, 4918-4922.
- Deforce,L., Tomizawa,K., Ito,N., Farrens,D., Song,P.S., and Furuya,M. (1991). In vitro assembly of apophytochrome and apophytochrome deletion mutants expressed in yeast with phycocyanobilin. *Proc. Natl. Acad. Sci. U. S. A.* 88, 10392-10396.
- Devlin,P.F., Patel,S.R., and Whitelam,G.C. (1998). Phytochrome E Influences Internode Elongation and Flowering Time in Arabidopsis. *Plant Cell* 10, 1479-1488.
- Devlin,P.F., Robson,P.R., Patel,S.R., Goosey,L., Sharrock,R.A., and Whitelam,G.C. (1999). Phytochrome D acts in the shade-avoidance syndrome in Arabidopsis by controlling elongation growth and flowering time. *Plant Physiol.* 119, 909-916.
- Dower,W.J., Miller,J.F., and Ragsdale,C.W. (1988). High efficiency transformation of *E.coli* by high voltage electroporation. *Nucleic Acids Res* 16, 6145.
- Eftink,M.R. (1991). Topics in Fluorescence Spectroscopy. J.R.Lakowicz, ed. Plenum Press, New York), pp. 52-126.
- Eilfeld,P. and Rüdiger,W. (1985). *Z Naturforsch* 40c, 109-114.
- El-DinEl-Assal,S., Alonso-Blanco,C., Peeters,A.J., Raz,V., and Koornneef,M. (2001). A QTL for flowering time in Arabidopsis reveals a novel allele of CRY2. *Natl. Genet.* 29, 435-440.
- Fankhauser,C., Yeh,K.C., Lagarias,J.C., Zhang,H., Elich,T.D., and Chory,J. (1999). Pks1, a Substrate Phosphorylated by Phytochrome That Modulates Light Signaling in Arabidopsis. *Science* 284, 1539-1541.
- Farrens,D., Holt,R., Rospendowski,B., Song,P.-S., and Cotton,T. (1989). Surface-enhanced resonance Raman Scattering (SERRS) Spectroscopy applied to phytochrome and its model compounds. II. Phytochrome and phycocyanin chromophores. *J. Am. Chem. Soc.* 111, 9162-9169.
- Farrens,D.L., Cordonnier,M.M., Pratt,L.H., and Song,P.S. (1992). The distance between the phytochrome chromophore and the N- terminal chain decreases during phototransformation. A novel fluorescence energy transfer method using labeled antibody fragments. *Photochem. Photobiol.* 56, 725-733.
- Fodor,S.P., Lagarias,J.C., and Mathies,R.A. (1990). Resonance Raman analysis of the Pr and Pfr forms of phytochrome. *Biochemistry* 29, 11141-11146.
- Foerstendorf,H., Benda,C., Gartner,W., Storf,M., Scheer,H., and Siebert,F. (2001). FTIR studies of phytochrome photoreactions reveal the C=O bands of the chromophore:

- consequences for its protonation states, conformation, and protein interaction. *Biochemistry* *40*, 14952-14959.
- Foerstendorf,H., Lamparter,T., Hughes,J., Gärtner,W., and Siebert,F. (2000). The photoreactions of recombinant phytochrome from the cyanobacterium *Synechocystis*: A low-temperature UV-Vis and FT-IR spectroscopic study. *Photochem. Photobiol.* *2000*, 655-661.
- Folta,K.M. and Spalding,E.P. (2001). Unexpected roles for cryptochrome 2 and phototropin revealed by high-resolution analysis of blue light-mediated hypocotyl growth inhibition. *Plant J* *26*, 471-478.
- Frankenberg,N., Mukougawa,K., Kohchi,T., and Lagarias,J.C. (2001). Functional genomic analysis of the hy2 family of ferredoxin-dependent bilin reductases from oxygenic photosynthetic organisms. *Plant Cell* *13*, 965-978.
- Furuya,M. and Song,P.S. (1994). Assembly and properties of holophytochrome. In *Photomorphogenesis in Plants*, R.E.Kendrick and G.H.M.Kronenberg, eds. (Dordrecht: Kluwer Academic Publishers), pp. 105-140.
- Gallagher,S., Short,T.W., Ray,P.M., Pratt,L.H., and Briggs,W.R. (1988). Light-mediated changes in two proteins found associated with plasmamembrane fractions from pea stem sections. *PNAS* *85*, 8003-8007.
- Gambetta,G.A. and Lagarias,J.C. (2001). Genetic engineering of phytochrome biosynthesis in bacteria. *Proc. Natl. Acad. Sci. U. S. A.* *98*, 10566-10571.
- García,D.M., Muro,P.M., I, Reyes,J.C., and Florencio,F.J. (2000). Light-dependent regulation of cyanobacterial phytochrome expression. *J Bacteriol* *2000*, 38-44.
- Garner,W.W. and Allard,H.A. (1920). Effect of the relative length of day and night and other factors of the environment on growth and reproduction in plants. *J. Agric. Res* *18*, 553-606.
- Gendreau,E., Hofte,H., Grandjean,O., Brown,S., and Traas,J. (1998). Phytochrome controls the number of endoreduplication cycles in the *Arabidopsis thaliana* hypocotyl. *Plant J.* *13*, 221-230.
- Gil,P., Kircher,S., Adam,E., Bury,E., Kozma,B.L., Schäfer,E., and Nagy,F. (2000). Photocontrol of subcellular partitioning of phytochrome-B:GFP fusion protein in tobacco seedlings. *Plant J* *2000*, 135-145.
- Giraud,E., Fardoux,J., Fourier,N., Hannibal,L., Genty,B., Bouyer,P., Dreyfus,B., and Vermeglio,A. (2002). Bacteriophytochrome controls photosystem synthesis in anoxygenic bacteria. *Nature* *417*, 202-205.
- Gong,W., Hao,B., Mansy,S.S., Gonzalez,G., Gilles-Gonzalez,M.A., and Chan,M.K. (1998). Structure of a biological oxygen sensor: a new mechanism for heme-driven signal transduction. *Proc. Natl. Acad. Sci. U. S. A.* *95*, 15177-15182.
- Goodner,B., Hinkle,G., Gattung,S., Miller,N., Blanchard,M., Quorollo,B., Goldman,B.S., Cao,Y., Askenazi,M., Halling,C., Mullin,L., Houmiel,K., Gordon,J., Vaudin,M., Iartchouk,O., Epp,A., Liu,F., Wollam,C., Allinger,M., Doughty,D., Scott,C., Lappas,C., Markelz,B., Flanagan,C., Crowell,C., Gurson,J., Lomo,C., Sear,C., Strub,G., Cielo,C., and Slater,S. (2001). Genome sequence of the plant pathogen and biotechnology agent *Agrobacterium tumefaciens* C58. *Science* *294*, 2323-2328.

- Govorunova,E.G., Jung,K.-H., Sineshchekov,O., and Spudich,J.L. (2004). Chlamydomonas sensory rhodopsins A and B: cellular content and role in photophobic responses. *Biophys J* 86, 2342-2349.
- Graham,L. and Gallop,P.M. (1994). Covalent protein crosslinks: general detection, quantitation, and characterization via modification with diphenylborinic acid. *Anal. Biochem* 217, 298-305.
- Grimm,R., Eckerskorn,C., Lottspeich,F., Zenger,C., and Rüdiger,W. (1988). Sequence analysis of proteolytic fragments of 124-kilodalton phytochrome from etiolated #Avena sativa# L.: Conclusions on the conformation of the native protein. *Planta* 174, 396-401.
- Guo, H. W., Duong, H., Ma, N., und Lin, C. T. The Arabidopsis Blue-Light Receptor Cryptochrome-2 Is a Nuclear- Protein Regulated by a Blue Light-Dependent Posttranscriptional Mechanism. *Plant Journal* Vol 19[3], 279-287. 1999.
- Guo,H.W., Yang,W.Y., Mockler,T.C., and Lin,C.T. (1998). Regulations of flowering time by Arabidopsis photoreceptors. *Science* 279, 1360-1363.
- Halliday,K.J., Thomas,B., and Whitelam,G.C. (1997). Expression of heterologous phytochromes A, B or C in transgenic tobacco plants alters vegetative development and flowering time. *Plant J.* 12, 1079-1090.
- Hamada,T., Tanaka,N., Noguchi,T., Kimura,N., and Hasunuma,K. (1996). Phytochrome regulates phosphorylation of a protein with characteristics of a nucleoside diphosphate kinase in the crude membrane fraction from stem sections of etiolated pea seedlings. *J. Photochem. Photobiol. B.* 33, 143-151.
- Hamazato,F., Shinomura,T., Hanzawa,H., Chory,J., and Furuya,M. (1997). Fluence and wavelength requirements for Arabidopsis CAB gene induction by different phytochromes. *Plant Physiol.* 115, 1533-1540.
- Harper,R.M., Stowe-Evans,E.L., Luesse,D.R., Muto,H., Tatematsu,K., Watahiki,M.K., Yamamoto,K., and Liscum,E. (2000). The NPH4 locus encodes the auxin response factor ARF7, a conditional regulator of differential growth in aerial Arabidopsis tissue. *Plant Cell* 12, 757-770.
- Haupt,W. and Häder,D.-P. (1994). Photomovement. In *Photomorphogenesis in plants*, R.E.Kendrick and G.H.M.Kronenberg, eds. (Dordrecht: Kluwer Academic Publishers), pp. 707-732.
- Hauser,B.A., Cordonnier-Pratt,M.M., Daniel-Vedele,F., and Pratt,L.H. (1995). The phytochrome gene family in tomato includes a novel subfamily. *Plant Mol. Biol.* 29, 1143-1155.
- Hellingwerf,K.J. (2000). Key issues in the photochemistry and signalling-state formation of photosensor proteins. *Journal-of-Photochemistry-and-Photobiology-B-Biology* 54, 94-102.
- Hellingwerf,K.J. (2002). The molecular basis of sensing and responding to light in microorganisms. *Antonie van Leeuwenhoek*, ed. (Netherlands: Kluwer Academie), pp. 51-59.
- Hennig,L., Poppe,C., Unger,S., and Schaefer,E. (1999). Control of hypocotyl elongation in Arabidopsis thaliana by photoreceptor. *Planta-Berlin*. April, 1999;208, 257-263.

- Herdman,M., Coursin,T., Rippka,R., Houmard,J., and Tandeau de Marsac,N. (2000). A new appraisal of the prokaryotic origin of eukaryotic phytochromes. *J. Mol. Evol.* *51*, 205-213.
- Heyne,K., Herbst,J., Stehlik,D., Esteban,B., Lamparter,T., Hughes,J., and Diller,R. (2002). Ultrafast dynamics of phytochrome from the cyanobacterium *Synechocystis*, reconstituted with phycocyanobilin and phycoerythrobilin. *Biophys. J.* *82*, 1004-1016.
- Hisada,A., Hanzawa,H., Weller,J.L., Nagatani,A., Reid,J.B., and Furuya,M. (2000). Light-induced nuclear translocation of endogenous pea phytochrome A visualized by immunocytochemical procedures. *Plant Cell* *2000*, 1063-1078.
- Ho,Y.S.J., Burden,L.M., and Hurley,J.H. (2000). Structure of the GAF domain, a ubiquitous signaling motif and a new class of cyclic GMP receptor. *EMBO J* *19*, 5288-5299.
- Hoecker,U., Tepperman,J.M., and Quail,P.H. (1999). SPA1, a WD-repeat protein specific to phytochrome A signal transduction. *Science* *284*, 496-499.
- Hoff,W.D., Matthijs,H.C.P., Schubert,H., Crielaard,W., and Hellingwerf,K.J. (1995). Rhodopsin(s) in eubacteria. *Biophysical Chemistry* *56*, 193-199.
- Huala,E., Oeller,P.W., Liscum,E., Han,I.S., Larsen,E., and Briggs,W.R. (1997). *Arabidopsis* NPH1: a protein kinase with a putative redox-sensing domain. *Science* *278*, 2120-2123.
- Huang,Z.J., Edery,I., and Rosbash,M. (1993). PAS is a dimerization domain common to *Drosophila* period and several transcription factors. *Nature* *364*, 259-262.
- Hübschmann,T., Börner,T., Hartmann,E., and Lamparter,T. (2001a). Characterisation of the Cph1 holo-phytochrome from *Synechocystis* sp. PCC 6803. *Eur. J. Biochem.* *268*, 2055-2063.
- Hübschmann,T., Jorissen,H.J.M.M., Börner,T., Gärtner,W., and Tandeau de Marsac,N. (2001b). Phosphorylation of proteins in the light-dependent signalling pathway of a filamentous cyanobacterium. *Eur. J. Biochem.* *268*, 3383-3389.
- Hudson,M., Ringli,C., Boylan,M.T., and Quail,P.H. (1999). The Far1 Locus Encodes a Novel Nuclear-Protein Specific to Phytochrome-A Signaling. *Genes Dev.* *13*, 2017-2027.
- Hughes,J., Lamparter,T., Mittmann,F., Hartmann,E., Gärtner,W., Wilde,A., and Börner,T. (1997). A prokaryotic phytochrome. *Nature* *386*, 663.
- Huq,E. and Tepperman,J.M.Q.P.H. (2000). GIGANTEA is a nuclear protein involved in phytochrome signaling in *Arabidopsis*. *PNAS* *97*, 9789-9794.
- Jancarik,J. and Kim,S.-H. (1991). Sparse matrix sampling: a screening method for crystallization of proteins. *J. Appl. Cryst.* *24*, 409-411.
- Jarillo,J.A., Capel,J., Tang,R.H., Yang,H.Q., Alonso,J.M., Ecker,J.R., and Cashmore,A.R. (2001). An *Arabidopsis* circadian clock component interacts with both CRY1 and phyB. *Nature* *410*, 487-490.
- Jensen,P.J., Hangarter,R.P., and Estelle,M. (1998). Auxin transport is required for hypocotyl elongation in light-grown but not dark-grown *Arabidopsis*. *Plant Physiol.* *116*, 455-462.

- Jiang,Z.Y., Swem,L.R., Rushing,B.G., Devanathan,S., Tollin,G., and Bauer,C.E. (1999). Bacterial Photoreceptor with Similarity to Photoactive Yellow Protein and Plant Phytochromes. *Science* 285, 406-409.
- Jones A M., Quail,P.H. (1986). Quarternary structure of 124-kilodalton phytochrome from *Avena sativa* L. *Biochemistry* 25, 2987-2995.
- Jones,A.M. and Edgerton,M.D. (1994). The anatomy of phytochrome, a unique photoreceptor in plants. *Semin. Cell Biol.* 5, 295-302.
- Jones,A.M. and Quail,P.H. (1989). Phytochrome structure: Peptide fragments from a amino-terminal domain involved in protein-chromophore interactions. *Planta* 178, 147-156.
- Jones,A.M., Vierstra,R.D., Daniels,S.M., and Quail,P. (1985). The role of separate molecular domains in the structure of phytochrome from etiolated *Avena sativa* L. *Planta* 164, 501-506.
- Jorissen,H.J., Braslavsky,S.E., Wagner,G., and Gartner,W. (2002a). Heterologous expression and characterization of recombinant phytochrome from the green alga *Mougeotia scalaris*. *Photochem. Photobiol.* 76, 457-461.
- Jorissen,H.J., Quest,B., Remberg,A., Coursin,T., Braslavsky,S.E., Schaffner,K., Tandeau de Marsac,N., and Gärtner,W. (2002b). Two independent, light-sensing two-component systems in a filamentous cyanobacterium. *Eur. J. Biochem.* 269, 2662-2671.
- Jung,K.-H., Trivedi,V.D., and Spudich,J.L. (2003). Demonstration of a sensory rhodopsin in eubacteria. *Mol Microbiol* 47, 1513-1522.
- Kagawa,T., Sakai,T., Suetsugu,N., Oikawa,K., Ishiguro,S., Kato,T., Tabata,S., Okada,K., and Wada,M. (2001). Arabidopsis NPL1: A phototropin homolog controlling the chloroplast highlight avoidance response. *Science* 291, 2138-2141.
- Kagawa,T. and Wada,M. (1999). Chloroplast-avoidance response induced by high-fluence blue light in prothallial cells of the fern *Adiantum capillus-veneris* as analyzed by microbeam irradiation. *Plant Physiol* 119, 917-924.
- Kagawa,T. and Wada,M. (2000). Blue light-induced chloroplast relocation in *Arabidopsis thaliana* as analyzed by microbeam irradiation. *Plant Cell Physiol* 41, 84-93.
- Kamiya,Y. and García-Martínez,J.L. (1999). Regulation of gibberellin biosynthesis by light. *Curr Opin Plant Biol* 2, 398-403.
- Kandori, H., Yoshihara, K., und Tokutomi, S. Primary Process of Phytochrome - Initial Step of Photomorphogenesis in Green Plants. *Journal of the American Chemical Society* Vol 114[Iss 27], 10958-10959. 1992.
- Kaneko,T., Sato,S., Kotani,H., Tanaka,A., Asamizu,E., Nakamura,Y., Miyajima,N., Hirosawa,M., Sugiura,M., Sasamoto,S., Kimura,T., Hosouchi,T., Matsuno,A., Muraki,A., Nakazaki,N., Naruo,K., Okumura,S., Shimpo,S., Takeuchi,C., Wada,T., Watanabe,A., Yamada,M., Yasuda,M., and Taba,S. (1996). Sequence analysis of the genome of the unicellular Cyanobacterium *Synechocystis* sp. strain PCC6803. II. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* 3, 109-136.



- Karniol, B. and Vierstra, R.D. (2003). The pair of bacteriophytochromes from *Agrobacterium tumefaciens* are histidine kinases with opposing photobiological properties. *Proc. Natl. Acad. Sci. U. S. A.* *100*, 2807-2812.
- Kawai, H., Kanegae, T., Christensen, S., Kiyosue, T., Sato, Y., Imaizumi, T., Kadota, A., and Wada, M. (2003). Responses of ferns to red light are mediated by an unconventional photoreceptor. *Nature* *421*, 287-290.
- Kay, S.A. (1997). PAS, present, and future: clues to the origins of circadian clocks [comment]. *Science* *276*, 753-754.
- Kehoe, D.M. and Grossman, A.R. (1997). New classes of mutants in complementary chromatic adaptation provide evidence for a novel four-step phosphorelay system. *J. Bacteriol.* *179*, 3914-3921.
- Kehoe, D.M. and Grossman, R. (1996). Similarity of a chromatic adaptation sensor to phytochrome and ethylene receptors. *Science* *273*, 1409-1412.
- Kendrick, R.E. and Kronenberg, G.H.M. (1994). *Photomorphogenesis in Plants*, 2nd edition. (Dordrecht: Kluwer Academic Publishers).
- Kim, L., Kircher, S., Toth, R., Adam, E., Schäfer, E., and Nagy, F. (2000). Light-induced nuclear import of phytochrome-A:GFP fusion proteins is differentially regulated in transgenic tobacco and Arabidopsis. *Plant J.* *22*, 125-133.
- Kinoshita, T., Michio Doi, M., Suetsugu, N., Kagawa, T., Wada, M., and Shimazaki, K.-I. (2001). phot1 and phot2 mediate blue light regulation of stomatal opening. *Nature* *414*, 656-660.
- Kircher, S., Kozma-Bognar, L., Kim, L., Adam, E., Harter, K., Schäfer, E., and Nagy, F. (1999). Light quality-dependent nuclear import of the plant photoreceptors phytochrome A and B. *Plant Cell* *11*, 1445-1456.
- Kiyosue, T. and Wada, M. (2000). LKP1 (LOV Kelch protein 1): A factor involved in the regulation of flowering time in Arabidopsis. *Plant J* *23*, 807-815.
- Kleiner, O., Kircher, S., Harter, K., and Batschauer, A. (1999). Nuclear localization of the Arabidopsis blue light receptor cryptochrome 2. *Plant J* *19*, 289-296.
- Kneip, C., Hildebrandt, P., Schlamann, W., Braslavsky, S.E., Mark, F., and Schaffner, K. (1999). Protonation state and structural changes of the tetrapyrrole chromophore during the P<sub>r</sub> P<sub>fr</sub> phototransformation of phytochrome. A Resonance Raman Spectroscopic study. *Biochemistry* *16*, 15185-15192.
- Kohchi, T., Mukougawa, K., Frankenberg, N., Masuda, M., Yokota, A., and Lagarias, J.C. (2001). The *Arabidopsis* hy2 gene encodes phytochromobilin synthase, a ferredoxin-dependent biliverdin reductase. *Plant Cell* *13*, 425-436.
- Kolukisaoglu, H.U., Marx, S., Wiegmann, C., Hanelt, S., and Schneider-Poetsch, H.A. (1995). Divergence of the phytochrome gene family predates angiosperm evolution and suggests that *Selaginella* and *Equisetum* arose prior to *Psilotum*. *J. Mol. Evol.* *41*, 329-337.
- Koorneef, M., Hanhart, C.J., and van der Veen, J.H. (1991). A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol. Gen. Genet.* *229*, 57-66.

- Koorneef,M., Rolff,E., and Spruit,C.J.P. (1980). Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* L. Heynh. *Z Pflanzenphysiologie* *100*, 147-160.
- Koorneef,M., Hanhart,C.J., and van der Veen,J.H. (1991). A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Mol. Gen. Genet.* *229*, 57-66.
- Koorneef,M., Rolff,E., and Spruit,C.J.P. (1980). Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* L. Heynh. *Z Pflanzenphysiologie* *100*, 147-160.
- Krishnamohan,A., Balaji,V., and Veluthambi,K. (2001). Efficient vir gene induction in *Agrobacterium tumefaciens* requires virA, virG, and vir box from the same Ti plasmid. *J Bacteriol* *183*, 4079-4089.
- Kufer,W. and Scheer,H. (1979). Chemical modification of biliprotein chromophores. *Z. Naturforsch. [C]*. *34*, 776-781.
- Kunkel,T., Tomizawa,K., I, Kern,R., Furuya,M., Chua,N.H., and Schäfer,E. (1993). In vitro formation of a photoreversible adduct of phycocyanobilin and tobacco apophytochrome B. *Eur. J. Biochem.* *215*, 587-594.
- Lagarias,D.M., Wu,S.H., and Lagarias,J.C. (1995). Atypical phytochrome gene structure in the green alga *Mesotaenium caldariorum*. *Plant Mol Biol* *29*, 1127-1142.
- Lagarias,J.C. and Mercurio,F.M. (1985). Structure function studies on phytochrome. Identification of light- induced conformational changes in 124-kDa *Avena* phytochrome in vitro. *J. Biol. Chem.* *260*, 2415-2423.
- Lagarias,J.C. and Rapoport,H. (1980). Chromopeptides from phytochrome. The structure and linkage of the Pr form of the phytochrome chromophore. *J. Am. Chem. Soc.* *102*, 4821-4828.
- Lamparter,T., Carrascal,M., Michael,N., Martínez,E., Rottwinkel,G., and Abián,J. (2004). The Biliverdin Chromophore Binds Covalently to a Conserved Cysteine Residue in the N-Terminus of *Agrobacterium* Phytochrome Agp1. *Biochemistry* *43*, 3659-3669.
- Lamparter,T., Michael,N., Caspani,O., Miyata,T., Shirai,K., and Inomata,K. (2003). Biliverdin Binds Covalently to *Agrobacterium* Phytochrome Agp1 via Its Ring A Vinyl Side Chain. *J. Biol. Chem.* *278*, 33786-33792.
- Lamparter,T., Esteban,B., and Hughes,J. (2001). Phytochrome Cph1 from the cyanobacterium *Synechocystis* PCC6803: purification, assembly, and quaternary structure. *Eur. J. Biochem.* *268*, 4720-4730.
- Lamparter,T. and Hughes,J. (2003). Phytochromes and phytochromes-like proteins in cyanobacteria. In *Comprehensive Series in Photosciences*, A.Batschauer (ED), ed. (Elsevier, Amsterdam).
- Lamparter,T., Michael,N., Mittmann,F., and Esteban,B. (2002). Phytochrome from *Agrobacterium tumefaciens* has unusual spectral properties and reveals an N-terminal chromophore attachment site. *Proc. Natl. Acad. Sci. U. S. A.* *99*, 11628-11633.

- Lamparter,T., Mittmann,F., Gärtner,W., Börner,T., Hartmann,E., and Hughes,J. (1997). Characterization of recombinant phytochrome from the cyanobacterium *Synechocystis*. *Proc. Natl. Acad. Sci. U. S. A.* *94*, 11792-11797.
- Landgraf,F.T., Forreiter,C., Hurtado,P.A., Lamparter,T., and Hughes,J. (2001). Recombinant holophytochrome in *Escherichia coli*. *FEBS Lett.* *508*, 459-462.
- Lapko,V.N., Jiang,X.Y., Smith,D.L., and Song,P.S. (1998). Surface Topography of Phytochrome A Deduced from Specific Chemical Modification with Iodoacetamide. *Biochemistry* *37*, 12526-12535.
- Lasceve,G., Leymarie,J., Olney,M.A., Liscum,E., Christie,J.M., Vavasseur,A., and Briggs,W.R. (1999). *Arabidopsis* contains at least four independent blue-light-activated signal transduction pathways. *Plant Physiol* *120*, 605-614.
- Li,L. and Lagarias,J.C. (1992). Phytochrome assembly. Defining chromophore structural requirements for covalent attachment and photoreversibility. *J. Biol. Chem.* *267*, 19204-19210.
- Li,L., Murphy,J.T., and Lagarias,J.C. (1995). Continuous fluorescence assay of phytochrome assembly in vitro. *Biochemistry* *34*, 7923-7930.
- Lin,C., Ahmad,M., Gordon,D., and Cashmore,A.R. (1995). Expression of an *Arabidopsis* cryptochrome gene in transgenic tobacco results in hypersensitivity to blue, UV-A, and green light. *Proc. Natl. Acad. Sci. U. S. A.* *92*, 8423-8427.
- Lin,C., Yang,H., Guo,H., Mockler,T., Chen,J., and Cashmore,A.R. (1998). Enhancement of blue-light sensitivity of *arabidopsis* seedlings by a blue light receptor cryptochrome 2. *Proc. Natl. Acad. Sci. U. S. A.* *95*, 2686-2690.
- Lin,C. (2000). Photoreceptors and regulation of flowering time. *Plant-Physiology-Rockville*. [print] May, *2000;123*, 39-50.
- Liscum,E. and Briggs,W.R. (1995b). Mutations in the NPH1 locus of *Arabidopsis* disrupt the perception of phototropic stimuli. *Plant Cell* *7*, 473-485.
- Liscum,E. and Briggs,W.R. (1995a). Mutations in the NPH1 locus of *Arabidopsis* disrupt the perception of phototropic stimuli. *Plant Cell* *7*, 473-485.
- Liscum,E. and Briggs,W.R. (1996). Mutations of *Arabidopsis* in potential transduction and response components of the phototropic signaling pathway. *Plant Physiol.* *112*, 291-296.
- Lissemore,J.L. and Quail,P.H. (1988). Rapid transcriptional regulation by phytochrome of the genes for phytochrome and chlorophyll a/b-binding protein in *Avena sativa*. *Mol. Cell Biol.* *8*, 4840-4850.
- Littlewood,T. and Evan,G.I. (1998). *Helix-Loop-Helix Transcriptions Factors*. Oxford University Press, New York.
- Lottspeich,F. and Zorbas,H. (1998). *Bioanalytik*. (Heidelberg, Berlin, Germany: Springer Verlag).
- Manabe K and Nakazawa M (1997). The structure and function of phytochrome A: The roles of the entire molecule and of its various parts. *J. Plant Res.* *110*, 109-122.

- Mancinelli,A. (1994). The physiology of phytochrome action. In *Photomorphogenesis in Plants*, 2nd edition, R.E.Kendrick and G.H.M.Kronenberg, eds. (Dordrecht, Netherlands: Kluwer Academic Publishers), pp. 211-269.
- Mandoli,D.F. and Briggs,W.R. (1981). Phytochrome control of two low-irradiance responses of etiolated oat seedlings. *Plant Physiol.* *63*, 733-739.
- Marchuk,D. (1991). *Nucleic Acids Res* 1154.
- Martinez-Garcia,J.F., Huq,E., and Quail,P.H. (2000). Direct targeting of light signals to a promoter element-bound transcription factor. *Science* 2000 May 5;288. (5467. ):859. -63. 288, 859-863.
- Mas,P., Devlin,P.F., Panda,S., and Kay,S.A. (2000). Functional interaction of phytochrome B and Cryptochrome 2. *Nature* *408*, 207-211.
- Mathews,S. and Sharrock,R.A. (1996). The phytochrome gene family in grasses (Poaceae): a phylogeny and evidence that grasses have a subset of the loci found in dicot angiosperms. *Mol. Biol. Evol.* *13*, 1141-1150.
- Mathews,S. and Sharrock,R.A. (1997). Photochrome gene diversity. *Plant Cell Environ.* *20*, 666-671.
- Matsushita,T., Mochizuki,N., and Nagatani,A. (2003). Dimers of the N-terminal domain of phytochrome B are functional in the nucleus. *Nature* *424*, 571-573.
- Matysik,J., Hildebrandt,P., Schlamann,W., Braslavsky,S.E., and Schaffner,K. (1995). Fourier-transform resonance Raman spectroscopy of intermediates of the phytochrome photocycle. *Biochemistry* *34*, 10497-10507.
- Mazzella,M.A., Cerdan,P.D., Staneloni,R.J., and Casal,J.J. (2001). Hierarchical coupling of phytochromes and cryptochromes reconciles stability and light modulation of *Arabidopsis* development. *Development* *128*, 2291-2299.
- McDonagh,A.F. (1979). *The porphyrins*. (New York: Academic Press), pp. 293-491.
- Miller,J., Dower,W., and Tompkins,L. (1988). High-voltage electroporation of bacteria: Genetic transformation of *Campylobacter jejuni* with plasmid DNA. *Proc. Natl. Acad. Sci. U. S. A.* *85*, 856-860.
- Mizutani,Y., Tokutomi,S., and Kitagawa,T. (1994). Resonance Raman spectra of the intermediates in phototransformation of large phytochrome: deprotonation of the chromophore in the bleached intermediate. *Biochemistry* *33*, 153-158.
- Mockler,T.C., Guo,H., Yang,H., Duong,H., and Lin,C. (1999). Antagonistic actions of *Arabidopsis* cryptochromes and phytochrome B in the regulation of floral induction. *Development* *126*, 2073-2082.
- Montgomery,B.L. and Lagarias,J.C. (2002). Phytochrome ancestry: sensors of bilins and light. *Trends Plant Sci.* *7*, 357-366.
- Mozley,D. and Thomas,B. (1995). Developmental and photobiological factors affecting photoperiodic induction in *Arabidopsis thaliana* Heynh. *Landsberg erecta*. *J. Exp. Bot.* *46*, 173-179.
- Muramoto,T., Kohchi,T., Yokota,A., Hwang,I., and Goodman,H.M. (1999). The *Arabidopsis* photomorphogenic mutant *hyl* is deficient in phytochrome

- chromophore biosynthesis as a result of a mutation in a plastid heme oxygenase. *Plant Cell* *11*, 335-348.
- Murphy, J.T. and Lagarias, J.C. (1997). The phytofluors: a new class of fluorescent protein probes. *Curr. Biol.* *7*, 870-876.
- Mustilli, A.C. and Bowler, C. (1997). Tuning in to the signals controlling photoregulated gene expression in plants. *EMBO J.* *16*, 5801-5806.
- Nagy, F. and Schäfer, E. (2000). Nuclear and cytosolic events of light-induced, phytochrome-regulated signaling in higher plants. *EMBO J* *19*, 157-163.
- Nakazawa, M., Hayashi, H., Yoshida, Y., and Manabe, K. (1993). Identification of surface-exposed parts of red-light- and far-red-light-absorbing forms of native pea phytochrome by limited proteolysis. *Plant Cell Physiol.* *34*, 83-91.
- Nakazawa, M. und Manabe, K. Differential Exposure of Tryptophan Residues in the Red and Far-Red Light Absorbing Forms of Phytochrome, as Revealed by Chemical Modification. *Plant Cell Physiol.* Vol 34[Iss 7], 1097-1105. 1993.
- Neff, M.M. and Chory, J. (1998). Genetic interactions between phytochrome A, phytochrome B, and cryptochrome 1 during arabidopsis development. *Plant Physiol.* *118*, 27-35.
- Neff, M.M., Fankhauser, C., and Chory, J. (2000). Light: an indicator of time and place. *Genes Dev* *14*, 257-271.
- Neff, M.M. and Van-Volkenburgh, E. (1994). Light-stimulated cotyledon expansion in Arabidopsis seedlings: The role of. *Plant Physiol.* *104*, 1027-1032.
- Neff, M.M., Nguyen, S.M., Malancharuvil, E.J., and Fujioka, S. (1999). BAS1: A gene regulating brassinosteroid levels and light responsiveness in. *Proceedings-of-the-National-Academy-of-Sciences-of-the-United-States-of* of-the.
- Neuhaus, G., Bowler, C., Kern, R., and Chua, N.H. (1993). Calcium/calmodulin-dependent and -independent phytochrome signal transduction pathways. *Cell* *73*, 937-952.
- Ni, M., Tepperman, J.M., and Quail, P.H. (1998). PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. *Cell* *95*, 657-667.
- Ni, M., Tepperman, J.M., and Quail, P.H. (1999). Binding of phytochrome B to its nuclear signaling partner Pif3 is reversibly induced by light. *Nature* *400*, 781-784.
- Nozue, K., Kanegae, T., Imaizumi, T., Fukuda, S., Okamoto, H., Yeh, K.C., Lagarias, J.C., and Wada, M. (1998). A phytochrome from the fern *Adiantum* with features of the putative photoreceptor NPH1. *Proc. Natl. Acad. Sci. U. S. A.* *95*, 15826-15830.
- Ogura, T., Tanaka, N., Yabe, N.T., Komatsu, S., and Hasunuma, K. (1999). Characterization of Protein Complexes Containing Nucleoside Diphosphate Kinase with Characteristics of Light Signal-Transduction Through Phytochrome in Etiolated Pea-Seedlings. *Photochem. Photobiol.* Vol 69, 397-403.
- Osterlund, M.T., Hardtke, C.S., Wei, N., and Deng, X.W. (2000). *Nature* *405*, 462-466.
- Otto, H., Lamparter, T., Borucki, B., Hughes, J., and Heyn, M.P. (2003). Dimerization and inter-chromophore distance of Cph1 phytochrome from *Synechocystis*, as

- monitored by fluorescence homo and hetero energy transfer. *Biochemistry* 42, 5885-5895.
- Park,C.-M., Bhoo,S.-H., and Song,P.-S. (2000a). Inter-domain crosstalk in the phytochrome molecules. *Cell & Developmental Biology* 11, 449-456.
- Park,C.M., Kim,J.I., Yang,S.S., Kang,J.G., Kang,J.H., Shim,J.Y., Chung,Y.H., Park,Y.M., and Song,P.S. (2000b). A second photochromic bacteriophytochrome from *Synechocystis* sp. PCC 6803: spectral analysis and down-regulation by light. *Biochemistry* 39, 10840-10847.
- Park,C.M., Shim,J.Y., Yang,S.S., Kang,J.G., Kim,J.I., Luka,Z., and Song,P.S. (2000c). Chromophore-apoprotein interactions in *Synechocystis* sp. PCC6803 phytochrome Cph1. *Biochemistry* 30, 6349-6356.
- Parker,W., Partis,M., and Song,P.S. (1992). N-terminal domain of *Avena* phytochrome: interactions with sodium dodecyl sulfate micelles and N-terminal chain truncated phytochrome. *Biochemistry* 31, 9413-9420.
- Parkinson,J.S. and Kofoid (1992). Communication modules in bacterial signaling proteins. *Annu. Rev. Genet.* 26, 71-112.
- Qin,M., Kuhn,R., Moran,S., and Quail,P.H. (1997). Overexpressed phytochrome C has similar photosensory specificity to phytochrome B but a distinctive capacity to enhance primary leaf expansion. *Plant J.* 12, 1163-1172.
- Quail,P.H. (1997). An emerging molecular map of the phytochromes. *Plant Cell Environ.* 20, 657-665.
- Quail,P.H. (2000). Phytochrome-interacting factors. *Semin. Cell Dev. Biol.* 11, 457-466.
- Quail,P.H. (2002). Phytochrome photosensory signalling networks. *Nat. Rev. Mol. Cell Biol.* 3, 85-93.
- Quail,P.H., Boylan,M.T., Parks,B.M., Short,T.W., Xu,Y., and Wagner,D. (1995). Phytochromes: Photosensory perception and signal transduction. *Science* 268, 675-680.
- Reed,J.W., Nagatani,A., Elich,T., Fagan,M., and Chory,J. (1994). Phytochrome A and phytochrome B have overlapping but distinct functions in *Arabidopsis* development. *Plant Physiol.* 104, 1139-1149.
- Reed,J.W., Nagpal,P., Poole,D.S., Furuya,M., and Chory,J. (1993). Mutations in the gene for the red/far-red light receptor phytochrome B alter cell elongation and physiological responses throughout *Arabidopsis* development. *Plant Cell* 5, 147-157.
- Rehorek, M. und Heyn, M. P. Binding of all-trans retinal to the purple membrane. Evidence for cooperativity and determination of the extinction coefficient. *Biochemistry* 18, 4977-4983. 1979.
- Remberg,A., Lindner,I., Lamparter,T., Hughes,J., Kneip,K., Hildebrandt,P., Braslavsky,S.E., Gärtner,W., and Schaffner,K. (1997). Raman spectroscopic and light-induced-kinetic characterization of a recombinant phytochrome of the cyanobacterium *Synechocystis*. *Biochemistry* 36, 13389-13395.

- Reymond P., Short T., Briggs,W.R., and Poff K L (1992). Light-inducible phosphorylation of a membrane protein plays an early role in signal reansduction for phototropism in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 89, 4718-4721.
- Rospendowski,B.N., Farrens,D.L., Cotton,T.M., and Song,P.S. (1989). Surface enhanced resonance Raman scattering (SERRS) as a probe of the structural differences between the Pr and Pfr forms of phytochrome. *FEBS Lett.* 258, 1-4.
- Rost,B. and Sander,C. (1993). Improved Prediction of Protein Secondary Structure by Use of Sequence Profiles and Neural Networks. *PNAS* 90, 7558-7562.
- Rost,B., Sander,C., and Schneider,R. (1994). PHD--an automatic mail server for protein secondary structure prediction. *Comput. Appl. Biosci.* 10, 53-60.
- Roux,S. (1994). Signal transduction in phytochrome responses. In *Photomorphogenesis in Plants*, 2nd edition (eds: Kendrick RE, Kronenberg GHM), Kluwer Academic Publishers , Netherlands, pp. 187-209.
- Rüdiger,W. and Thümmler,F. (1994). The phytochrome chromophore. In *Photomorphogenesis in plants*, 2nd edition, R.E.Kendrick and G.H.M.Kronenberg, eds. (Dordrecht: Kluwer Academic Publishers), pp. 51-69.
- Sage,L.C. (1992). *Pigment of the Imagination - A History of Phytochrome Research.* Academic Press).
- Saijo,Y., Sullivan,J.A., Wang,H., Yang,J., Shen,Y., Rubio,V., Ma,L., Hoecker,U., and Deng,X.W. (2003). The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. *Genes-and-Development.* [print] August 17, 2642-2647.
- Sakai,T., Wada,T., Ishiguro,S., and Okada,K. (2000). RPT2. A signal transducer of the phototropic response in *Arabidopsis*. *Plant Cell* 12, 225-236.
- Sakai,T., Kagawa,T., Kasahara,M., Swartz,T.E., Christie,J.M., Briggs,W.R., Wada,M., and Okada,K. (2001). *Arabidopsis* *nph1* and *npl1*: blue light receptors that mediate both phototropism and chloroplast relocation. *Proc. Natl. Acad. Sci. U. S. A.* 98, 6969-6974.
- Sakamoto,K. and Nagatani,A. (1996). Nuclear localization activity of phytochrome B. *Plant J.* 10, 859-868.
- Salomon,M., Christie,J.M., Knieb,E., Lempert,U., and Briggs,W.R. (2000). Photochemical and mutational analysis of the FMN-binding domains of the plant blue light receptor, phototropin. *Biochemistry* 39, 9401-9410.
- Sambrook,J. and Russell,A.J. (2001).
- Sambrook,J., Fritsch,E.F., and Maniatis,T. (1989). *Molecular cloning, a laboratory manual.* CSH Press).
- Sanger,F., Nicklen,S., and Coulsen,A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. U. S. A.* 74, 5463-5467.
- Schendel,R. and Rüdiger,W. (1988). Electrophoresis and electrofocussing of phytochrome from etiolated *Avena sativa*. *Z Naturforsch* 44c, 12-18.
- Schendel,R., Tong,Z., and Rüdiger,W. (1989). Partial proteolysis of rice phytochrome: comparison with oat phytochrome. *Z. Naturforsch. [C].* 44, 757-764.

- Schmitz,O., Katayama,M., Williams,S.B., Kondo,T., and Golden,S.S. (2000). CikA, a bacteriophytochrome that resets the cyanobacterial circadian clock. *Science* 289, 765-768.
- Schneider-Poetsch,H.A., Braun,B., Marx,S., and Schaumburg,A. (1991). Phytochromes and bacterial sensor proteins are related by structural and functional homologies. Hypothesis on phytochrome-mediated signal-transduction. *FEBS Lett.* 281, 245-249.
- Sharrock,R.A. and Quail,P.H. (1989). Novel phytochrome sequences in *Arabidopsis thaliana*: structure, evolution, and differential expression of a plant regulatory photoreceptor family. *Genes Dev.* 3, 1745-1757.
- Shinomura,T., Nagatani,A., Chory,J., and Furuya,M. (1994). The induction of seed germination in *Arabidopsis thaliana* is regulated principally by phytochrome B and secondarily by phytochrome-A. *Plant Physiol.* 104, 363-371.
- Shinomura,T., Nagatani,A., Hanzawa,H., Kubota,M., Watanabe,M., and Furuya,M. (1996). Action spectra for phytochrome A- and B-specific photoinduction of seed germination in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 93, 8129-8133.
- Shinomura,T., Uchida,K., and Furuya,M. (2000). Elementary processes of photoperception by phytochrome A for high. *Plant Physiol* 2000, 147-156.
- Short,T.W. and Briggs,W.R. (1994). The transduction of blue light signals in higher plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 45, 143-171.
- Sineshchekov,O.A., Jung,K.H., and Spudich,J.L. (2002a). Two rhodopsins mediate phototaxis to low- and high-intensity light in *Chlamydomonas reinhardtii*. *Proc. Natl. Acad. Sci. U. S. A.* 99, 8689-8694.
- Sineshchekov,V., Hughes,J., Hartmann,E., and Lamparter,T. (1998a). Fluorescence and photochemistry of recombinant phytochrome from the cyanobacterium *Synechocystis*. *Photochem. Photobiol.* Vol 67, 263-267.
- Sineshchekov,V., Hughes,J., Hartmann,E., and Lamparter,T. (1998b). Fluorescence and photochemistry of recombinant phytochrome from the cyanobacterium *Synechocystis*. *Photochem Photobiol* 67, 263-267.
- Sineshchekov,V., Koppel,L., Esteban,B., Hughes,J., and Lamparter,T. (2002b). Fluorescence investigation of the recombinant cyanobacterial phytochrome (Cph1) and its C-terminally truncated monomeric species (Cph1Delta2): implication for holoprotein assembly, chromophore-apoprotein interaction and photochemistry. *J Photochem. Photobiol. B.* 67, 39-50.
- Sineshchekov,V.A. (1995). Photobiophysics and photobiochemistry of the heterogeneous phytochrome system. *Biochim. Biophys. Acta* 1228, 125-164.
- Singh,B.R., Chai,Y.G., Song,P.S., Lee,J., and Robinson,G.W. (1988). A photoreversible conformational change in 124 kDa *Avena* phytochrome. *Biochim. Biophys. Acta* 936, 395-405.
- Singh,B.R., Choi,J., Kwon,T.I., and Song,P.S. (1989). Use of bilirubin oxidase for probing chromophore topography in tetrapyrrole proteins. *J. Biochem. Biophys. Methods* 18, 135-147.



- Singh, B.R. and Song, P.-S. (1990). A differential molecular topography of the Pr and Pfr forms of native oat phytochrome as probed by fluorescence quenching. *Planta* *181*, 263-267.
- Singh, B.R. and Song, P.S. (1989). Interactions between native oat phytochrome and tetrapyrroles. *Biochim. Biophys. Acta* *996*, 62-69.
- Smith, H. (1970). Phytochrome and photomorphogenesis in plants. *Nature* *227*, 665-668.
- Smith, H. (2000). Phytochromes and light signal perception by plants--an emerging synthesis. *Nature* *407*, 585-591.
- Somers, D.E. and Quail, P.H. (1995). Phytochrome-mediated light regulation of PHYA- and PHYB- GUS transgenes in *Arabidopsis thaliana* seedlings. *Plant Physiol.* *107*: 2, 523-534.
- Somers, D.E., Schultz, T.F., Milnamow, M., and Kay, S.A. (2000). ZEITLUPE encodes a novel clock-associated PAS protein from *Arabidopsis*. *Cell* *101*, 319-329.
- Song, P.S. (1988). The molecular topography of phytochrome: chromophore and apoprotein. *J. Photochem. Photobiol. B.* *2*, 43-57.
- Spector, T. (1978). Refinement of the coomassie blue for protein quantitation. *Anal. Biochem.* *86*, 142-146.
- Sprenger, W.W., Hoff, W.D., Armitage, J.P., and Hellingwerf, K.J. (1993). The eubacterium *Ectothiorhodospira halophila* is negatively phototactic, with a wavelength dependence that fits the absorption spectrum of the photoactive yellow protein. *J. Bacteriol.* *175*, 3096-3104.
- Spudich, J.L., Yang, C.-S., Jung, K.-H., and Spudich, E.N. (2000). Retinylidene Proteins: Structures and Functions from Archaea to Humans. *Annu. Rev. Cell Dev. Biol.* *16*, 365-392.
- Stock, A. M., Robinson, V. L., und Goudreau, P. N. Two-component signal transduction. *Annu. Rev. Biochem.* *69*, 183-215. 2000.
- Stowe-Evans, E.L., Harper, R.M., Motchoulski, A.V., and Liscum, E. (1998). NPH4, a conditional modulator of auxin-dependent differential growth responses in *Arabidopsis*. *Plant Physiol.* *118*, 1265-1275.
- Swartz, T.E., Corchnoy, S.B., Christie, J.M., Lewis, J.W., Szundi, I., Briggs, W.R., and Bogomolni, R.A. (2001). The photocycle of a flavin-binding domain of the blue light photoreceptor phototropin. *J Biol Chem.* *276*, 36493-36500.
- Sweere, U., Eichenberg, K., Lohrmann, J., Mira-Rodado, V., Baurle, I., Kudla, J., Nagy, F., Schäfer, E., and Harter, K. (2001). Interaction of the response regulator ARR4 with phytochrome B in modulating red light signaling. *Science* *294*, 1108-1111.
- Tanaka, N., Ogura, T., Noguchi, T., Hirano, H., Yabe, N., and Hasunuma, K. (1998). Phytochrome-mediated light signals are transduced to nucleoside diphosphate kinase in *Pisum sativum* L. cv. Alaska. *J. Photochem. Photobiol. B* *45*, 113-121.
- Taylor, B. L. und Zhulin, I. B. Pas Domains - Internal Sensors of Oxygen, Redox Potential, and Light. *Microbiology and Molecular Biology Reviews* Vol 63[2], 479. 1999.

- Terauchi,K., Montgomery,B.L., Grossman,A.R., Lagarias,J.C., and Kehoe,D.M. (2003). RcaE is a Complementary Chromatic Adaptation Photoreceptor Required for Green and Red Light Responsiveness. *Mol. Microbiol.* 51, 567-577.
- Terry,M.J. (1997). Phytochrome chromophore-deficient mutants. *Plant Cell Environ.* 20, 740-745.
- Terry,M.J., Maines,M.D., and Lagarias,J.C. (1993). Inactivation of phytochrome- and phycobiliprotein-chromophore precursors by rat liver biliverdin reductase. *J. Biol. Chem.* 268, 26099-26106.
- Terry,M.J., McDowell,M.T., and Lagarias,J.C. (1995). (3Z)- and (3E)-phytochromobilin are intermediates in the biosynthesis of the phytochrome chromophore. *J. Biol. Chem.* 270, 11111-11118.
- Thompson,J.D., Gibson,T.J., Plewniak,F., Jeanmougin,F., and Higgins,D.G. (1997). The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24, 4876-4882.
- Thümmler,F. and Rüdiger,W. (1983). Models for the photoreversibility of phytochrome. Z,E isomerisation of chromopeptides from phycocyanin and phytochrome. *Tetrahedron* 39, 1943-1951.
- Toyama, A., Nakazawa, M., Manabe, K., Takeuchi, H., und Harada, I. Ultraviolet Resonance Raman-Spectra of Phytochrome - A Comparison of the Environments of Tryptophan Side-Chains Between Red-Light Absorbing and Far Red-Light Absorbing Forms. *Photochem.Photobiol.* Vol 57[Iss 2], 391-395. 1993.
- Tzfira,T. and Citovsky,V. (2002). Partners-in-infection:host proteins involved in the transformation of plant cells by *Agrobacterium*. *Trends Cell Biol* 12, 121-129.
- van Thor,J.J., Borucki,B., Crielaard,W., Otto,H., Lamparter,T., Hughes,J., Hellingwerf,K.J., and Heyn,M.P. (2001). Light-induced proton release and proton uptake reactions in the cyanobacterial phytochrome Cph1. *Biochemistry* 40, 11460-11471.
- Vierstra,R.D. and Davis,S.J. (2000). Bacteriophytochromes: new tools for understanding phytochrome signal transduction. *Semin. Cell Dev. Biol.* 11, 511-521.
- Vierstra,R.D., Quail,P.H., Hahn,T.R., and Song,P.S. (1987). Comparison of the protein conformations between different forms (Pr and Pfr) of native (124 kDa) and degraded (118/114 kDa) phytochromes from *Avena sativa*. *Photochem. Photobiol.* 45, 429-432.
- Vince-Prue,D. (1994). The duration of light and photoperiodic responses. In *Photomorphogenesis in plants*, R.E.Kendrick and G.H.M.Kronenberg, eds. (Dordrecht, The Netherlands: Kluwer Academic), pp. 447-490.
- Vinograd,J. and Lebowitz,J. (1966). Physical and topological properties of circular DNA. *J. Gen. Physiol.* 49, 103-125.
- Wagner,D., Fairchild,C.D., Kuhn,R.M., and Quail,P.H. (1996). Chromophore-bearing NH<sub>2</sub>-terminal domains of phytochromes A and B determine their photosensory specificity and differential light lability. *Proc. Natl. Acad. Sci. U. S. A.* 93, 4011-4015.

- Wang,H., Ma,L.G., Li,J.M.Z.H.Y., and Deng,X.W. (2001). Direct interaction of Arabidopsis cryptochromes with COP1 in light control development. *Science* 294, 154-158.
- Wang,X. and Iino,M. (1998). Interaction of cryptochrome 1, phytochrome, and ion fluxes in blue- light-induced shrinking of arabidopsis hypocotyl protoplasts. *Plant Physiol.* 117, 1265-1279.
- Wang,Y.-C., Stewart,S.J., Cordonnier,M.-M., and Pratt,L. (1991). Avena sativa L. contains three phytochromes, only one of which is abundant in etiolated tissue. *Planta* 184, 96-104.
- Weller,J.L., Perrotta,G., Schreuder,M.E., van Tuinen,A., Koornneef,M., Giuliano,G., and Kendrick,R.E. (2001). Genetic dissection of blue-light sensing in tomato using mutants deficient in cryptochrome 1 and phytochromes A, B1 and B2. *Plant J* 25, 427-440.
- Wells,T.A., Nakazawa,M., Manabe,K., and Song,P.S. (1994). A conformational change associated with the phototransformation of Pisum phytochrome A as probed by fluorescence quenching. *Biochemistry* 33, 708-712.
- West,A.H. and Stock,A.M. (2001). Histidine kinases and response regulator proteins in two-component signaling systems. *Trends. Biochem Sci.* 26, 369-376.
- Whitelam,G.C., Johnson,E., Peng,J., Carol,P., Anderson,M.L., Cowl,J.S., and Harberd,N.P. (1993). Phytochrome A null mutants of *Arabidopsis* display a wild-type phenotype in white light. *Plant Cell* 5, 757-768.
- Wilde,A., Churin,Y., Schubert,H., and Börner,T. (1997). Disruption of a Synechocystis sp. PCC 6803 gene with partial similarity to phytochrome genes alters growth under changing light qualities. *FEBS Lett.* 406, 89-92.
- Wood,D.W., Setubal,J.C., Kaul,R., Monks,D.E., Kitajima,J.P., Okura,V.K., Zhou,Y., Chen,L., Wood,G.E., Almeida,N.F.J., Woo,L., Chen,Y., Paulsen,I.T., Eisen,J.A., Karp,P.D., Bovee,D.S., Chapman,P., Clendenning,J., Deatherage,G., Gillet,W., Grant,C., Kutayavin,T., Levy,R., Li,M.J., McClelland,E., Palmieri,A., Raymond,C., Rouse,G., Saenphimmachak,C., Wu,Z., Romero,P., Gordon,D., Zhang,S., Yoo,H., Tao,Y., Biddle,P., Jung,M., Krespan,W., Perry,M., Gordon-Kamm,B., Liao,L., Kim,S., Hendrick,C., Zhao,Z.Y., Dolan,M., Chumley,F., Tingey,S.V., Tomb,J.F., Gordon,M.P., Olson,M.V., and Nester,E.W. (2001). The genome of the natural genetic engineer *Agrobacterium tumefaciens* C58. *Science* 294, 2317-2323.
- Wu,S.H. and Lagarias,J.C. (1997). The phytochrome photoreceptor in the green alga *Mesotaenium caldariorum*: Implication for a conserved mechanism of phytochrome action. *Plant Cell Environ.* 20, 691-699.
- Wu,S.H. and Lagarias,J.C. (2000). Defining the bilin lyase domain: lessons from the extended phytochrome superfamily. *Biochemistry* 39, 13487-13495.
- Wu,S.H., McDowell,M.T., and Lagarias,J.C. (1997). Phycocyanobilin is the natural precursor of the phytochrome chromophore in the green alga *Mesotaenium caldariorum*. *J. Biol. Chem.* 272, 25700-25705.
- Yamaguchi,R., Nakamura,M., Mochizuki,N., Kay,S.A., and Nagatani,A. (1999). Light-dependent translocation of a phytochrome B-GFP fusion protein to the nucleus in transgenic Arabidopsis. *J Cell Biol* 145, 437-445.

- Yang,H.Q., Wu,Y.J., Tang,R.H., Liu,D., Liu,Y., and Cashmore,A.R. (2000). The C termini of Arabidopsis cryptochromes mediate a constitutive light response. *Cell* 103, 815-827.
- Yeh,K.C. and Lagarias,J.C. (1998). Eukaryotic phytochromes: Light-regulated serine/threonine protein kinases with histidine kinase ancestry. *Proc. Natl. Acad. Sci. U. S. A.* 95, 13976-13981.
- Yeh,K.C., Wu,S.H., Murphy,J.T., and Lagarias,J.C. (1997). A cyanobacterial phytochrome two-component light sensory system. *Science* 277, 1505-1508.
- Yoshihara,S., Suzuki,F., Fujita,H., Geng,X.X., and Ikeuchi,M. (2000). Novel putative photoreceptor and regulatory genes required for the phototactic movement of the unicellular motile cyanobacterium *Synechocystis* PCC 6803. *Plant Cell Physiol.* 41, 1299-1304.
- Zambre,M., Terryn,N., De Clercq,J., De Buck,S., Dillen,W., Van Montagu,M., Van Der Straeten,D., and Angenon,G. (2003). Light strongly promotes gene transfer from *Agrobacterium tumefaciens* to plant cells. *Planta* 216, 580-586.
- Zeidler,M., Lamparter,T., Hughes,J., Hartmann,E., Remberg,A., Braslavsky,S., Schaffner,K., and Gärtner,W. (1998). Recombinant phytochrome of the moss *Ceratodon purpureus*: heterologous expression and kinetic analysis of Pr $\rightarrow$ Pfr conversion. *Photochem. Photobiol.* 68, 857-863.
- Zeiger,E. (2000). Sensory transduction of blue light in guard cells. *Trends Plant Sci* 5, 183-185.
- Zeiger,E. and Helper,P. (1977). Light and stomatal function: Blue light stimulates swelling of guard cell protoplasts. *Science* 196, 887-889.
- Zhu,J., Oger,P.M., Schrammeijer,B., Hooykaas,P.J., Farrand,S.K., and Winans,S.C. (2000). The bases of crown gall tumorigenesis. *J Bacteriol* 182, 3885-3895.
- Zhulin,I.B., Taylor,B.L., and Dixon,R. (1997). PAS domain S-boxes in Archaea, Bacteria and sensors for oxygen and redox. *Trends. Biochem. Sci.* 22, 331-333.
- Zimmermann,S., Baumann,A., Jaekel,K., Marbach,I., Engelberg,D., and Frohnmeyer,H. (1999). UV-responsive genes of arabidopsis revealed by similarity to the Gcn4-mediated UV response in yeast. *J. Biol. Chem.* 274, 17017-17024.

