

8. Literatur

- [1] Abosh D, Rosser B, Kaita K. Outcomes following liver transplantation for patient with alcohol-versus nonalcohol-induced liver disease: Can J Gastroenterology 2000; 14(10): 851- 5.
- [2] Abrignani S. Antigen-independent activation of resting T cells in the liver of patients with chronic hepatitis. Dev Biol Stand 1998; 92: 191- 4.
- [3] Adachi Y, Bradford BU, Gao W. Inactivation of kupffer cells prevents early alcohol-induced liver injury. Hepatology 1994; 20: 453- 60.
- [4] Adachi Y, Moore LE, Bradford BU. Antibiotics prevent liver injury in rats following long-term exposure to ethanol. Gastroenterology 1995; 108: 218- 24.
- [5] Adinoff B, Martin PR, Bone GHA. Hypothalamic-Pituitary-Adrenal Axis functioning and cerebrospinal fluid corticotropin releasing hormone and corticotropin levels in alcoholics after recent and long-term abstinence. Arch Gen Psychiatry 1990; 47: 325- 30.
- [6] Adinoff B, Risher-Flowers D, De Jong J. Disturbances of HPA-axis functioning during ethanol withdrawal in six men. Am J Psychiatry 1991; 148(8): 1023- 5.
- [7] Allemann PI. Marker für übermässigen Alkoholkonsum (Screening). Therapeutische Umschau Verlag Hans Huber, Bern 2000; 57(4):185-90.
- [8] Anderson FH, Zeng L, Rock NR. An assessment of the clinical utility of serum ALT and AST in chronic hepatitis. Hepatology Research 2000; 18(1): 63- 71.
- [9] Anthony PP, Ishak KG, Nayak NC. The morphology of cirrhosis: J Clin Pathol 1978; 31: 395- 14.
- [10] Armstrong L, Jordan N, Millar A. Interleukin-10 regulation of tumor necrosis factor alpha from human alveolar macrophages and peripheral blood monocytes. Thorax 1996; 51: 143- 49.
- [11] Basista MH, Gavaler J, Stieffenhofer A. Effect of ethanol on Kupffer cell funktion. Alcohol Clin Exp Res 1993; 17(3): 556- 60.
- [12] Batey RG, Clancy RL, Pang GT. Alcoholic hepatitis as a T-cell mediated disorder: an hypothesis. Alcohol Clin Exp Res 1999; 23(7): 1207- 09.
- [13] Baxter F. Septic Shock. Can J Anaesth 1997; 44: 59- 72.
- [14] Biffl WL, Moore EE, Moore F. Interleukin-6 in the injured patient: marker of injury or mediator of inflammation? Ann Surg 1996;224(5):647-64.
- [15] Bode C, Bode JC, Hahn EG, Rossol S, Schäfer C, Schuppmann. Alkohol und Leber. In: Singer MV, Teysen S (Hrsg) Kompendium Alkohol Folgekrankheiten, Klinik, Diagnostik, Therapie. Springer, Berlin Heidelberg New York Toxyo,199-249.
- [16] Bode JC, Heidelbach R, Durr HK. Jejunal microflora in patients with chronic alcoholic abuse: Hepatogastroenterology 1984; 31: 30- 34.
- [17] Bone RC. Toward a theory regarding the pathogenesis of the systemic inflammatory response syndrome: What we do and do not know about cytokine regulation. Crit Care Med 1996; 24: 163- 72.
- [18] Borelli E, Sutter PM, Roux. Lombard P. Interleukin-6 and tumor necrosis factor plasma levels in the development of multiple organ failure. Ares Serono Symp. Abstract Book, Montreux 1991; p49.
- [19] Boyer N, Marcellin P. Pathogenesis, diagnosis and management of hepatitis C. J of Hepatology 2000; 32: 98-112.
- [20] Breuer JP, Neumann T, Heinz A, Kox WJ, Spies C. Der Alkoholkranke als Patient. Wien Klin Wochenschr Springer-Verlag 2003; 115(17-18):618-33.

- [21] Damas P, Reuter A, Gysen P, Demonty J, Lamy M, Franchimont P. Tumor necrosis factor- alpha and interleukin-1 serum levels during severe sepsis in humans. Crit Care Med 1989; 17: 975- 78.
- [22] De Bleser PJ, Braet F, Lovisetti P. Cell biology of liver endothel and Kupffer cells. Gut 1994; 35: 1509- 16.
- [23] De Waal Malefy R, Abrams J, Bennett B. Interleukin-10 inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes: J Exp Med 1991; 174: 1209- 20.
- [24] Decker T, Lohmann-Matthes ML, Karck U. Comparative study of cytotoxicity, TNF, and prostaglandin release after stimulation of rat Kupffer cells, murine Kupffer cells, and murine inflammatory liver macrophages. J Leukoc Biol 1989; 45:139- 46.
- [25] Del Arbol JL, Aguirre JC, Raya J. Plasma concentrations of β -Endorphin, Adrenocorticotropic Hormone, and Cortisol in drinking and abstinent chronic alcoholics. Alcohol 1995; 12(6): 525- 29.
- [26] Deviere J, Content J, Denys C. High interleukin-6 levels and increased production by leucocytes in alcoholic liver cirrhosis. Correlation with IgA serum levels and lymphokines production. Clin Exp Immunol 1989; 77: 221- 25.
- [27] Diez Ruiz A, Santos Perez JL. Tumor necrosis factor, interleukin-1 and interleukin-6 in alcoholic cirrhosis. Alcohol Alcohol 1993 ;28 :319.
- [28] Ehrenreich H, Schuck J, Stender N. Endocrine and hemodynamic effects of stress versus systemic CRF in alcoholics during early and medium term abstinence. Alcohol Clin Exp Res 1997; 21(7): 1285. 93.
- [29] Engelberts I, Möller A, Leeuwenberg JFM, van der Linden CJ, Buurman WA. Administration of tumor necrosis factor α inhibitors after exposure to TNF α prevents development of the maximal biological effect: An argument for clinical treatment with TNF α -inhibitors. J Surg Res 1992; 53: 510- 14.
- [30] Ewing JA. Detecting Alkoholism,The Cage Questionnaire.JAMA1984; 252:1905- 07.
- [31] Farges O, Saliba F, Farhamant H, et al. Incidence of rejection and infection after liver transplantation as a function of primary disease: possible influence of alcohol and polyclonal immunoglobulins. Hepatology 1996;23:240.
- [32] Faunce DE, Gregory MS, Kovacs EJ. Glucocorticoids protect against suppression of T cell responses in a murine model of acute ethanol exposure and thermal injury by regulating IL-6. J Leuk Biol 1998; 64: 724- 32.
- [33] Fernandez-Checa JC, Kaplowitz N, Garcia Ruiz C. GSH transport in mitochondria: defense against TNF-induced oxidative stress and alcohol-induced defect. Am J Physiol Gastrointest Liver Physiol 1997; 273: 7-17.
- [34] Fiorentino F, Zlotnik A, Mosmann TR. IL-10 inhibits cytokine production by activated macrophages. J Immunol 1991; 147(11): 3815- 22.
- [35] Friedman SL. Molecular regulation of hepatic fibrosis, an integrated cellular response to tissue injury. J Biol Chem 2000; 2247- 50.
- [36] Garner JS, Jarvis WR, Emori TG. CDC definitions for nosocomial infections. Am J Infection Control 1988; 16: 128- 40.
- [37] Goebel KM, Goebel FD, Schubotz R, Schneider J. Red cell metabolic and membrane feature in haemolytic anaemia of alcoholic liver disease (Zieve's syndrome). Br. J Haematol. 1977; 35(4):573-83.
- [38] Goto M, Lemasters JJ, Thurman RG. Activation of voltage- dependent calcium channels in kupffer cells by chronic treatment with alcohol in the rat. The Journal of pharmacology and experimental therapeutics 1993; 267(1): 1264- 68.
- [39] Grant BF. DSM-III -R and proposed DSM- IV alcohol abuse and dependence, United states 1988: A nosological study. Alcohol Clin Exp Res 1988; 16: 1068- 77.

- [40] Hanck C, Glatzel M, Singer MV. Gene expression of TNF receptor in peripheral blood mononuclear cells of patient with alcoholic cirrhosis. *J Hepatol* 2000; 32(1): 51- 7.
- [41] Hayashi M, Keeffe EB, Krams SM, et al. Allograft rejection after liver transplantation for autoimmune liver disease. *Liver Transplantation* 1998; 4:208.
- [42] Heijnen CJ, Kavelaars A, Ballieux RE. Beta-endorphin: cytokine and neuropeptide. *Immunol Rev* 1991; 119: 41- 63.
- [43] Heinrich PC, Castell JV, Andus T. Interleukin-6 and the acute phase response. *Biochem Journal* 1990; 265: 621- 36.
- [44] Hempel L, Körholz D, Bönig H. Interleukin-10 directly inhibits the interleukin-6 production in T-cells. *Scand J Immunol* 1995; 41: 462- 66.
- [45] Herold G, Herold 2004 Köln;462.
- [46] Heumann D, Gallay P, Barras C. Control of LPS binding and LPS-induced tumor necrosis factor secretion in human peripheral blood monocytes. *J Immunol* 1992; 148: 3505- 12.
- [47] Honchel R, Ray Mukunda B, Marsano L. Tumor Necrosis Factor in alcohol enhanced endotoxin liver injury. *Alcohol Clin Exp Res* 1992; 16(4): 665- 69.
- [48] Kakumu S, Okumura A, Ishikawa T, Iwata K, Yano M, Yoshioka K. Production of interleukin 10 and 12 by peripheral blood mononuclear cells in chronic hepatitis C virus infection. *Clin Exp Immunol* 1997;108:38.
- [49] Kavakami M, Switzer BR, Herzog SR. Immune suppression after acute ethanol ingestion and thermal injury. *J Surg Res* 1991; 51(3): 210- 05.
- [50] Khoruts A, Stahnke L, McClain GJ. Circulating tumor necrosis factor, IL-1 and IL-6 concentration in chronic alcoholic patients. *Hepatology* 1991; 13: 267- 76.
- [51] Kievolt-Glaser JK, Preacher KJ, MacCallum RC. Chronic stress and age-related increases in the proinflammatory cytokine IL-6. *PNAS* 2003; 100(15): 9090- 95.
- [52] Kirmaz C, Terzioglu E, Topalak O,et al. Serum tumor growth factor-beta1 levels in patient with cirrhosis, chronic hepatitis B and chronic hepatitis C. *Eur Cytokine Netw* 2004;15:112.
- [53] Kishimoto T. The biology of Interleukin-6. *Blood* 1989; 74(1): 1-10.
- [54] Kjell O, Björk P, Bergenfeldt M, Hageman R, Thomsen RC. Interleukin-1 receptor antagonist reduced mortality from endotoxin shock. *Nature* 1990; 348: 550- 52.
- [55] Klava A, Windsor ACJ,Farmery SM. IL-10. *Arch Surg* 1997; 132:425- 29.
- [56] Knolle P, Schlaak J, Uhrig A. Human Kupffer cells secrete IL-10 in response to LPS challenge. *J Hepatology* 1995; 22: 226- 29.
- [57] Knolle P, Uhrig A, Hegenbarth S. IL-10 down-regulates t cell activation by antigen-presenting liver sinusoidal endothelial cells through decreased antigen uptake via the maaose receptor and lowered surface expression of accessory molecules. *Clin Exp Immunol* 1998; 114: 427- 33.
- [58] Koivisto T; Mishin VM, Mak KM. Induction of cytochrome P-450E1 by ethanol in ret kupffer cells. *Alcohol Clin Exp Res* 1996; 20:207- 12.
- [59] Krom RA. Liver transplantation and alcohol: who should get transplants? *Hepatology* 1994; 20: 28-32.
- [60] Louis H, Le Moine O, Peny MO. Hepatoprotective role of IL-10 in galactosamine/Lipopolysaccharide mouse liver injury. *Gastroenterology* 1997; 112: 935- 42.
- [61] Lumeng L, Li TK. Vitamin B6 metabolism in chronic alcohol abuse. Pyridoxal phosphate synthesis and degradation in human erythrocytes. *J Clin Invest* 1974; 53: 693- 97.
- [62] Malik P, Bogetti D, Sileri P, Testa G, Blumenthal SA, Wiley TE, Layden TJ, Benedetti E. Spur cell anemia in alkoholic cirrhosis: cure by orthopic liver transplantation and recurrence after liver graft failure. *Int Surg*. 2002; 87(4):201-04.

- [63] Marchesi C, Chiodera P, Ampollini P. Beta-endorphin, adrenocorticotrophic hormone and cortisol secretion in abstinent alcoholics. *Psychiatry research* 1997; 72: 187- 94.
- [64] Maruyama S, Hirayama C, Yamamoto S. Red blood cell status in alcoholic and non-alcoholic liver disease. *J Lab Clin Med* 2001; 138(5): 332- 7.
- [65] Massignon D, Lepape A, Bienvenu J, Barbier Y, Boileau C, Coeur P. Coagulation/fibrinolysis balance in septic shock related to cytokines and clinical state. *Haemostasis* 1994; 24: 36- 48.
- [66] Matloff DS, Selinger MJ, Kaplan MM. Hepatic Transaminase Activity in Alcoholic liver disease: *Gastroenterology* 1980; 78: 1389- 92.
- [67] Mc Guinness PH, Painter D, Davies S. Increases in intrahepatic CD68 positive cells, MAC387 positive cells, and inflammatory cytokines in chronic hepatitis C infection: *Gut* 2000; 46(2): 260- 9.
- [68] Mc Clain C, Hill D, Schmidt J. Cytokines and alcoholic liver disease. *Semin Liver Dis* 1993; 13: 170- 82.
- [69] Mc Cuskey RS, Nishida J, Eguchi H. Role of endotoxin in the hepatic microvascular inflammatory response to ethanol. *J Gastroenterol Hepatol* 1995; 10(1)Suppl: 18-23.
- [70] Mc Sween RNM, Burt AD. Histologic spectrum of alcoholic liver disease. *Semin Liver Dis* 1986; 6: 221- 32.
- [71] Moore RD, Bone LR, Geller G, Mamon JA, Stokes EJ, Levine DM. Prevalence, detection and treatment of alcoholism in hospitalised patients. *JAMA* 1989; 261: 403- 07.
- [72] Nalpas B, Vassault A, Charpin S. Mitochondrial Aspartate Aminotransferase as a Marker of chronic alcoholism: Diagnostic Value and Interpretation in a Liver Unit. *Hepatology* 1986; 6(4): 608- 14.
- [73] Napoli J, Bishop GA, McGuinness PH, Painter DM. Progresive liver injury in chronic Hepatitis C infection correlates with increased intrahepatic expression of Th1-associated cytokines. *Hepatology* 1996; 24: 759- 65.
- [74] Napolitano LM, Koruda MJ, Zimmermann K, McCowan K, Chang J, Meyer AA. Chronic ethanol intake and burn injury: evidence for synergistic alteration in gut and immune integrity. *J Trauma* 1995; 38: 198-07.
- [75] Neidhardt R, Keel M, Steckholzer U. Relationship of interleukin-10 plasma levels to serverity of injury and clinical outcome in injured patients. *The Journal of Trauma: Injury, infection, and critical care* 1997; 42(5): 863- 71.
- [76] Neta R, Oppenheim JJ. Why should internists be interested in interleukin-1. *Ann Intern Med* 1988; 109(1).
- [77] Neuberger J. Incidence, timing, and risk factors for acute and chronic rejection. *Liver Transpl Surg* 1999;5:30.
- [78] Oyanagi Y, Takahashi T, Matsui S. Enhanced expression of interleukin-6 in chronic hepatitis C. *Liver* 1999; 19: 464- 72.
- [79] Palluy O, Morliere L, Gris JC, Bonne C, Modat G. Hypoxia/reoxygenetion stimulates endothelium to promote neutrophil adhesion. *Free Rad Biol Med* 1992; 13: 21- 30.
- [80] Pereira SP, Williams R. Alcohol relapse and functional outcome after liver transplantation for alcoholic liver disease. *Liver Transplantation* 2001; 7(3): 1-4.
- [81] Platz KP, Mueller AR, Spree E et al. Liver transplantation for alcoholic cirrhosis. *Transpl Int* 2000;13:127.
- [82] Preuss R, Renz- Polster H, Reuter H, Wellhoehner P. 7. Kapitel: Leber, Galle, Pankreas. In: Renz- Polster H, Braun J. Basislehrbuch Innere Medizin. 2. korrigierte Auflage, Urban und Fischer Verlag München, Jena 1999; 601- 52.
- [83] Preuss R; Reuter H, Wellhoehner P. 7. Kapitel: Child-Pugh-Score zur Prognoseabschätzung bei Leberzirrhose. Basislehrbuch Innere Medizin. 2. korrigierte Auflage, Urban und Fischer Verlag München, Jena 1999; 641.

- [84] Pruett SB, Collier SD, Wu WJ. Ethanol induced activation of the HPA-axis in a mouse model. *Life Sci* 1998; 63(13): 1137- 46.
- [85] Reinert, Duane F, Allen, John P. The Alcohol Use Disorders Identification Test (AUDIT): A Review of Recent Research. *Alcohol Clin Exp Res*, 2002; 26(2):272-9.
- [86] Rosser BG, Gores GJ. Liver cell necrosis: cellular mechanisms and clinical implications. *Gastroenterology* 1995; 108: 252- 75.
- [87] Rubin E, Lieber C. Alcohol induced hepatic injury in nonalcoholic volunteers. *N Engl J Med* 1968; 278: 869- 72.
- [88] Rudell WAS, Axon ATR, Findlay JM. Effect of cimetidine on the gastric bacterial flora. *Lancet* 1980; 1: 672- 74.
- [89] Sander M, Irwin M, Sinha P, Naumann E, Kox WJ, Spies CD. Suppression of interleukin-6 to interleukin-10 in chronic alcoholics:association with postoperative infections. *Intensive Care Med* 2002;28:285.
- [90] Sander M, Naumann E, Irwin M. Decreased interleukin6/10 ratios in chronic alcoholics and postoperative infections in the intensive care unit. *Intensive Care Med* 2002 15A
- [91] Sarkar DK. The neuroendokrino-immune axis of alcoholics. *Alcohol Clin Exp Res* 1996;20(Suppl.): 256- 59.
- [92] Saunders JB, Aasland OG, Babor TF. Development of the alcohol use disorder identification test. *Addiction* 88 1993; 791- 03.
- [93] Schäfer M, Mousa SA, Stein C. CRF in antinociception and inflammation. *Eur J Pharmacol* 1997; 323(1): 1-10. Review.
- [94] Scharf RE, Aul C. Alcohol-induced disorder of the hematopoietic system. *Gastroenterol*. 1988; Suppl 3:75-83.
- [95] Schulze-Osthoff K, Beyaert R, Vandevorde V. Depletion of the mitochondrial electron transport abrogates the cytotoxic and gene-inductive effects of TNF. *EMBO J* 1993; 12(8): 3095- 104.
- [96] Shong M, Ro HK, Kim YK, Yoo CJ, Lee JH, Song CU, Cho BY. The cytokines, interleukin-1 beta, interleukin-6 and interferon- gamma upregulate the expression of intercellular adhesion molecule-1 (ICAM-1) in rat thyroid cell line, FRTL-5. *Korean J Intern Med* 1994; 8: 88- 92.
- [97] Song C, Lin A, De Long R. Cytokines in detoxified patients with chronic alcoholism without liver disease: increased monocytic cytokine production. *Biological Psychiatry* 1999. 45(9): 1212- 16.
- [98] Sorensen TIA, Orholm M, Bensten KD, Hoybye G, Eghoje K, Christoffersen P. Prospective evalution of alcohol abuse and alcoholic liver disease in men as predictors of development of cirrhosis. *Lancet* 1984; 2: 241- 47.
- [99] Spies C, Neuner B, Neumann T, Blum S. Intercurrent complication in chronic alcoholic men admitted to the intensive care unit following trauma. *Intensive Care Med* 1996; 22: 286- 93.
- [100] Spies C, Nordmann A, Brummer G, Marcs C. Intensive care unit stay is prolonged in chronic alcoholic men following tumor resection of the upper digestive tract. *Acta Anaesthesiol Scand* 1996; 40: 649- 56.
- [101] Spies C, Rommelspacher H. Alcohol withdrawal in the surgical patient: prevention and treatment. *Anesth Analg* 1999; 88(4): 946-54.
- [102] Spies C, Tonnesen H, Andreasson S, Helander A, Conigrave K. Perioperative morbidity and mortality in chronic alcoholic patients. *Alcohol Clin Exp Res* 2001; 25: 164S-70S.

- [103] Spies C, von Dossow V, Eggers V, Jetschmann G, El-Hilali R, Egert J, Fischer M, Schröder T, Höflich C, Sinha P, Paschen C, Mirsalim P, Brunsch R, Hopf J, Marks C, Wernecke KD, Pragst F, Ehrenreich H, Müller C, Tonnesen H, Oelkers W, Rohde W, Stein C, Kox WJ. Altered Cell-mediated Immunity and Increased Postoperative Infection Rate in Long-term Alcoholic Patients. *Anaesthesiology* 2004; 100: 1088-100.
- [104] Stefanini GF, Biselli M, Grazi GL. *Hepatogastroenterology* 1997; 44(7): 1356- 9.
- [105] Szabo G, Mandrekar P, Girouard L. Regulation of human monocyte functions by acute ethanol treatment: Decreased Tumor necrosis factor alpha, Interleukin-1beta and elevated Interleukin-10, and Transforming growth factor beta production. *Alcohol Clin Exp Res* 1996; 20(5): 900- 07.
- [106] Szabo G. Consequences of alcohol consumption on host defence. *Alcohol Alcohol* 1999; 34(6): 830- 41. Review.
- [107] Szuster-Cieselska A, Daniluk J, Kanderfer-Zerszen M. Serum levels of cytokines in alcoholic liver cirrhosis and pancreatitis. *Arch Immunol Ther Exp (Warsz)* 2000; 48(4): 301-7.
- [108] Tonnessen H, Rosenberg J, Nielson HJ. Effect of preoperative abstinence on poor postoperative outcome in alcohol misusers: randomised controlled trial. *British Medical Journal* 1999; 318: 1311- 16.
- [109] Tonnessen H, Petersoen KR, Hoigaard L. Postoperative morbidity among symptom-free alcohol misusers. *Lancet* 1992; 340: 334- 37.
- [110] Tracey KJ. Tumour Necrosis Factor- Alpha. In: Thomson A. *The Cytokine Handbook*. 2. Auflage, Department of Surgery and Molecular genetics and Biochemistry, School of Medicine, University of Pittsburgh, USA 1994; 289- 04.
- [111] Van Gool J, Van Vugt H, Helle M. The relation among stress, adrenalin, interleukin-6 and acute phase proteins in the rat. *Clinical Immunology and Immunopathology* 1990; 57: 200- 10.
- [112] Videla L, Bernstein J, Israel Y. Metabolic alteration produced in the liver by chronic alcohol administration. *Biochem J* 1973; 134: 507- 14.
- [113] Vincent JL, Bihari D. Sepsis, severe sepsis or sepsis syndrom: need for clarification. *Intensiv Care Med* 1992; 18: 255- 57.
- [114] Von Bardeleben U, Heuser I, Holsboer F. Human CRH stimulation response during acute withdrawal and after medium-term abstinence from alcohol abuse. *Psychoneuroendocrinology* 1989; 14(6): 441- 49.
- [115] Von Dauciuc I, Alappat JM, De Souza. Effect of acut and chronic alcohol administration to rats on the expression of interleukin-6 cell-surface receptors of hepatic parenchymal and nonparenchymal cells. *Alcohol Clin Exp Res* 1994; 18(5): 1207-14.
- [116] Von Deauciuc L. Alcohol and cytokine networks. *Alcohol* 1997; 14(5): 421- 30.
- [117] Von Heymann C, Langenkamp J, Dubisz N. Posttraumatic immune modulation in chronic alcoholics is associated with multiple organ dysfunction syndrome. *Trauma* 2002; 52(1): 95- 103.
- [118] Waage A, Slupphang G, Shalaby R. Glucocorticoids inhibit the production of IL-6 from monocytes, endothelial cells and fibroblasts. *Eur J Immunol* 1990; 20: 2439- 43.
- [119] Wang P, Wu P, Siegel MI. IL-10 inhibits NFkB activation in human monocytes. *The american society for biochemistry and molecular biology* 1995; 270(16): 9558- 63.
- [120] Wilckens T, De Rijk R. Glucocorticoids and immune function: unknown dimensions and new frontiers: *Immunology* 1997; 18(9): 418- 24.
- [121] Winwood PJ, Arthur MJP. Kupffer cells: their activation and role in animal models of liver injury and human liver disease. *Seminars in liver disease* 1993; 13(1): 50- 59.

- [122] World health Organisation. The ICD- 10 classification of mental and behavioural disorders In:. Clinical discriptions and diagnostic guidelines (10th revision). (Monographie) Geneva. World Health Organisation 1992.
- [123] Yoshidome H, Kato A, Edwards MJ. IL-10 suppresses hepatic ischemia/reperfusion injury in mice: implications of a central role for nuklear factor kB. Hepatology 1999; 30: 203- 08.
- [124] Yoshioka K, Kakumu S, Arao M. Tumor necrosis factor alpha production by peripheral blood mononuclear cells of patients with chronic liver disease. Hepatology 1989; 10(5): 769- 73.