

## 8. Literaturverzeichnis

Abraham, E.J., Morris-Hardemann, J.N., Swenson, L.M., Knoppel, E.L., Ramanathan, B., Wright, K.J., Grieger, D.M., Minton, J.E. (1998)

Pituitary function in the acute phase response in domestic farm animals: cytokines, prostaglandins and secretion of ACTH.

Domestic animal endocrinology, Vol. 15 (5), 389-396.

Ahlers, D., J. Aurich, E. Bleckmann, A. Bieneck, M. Emmert, M. Hoedemaker, J. Lange, S. Mansfeld, P. Scheibl, W. Zaremba, and H. Zerbe. (2000).

Verlauf des Puerperiums und das weitere Schicksal von Rindern ohne und mit Retention secundinarum nach Abkalbung mit tierärztlicher Geburtshilfe und intrauteriner Applikation von Ampicillin-Cloxacillin- oder Tetracyclin-haltigen Uterusstäben.

Tierärztl. Umschau 55:479-488.

Aiello, S.E. (Hrsg.) (1998)

Anti-inflammatory agents.

in: The Merck veterinary manual. 8<sup>th</sup> edition, 1817-1831.

Aiumlamai, S., Odensvik, K., Stabenfeldt, K., Kindahl, H. (1990)

Regulation of prostaglandin biosynthesis with flunixin meglumine in the bovine species.

J. Vet. Med. A, 37, 16-22.

Alsemgeest, S.P., Kalsbeek, H.C., Wensing, T., Koeman, J.P., van Ederen, A.M., Gruys, E. (1994)

Concentrations of serum amyloid-A (SAA) and haptoglobin (HP) as parameters of inflammatory diseases in cattle.

Vet. Q. (1), 21-23.

Amiridis, G.S., Leontides, L., Tassos, E., Kostoulas, P., Fthenakis, G.C. (2001)

Flunixin meglumine accelerates uterine involution and shortens the calving-to-first-oestrus interval in cows with puerperal metritis.

J. vet. Pharmacol. Therap., 24, 365-367.

Anderson, K.L., Neff-Davis, C.A., Davis, L.E., Bass, V.D. (1990)

Pharmakokinetik of flunixin meglumine in lactating cattle after single multiple intramuscular and intravenous administrations.

Am J Vet Res, Vol 51, No. 9, 1464-1467.

Aurich, J. (1995)

Der Endometritis Pyometra Komplex bei Großtieren.

in: J. Aurich, A. Kuntze, (Hrsg.): Vet Spezial, Verlag Gustav Fischer Jena und Stuttgart, S. 51-107.

Aurich, J.E. ,und Grunert, E. (1996)

Pathogenese und Therapie postpuerperaler Endometritiden beim Rind.

Praktischer Tierarzt, Coll. vet. XXVI, 28-30.

Bartlett, P.C., John, H.K., Wilke, M.A., Kaneene, J.B., Mather E.C. (1986)

Metritis complex in Michigan Holstein-Friesian cattle: Incidence, descriptive epidemiology and estimated economic impact

Preventive Veterinary Medicine,4 (1986), 235-248

Baumann, H., Gauldie, J. (1994)

The acute phase response.

Immunology today, Vol. 15, No. 2, 74-80.

Bekana, M., Jonsson, P., Ekman, T., Kindahl, H. (1994)

Intrauterine bacteril findings in postpartum cows with retained fetal membranes.

J Vet Med, 41, 663-670.

Bekana, M., Odensvik, K., Kindahl, H. (1996)

Prostaglandin F<sub>2α</sub> metabolite and progesterone profiles in postpartum cows with retained fetal membranes.

Acta vet. scand. 37, 171-185.

- Bonnett, B.N., Martin, S.W., Gannon, V.P., Miller, R.B., Etherington, W.G. (1991)  
Endometrial biopsy in Holstein-Frisian dairy cows. III. Bacteriological analysis and correlations with histological findings.  
Can J Vet Res, 55(2), 168-173.
- Borsberry, S., Dobson, H. (1989)  
Periparturient diseases and their effect on reproductive performance in five dairy herds.  
Vet Rec, 124, 217-219.
- Bostedt, H., Schels, H., Günzler, D. (1979)  
Klinische und bakteriologische Befunde am Genitaltrakt von Rindern nach gestörten Geburten in den ersten drei Wochen des Puerperiums.  
Zbl. Vet. Med. B, 26, 397-412.
- Bretzlaff, K. (1987)  
Rationale for the treatment of endometritis in the dairy cow.  
Food animal practice, 3, 593-607.
- Chenault, J.R., McAllister, J.F., Chester, S.T., Dame, K.J. Kausche, F.M. (2001)  
Efficacy of ceftiofur hydrochloride administered parenterally for five consecutive days for treatment of acute post-partum metritis in dairy cows  
AABP- Proc. 34, 137-138
- Cheng, Z., McKeller, Q., Nolan, A. (1998)  
Pharmacokinetic studies of flunixin meglumine and phenylbutazone in plasma exsudate and transudate in sheep.  
J. vet. Pharmacol. Therap., 21, 315-21.
- Cheryk, L.A., Hooper-McGrevy, K.E., Gentry, P.A. (1998)  
Alterations in bovine platelet function and acute phase proteins induced by Pasteurella haemolytica A1.  
Can J Vet Res, 62, 1-8.

Clauss, A. (1957)

Gerinnungsphysiologische Schnellmethode zur Bestimmung des Fibrinogens.  
Acta haemat. 17, 237-246.

Cohen, R.O., Colodner, R., Ziv, G., Keness, J. (1996)

Isolation and microbial susceptibility of obligate anaerobic bacteria recovered from the uteri of dairy cows with retained fetal membranes and postparturient endometritis.  
Zentralbl Veterinärmed B, 43(4), 193-199.

Conner, J.G., Eckersall, P.D, Wiseman, A., Aitchinson, T.C., Douglas, T.A. (1988)

Bovine acute phase response following turpentine injection.  
Res Vet Sci, 44, 82-88.

Conner, J.G., Eckersall, P.D., Doherty, M., Douglas, T.A. (1986)

Acute phase response and mastitis in the cow.  
Res Vet Sci, 41, 126-128.

De Kruif, A., Gunnink, J. W., und de Bois, C. H. W. (1982)

Onderzoek en behandeling van endometritis post partum bij het rund.  
Tijdschr. Diergeneesk. 107, 117 – 125

De Kruif, A., Mansfeld, R., Hoedemaker, M. (1998)

Tierärztliche Bestandsbetreuung beim Milchrind.  
Ferdinand Enke Verlag, Stuttgart, 1998, 61-63

Deluyker, H.A., Gay, J.M., Weaver, L.D., Azari, A.S. (1991)

Change of milk yield with clinical diseases for a high producing dairy herd.  
J Dairy Sci, 74(2), 437-445.

Dohmen, M.J., Joop, K., Sturk, A., Bols, P.E., Lohuis, J.A. (2000)

Relationship between intrauterine bacterial contamination, endotoxin levels and the development of endometritis in postpartum cows with dystocia or retained placenta.  
Theriogenology, 54(7), 1019-1032.

Drillich M., Beetz, O., Pfützner, A., Sabin, M., Sabin H.-J., Kutzer, P., Nattermann, H., Heuwieser, W. (2001)  
Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows  
J. Dairy Sci. 84, 2010-2017

Drillich, M. (1999)  
Vergleich des strategischen Einsatzes von Prostaglandin F2a mit konventionellen Methoden des Fruchtbarkeitsmanagements in zwei Milchviehbetrieben  
Berlin, Freie Universität, Fachbereich Veterinärmedizin, Diss.

Eckersall, P.D., Conner, J.G. (1988)  
Bovine and canine acute-phase-proteins.  
Veterinary Research Communications, 12, 169-178.

Edwards, J.L., Tozer, P.R. (2003)  
Using activity and milk yield as predictors of fresh cow disorders.  
[http://www.aipl-arsusda.gov/publish/other/2003/adsa03\\_jle.html](http://www.aipl-arsusda.gov/publish/other/2003/adsa03_jle.html).

Ek, N. (1972)  
The quantitative determination of fibrinogen in normal bovine plasma and cows with inflammatory conditions.  
Acta Vet. Scand., 13, 175-184.

Erb, H. N., Smith, R. D., Oltenacu, P. A., Guard, C. L., Hillman, R. B., Powers, P. A., Smith, M.C., und White, M. E. (1985)  
Path model of reproductive disorders and performance, milk fever, mastitis, milk yield, and culling in Holstein cows.  
J. Dairy Sci. 68, 3337 – 3349

Espinasse, J., Thouvenot, J.P., Dalle, S., Garcia, J., Schelcher, F., Salat, O., Valarcher, J.F., Daval., S. (1994)  
Comparative study of the action of flunixin meglumine and tolfenamic acid on prostaglandin E2 synthesis in bovine inflammatory exudate.  
J Vet Pharmacol Ther, 17(4), 271-274.

Esslemont, R.J., Kossaibati, M.A. (1997)

Culling in 50 dairy herds in England.

Vet Rec, 140, 36-39.

Franklyn, B.G. (1984)

Plasma fibrinogen measurement: prognostic value in calf bronchopneumonia.

Zbl. Vet. Med. A, 31, 361-369.

Gilbert, R.O. (1992)

Bovine endometritis: the burden of proof.

Cornell Vet 82, 11-14

Godson, D.L., Baca-Estrada, M.E., van Kessel, A.G., Hughes, P.A., Morsy, M.A., vna

Donkergoed, J., Harland, R.J., Shuster, D.E., Daley, M.J., Babiuk, L.A. (1995)

Regulation of bovine acute phase responses by recombinant Interleukin-1 $\beta$ .

Can J Vet Res, 59, 249-255.

Golbs, S., Scherkl, R. (1996)

Pharmakologie der Entzündung und der Allergie.

in: Frey, H.H., Löscher, W.: Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin, Ferdinand Enke Verlag, Stuttgart, 424-445.

Gröhn, Y. T., Erb, H. N., McCulloch, C. E., und Saloniemi, H. S. (1990)

Epidemiology of reproductive disorders in dairy cattle: Association among host characteristics, disease and production.

Prev. Vet. Med. 8, 25 – 39

Gustafsson, B. K. (1984)

Therapeutic strategies involving antimicrobial treatment of the uterus in large animals.

J. Am. Vet. Med. Assoc. 185, 1194 – 1198

Gymnich, S., Deszczka, Wimmers, K., Bidlingmaier, M., Schellander, K., Petersen, B. (2001)  
Haptoglobin as an indicator for animal welfare: Effects of different hygienic conditions and  
transport stress on haptoglobin plasma concentration.

11<sup>th</sup> ICPD, 12-16 August 2001, KLV Denmark.

Heinrich, P.C., Castell, J.V., Andus, T. (1990)

Interleukin-6 and the acute phase response.

Biochem. J., 265, 621-636.

Heuer, C., Schukken, Y.H., Dobbelaar, P. (1999)

Postpartum body condition score and results from the first test day milk as predictors of  
disease, fertility, yield and culling in commercial dairy herds.

J Dairy Sci, 82, 295-304.

Heuwieser, W., Tenhagen, B.A., Tischer, M., Lühr, J., Blum, H. (2000)

Effect of three programs for the treatment of endometritis on the reproductive performance of  
a dairy herd.

Vet Rec, 146, 338-341.

Higgins, A.J., Lees, P. (1984)

The acute inflammatory process, arachidonic acid metabolism and the mode of action of  
antiinflammatory drugs.

Equine vet. J., 16(3), 163-175.

Hirvonen, J. (2000)

Acute phase response in dairy cattle.

<http://www.thesis.helsinki.fi/julkaisut/ela/kliin/muut/hirvonen/introduction/html>.

Hirvonen, J., Huszenicza, G., Kulcsár, Pyörälä, S. (1998)

Acute phase response in dairy cows with acute postpartum metritis.

Theriogenology, 51, 1071-1083.

Horadagoda, N.U., Knox, K.M.G., Gibbs, H.A., Reid, S.W.J., Horadagoda, A., Edwards, S.E.R., Eckersall, P.D. (1999)

Acute phase proteins in cattle: discrimination between acute and chronic inflammation.  
Vet Rec, 144, 437-441.

Hultén, C., Sandgren, B., Skiöldebrand, E., Klingeborn, B., Marhaug, G., Forsberg, M. (1999)  
The acute phase protein Serum Amyloid A (SAA) as an inflammatory marker in equine influenza virus infection.

Acta vet. scand., 40, 323-333.

Huszenicza, G., Fodor, M., Kulcsar, M., Dohmen, M.J.W., Vamos, M., Porkolab, L., Kegl, T., Bartyik, J., Lohuis, J.A.C.M., Janosi, S., Szita, G. (1999)

Uterine bacteriology, resumption of cyclic ovarian activity and fertility in postpartum cows kept in large – scale herds.

Reprod Dom Anim, 34, 237-245.

Jarlov, N., Andersen, P.H., Hesselholt, M. (1992)

Pathophysiology of experimental bovine endotoxemia: endotoxin induced synthesis of prostaglandins and thromboxane and the modulatory effect of some non-steroidal anti-inflammatory drugs.

Acta Vet Scand, 33(1), 1-8.

Kask, K., Kindahl, H., Gustaffson, H. (1998)

Bacteriological and histological investigation of the postpartum bovine uterus in two estonian dairy herds.

Acta vet. scand., 39, 423-432.

Kietzmann, M. (1999)

Arzneimitteltherapie bei uterinen Erkrankungen.

GroßtierVet, April 1999, 6-10.



Knutti, B., Küpfer, U., Busato, A. (2000)

Reproduktive efficiency of cows with endometritis after treatment with intrauterine infusions or prostaglandin injections, or no treatment.

J. Vet. Med. 47, 609-615

Königsson, K., Gustafsson, H., Gunnarsson, A., Kindahl, H. (2001)

Clinical and bacteriological aspects on the use of tetracycline and flunixin in primiparous cows with induced retained placenta and post partal endometritis.

Reprod Domest Anim., 36(5), 247-256.

Kristula, M., Smith, B., (2001)

Use of daily postpartum temperatures to select dairy cows for treatment with systemic antibiotics.

The AABP Proceedings-Vol. 34, 182

Landoni, M.F., Cunningham, F.M., Lees, P. (1995)

Comparative pharmacodynamics of flunixin, ketoprofen and tolfenaamic acid in calves.

Vet Rec, 137(17), 428-431.

LeBlanc, S.J., Duffield, T.F., Leslie, K.E., Bateman, K.G., Keefe, G.P., Walton, J.S., Johnson, W.H. (2002)

The effect of treatment of clinical endometritis on reproductive performance in dairy cows.

J. Dairy Sci. 85, 2237-2249.

Lee, L. A., Ferguson, J. D., und Galligan, D. T. (1989)

Effect of disease on days open assessed by survival analysis.

J. Dairy Sci. 72, 1020 – 1026

Lees, P. (1991)

General aspects of inflammation.

Flem. Vet. J. 62, Suppl. 1, 43-54.

Leslie, K.E. (1983):

The events of normal and abnormal postpartum reproductive endocrinology and uterine involution in dairy cows: a review.

Can Vet J 24, 67-71

Lewis, G.S. (1997)

Uterine health and disorders.

J Dairy Sci, 80, 984-994.

Lowder, M.Q. (1993)

Diagnosing and treating bovine postpartum endometritis.

Vet. Med., 474-480.

Mackiewicz, A. (2001)

Application of acute-phase-proteins in human diagnosis.

2<sup>nd</sup> European Colloquium on Acute Phase Proteins, Bonn.

Mansfeld, R., De Kruif, A., Hoedemaker, M., und Heuwieser, W. (1999)

Fruchtbarkeitsüberwachung auf Herdenbasis.

in: Grunert, E., und de Kruif, A. (Hrsg.): Fertilitätsstörungen beim weiblichen Rind.

3. Auflage, Parey Buchverlag, Berlin, 337 – 350

Markusfeld, D. (1984)

Factors responsible for post parturient metritis in dairy cattle.

Vet. Rec. 114, 539-542.

Markusfeld, O., Galon, N., Ezra, E. (1997)

Body condition score, health, yield and fertility in dairy cows.

Vet Rec, 141(3), 67-72.

Martinez, J., und Thibier, M. (1984)

Reproductive disorders in dairy cattle. Respective influence of herds, seasons, milk yield and parity.

Theriogenology 21, 569-581.

- Mateus, L., da Costa, L.L., Bernardo, F., Silva, J.R. (2002)  
Influence of puerperal uterine infection on uterine involution and postpartum ovarian activity in dairy cows.  
Reprod Domest Anim, 37(1), 31-35.
- McNair, J., Kennedy, G., Bryson, D.G., Reilly, G.A.C., McDowell, S.W.J., Mackie, D.P. (1997)  
Evaluation of a competitive immunoassay for the detection of bovine haptoglobin.  
Res Vet Sci, 63, 145-149.
- McSherry, B.J., Horney, F.D., deGroot, J.J. (1970)  
Plasma fibrinogen levels in normal and sick cows.  
Can. J. comp. Med. 34, 191-197.
- Messow, C. (1959)  
Fibrinogen und Fibrinolyse.  
Dtsch. tierärztl. Wschr., 1.August 1959, Nr. 15, 401-406.
- Montes, A. J., und Pugh, D. G. (1993)  
Clinical approach to postpartum metritis.  
Compend. Contin. Educ. Pract. Vet. 15, 1131 – 1137
- Morimatsu, M., Syuto, B., Shimada, N., Fujinaga, T., Yamamoto, S., Saito, M., Naiki, M. (1991)  
Isolation and characterisation of bovine haptoglobin from acute phase sera.  
Journal of Biological Chemistry, 266, No. 18, 11833-11837.
- Noakes, D.E., Till, D., und Smith, D.R. (1989)  
Bovine uterine flora post partum: A comparison of swabbing and biopsy.  
Vet. Rec. 124, 563-564.
- Noakes, D.E., Wallace, L., und Smith, G.R. (1991)  
Bacterial flora of the uterus of cows after calving on two hygienically contrasting farms.  
Vet. Rec. 128, 440-442.

Odensvik, K., Fredriksson, G. (1993)

The effect of intensive flunixin treatment during the postpartum period in the bovine.  
J. Vet. Med. A 40, 561-568.

Odensvik, K., Magnusson, U. (1996)

Effect of oral administration of flunixin meglumine on the inflammatory response to endotoxin in heifers.

Am J Vet Res, 57, 201-204.

Okker, H., Schmitt, E.J., Vos, P.L.A.M., Scherpenisse, P., Bergwerff, A.A., Jonker, F.H. (2002)

Pharmacokinetics of ceftiofur in plasma and uterine secretions and tissues after subcutaneous postpartum administration in lactating dairy cows.

J. vet. Pharmacol. Therap., 25, 33-38.

Olson, J. D. (1996)

Metritis / Endometritis: Medically sound treatment.

The Bovine Proceedings 29, 8 – 14

Olson, J.D., Ball, L., Mortimer, R., Farin, P.W., Adney, W.S., und Huffmann, E.M. (1984)

Aspects of bacteriology and endocrinology of cows with pyometra and retained fetal membranes.

Am. J. Vet. Res. 45, 2251-2255.

Olson, J.D., Bretzlaff, K., Mortimer, R.G., und Ball, L. (1986)

The metritis-pyometra complex.

In: J.D. Olson (Hrsg.): Current therapy in Theriogenology 2. Auflage, W. B. Saunders C., Philadelphia, S. 227-236.

Paisley, L. G., Mickelsen, W. D., und Anderson, P. B. (1986)

Mechanism and therapy for retained fetal membranes and uterine infections of cows: A review.

Theriogenology 25, 353 – 381

Panndorf, H., Richter, H., Dittrich, B. (1976)

Haptoglobin bei Haussäugetieren.

Arch. exper. Vet.med. Bd. 30, H.2, 193-202.

Pennington, J.A., Albright, J.L., Callahan, C.J. (1986)

Relationship of sexual activities in estrous cows different frequencies of observation and pedometer measurements.

J. Dairy Sci. 69, 2925-2934.

Pineiro, C., Lorenzo, E., Pineiro, M., Lampraeve, F., Alava, M.A. (2001)

Effect of transport and farm sanitary conitions in the concentration of acute phase plasma proteins Pig-Map and haptoglobin.

2<sup>nd</sup> European Colloquium on Acute Phase Proteins, Bonn.

Playfair, J.H.L., Baron, D. (1995).

Akute Entzündungen.

in: Playfair, J.H.L., Baron, D. (Hrsg.): Immunologie. 5. Auflage, Blackwell

Wissenschaftsverlag, Berlin, 18-19.

Playfair, J.H.L., Baron, D. (1995).

Das Zytokin – Netzwerk.

in: Playfair, J.H.L., Baron, D. (Hrsg.): Immunologie. 5. Auflage, Blackwell

Wissenschaftsverlag, Berlin, 50-51.

Radostits, O. M., Leslie, K. E., und Fetrow, J. (1994)

Maintaining reproductive efficiency in dairy cattle.

in: Radostits, O. M., Leslie, K. E., und Fetrow, J. (Hrsg.): Herd health: Food animal production medicine.

2. Auflage, W. B. Saunders Company, Philadelphia, 141 – 158

Rajala, P.J., Gröhn, Y.T. (1998)

Effects of dystocia, retained placenta and metritis on milk yield in dairy cows.

J Dairy Sci, 81, 3172-3181.

Salonen, M., Hirvonen, J., Pyörälä, S., Sankari, S., Sandholm, M. (1996)

Quantitative determination of bovine serum haptoglobin in experimentally induced Escherichia coli mastitis.

Res Vet Sci, 60 (1), 88-91.

Sandals, W.C.D., Curtis, R.A., Cote, J.F., und Martin, S.W. (1979)

The effect of retained placenta and metritis complex on reproductive performance in dairy cattle. A case control study.

Can. vet. J. 20, 131-135.

Schmitt, E., Boucher, J.F., Van den Eede, Ch., Chenault, J.R. (2001)

Comparison of clinical efficacy of ceftiofur to oxytetracycline for treatment of acute puerpera metritis in cows

The bovine practitioner Sept 2001, 199

Selinger, P., Mazurova, J. (1976)

Results of bacteriological and cytological studies of uterine lavages in cows from herds with fertility disorders.

Vet Med (Praha), 21(5), 309-318.

Sheldon, I.M., Noakes, D.E. Rycroft, A., Dobson, H. (2001)

Acute phase protein responses to uterine bacterial contamination in cattle after calving.

Vet Rec, 148, 172-175.

Simerl, N.A., Wilcox, C.J., Thatcher, W.W., Martin, F.G. (1991)

Prepartum and peripartum reproductive performance of dairy heifers freshening at young ages.

J Dairy Sci, 74, 1724-1729.

Skinner, J.G., Brown, R.A., Roberts, L. (1991)

Bovine haptoglobin response in clinically defined field conditions.

Vet. Rec., February 16, 147-149.

Skinner, J.G., Roberts, L. (1994)

Haptoglobin as an indicator of infection in sheep.

Vet Rec, 134, 33-36.

Smith, B.I., Donovan, G.A., Risco, C., Littell, R., Young, C., Stanker, L.H., Elliot, J. (1998)

Comparison of various antibiotic treatments for cows diagnosed with toxic puerperal metritis

J. Dairy Sci.81, 1555-1562

Smith, B.I., Risco, C.A. (2002)

Clinical manifestation of postpartum metritis in dairy cattle.

Compendium, Vol. 24, No. 8, 56-63.

Spooner, R.L., Miller, J.K. (1971)

The measurement of haemoglobin reactive protein in ruminants as an aid to the diagnosis of acute inflammation.

Vet. Rec., 88, 2-4.

Steffan, J., Adriamanga, S. (1983)

Treatment of metritis with antibiotics or prostaglandin  $F_{2\alpha}$  and influence of ovarian cyclicity in dairy cows.

Am J Vet Res, Vol. 45, No.6, 1090-1094.

Stevenson, J.S., Call, E.P. (1988)

Reproductive disorders in the periparturient dairy cow.

J Dairy Sci, 71, 2572-2583.

Tenhagen, B.A., Edinger, D., Heuwieser, W. (1999)

Einfluss von Schweregeburten bei Erstkalbinnen auf den Verlauf der ersten Laktation.

Tierärztl. Umschau, 54, 617-623.

Tenhagen, B.-A., Tischer, M., Heuwieser, W., und Blum, H. (1998)

Use of sensitivity analysis to economically evaluate reproductive management programs in dairy herds.

Pages 663-668 in Proc. 20th World Buiatrics Congr., Aydney, Australia.

Tischer, M. (1998)

Vergleich von intrauterinen Arzneimittelapplikationen mit einem strategischen Prostaglandinprogramm zur Behandlung von chronischen Endometritiden in einer Milchviehherde  
Berlin, Freie Universität, Fachbereich Veterinärmedizin, Diss.

Vandeplassche, M. (1981)

Neue vergleichende Aspekte der Involution und der puerperalen Metritis bei Stute, Kuh und Sau.  
Mh. Vet. Med., 36, 804-807.

Voet, D., Voet, J.G. (1994)

Arachidonsäurestoffwechsel: Prostaglandine, Prostacycline, Thromboxane und Leukotriene.  
in: Voet, D., Voet, J.G.: Biochemie, Verlagsgesellschaft mbH, Weinheim, 659-668.

Waelchli, R.O., Thun, R., Stocker, H. (1999)

Effect of flunixin meglumine on placental expulsion in dairy cattle after a caesarean.  
Vet Rec, 144, 702-703.

Whitacre, M.D. (1992)

Intrauterine infusion in the post partum dairy cow.  
Vet. Med. 4, 376-381.

Whitaker, D.A., Kelly, J.M., Smith, S. (2000)

Disposal and disease rates in 340 British dairy herds.  
Vet Rec, 146, 363-367.

Young, C.R., Eckersall, P.D., Saini, P.K., Stanker, L.H. (1995)

Validation of immunoassays for bovine haptoglobin.  
Veterinary Immunology and Immunopathology, 49, 1-13.

Young, C.R., Wittum, T.E., Stanker, L.H., Perino, L.J., Griffin, D.D., Littledike, E.T. (1996)

Serum haptoglobin concentrations in a population of feedlot cattle.  
AJVR, Vol. 57, No.2, 138-141.



Zhou, Ch., Boucher, J.F., Dame, K.J. (2001)

A multi-location clinical study of Ceftiofur for the treatment of postpartum cows with elevated temperatur and other complications

AABP-Proc. 2001