

## 7 Literatur

1. Abolmaali, N.D., Hietschold, V., Appold, S., Ebert, W., Vogl, T.J.: Gadomer-17-enhanced 3D navigator-echo MR angiography of the pulmonary arteries in pigs. *Eur Radiol* (2002) 12: 692-697
2. Akritas, M.G., Arnold, S.F., Brunner, E.: Nonparametric hypothesis and rank statistics for unbalanced factorial designs. *J Am Stat Assoc* (1997) 92: 258-265
3. Allkemper, T., Bremer, C., Matuszewski, L., Ebert, W., Reimer, P.: Contrast-enhanced blood-pool MR angiography with optimized iron oxides: effect of size and dose on vascular contrast enhancement in rabbits. *Radiology* (2002) 223: 432-438
4. Amano, Y., Herfkens, R.J., Shifrin, R.Y., Alley, M.T., Pelc, N.J.: Three-dimensional cardiac cine magnetic resonance imaging with an ultrasmall superparamagnetic iron oxide blood pool agent (NC100150). *J Magn Reson Imaging* (2000) 11: 81-86
5. Ames, B.N.: Identifying environmental chemicals causing mutations and cancer. *Science* (1979) 204: 587-593.
6. Anzai, Y., Piccoli, C.W., Outwater, E.K., Stanford, W., Bluemke, D.A., Nurenberg, P., Saini, S., Maravilla, K.R., Feldman, D.E., Schmiedl, U.P., Brunberg, J.A., Francis, I.R., Harms, S.E., Som, P.M., Tempany, C.M.: Evaluation of neck and body metastases to nodes with ferumoxtran 10-enhanced MR imaging: phase III safety and efficacy study. *Radiology* (2003) 228: 777-788
7. Anzai, Y., Prince, M.R., Chenevert, T.L., Maki, J.H., Londy, F., London, M., McLachlan, S.J.: MR angiography with an ultrasmall superparamagnetic iron oxide blood pool agent. *J Magn Reson Imaging* (1997) 7: 209-214
8. Ardenne von, M., Kirsch, R.: Zur Methodik der Extrem-Hyperthermie, insbesondere bei der Krebs-Mehrschritt-Chemotherapie. *Dtsch Gesundheitsw* (1965) 20: 1935-1940
9. Ashley, D.L., Goldstein, J.H.: The application of dextran-magnetite as a relaxation agent in the measurement of erythrocyte water exchange using pulsed nuclear magnetic resonance spectroscopy. *Biochem Biophys Res Commun* (1980) 97: 114-120
10. Ba-Ssalamah, A., Heinz-Peer, G., Schima, W., Schibany, N., Schick, S., Prokesch, R.W., Kaider, A., Teleky, B., Wrba, F., Lechner, G.: Detection of focal hepatic lesions: comparison of unenhanced and SHU 555 A-enhanced MR imaging versus biphasic helical CTAP. *J Magn Reson Imaging* (2000) 11: 665-672
11. Bartolotta, T.V., Midiri, M., Galia, M., Carcione, A., De Maria, M., Lagalla, R.: Benign hepatic tumors: MRI features before and after administration of superparamagnetic contrast media. *Radiol Med* (2001) 101: 219-229
12. Bean, C.P., Jacobs, I.S.: Magnetic granulometry and superparamagnetism. *J Appl Phys* (1956) 27: 1448-1452
13. Beckmann, N., Joergensen, J., Bruttel, K., Rudin, M., Schuurman, H.J.: Magnetic resonance imaging for the evaluation of rejection of a kidney allograft in the rat. *Transpl Int* (1996) 9: 175-183
14. Bedaux, W.L., Hofman, M.B., Wielopolski, P.A., de Cock, C.C., Hoffmann, V., Oudkerk, M., de Feyter, P.J., van Rossum, A.C.: Three-dimensional magnetic resonance coronary angiography using a new blood pool contrast agent: initial experience. *J Cardiovasc Magn Reson* (2002) 4: 273-282
15. Bee, A., Massart, R., Neveu, S.: Synthesis of very fine maghemite particles. *J Magn Mater* (1995) 149: 6-9

16. Beets-Tan, R.G., Van Engelshoven, J.M., Greve, J.W.:  
Hepatic adenoma and focal nodular hyperplasia: MR findings with superparamagnetic iron oxide-enhanced MRI. *Clin Imaging* (1998) 22: 211-215
17. Bellin, M.F., Roy, C., Kinkel, K., Thoumas, D., Zaim, S., Vanel, D., Tuchmann, C., Richard, F., Jacqmin, D., Delcourt, A., Challier, E., Leuret, T., Cluzel, P.:  
Lymph node metastases: safety and effectiveness of MR imaging with ultrasmall superparamagnetic iron oxide particles--initial clinical experience.  
*Radiology* (1998) 207: 799-808
18. Bilder, G.E.:  
Studies on immune competence in the rat: changes with age, sex, and strain.  
*J Gerontol* (1975) 30: 641-646
19. Bjerner, T., Ericsson, A., Wikstrom, G., Johansson, L., Nilsson, S., Ahlstrom, H., Hemmingsson, A.:  
Evaluation of nonperfused myocardial ischemia with MRI and an intravascular USPIO contrast agent in an ex vivo pig model. *J Magn Reson Imaging* (2000) 12: 866-872
20. Bjerner, T., Johansson, L., Ericsson, A., Wikstrom, G., Hemmingsson, A., Ahlstrom, H.:  
First-pass myocardial perfusion MR imaging with outer-volume suppression and the intravascular contrast agent NC100150 injection: preliminary results in eight patients.  
*Radiology* (2001) 221: 822-826
21. Bluemke, D.A., Stillman, A.E., Bis, K.G., Grist, T.M., Baum, R.A., D'Agostino, R., Malden, E.S., Pierro, J.A., Yucel, E.K.:  
Carotid MR angiography: phase II study of safety and efficacy for MS-325.  
*Radiology* (2001) 219: 114-122
22. Bluemke, D.A., Weber, T.M., Rubin, D., de Lange, E.E., Semelka, R., Redvanly, R.D., Chezmar, J., Outwater, E., Carlos, R., Saini, S., Holland, G.A., Mammone, J.F., Brown, J.J., Milestone, B., Javitt, M.C., Jacobs, P.:  
Hepatic MR imaging with ferumoxides: multicenter study of safety and effectiveness of direct injection protocol. *Radiology* (2003) 228: 457-464
23. Bock, J.C., Pison, U., Kaufmann, F., Felix, R.:  
Gd-DTPA-polylysine-enhanced pulmonary time-of-flight MR angiography.  
*J Magn Reson Imaging* (1994) 4: 473-476
24. Bogdanov, A.A., Jr., Weissleder, R., Frank, H.W., Bogdanova, A.V., Nossif, N., Schaffer, B.K., Tsai, E., Papisov, M.I., Brady, T.J.:  
A new macromolecule as a contrast agent for MR angiography: preparation, properties, and animal studies. *Radiology* (1993) 187: 701-706
25. Borisch, I., Horn, M., Butz, B., Zorger, N., Draganski, B., Hoelscher, T., Bogdahn, U., Link, J.:  
Preoperative evaluation of carotid artery stenosis: comparison of contrast-enhanced MR angiography and duplex sonography with digital subtraction angiography.  
*Am J Neuroradiol* (2003) 24: 1117-1122
26. Both, M., Reinhold-Keller, E., Muller-Hulsbeck, S., Biederer, J., Gross, W.L., Heller, M., Reuter, M.:  
Entzündliches Aortenbogensyndrom: Stenosedagnostik mittels kontrastmittelverstärkter 3D-MR-Angiographie im Vergleich mit der DSA. *Fortschr Röntgenstr* (2004) 176: 48-55
27. Botnar, R.M., Stuber, M., Danias, P.G., Kissinger, K.V., Manning, W.J.:  
Improved coronary artery definition with T2-weighted, free-breathing, three-dimensional coronary MRA. *Circulation* (1999) 99: 3139-3148
28. Brain, J.D., Molina, R.M., DeCamp, M.M., Warner, A.E.:  
Pulmonary intravascular macrophages: their contribution to the mononuclear phagocyte system in 13 species. *Am J Physiol* (1999) 276: L146-154
29. Bremer, C., Allkemper, T., Baermig, J., Reimer, P.:  
RES-specific imaging of the liver and spleen with iron oxide particles designed for blood pool MR-angiography. *J Magn Reson Imaging* (1999) 10: 461-467

30. Briggs, R.W., Wu, Z., Mladinich, C.R.J., Stoupis, C., Gauger, J., Liebig, T., Ros, P.R., Ballinger, J.R., Kubilis, P.:  
In vivo animal tests of an artifact-free contrast agent for gastrointestinal MRI.  
*Magn Reson Imaging* (1997) 15: 559-566
31. Brouwer, A., Knook, D.L.:  
The reticuloendothelial system and aging: a review. *Mech Ageing Dev* (1983) 21: 205-228
32. Brunner, E., Langer, F.:  
Nonparametric analysis of longitudinal data.  
Munich - Vienna: Oldenbourg Publishers, (1999).
33. Bulte, J.M., Vymazal, J., Brooks, R.A., Pierpaoli, C., Frank, J.A.:  
Frequency dependence of MR relaxation times. II. Iron oxides.  
*J Magn Reson Imaging* (1993) 3: 641-648
34. Bulte, J.W.M., Douglas, T., Witwer, B., Zhang, S.-C., Strable, E., Lewis, B.K., Zywicke, H., Miller, B., van Gelderen, P., Moskowitz, B.M., Duncan, L.D., Frank, J.A.:  
Magnetodendrimers allow endosomal magnetic labeling and in vivo tracking of stem cells.  
*Nat Biotechnol* (2001) 19: 1141-1147
35. Bulte, J.W.M., Duncan, I.D., Frank, J.A.:  
In vivo magnetic resonance tracking of magnetically labeled cells after transplantation.  
*J Cereb Blood Flow Metab* (2002) 22: 899-907
36. Bunce, N.H., Keegan, J., Gatehouse, P.D., Moon, J.C., Firmin, D.N., Hoffmann, V., Pennell, D.J.:  
Initial experience with the intravascular contrast agent NC100150-injection (Clariscan) for breath-hold and navigator-gated magnetic resonance coronary artery imaging.  
*J Magn Reson Imaging* (2002) 16: 217-223
37. Butler, R.F.:  
Paleomagnetism: Magnetic Domains to Geologic Terranes. Blackwell Scientific Publications, Electronic Edition [www.earth.rochester.edu/butlerbook/tocpref.pdf](http://www.earth.rochester.edu/butlerbook/tocpref.pdf) (2004)
38. Butz, B., Dorenbeck, U., Borisch, I., Zorger, N., Lenhart, M., Feuerbach, S., Link, J.:  
High-resolution contrast-enhanced magnetic resonance angiography of the carotid arteries using fluoroscopic monitoring of contrast arrival: diagnostic accuracy and interobserver variability. *Acta Radiol* (2004) 45: 164-170
39. Cantrell, W., Elko, E.E.:  
Effect of age on phagocytosis of carbon in the rat. *Exp Parasitol* (1973) 34: 337-343
40. Cavagna, F.M., Lorusso, V., Anelli, P.L., Maggioni, F., de Haen, C.:  
Preclinical profile and clinical potential of gadocoletic acid trisodium salt (B22956/1), a new intravascular contrast medium for MRI. *Acad Radiol* (2002) 9 Suppl 2: S491-494
41. Cerdan, S., Lotscher, H.R., Kunnecke, B., Seelig, J.:  
Monoclonal antibody-coated magnetite particles as contrast agents in magnetic resonance imaging of tumors. *Magn Reson Med* (1989) 12: 151-163
42. Chan, T.W., Eley, C., Liberti, P., So, A., Kressel, H.Y.:  
Magnetic resonance imaging of abscesses using lipid-coated iron oxide particles.  
*Invest Radiol* (1992) 27: 443-449
43. Clarke, S.E., Weinmann, H.J., Dai, E., Lucas, A.R., Rutt, B.K.:  
Comparison of two blood pool contrast agents for 0.5-T MR angiography: experimental study in rabbits. *Radiology* (2000) 214: 787-794
44. Clement, O., Rety, F., Cuenod, C.A., Siauve, N., Carnot, F., Bordat, C., Siche, M., Frija, G.:  
MR lymphography: evidence of extravasation of superparamagnetic nanoparticles into the lymph. *Acad Radiol* (1998) 5 Suppl 1: S170-172
45. Clement, O., Schouman-Claeys, E., Frija, G.:  
Ferrite particles (AMI25): influence of cirrhosis and chemotherapy on magnetic resonance imaging of metastasis. *Invest Radiol* (1990) 25: S61-S62
46. Constable, R.T., Kay, I., Smith, M.R., Henkelman, R.M.:  
High quality zoomed MR images. *J Comput Assist Tomogr* (1989) 13: 179-181

47. Corot, C., Violas, X., Robert, P., Gagneur, G., Port, M.: Comparison of different types of blood pool agents (P792, MS325, USPIO) in a rabbit MR angiography-like protocol. *Invest Radiol* (2003) 38: 311-319
48. Cosottini, M., Pingitore, A., Puglioli, M., Michelassi, M.C., Lupi, G., Abbruzzese, A., Calabrese, R., Lombardi, M., Parenti, G., Bartolozzi, C.: Contrast-enhanced three-dimensional magnetic resonance angiography of atherosclerotic internal carotid stenosis as the noninvasive imaging modality in revascularization decision making. *Stroke* (2003) 34: 660-664
49. Cronberg, C.N., Sjoberg, S., Albrechtsson, U., Leander, P., Lindh, M., Norgren, L., Danielsson, P., Sonesson, B., Larsson, E.M.: Peripheral arterial disease. Contrast-enhanced 3D MR angiography of the lower leg and foot compared with conventional angiography. *Acta Radiol* (2003) 44: 59-66
50. D'Arienzo, A., Scaglione, G., Vicinanza, G., Manguso, F., Bennato, R., Belfiore, G., Imbriaco, M., Mazzacca, G.: Magnetic resonance imaging with ferumoxil, a negative superparamagnetic oral contrast agent, in the evaluation of ulcerative colitis. *Am J Gastroenterol* (2000) 95: 720-724
51. Daldrup, H.E., Link, T.M., Blasius, S., Strozyk, A., Konemann, S., Jurgens, H., Rummeny, E.J.: Monitoring radiation-induced changes in bone marrow histopathology with ultra-small superparamagnetic iron oxide (USPIO)-enhanced MRI. *J Magn Reson Imaging* (1999) 9: 643-652
52. Daldrup-Link, H.E., Brasch, R.C.: Macromolecular contrast agents for MR mammography: current status. *Eur Radiol* (2003) 13: 354-365
53. Daldrup-Link, H.E., Reinlander, C., Link, T.M., Richter, K.J., Konemann, S., Rummeny, E.J.: Experimentelle Untersuchungen zur Wertigkeit von SPIO für die MRT des Knochenmarkes vor und nach Ganzkörperbestrahlung. *Fortschr Röntgenstr* (2001) 173: 547-553
54. Daldrup-Link, H.E., Rummeny, E.J., Ihssen, B., Kienast, J., Link, T.M.: Iron-oxide-enhanced MR imaging of bone marrow in patients with non-Hodgkin's lymphoma: differentiation between tumor infiltration and hypercellular bone marrow. *Eur Radiol* (2002) 12: 1557-1566
55. del Frate, C., Bazzocchi, M., Morteale, K.J., Zuiani, C., Londero, V., Como, G., Zanardi, R., Ros, P.R.: Detection of liver metastases: comparison of gadobenate dimeglumine-enhanced and ferumoxides-enhanced MR imaging examinations. *Radiology* (2002) 225: 766-772
56. Deserno, W.M., Harisinghani, M.G., Taupitz, M., Jager, G.J., Witjes, J.A., Mulders, P.F., Hulsbergen Van De Kaa, C.A., Kaufmann, D., Barentsz, J.O.: Urinary Bladder Cancer: Preoperative Nodal Staging with Ferumoxtran-10-enhanced MR Imaging. *Radiology* (2004)
57. Dirksen, M.S., Lamb, H.J., van der Geest, R., de Roos, A.: Toward comparability of coronary magnetic resonance angiography: proposal for a standardized quantitative assessment. *Eur Radiol* (2003) 13: 2353-2357
58. Dong, Q., Hurst, D.R., Weinmann, H.J., Chenevert, T.L., Londy, F.J., Prince, M.R.: Magnetic resonance angiography with gadomer-17. An animal study original investigation. *Invest Radiol* (1998) 33: 699-708
59. Dromain, C., de Baere, T., Baudin, E., Galline, J., Ducreux, M., Boige, V., Duvillard, P., Laplanche, A., Caillet, H., Lasser, P., Schlumberger, M., Sigal, R.: MR imaging of hepatic metastases caused by neuroendocrine tumors: comparing four techniques. *Am J Roentgenol* (2003) 180: 121-128
60. Edelman, R.R., Manning, W.J., Burstein, D., Paulin, S.: Coronary arteries: breath-hold MR angiography. *Radiology* (1991) 181: 641-643

61. Elste, V., Wagner, S., Taupitz, M., Pfefferer, D., Kresse, M., Hamm, B., Berg, R., Wolf, K.J., Semmler, W.:  
Magnetic resonance lymphography in rats: effects of muscular activity and hyperthermia on the lymph node uptake of intravenously injected superparamagnetic iron oxide particles. *Acad Radiol* (1996) 3: 660-666
62. Fain, S.B., Riederer, S.J.:  
Dependence of venous enhancement on the field of view in 3D contrast-enhanced MRA using the elliptical centric view order. *Magn Reson Med* (2001) 45: 1134-1141
63. Fink, C., Hallscheidt, P.J., Hosch, W.P., Ott, R.C., Wiesel, M., Kauffmann, G.W., Dux, M.:  
Preoperative evaluation of living renal donors: value of contrast-enhanced 3D magnetic resonance angiography and comparison of three rendering algorithms. *Eur Radiol* (2003) 13: 794-801
64. Fleige, G., Nolte, C., Synowitz, M., Seeberger, F., Kettenmann, H., Zimmer, C.:  
Magnetic labeling of activated microglia in experimental gliomas. *Neoplasia* (2001) 3: 489-499
65. Fleige, G., Seeberger, F., Laux, D., Kresse, M., Taupitz, M., Pilgrim, H., Zimmer, C.:  
In Vitro Characterization of Two Different Ultrasmall Iron Oxide Particles for Magnetic Resonance Cell Tracking. *Invest Radiol* (2002) 37: 482-488
66. Forth, W., Henschler, D., Rummel, W.:  
*Pharmakologie und Toxikologie: Bibliographisches Institut, Mannheim, Wien, Zürich, (1983).*
67. Frahm, J., Haase, A., Matthaei, D.:  
Rapid three-dimensional MR imaging using the FLASH technique. *J Comput Assist Tomogr* (1986) 10: 363-368
68. Franklin, R.J., Blaschuk, K.L., Bearchell, M.C., Prestoz, L.L., Setzu, A., Brindle, K.M., French-Constant, C.:  
Magnetic resonance imaging of transplanted oligodendrocyte precursors in the rat brain. *Neuroreport* (1999) 10: 3961-3965
69. Goldfarb, J.W., Edelman, R.R.:  
Coronary arteries: breath-hold, gadolinium-enhanced, three-dimensional MR angiography. *Radiology* (1998) 206: 830-834
70. Grandin, C., Van Beers, B.E., Robert, A., Gigot, J.F., Geubel, A., Pringot, J.:  
Benign hepatocellular tumors: MRI after superparamagnetic iron oxide administration. *J Comput Assist Tomogr* (1995) 19: 412-418
71. Groman, E.V., Paul, K.G., Frigo, T.B., Bengel, H., Lewis, J.M.:  
Polyol and polyether iron oxide complexes as pharmacological and/or MRI contrast agents. Patent: US 2003-410527, US 2003232084 (2003)
72. Halliwell, B., Gutteridge, J.M.:  
Role of free radicals and catalytic metal ions in human disease: an overview. *Methods Enzymol* (1990) 186: 1-85
73. Hamm, B., Mahfouz, A.E., Taupitz, M., Mitchell, D.G., Nelson, R., Halpern, E., Speidel, A., Wolf, K.J., Saini, S.:  
Liver metastases: improved detection with dynamic gadolinium-enhanced MR imaging? *Radiology* (1997) 202: 677-682
74. Hamm, B., Reichel, M., Vogl, T., Taupitz, M., Wolf, K.J.:  
Superparamagnetische Eisenpartikel. Klinische Ergebnisse in der MR-Diagnostik von Lebermetastasen. *Fortschr Röntgenstr* (1994) 160: 52-58
75. Hamm, B., Staks, T., Taupitz, M., Maibauer, R., Speidel, A., Huppertz, A., Frenzel, T., Lawaczek, R., Wolf, K.J., Lange, L.:  
Contrast-enhanced MR imaging of liver and spleen: first experience in humans with a new superparamagnetic iron oxide. *J Magn Reson Imaging* (1994) 4: 659-668
76. Hamm, B., Taupitz, M., Hussmann, P., Wagner, S., Wolf, K.J.:  
MR lymphography with iron oxide particles: dose-response studies and pulse sequence optimization in rabbits. *Am J Roentgenol* (1992) 158: 183-190

77. Hamm, B., Thoeni, R.F., Gould, R.G., Bernardino, M.E., Luning, M., Saini, S., Mahfouz, A.E., Taupitz, M., Wolf, K.J.:  
Focal liver lesions: characterization with nonenhanced and dynamic contrast material-enhanced MR imaging. *Radiology* (1994) 190: 417-423
78. Harisinghani, M.G., Barentsz, J., Hahn, P.F., Deserno, W.M., Tabatabaei, S., van de Kaa, C.H., de la Rosette, J., Weissleder, R.:  
Noninvasive detection of clinically occult lymph-node metastases in prostate cancer. *N Engl J Med* (2003) 348: 2491-2499
79. Harisinghani, M.G., Saini, S., Slater, G.J., Schnall, M.D., Rifkin, M.D.:  
MR imaging of pelvic lymph nodes in primary pelvic carcinoma with ultrasmall superparamagnetic iron oxide (Combidex): preliminary observations. *J Magn Reson Imaging* (1997) 7: 161-163
80. Harisinghani, M.G., Saini, S., Weissleder, R., Hahn, P.F., Yantiss, R.K., Tempany, C., Wood, B.J., Mueller, P.R.:  
MR lymphangiography using ultrasmall superparamagnetic iron oxide in patients with primary abdominal and pelvic malignancies: radiographic-pathologic correlation. *Am J Roentgenol* (1999) 172: 1347-1351
81. Hasegawa, M., Hokukoku, S.:  
Magnetic iron oxide-dextran complex and process for its production.  
Patent: US 4.101.435 (1978)
82. Helbich, T.H., Gossman, A., Mareski, P.A., Raduchel, B., Roberts, T.P., Shames, D.M., Muhler, M., Turetschek, K., Brasch, R.C.:  
A new polysaccharide macromolecular contrast agent for MR imaging: biodistribution and imaging characteristics. *J Magn Reson Imaging* (2000) 11: 694-701
83. Hemmingsson, A., Carlsten, J., Ericsson, A., Klaveness, J., Sperber, G.O., Thuomas, K.A.:  
Relaxation enhancement of the dog liver and spleen by biodegradable superparamagnetic particles in proton magnetic resonance imaging. *Acta Radiol* (1987) 28: 703-705
84. Herborn, C.U., Barkhausen, J., Paetsch, I., Hunold, P., Mahler, M., Shamsi, K., Nagel, E.:  
Coronary arteries: contrast-enhanced MR imaging with SH L 643A--experience in 12 volunteers. *Radiology* (2003) 229: 217-223
85. Hergt, R., Andra, W., d'Ambly, C.G., Hilger, I., Kaiser, W.A., Richter, U., Schmidt, H.G.:  
Physical limits of hyperthermia using magnetite fine particles. *IEEE Trans Magn* (2002) 34: 3745-3754
86. Hoffman, H.T., Quets, J., Toshiaki, T., Funk, G.F., McCulloch, T.M., Graham, S.M., Robinson, R.A., Schuster, M.E., Yuh, W.T.:  
Functional magnetic resonance imaging using iron oxide particles in characterizing head and neck adenopathy. *Laryngoscope* (2000) 110: 1425-1430
87. Hofman, M.B., Henson, R.E., Kovacs, S.J., Fischer, S.E., Lauffer, R.B., Adzamlı, K., De Becker, J., Wickline, S.A., Lorenz, C.H.:  
Blood pool agent strongly improves 3D magnetic resonance coronary angiography using an inversion pre-pulse. *Magn Reson Med* (1999) 41: 360-367
88. Huber, M.E., Paetsch, I., Schnackenburg, B., Bornstedt, A., Nagel, E., Fleck, E., Boesiger, P., Maggioni, F., Cavagna, F.M., Stuber, M.:  
Performance of a new gadolinium-based intravascular contrast agent in free-breathing inversion-recovery 3D coronary MRA. *Magn Reson Med* (2003) 49: 115-121
89. Hudgins, P.A., Anzai, Y., Morris, M.R., Lucas, M.A.:  
Ferumoxtran-10, a superparamagnetic iron oxide as a magnetic resonance enhancement agent for imaging lymph nodes: a phase 2 dose study. *Am J Neuroradiol* (2002) 23: 649-656
90. ICH: ICH harmonised tripartite guideline for good clinical practice.  
Brookwood Medical Publications [www.ich.org](http://www.ich.org)

91. Ichikawa, T., Arbab, A.S., Araki, T., Touyama, K., Haradome, H., Hachiya, J., Yamaguchi, M., Kumagai, H., Aoki, S.:  
Perfusion MR imaging with a superparamagnetic iron oxide using T2-weighted and susceptibility-sensitive echoplanar sequences: evaluation of tumor vascularity in hepatocellular carcinoma. *Am J Roentgenol* (1999) 173: 207-213
92. Idee, J.M., Berthommier, C., Goulas, V., Corot, C., Santus, R., Hermine, C., Schaefer, M., Bonnemain, B.:  
Haemodynamic effects of macrocyclic and linear gadolinium chelates in rats: role of calcium and transmetallation. *Biometals* (1998) 11: 113-123
93. Johansson, L.O., Bjornerud, A., Ahlstrom, H.K., Ladd, D.L., Fujii, D.K.:  
A targeted contrast agent for magnetic resonance imaging of thrombus: implications of spatial resolution. *J Magn Reson Imaging* (2001) 13: 615-618
94. Johansson, L.O., Fischer, S.E., Lorenz, C.H.:  
Benefit of T1 reduction for magnetic resonance coronary angiography: a numerical simulation and phantom study. *J Magn Reson Imaging* (1999) 9: 552-556
95. Johansson, L.O., Nolan, M.M., Taniuchi, M., Fischer, S.E., Wickline, S.A., Lorenz, C.H.:  
High-resolution magnetic resonance coronary angiography of the entire heart using a new blood-pool agent, NC100150 injection: comparison with invasive x-ray angiography in pigs. *J Cardiovasc Magn Reson* (1999) 1: 139-143
96. Jordan, A., Scholz, R., Wust, P., Fahling, H., Krause, J., Wlodarczyk, W., Sander, B., Vogl, T., Felix, R.:  
Effects of magnetic fluid hyperthermia (MFH) on C3H mammary carcinoma in vivo. *Int J Hyperthermia* (1997) 13: 587-605
97. Jordan, A., Wust, P., Scholz, R., Tesche, B., Fahling, H., Mitrovics, T., Vogl, T., Cervos-Navarro, J., Felix, R.:  
Cellular uptake of magnetic fluid particles and their effects on human adenocarcinoma cells exposed to AC magnetic fields in vitro. *Int J Hyperthermia* (1996) 12: 705-722
98. Jung, C.W., Jacobs, P.:  
Physical and chemical properties of superparamagnetic iron oxide MR contrast agents: ferumoxides, ferumoxtran, ferumoxsil. *Magn Reson Imaging* (1995) 13: 661-674
99. Kang, H.W., Josephson, L., Petrovsky, A., Weissleder, R., Bogdanov, A., Jr.:  
Magnetic Resonance Imaging of Inducible E-Selectin Expression in Human Endothelial Cell Culture. *Bioconjug Chem* (2002) 13: 122-127
100. Kanno, S., Lee, P.C., Dodd, S.J., Williams, M., Griffith, B.P., Ho, C.:  
A novel approach with magnetic resonance imaging used for the detection of lung allograft rejection. *J Thorac Cardiovasc Surg* (2000) 120: 923-934
101. Kanno, S., Wu, Y.J., Lee, P.C., Dodd, S.J., Williams, M., Griffith, B.P., Ho, C.:  
Macrophage accumulation associated with rat cardiac allograft rejection detected by magnetic resonance imaging with ultrasmall superparamagnetic iron oxide particles. *Circulation* (2001) 104: 934-938
102. Kato, N., Ihara, S., Tsujimoto, T., Miyazawa, T.:  
Effect of resovist on rats with different severities of liver cirrhosis. *Invest Radiol* (2002) 37: 292-298
103. Kehagias, D.T., Gouliamos, A.D., Smyrniotis, V., Vlahos, L.J.:  
Diagnostic efficacy and safety of MRI of the liver with superparamagnetic iron oxide particles (SH U 555 A). *J Magn Reson Imaging* (2001) 14: 595-601
104. Kellar, K.E., Fujii, D.K., Gunther, W.H., Briley-Saebo, K., Spiller, M., Koenig, S.H.:  
'NC100150', a preparation of iron oxide nanoparticles ideal for positive-contrast MR angiography. *MAGMA* (1999) 8: 207-213
105. Kim, D.K., Zhang, Y., Voit, W., Rao, K.V., Kehr, J., Bjelke, B., Muhammed, M.:  
Superparamagnetic iron oxide nanoparticles for bio-medical applications. *Scripta Mater* (2001) 44: 1713-1717

106. Kirschvink, J.L., Kobayashi-Kirschvink, A., Woodford, B.J.:  
Magnetite biomineralization in the human brain. *Proc Natl Acad Sci* (1992) 89: 7683-7687
107. Klein, C., Schalla, S., Schnackenburg, B., Bornstedt, A., Hoffmann, V., Fleck, E., Nagel, E.:  
Improvement of image quality of non-invasive coronary artery imaging with magnetic resonance by the use of the intravascular contrast agent Clariscan trade mark (NC100150 injection) in patients with coronary artery disease.  
*J Magn Reson Imaging* (2003) 17: 656-662
108. Knollmann, F.D., Bock, J.C., Rautenberg, K., Beier, J., Ebert, W., Felix, R.:  
Differences in predominant enhancement mechanisms of superparamagnetic iron oxide and ultrasmall superparamagnetic iron oxide for contrast-enhanced portal magnetic resonance angiography: preliminary results of an animal study original investigation.  
*Invest Radiol* (1998) 33: 637-643
109. Knollmann, F.D., Böck, J.C., Teltenkötter, S., Włodarczyk, W., Mühler, A., Vogl, T.J., Felix, R.:  
Evaluation of portal MR angiography using superparamagnetic iron oxide.  
*J Magn Reson Imaging* (1997) 7: 191-196
110. Knuesel, P.R., Nanz, D., Wolfensberger, U., Saranathan, M., Lehning, A., Luescher, T.F., Marincek, B., Von Schulthess, G.K., Schwitler, J.:  
Multislice breath-hold spiral magnetic resonance coronary angiography in patients with coronary artery disease: Effect of intravascular contrast medium.  
*J Magn Reson Imaging* (2002) 16: 660-667
111. Kobozev, N.I., Evdokimov, V.B., Zubovich, I.A., Mal'tsev, A.N.:  
Magnetochemistry of active centers. I. Magnetic and catalytic properties of dilute layers.  
*Zhur Fiz Khim* (1952) 26: 1349-1373
112. Krause, W., Gerlach, S., Muschick, P.:  
Prevention of the hemodynamic effects of iopromide-carrying liposomes in rats and pigs.  
*Invest Radiol* (2000) 35: 493-503
113. Kreft, B.P., Tanimoto, A., Leffler, S., Finn, J.P., Oksendal, A.N., Stark, D.D.:  
Contrast-enhanced MR imaging of diffuse and focal splenic disease with use of magnetic starch microspheres.  
*J Magn Reson Imaging* (1994) 4: 373-379
114. Kresse, M., Wagner, S., Pfeufferer, D., Lawaczek, R., Elste, V., Semmler, W.:  
Targeting of ultrasmall superparamagnetic iron oxide (USPIO) particles to tumor cells in vivo by using transferrin receptor pathways. *Magn Reson Med* (1998) 40: 236-242
115. Krieg, F.M., Andres, R.Y., Winterhalter, K.H.:  
Superparamagnetically labelled neutrophils as potential abscess-specific contrast agent for MRI. *Magn Reson Imaging* (1995) 13: 393-400
116. Kroencke, T.J., Wasser, M.N., Pattynama, P.M., Barentsz, J.O., Grabbe, E., Marchal, G., Knopp, M.V., Schneider, G., Bonomo, L., Pennell, D.J., del Maschio, A., Hentrich, H.R., Dapra, M., Kirchin, M.A., Spinazzi, A., Taupitz, M., Hamm, B.:  
Gadobenate dimeglumine-enhanced MR angiography of the abdominal aorta and renal arteries. *Am J Roentgenol* (2002) 179: 1573-1582
117. Kronick, P., Gilpin, R.W.:  
Use of superparamagnetic particles for isolation of cells.  
*J Biochem Biophys Meth* (1986) 12: 73-80
118. La Noce, A., Stoelben, S., Scheffler, K., Hennig, J., Lenz, H.M., La Ferla, R., Lorusso, V., Maggioni, F., Cavagna, F.:  
B22956/1, a new intravascular contrast agent for MRI: first administration to humans--preliminary results. *Acad Radiol* (2002) 9 Suppl 2: S404-406
119. Larsson, E.M., Sunden, P., Olsson, C.G., Debatin, J., Duerinckx, A.J., Baum, R., Hahn, D., Ebner, F.:  
MR venography using an intravascular contrast agent: results from a multicenter phase 2 study of dosage. *Am J Roentgenol* (2003) 180: 227-232

120. Lauffer, R.B., Parmelee, D.J., Dunham, S.U., Ouellet, H.S., Dolan, R.P., Witte, S., McMurry, T.J., Walovitch, R.C.:  
MS-325: albumin-targeted contrast agent for MR angiography.  
*Radiology* (1998) 207: 529-538
121. Lawaczeck, R., Bauer, H., Frenzel, T., Hasegawa, M., Ito, Y., Kito, K., Miwa, N., Tsutsui, H., Vogler, H., Weinmann, H.J.:  
Magnetic iron oxide particles coated with carboxydextran for parenteral administration and liver contrasting. Pre-clinical profile of SH U555A. *Acta Radiol* (1997) 38: 584-597
122. Lefebure, S., Dubois, E., Cabuil, V., Neveu, S., Massart, R.:  
Monodisperse magnetic nanoparticles: preparation and dispersion in water and oils.  
*J Mater Res* (1998) 13: 2975-2981
123. Leiner, T., Ho, K.Y., Ho, V.B., Bongartz, G., Mali, W.P., Rasch, W., Van Engelshoven, J.M.:  
Multicenter phase-II trial of safety and efficacy of NC100150 for steady-state contrast-enhanced peripheral magnetic resonance angiography. *Eur Radiol* (2003) 13: 1620-1627
124. Lenhart, M., Framme, N., Volk, M., Strotzer, M., Manke, C., Nitz, W.R., Finkenzeller, T., Feuerbach, S., Link, J.:  
Time-resolved contrast-enhanced magnetic resonance angiography of the carotid arteries: diagnostic accuracy and inter-observer variability compared with selective catheter angiography. *Invest Radiol* (2002) 37: 535-541
125. Leveille-Webster, C.R., Rogers, J., Arias, I.M.:  
Use of an asialoglycoprotein receptor-targeted magnetic resonance contrast agent to study changes in receptor biology during liver regeneration and endotoxemia in rats.  
*Hepatology* (1996) 23: 1631-1641
126. Lewin, M., Carlesso, N., Tung, C.H., Tang, X.W., Cory, D., Scadden, D.T., Weissleder, R.:  
Tat peptide-derivatized magnetic nanoparticles allow in vivo tracking and recovery of progenitor cells. *Nat Biotechnol* (2000) 18: 410-414
127. Li, D., Dolan, R.P., Walovitch, R.C., Lauffer, R.B.:  
Three-dimensional MRI of coronary arteries using an intravascular contrast agent.  
*Magn Reson Med* (1998) 39: 1014-1018
128. Li, D., Kaushikkar, S., Haacke, E.M., Woodard, P.K., Dhawale, P.J., Kroeker, R.M., Laub, G., Kuginuki, Y., Gutierrez, F.R.:  
Coronary arteries: three-dimensional MR imaging with retrospective respiratory gating.  
*Radiology* (1996) 201: 857-863
129. Li, D., Zheng, J., Weinmann, H.J.:  
Contrast-enhanced MR imaging of coronary arteries: comparison of intra- and extravascular contrast agents in swine. *Radiology* (2001) 218: 670-678
130. Loewe, C., Schillinger, M., Haumer, M., Loewe-Grgurin, M., Lammer, J., Thurnher, S., Fuchsjager, M., Cejna, M.:  
MRA versus DSA in the assessment of occlusive disease in the aortic arch vessels: accuracy in detecting the severity, number, and length of stenoses.  
*J Endovasc Ther* (2004) 11: 152-160
131. Lonnemark, M., Hemmingsson, A., Bach-Gansmo, T., Ericsson, A., Oksendal, A., Nyman, R., Moxnes, A.:  
Effect of superparamagnetic particles as oral contrast medium at magnetic resonance imaging. A phase I clinical study. *Acta Radiol* (1989) 30: 193-196
132. Lorenz, C.H., Flacke, S., Chia, J.M., Taniuchi, M., Allen, J.S., McLEAn, M., Setser, R.M., Chan, T., Weiskoff, R.:  
Steady-State Detection of Ischemia with AngioMARK.  
*Proc Int Soc Mag Reson Med* 8 (2000) 40
133. Lorenz, C.H., Johansson, L.O.:  
Contrast-enhanced coronary MRA.  
*J Magn Reson Imaging* (1999) 10: 703-708

134. Lucidarme, O., Baleston, F., Cadi, M., Bellin, M.F., Charlotte, F., Ratzu, V., Grenier, P.A.: Non-invasive detection of liver fibrosis: Is superparamagnetic iron oxide particle-enhanced MR imaging a contributive technique? *Eur Radiol* (2003) 13: 467-474
135. Mack, M.G., Balzer, J.O., Straub, R., Eichler, K., Vogl, T.J.: Superparamagnetic iron oxide-enhanced MR imaging of head and neck lymph nodes. *Radiology* (2002) 222: 239-244
136. Mallouhi, A., Schocke, M., Judmaier, W., Wolf, C., Dessl, A., Czermak, B.V., Waldenberger, P., Jaschke, W.R.: 3D MR angiography of renal arteries: comparison of volume rendering and maximum intensity projection algorithms. *Radiology* (2002) 223: 509-516
137. Manning, W.J., Li, W., Boyle, N.G., Edelman, R.R.: Fat-suppressed breath-hold magnetic resonance coronary angiography. *Circulation* (1993) 87: 94-104
138. Marchessault, R.H., Ricard, S., Rioux, P.: In Situ synthesis of ferrites in lignocellulosis. *Carbohydr Res* (1992) 224: 133-139
139. Maron, D.M., Ames, B.N.: Revised methods for the Salmonella mutagenicity test. *Mutat Res* (1983) 113: 173-215.
140. Massart, R., Dubois, E., Cabuil, V., Hasmonay, E.: Preparation and properties of monodisperse magnetic fluids. *J Magn Magn Mater* (1995) 149: 1-5
141. McLachlan, S.J., Morris, M.R., Lucas, M.A., Fisco, R.A., Eakins, M.N., Fowler, D.R., Scheetz, R.B., Olukotun, A.Y.: Phase I clinical evaluation of a new iron oxide MR contrast agent. *J Magn Reson Imaging* (1994) 4: 301-307
142. Meaney, J.F., Johansson, L.O., Ahlstrom, H., Prince, M.R.: Pulmonary magnetic resonance angiography. *J Magn Reson Imaging* (1999) 10: 326-338
143. Meyer, C.H., Hu, B.S., Nishimura, D.G., Macovski, A.: Fast spiral coronary artery imaging. *Magn Reson Med* (1992) 28: 202-213
144. Misselwitz, B., Schmitt-Willich, H., Ebert, W., Frenzel, T., Weinmann, H.J.: Pharmacokinetics of Gadomer-17, a new dendritic magnetic resonance contrast agent. *MAGMA* (2001) 12: 128-134
145. Mohrs, O.K., Oberholzer, K., Krummenauer, F., Bernhard, S., Kalden, P., Neufang, A., Thelen, M., Kreitner, K.F.: Ergebnisse der kontrastverstärkten MR-Angiographie der aortoiliakalen Gefäße mit einem 1-molaren Kontrastmittel bei 1,0 T: Vergleich zur i. a. DSA. *Fortschr Röntgenstr* (2004) 176: 985-991
146. Moroz, P., Jones, S.K., Gray, B.N.: Tumor response to arterial embolization hyperthermia and direct injection hyperthermia in a rabbit liver tumor model. *J Surg Oncol* (2002) 80: 149-156
147. Moroz, P., Jones, S.K., Winter, J., Gray, B.N.: Targeting liver tumors with hyperthermia: Ferromagnetic embolization in a rabbit liver tumor model. *J Surg Oncol* (2001) 78: 22-29
148. Mühler, A., Freise, C.E., Kuwatsuru, R., Rosenau, W., Liu, T., Mintorovitch, J., Clement, O., Vexler, V., Mahboubi, S., Lang, P.: Acute liver rejection: evaluation with cell-directed MR contrast agents in a rat transplantation model. *Radiology* (1993) 186: 139-146
149. Mühler, A., Saeed, M., Brasch, R.C., Higgins, C.B.: Amelioration of cardiodepressive effects of gadopentetate dimeglumine with addition of ionic calcium. *Radiology* (1992) 184: 159-164
150. Mühler, A., Zhang, X., Wang, H., Lawaczeck, R., Weinmann, H.J.: Investigation of mechanisms influencing the accumulation of ultrasmall superparamagnetic iron oxide particles in lymph nodes. *Invest Radiol* (1995) 30: 98-103

151. Müller, M., Reimer, P., Wiedermann, D., Allkemper, T., Marx, C., Tombach, B., Rummeny, E.J., Shamsi, K., Balzer, T., Peters, P.E.:  
T1-gewichtete dynamische MRT mit neuen superparamagnetischen Eisenoxidpartikeln (Resovist): Ergebnisse einer Studie am Phantom sowie an 25 Patienten.  
*Fortschr Röntgenstr* (1998) 168: 228-236
152. Müller, R.D., Vogel, K., Neumann, K., Hirche, H., Barkhausen, J., Stoblen, F., Henrich, H., Langer, R.:  
SPIO-MR imaging versus double-phase spiral CT in detecting malignant lesions of the liver.  
*Acta Radiol* (1999) 40: 628-635
153. Noguchi, Y., Murakami, T., Kim, T., Hori, M., Osuga, K., Kawata, S., Kumano, S., Okada, A., Sugiura, T., Nakamura, H.:  
Detection of hepatocellular carcinoma: comparison of dynamic MR imaging with dynamic double arterial phase helical CT. *Am J Roentgenol* (2003) 180: 455-460
154. Normann, S.J.:  
Reticuloendothelial system function. VI. Experimental alterations influencing the correlation between portal blood flow and colloid clearance. *J Reticuloendothel Soc* (1973) 13: 47-60
155. Oswald, P., Clement, O., Chambon, C., Schouman-Claeys, E., Frija, G.:  
Liver positive enhancement after injection of superparamagnetic nanoparticles: respective role of circulating and uptaken particles. *Magn Reson Imaging* (1997) 15: 1025-1031
156. Padfield, J.: Annual Report & Accounts 2001.  
<http://www.amersham.com/investors/AR01/rep15.html> (2001)
157. Paetsch, I., Huber, M.E., Bornstedt, A., Schnackenburg, B., Boesiger, P., Stuber, M., Fleck, E., Cavagna, F., Nagel, E.:  
Improved three-dimensional free-breathing coronary magnetic resonance angiography using gadoleptic acid (B-22956) for intravascular contrast enhancement.  
*J Magn Reson Imaging* (2004) 20: 288-293
158. Pannu, H.K., Wang, K.P., Borman, T.L., Bluemke, D.A.:  
MR imaging of mediastinal lymph nodes: evaluation using a superparamagnetic contrast agent. *J Magn Reson Imaging* (2000) 12: 899-904
159. Panting, J.R., Taylor, A.M., Gatehouse, P.D., Keegan, J., Yang, G.Z., McGill, S., Francis, J.M., Burman, E.D., Firmin, D.N., Pennell, D.J.:  
First-pass myocardial perfusion imaging and equilibrium signal changes using the intravascular contrast agent NC100150 injection. *J Magn Reson Imaging* (1999) 10: 404-410
160. Patek, P.R., De Mignard, A., Bernick, S.:  
Age changes in structure and responses of reticuloendothelial cells of rat liver.  
*J Reticuloendothel Soc* (1967) 4: 211-218
161. Päufer, S., Reszka, R., Wagner, S., Wolf, K.J., Buhr, H.J., Berger, G.:  
Liposome-encapsulated superparamagnetic iron oxide particles as markers in an MRI-guided search for tumor-specific drug carriers.  
*Anticancer Drug Des* (1997) 12: 125-135
162. Pilgrim, H.:  
Stabilized suspension of magnetic particles and its preparation and use in NMR diagnosis.  
Patent: DE 3709851 (1988)
163. Pilgrim, H.:  
Superparamagnetic particles with increased R1 relaxivity, their production and their use.  
Patent: WO 9735200 (1997)
164. Pilgrim, H.:  
Superparamagnetic particles with increased R1 relaxivity, process for producing said particles and use thereof. Patent: EP 888545, JP 2000-507197 (2000)
165. Port, M., Corot, C., Raynal, I., Idee, J.M., Dencausse, A., Lancelot, E., Meyer, D., Bonnemain, B., Lautrou, J.:  
Physicochemical and biological evaluation of P792, a rapid-clearance blood-pool agent for magnetic resonance imaging. *Invest Radiol* (2001) 36: 445-454

166. Port, M., Corot, C., Rousseaux, O., Raynal, I., Devoldere, L., Idee, J.M., Dencausse, A., Le Greneur, S., Simonot, C., Meyer, D.: P792:  
a rapid clearance blood pool agent for magnetic resonance imaging: preliminary results. *MAGMA* (2001) 12: 121-127
167. Potenza, B.B., Boscardin, M., de Michele, V., Gramaccioli, C.M., Quaroni, A., Roseo, I., Sosso, F.:  
Das grosse Buch der Mineralien und Edelsteine.  
Klagenfurt: Neuer Kaiser Verlag - Gesellschaft m.b.H., (2002).
168. Prince, M.R.:  
Gadolinium-enhanced MR aortography. *Radiology* (1994) 191: 155-164
169. Prince, M.R., Yucel, E.K., Kaufman, J.A., Harrison, D.C., Geller, S.C.:  
Dynamic gadolinium-enhanced three-dimensional abdominal MR arteriography. *J Magn Reson Imaging* (1993) 3: 877-881
170. Prince, M.R., Zhang, H.L., Chabra, S.G., Jacobs, P., Wang, Y.:  
A pilot investigation of new superparamagnetic iron oxide (ferumoxylol) as a contrast agent for cardiovascular MRI. *J X-Ray Science Tech* (2003) 11: 231-240
171. Reimer, P., Balzer, T.:  
Ferucarbotran (Resovist): a new clinically approved RES-specific contrast agent for contrast-enhanced MRI of the liver: properties, clinical development, and applications. *Eur Radiol* (2003) 13: 1266-1276
172. Reimer, P., Jahnke, N., Fiebich, M., Schima, W., Deckers, F., Marx, C., Holzknacht, N., Saini, S.:  
Hepatic lesion detection and characterization: value of nonenhanced MR imaging, superparamagnetic iron oxide-enhanced MR imaging, and spiral CT-ROC analysis. *Radiology* (2000) 217: 152-158
173. Reimer, P., Marx, C., Rummeny, E.J., Muller, M., Lentschig, M., Balzer, T., Dietl, K.H., Sulkowski, U., Berns, T., Shamsi, K., Peters, P.E.:  
SPIO-enhanced 2D-TOF MR angiography of the portal venous system: results of an intraindividual comparison. *J Magn Reson Imaging* (1997) 7: 945-949
174. Reimer, P., Muller, M., Marx, C., Wiedermann, D., Muller, R., Rummeny, E.J., Ebert, W., Shamsi, K., Peters, P.E.:  
T1 effects of a bolus-injectable superparamagnetic iron oxide, SH U 555 A: dependence on field strength and plasma concentration--preliminary clinical experience with dynamic T1-weighted MR imaging. *Radiology* (1998) 209: 831-836
175. Reimer, P., Weissleder, R., Brady, T.J., Yeager, A.E., Baldwin, B.H., Tennant, B.C., Wittenberg, J.:  
Experimental hepatocellular carcinoma: MR receptor imaging. *Radiology* (1991) 180: 641-645
176. Reimer, P., Weissleder, R., Lee, A.S., Buettner, S., Wittenberg, J., Brady, T.J.:  
Asialoglycoprotein receptor function in benign liver disease: evaluation with MR imaging. *Radiology* (1991) 178: 769-774
177. Reimer, P., Weissleder, R., Lee, A.S., Wittenberg, J., Brady, T.J.:  
Receptor imaging: application to MR imaging of liver cancer. *Radiology* (1990) 177: 729-734
178. Reimer, P., Weissleder, R., Shen, T., Knoefel, W.T., Brady, T.J.:  
Pancreatic receptors: initial feasibility studies with a targeted contrast agent for MR imaging. *Radiology* (1994) 193: 527-531
179. Remonda, L., Senn, P., Barth, A., Arnold, M., Lovblad, K.O., Schroth, G.:  
Contrast-enhanced 3D MR angiography of the carotid artery: comparison with conventional digital subtraction angiography. *Am J Neuroradiol* (2002) 23: 213-219
180. Remsen, L.G., McCormick, C.I., Roman-Goldstein, S., Nilaver, G., Weissleder, R., Bogdanov, A., Hellstrom, I., Kroll, R.A., Neuwelt, E.A.:  
MR of carcinoma-specific monoclonal antibody conjugated to monocrySTALLINE iron oxide nanoparticles: the potential for noninvasive diagnosis. *Am J Neuroradiol* (1996) 17: 411-418

181. Renshaw, P.F., Owen, C.S., Evans, A.E., Leigh, J.S., Jr.:  
Immunospecific NMR contrast agents. *Magn Reson Imaging* (1986) 4: 351-357
182. Rety, F., Clement, O., Siauve, N., Cuenod, C.A., Carnot, F., Sich, M., Buisine, A., Frija, G.:  
MR lymphography using iron oxide nanoparticles in rats: pharmacokinetics in the lymphatic system after intravenous injection. *J Magn Reson Imaging* (2000) 12: 734-739
183. Rohl, L., Ostergaard, L., Simonsen, C.Z., Vestergaard Poulsen, P., Sorensen, L., Bjornerud, A., Saebo, K.B., Gyldensted, C.:  
NC100150-enhanced 3D-SPGR MR angiography of the common carotid artery in a pig vascular stenosis model. Quantification of stenosis and dose optimization. *Acta Radiol* (1999) 40: 282-290
184. Rozenman, Y., Zou, X.M., Kantor, H.L.:  
Magnetic resonance imaging with superparamagnetic iron oxide particles for the detection of myocardial reperfusion. *Magn Reson Imaging* (1991) 9: 933-939
185. Rubin, D.L., Desser, T.S., Semelka, R., Brown, J., Nghiem, H.V., Stevens, W.R., Bluemke, D., Nelson, R., Fultz, P., Reimer, P., Ho, V., Kristy, R.M., Pierro, J.A.:  
A multicenter, randomized, double-blind study to evaluate the safety, tolerability, and efficacy of OptiMARK (gadoversetamide injection) compared with Magnevist (gadopentetate dimeglumine) in patients with liver pathology: results of a Phase III clinical trial. *J Magn Reson Imaging* (1999) 9: 240-250
186. Ruehm, S.G., Corot, C., Vogt, P., Kolb, S., Debatin, J.F.:  
Magnetic resonance imaging of atherosclerotic plaque with ultrasmall superparamagnetic particles of iron oxide in hyperlipidemic rabbits. *Circulation* (2001) 103: 415-422
187. Saeed, M., Wendland, M.F., Engelbrecht, M., Sakuma, H., Higgins, C.B.:  
Value of blood pool contrast agents in magnetic resonance angiography of the pelvis and lower extremities. *Eur Radiol* (1998) 8: 1047-1053
188. Sahani, D., Saini, S., Sharma, R., O'Malley, M., Hahn, P.:  
Dynamic T1-weighted ferumoxides enhanced MRI for imaging liver hemangiomas: preliminary observations. *Abdom Imaging* (2001) 26: 166-170
189. Saini, S., Edelman, R.R., Sharma, P., Li, W., Mayo Smith, W., Slater, G.J., Eisenberg, P.J., Hahn, P.F.:  
Blood-pool MR contrast material for detection and characterization of focal hepatic lesions: initial clinical experience with ultrasmall superparamagnetic iron oxide (AMI-227). *Am J Roentgenol* (1995) 164: 1147-1152
190. Saini, S., Sharma, R., Baron, R.L., Turner, D.A., Ros, P.R., Hahn, P.F., Small, W.C., Delange, E.E., Stillman, A.E., Edelman, R.R., Runge, V.M., Outwater, E.K.:  
Multicentre dose-ranging study on the efficacy of USPIO ferumoxtran-10 for liver MR imaging. *Clin Radiol* (2000) 55: 690-695
191. Saini, S., Stark, D.D., Hahn, P.F., Bousquet, J.C., Introcasso, J., Wittenberg, J., Brady, T.J., Ferrucci, J.T., Jr.:  
Ferrite particles: a superparamagnetic MR contrast agent for enhanced detection of liver carcinoma. *Radiology* (1987) 162: 217-222
192. Saini, S., Stark, D.D., Hahn, P.F., Wittenberg, J., Brady, T.J., Ferrucci, J.T.:  
Ferrite particles: a superparamagnetic MR contrast agent for the reticuloendothelial system. *Radiology* (1987) 162: 211-216
193. Sako, M., Hirota, S.:  
Embolotherapy of hepatomas using ferromagnetic microspheres, its clinical evaluation and the prospect of its use as a vehicle in chemoembolo-hyperthermic therapy. *Gan to Kagaku Ryoho* (1986) 13: 1618-1624
194. Sandstede, J.J., Pabst, T., Wacker, C., Wiesmann, F., Hoffmann, V., Beer, M., Kenn, W., Bauer, W., Hahn, D.:  
Breath-hold 3D MR coronary angiography with a new intravascular contrast agent (feruglose)--first clinical experiences. *Magn Reson Imaging* (2001) 19: 201-205

195. Sardanelli, F., Molinari, G., Zandrino, F., Balbi, M.:  
Three-dimensional, navigator-echo MR coronary angiography in detecting stenoses of the major epicardial vessels, with conventional coronary angiography as the standard of reference. *Radiology* (2000) 214: 808-814
196. Schafer, F.K., Schafer, P.J., Jahnke, T., Walluscheck, K., Priebe, M., Hentsch, A., Heller, M., Brossmann, J.:  
Erste klinische Erfahrungen zur Wertigkeit der kontrastverstärkten Magnetresonanztomographie der Becken-Bein-Strombahn mit dem 1,0 molaren Gadobutrol - Vergleich zur i. a. DSA. *Fortschr Röntgenstr* (2003) 175: 556-564
197. Schaffer, B.K., Linker, C., Papisov, M., Tsai, E., Nossiff, N., Shibata, T., Bogdanov, A., Jr., Brady, T.J., Weissleder, R.:  
MION-ASF: biokinetics of an MR receptor agent. *Magn Reson Imaging* (1993) 11: 411-417
198. Schellenberger, E.A., Bogdanov, A., Jr., Hogemann, D., Tait, J., Weissleder, R., Josephson, L.:  
Annexin V-CLIO: a nanoparticle for detecting apoptosis by MRI. *Mol Imaging* (2002) 1: 102-107
199. Scherer, F., Anton, M., Schillinger, U., Henke, J., Bergemann, C., Kruger, A., Gansbacher, B., Plank, C.:  
Magnetofection: enhancing and targeting gene delivery by magnetic force in vitro and in vivo. *Gene Ther* (2002) 9: 102-109
200. Schmiedl, U., Brasch, R.C., Ogan, M.D., Moseley, M.E.:  
Albumin labeled with Gd-DTPA. An intravascular contrast-enhancing agent for magnetic resonance blood pool and perfusion imaging. *Acta Radiol Suppl* (1990) 374: 99-102
201. Schmitz, S.A., Albrecht, T., Jensen, K., Wolf, K.J.:  
SPIO-unterstützte MR-Angiographie des portalvenösen Systems. *Fortschr Röntgenstr* (2000) 172: 51-54
202. Schmitz, S.A., Albrecht, T., Wolf, K.J.:  
MR angiography with superparamagnetic iron oxide: feasibility study. *Radiology* (1999) 213: 603-607
203. Schmitz, S.A., Coupland, S.E., Gust, R., Winterhalter, S., Wagner, S., Kresse, M., Semmler, W., Wolf, K.J.:  
Superparamagnetic iron oxide-enhanced MRI of atherosclerotic plaques in Watanabe heritable hyperlipidemic rabbits. *Invest Radiol* (2000) 35: 460-471
204. Schmitz, S.A., Taupitz, M., Wagner, S., Wolf, K.J., Beyersdorff, D., Hamm, B.:  
Magnetic resonance imaging of atherosclerotic plaques using superparamagnetic iron oxide particles. *J Magn Reson Imaging* (2001) 14: 355-361
205. Schmitz, S.A., Winterhalter, S., Schiffler, S., Gust, R., Wagner, S., Kresse, M., Coupland, S.E., Semmler, W., Wolf, K.J.:  
USPIO-enhanced direct MR imaging of thrombus: preclinical evaluation in rabbits. *Radiology* (2001) 221: 237-243
206. Schnorr, J., Taupitz, M., Wagner, S., Pilgrimm, H., Hansel, J., Hamm, B.:  
Age-related blood half-life of particulate contrast material: experimental results with a USPIO in rats. *J Magn Reson Imaging* (2000) 12: 740-744
207. Schnorr, J., Wagner, S., Abramjuk, C., Wojner, I., Schink, T., Kroencke, T.J., Schellenberger, E.A., Hamm, B., Pilgrimm, H., Taupitz, M.:  
Comparison of the Iron-Oxide-Based Blood-Pool Contrast Medium VSOP-C184 with Gadopentetate Dimeglumine for First-Pass MRA of the Aorta and Renal Arteries in Pigs. *Invest Radiol* (2004) 39: 546-553
208. Schnorr, J., Wagner, S., Ebert, W., Heyer, C., Laub, G., Kivelitz, D., Abramjuk, C., Hamm, B., Taupitz, M.:  
MR-Angiographie der Koronararterien: Vergleich des Blutpoolkontrastmittels Gadomer mit Gd-DTPA am Schwein. *Fortschr Röntgenstr* (2003) 175: 822-829
209. Schulze, E., Ferrucci, J.T., Jr., Poss, K., Lapointe, L., Bogdanova, A., Weissleder, R.:  
Cellular uptake and trafficking of a prototypical magnetic iron oxide label in vitro. *Invest Radiol* (1995) 30: 604-610

210. Semelka, R.C., Cance, W.G., Marcos, H.B., Mauro, M.A.:  
Liver metastases: comparison of current MR techniques and spiral CT during arterial portography for detection in 20 surgically staged cases. *Radiology* (1999) 213: 86-91
211. Semelka, R.C., Martin, D.R., Balci, C., Lance, T.:  
Focal liver lesions: comparison of dual-phase CT and multisequence multiplanar MR imaging including dynamic gadolinium enhancement. *J Magn Reson Imaging* (2001) 13: 397-401
212. Seneterre, E., Weissleder, R., Jaramillo, D., Reimer, P., Lee, A.S.,  
Brady, T.J., Wittenberg, J.: Bone marrow: ultrasmall superparamagnetic iron oxide for MR imaging. *Radiology* (1991) 179: 529-533
213. Shen, T., Weissleder, R., Papisov, M., Bogdanov, A., Jr., Brady, T.J.:  
Monocrystalline iron oxide nanocompounds (MION): physicochemical properties. *Magn Reson Med* (1993) 29: 599-604
214. Sigal, R., Vogl, T., Casselman, J., Moulin, G., Veillon, F., Hermans, R., Dubrulle, F., Viala, J.,  
Bosq, J., Mack, M., Depondt, M., Mattelaer, C., Petit, P., Champsaur, P., Riehm, S.,  
Dadashitazehozhi, Y., de Jaegere, T., Marchal, G., Chevalier, D., Lemaitre, L., Kubiak, C.,  
Helmberger, R., Halimi, P.:  
Lymph node metastases from head and neck squamous cell carcinoma: MR imaging with ultrasmall superparamagnetic iron oxide particles (Sinerem MR) -- results of a phase-III multicenter clinical trial. *Eur Radiol* (2002) 12: 1104-1113
215. Spuentrup, E., Buecker, A., Stuber, M., Botnar, R., Nguyen, T.H., Bornert, P.,  
Kolker, C., Gunther, R.W.:  
Navigator-gated coronary magnetic resonance angiography using steady-state-free-precession: comparison to standard T2-prepared gradient-echo and spiral imaging. *Invest Radiol* (2003) 38: 263-268
216. Spuentrup, E., Stuber, M., Botnar, R.M., Kissinger, K.V., Manning, W.J.:  
Real-time motion correction in navigator-gated free-breathing double-oblique submillimeter 3D right coronary artery magnetic resonance angiography. *Invest Radiol* (2002) 37: 632-636
217. Stark, D.D., Bradley, W.G.:  
*Magnetic Resonance Imaging*. St. Louis: Mosby-Year Book Inc, (1992).
218. Stark, D.D., Weissleder, R., Elizondo, G., Hahn, P.F., Saini, S., Todd, L.E.,  
Wittenberg, J., Ferrucci, J.T.:  
Superparamagnetic iron oxide: clinical application as a contrast agent for MR imaging of the liver. *Radiology* (1988) 168: 297-301
219. Stillman, A.E., Wilke, N., Li, D., Haacke, M., McLachlan, S.:  
Ultrasmall superparamagnetic iron oxide to enhance MRA of the renal and coronary arteries: studies in human patients. *J Comput Assist Tomogr* (1996) 20: 51-55
220. Stroszczyński, C., Gaffke, G., Gretschel, S., Rambow, A., Jost, D., Schlecht, I.,  
Schneider, U., Schicke, B., Hohenberger, P., Gebauer, B., Felix, R.:  
Differenzierung von Leberläsionen mit SPIO-gestützten T1-w und T2-w MRT-Aufnahmen: Eine ROC-Analyse. *Fortschr Röntgenstr* (2003) 175: 1368-1375
221. Stuber, M., Botnar, R.M., Danias, P.G., McConnell, M.V., Kissinger, K.V.,  
Yucel, E.K., Manning, W.J.:  
Contrast agent-enhanced, free-breathing, three-dimensional coronary magnetic resonance angiography. *J Magn Reson Imaging* (1999) 10: 790-799
222. Stuber, M., Botnar, R.M., Danias, P.G., Sodickson, D.K., Kissinger, K.V., Van Cauwen, M.,  
De Becker, J., Manning, W.J.:  
Double-oblique free-breathing high resolution three-dimensional coronary magnetic resonance angiography. *J Am Coll Cardiol* (1999) 34: 524-531
223. Suwa, T., Ozawa, S., Ueda, M., Ando, N., Kitajima, M.:  
Magnetic resonance imaging of esophageal squamous cell carcinoma using magnetite particles coated with anti-epidermal growth factor receptor antibody. *Int J Cancer* (1998) 75: 626-634

224. Suzuki, M., Honda, H., Kobayashi, T., Wakabayashi, T., Yoshida, J., Takahashi, M.: Development of a target-directed magnetic resonance contrast agent using monoclonal antibody-conjugated magnetic particles. *Noshuyo Byori* (1996) 13: 127-132
225. Swan, J.S., Kennell, T.W., Acher, C.W., Heisey, D.M., Grist, T.M., Korosec, F.R., Hagenauer, M.E.: Magnetic resonance angiography of aorto-iliac disease. *Am J Surg* (2000) 180: 6-12
226. Tacke, J., Adam, G., Classen, H., Muhler, A., Prescher, A., Gunther, R.W.: Dynamic MRI of a hypovascularized liver tumor model: comparison of a new blood pool contrast agent (24-gadolinium-DTPA-cascade-polymer) with gadopentetate dimeglumine. *J Magn Reson Imaging* (1997) 7: 678-682
227. Takahama, K., Amano, Y., Hayashi, H., Ishihara, M., Kumazaki, T.: Detection and characterization of focal liver lesions using superparamagnetic iron oxide-enhanced magnetic resonance imaging: comparison between ferumoxides-enhanced T1-weighted imaging and delayed-phase gadolinium-enhanced T1-weighted imaging. *Abdom Imaging* (2003) 28: 525-530
228. Tang, J., Myers, M., Bosnick, K.A., Brus, L.E.: Magnetite Fe<sub>3</sub>O<sub>4</sub> Nanocrystals: Spectroscopic Observation of Aqueous Oxidation Kinetics. *J Phys Chem B* (2003) 107: 7501-7506
229. Tanimoto, A., Yuasa, Y., Hiramatsu, K.: Enhancement of phase-contrast MR angiography with superparamagnetic iron oxide. *J Magn Reson Imaging* (1998) 8: 446-450
230. Taupitz, M., Schmitz, S., Hamm, B.: Superparamagnetische Eisenoxidpartikel: Aktueller Stand und zukünftige Entwicklung. *Fortschr Röntgenstr* (2003) 175: 752-765
231. Taupitz, M., Schnorr, J., Abramiuk, C., Wagner, S., Pilgrimm, H., Hünigen, H., Hamm, B.: New Generation of Monomer-Stabilized Very Small Superparamagnetic Iron Oxide Particles (VSOP) as Contrast Medium for MR Angiography - Preclinical Results in Rats and Rabbits. *J Magn Reson Imaging* (2000) 12: 905-911
232. Taupitz, M., Schnorr, J., Wagner, S., Abramjuk, C., Pilgrimm, H., Kivelitz, D., Schink, T., Hansel, J., Laub, G., Hunigen, H., Hamm, B.: Coronary MR angiography: experimental results with a monomer-stabilized blood pool contrast medium. *Radiology* (2002) 222: 120-126
233. Taupitz, M., Wagner, S., Hamm, B., Binder, A., Pfefferer, D.: Interstitial MR lymphography with iron oxide particles: results in tumor-free and VX2 tumor-bearing rabbits. *Am J Roentgenol* (1993) 161: 193-200
234. Taupitz, M., Wagner, S., Schnorr, J., Dewey, M., Rogalla, P., Krug, L., Laub, G., Kivelitz, D., Pilgrimm, H., Kravec, I., Hamm, B.: MR angiography of the entire coronary system with a very small iron oxide blood-pool contrast medium: results of a phase IB clinical trial. *Proc Int Soc Mag Reson Med* 11 (2003) 725
235. Taylor, A.M., Panting, J.R., Keegan, J., Gatehouse, P.D., Amin, D., Jhooti, P., Yang, G.Z., McGill, S., Burman, E.D., Francis, J.M., Firmin, D.N., Pennell, D.J.: Safety and preliminary findings with the intravascular contrast agent NC100150 injection for MR coronary angiography. *J Magn Reson Imaging* (1999) 9: 220-227
236. Taylor, A.M., Panting, J.R., Keegan, J., Gatehouse, P.D., Jhooti, P., Yang, G.Z., McGill, S., Francis, J.M., Burman, E.D., Firmin, D.N., Pennell, D.J.: Use of the intravascular contrast agent NC100150 injection in spin-echo and gradient-echo imaging of the heart. *J Cardiovasc Magn Reson* (1999) 1: 23-32
237. Tiefenauer, L.X., Kuehne, G., Andres, R.Y.: Antibody-magnetite nanoparticles: In vitro characterization of a potential tumor-specific contrast agent for magnetic resonance imaging. *Bioconj Chem* (1993) 4: 347-352

238. Tombach, B., Reimer, P., Mahler, M., Ebert, W., Pering, C., Heindel, W.:  
First-pass and equilibrium phase MRA following intravenous bolus injection of SH U 555 C:  
Phase I clinical trial in elderly volunteers with risk factors for arterial vascular disease.  
*Acad Radiol* (2002) 9 Suppl 2: S425-427
239. Turetschek, K., Huber, S., Floyd, E., Helbich, T., Roberts, T.P., Shames, D.M., Tarlo, K.S.,  
Wendland, M.F., Brasch, R.C.:  
MR imaging characterization of microvessels in experimental breast tumors by using a  
particulate contrast agent with histopathologic correlation. *Radiology* (2001) 218: 562-569
240. Turetschek, K., Huber, S., Helbich, T., Floyd, E., Tarlo Kirk, S., Roberts Timothy, P.L.,  
Shames David, M., Wendland Michael, F., Brasch Robert, C.:  
Dynamic MRI enhanced with albumin-(Gd-DTPA)<sub>30</sub> or ultrasmall superparamagnetic iron  
oxide particles (NC100150 injection) for the measurement of microvessel permeability in  
experimental breast tumors. *Acad Radiol* (2002) 9 Suppl 1: S112-114
241. Turetschek, K., Roberts, T.P., Floyd, E., Preda, A., Novikov, V., Shames, D.M.,  
Carter, W.O., Brasch, R.C.:  
Tumor microvascular characterization using ultrasmall superparamagnetic iron oxide  
particles (USPIO) in an experimental breast cancer model.  
*J Magn Reson Imaging* (2001) 13: 882-888
242. Urhahn, R., Adam, G., Busch, N., Chen, J.H., Euringer, W., Gunther, R.W.:  
Superparamagnetische Eisenoxidpartikel: Welchen Stellenwert hat der T1-Effekt für die MR-  
Diagnostik fokaler Leberläsionen? *Fortschr Röntgenstr* (1996) 165: 364-370
243. Vavrik, J., Rohmoser, G.M., Madani, B., Ersek, M., Tscholakoff, D., Bucek, R.A.:  
Comparison of MR angiography versus digital subtraction angiography as a basis for  
planning treatment of lower limb occlusive disease. *J Endovasc Ther* (2004) 11: 294-301
244. Vogl, T.J., Hammerstingl, R., Pegios, W., Hamm, B., Eibl-Eibesfeldt, A., Woessmer, B.,  
Lissner, J., Felix, R.:  
Wertigkeit des leberspezifischen superparamagnetischen Kontrastmittels AMI-25 für die  
Detektion und Differential-diagnose lebereigener Tumoren versus Metastasen.  
*Fortschr Röntgenstr* (1994) 160: 319-328
245. Wacker, F.K., Reither, K., Ebert, W., Wendt, M., Lewin, J.S., Wolf, K.J.:  
MR Image-guided Endovascular Procedures with the Ultrasmall Superparamagnetic Iron  
Oxide SH U 555 C as an Intravascular Contrast Agent: Study in Pigs.  
*Radiology* (2003) 226: 459-464
246. Wagenseil, J.E., Johansson, L.O., Lorenz, C.H.:  
Characterization of t1 relaxation and blood-myocardial contrast enhancement of NC100150  
injection in cardiac MRI. *J Magn Reson Imaging* (1999) 10: 784-789
247. Wagner, H.N., Jr., Migita, T., Solomon, N.:  
Effect of age on reticuloendothelial function in man. *J Gerontol* (1966) 21: 57-62
248. Wagner, S., Pfefferer, D., Ebert, W., Kresse, M., Taupitz, M., Hamm, B.,  
Lawaczek, R., Wolf, K.J.:  
Intravenous MR lymphography with superparamagnetic iron oxide particles: Experimental  
studies in rats and rabbits. *Eur Radiol* (1995) 5: 640-646
249. Wagner, S., Schnorr, J., Pilgrim, H., Hamm, B., Taupitz, M.:  
Monomer-coated very small superparamagnetic iron oxide particles as contrast medium for  
magnetic resonance imaging: Preclinical in vivo characterization.  
*Invest Radiol* (2002) 37: 167-177
250. Wang, Y., Rossman, P.J., Grimm, R.C., Riederer, S.J., Ehman, R.L.:  
Navigator-echo-based real-time respiratory gating and triggering for reduction of respiration  
effects in three-dimensional coronary MR angiography. *Radiology* (1996) 198: 55-60
251. Wang, Y.X., Hussain, S.M., Krestin, G.P.:  
Superparamagnetic iron oxide contrast agents: physicochemical characteristics and  
applications in MR imaging. *Eur Radiol* (2001) 11: 2319-2331

252. Watanabe, Y.:  
Serial inbreeding of rabbits with hereditary hyperlipidemia (WHHL-rabbit).  
*Atherosclerosis* (1980) 36: 261-268
253. Watts, R., Wang, Y., Redd, B., Winchester, P.A., Kent, K.C., Bush, H.L., Prince, M.R.:  
Recessed elliptical-centric view-ordering for contrast-enhanced 3D MR angiography of the carotid arteries. *Magn Reson Med* (2002) 48: 419-424
254. Weinmann, H.J., Laniado, M., Mutzel, W.:  
Pharmacokinetics of GdDTPA/dimeglumine after intravenous injection into healthy volunteers. *Physiol Chem Phys Med NMR* (1984) 16: 167-172
255. Weishaupt, D., Hilfiker, P.R., Schmidt, M., Debatin, J.F.:  
Pulmonary hemorrhage: imaging with a new magnetic resonance blood pool agent in conjunction with breathheld three-dimensional magnetic resonance angiography. *Cardiovasc Intervent Radiol* (1999) 22: 321-325
256. Weissleder, R.:  
Target-specific superparamagnetic MR contrast agents.  
*Magn Reson Med* (1991) 22: 209-212
257. Weissleder, R., Cheng, H.C., Bogdanova, A., Bogdanov, A.J.:  
Magnetically labeled cells can be detected by MR imaging.  
*J Magn Reson Imaging* (1997) 7: 258-263
258. Weissleder, R., Elizondo, G., Josephson, L., Compton, C.C., Fretz, C.J., Stark, D.D., Ferrucci, J.T.:  
Experimental lymph node metastases: enhanced detection with MR lymphography. *Radiology* (1989) 171: 835-839
259. Weissleder, R., Elizondo, G., Wittenberg, J., Lee, A.S., Josephson, L., Brady, T.J.:  
Ultrasmall superparamagnetic iron oxide: an intravenous contrast agent for assessing lymph nodes with MR imaging. *Radiology* (1990) 175: 494-498
260. Weissleder, R., Elizondo, G., Wittenberg, J., Rabito, C.A., Bengele, H.H., Josephson, L.:  
Ultrasmall superparamagnetic iron oxide: characterization of a new class of contrast agents for MR imaging. *Radiology* (1990) 175: 489-493
261. Weissleder, R., Lee, A.S., Fischman, A.J., Reimer, P., Shen, T., Wilkinson, R., Callahan, R.J., Brady, T.J.:  
Polyclonal human immunoglobulin G labeled with polymeric iron oxide: antibody MR imaging. *Radiology* (1991) 181: 245-249
262. Weissleder, R., Lee, A.S., Khaw, B.A., Shen, T., Brady, T.J.:  
Antimyosin-labeled monocrystalline iron oxide allows detection of myocardial infarct: MR antibody imaging. *Radiology* (1992) 182: 381-385
263. Weissleder, R., Moore, A., Mahmood, U., Bhorade, R., Benveniste, H., Chiocca, E.A., Basilion, J.P.:  
In vivo magnetic resonance imaging of transgene expression. *Nat Med* (2000) 6: 351-354
264. Weissleder, R., Reimer, P., Lee, A.S., Wittenberg, J., Brady, T.J.:  
MR receptor imaging: ultrasmall iron oxide particles targeted to asialoglycoprotein receptors. *Am J Roentgenol* (1990) 155: 1161-1167
265. Weissleder, R., Stark, D.D., Engelstad, B.L., Bacon, B.R., Compton, C.C., White, D.L., Jacobs, P., Lewis, J.:  
Superparamagnetic iron oxide: pharmacokinetics and toxicity. *Am J Roentgenol* (1989) 152: 167-173
266. Wesbey, G.E., Engelstad, B.L., Brasch, R.C.:  
Paramagnetic pharmaceuticals for magnetic resonance imaging. *Physiol Chem Phys Med NMR* (1984) 16: 145-155
267. Widder, D.J., Greif, W.L., Widder, K.J., Edelman, R.R., Brady, T.J.:  
Magnetite albumin microspheres: a new MR contrast material. *Am J Roentgenol* (1987) 148: 399-404

268. Wielopolski, P.A., van Geuns, R.J., de Feyter, P.J., Oudkerk, M.:  
Breath-hold coronary MR angiography with volume-targeted imaging.  
*Radiology* (1998) 209: 209-219
269. Wiener, E.C., Brechbiel, M.W., Brothers, H., Magin, R.L., Gansow, O.A., Tomalia, D.A.,  
Lauterbur, P.C.:  
Dendrimer-based metal chelates: a new class of magnetic resonance imaging contrast  
agents. *Magn Reson Med* (1994) 31: 1-8
270. Wilhelm, C., Billotey, C., Roger, J., Pons, J.N., Bacri, J.C., Gazeau, F.:  
Intracellular uptake of anionic superparamagnetic nanoparticles as a function of their surface  
coating. *Biomaterials* (2003) 24: 1001-1011
271. Willmann, J.K., Wildermuth, S., Pfammatter, T., Roos, J.E., Seifert, B., Hilfiker, P.R.,  
Marincek, B., Weishaupt, D.:  
Aortoiliac and renal arteries: prospective intraindividual comparison of contrast-enhanced  
three-dimensional MR angiography and multi-detector row CT angiography.  
*Radiology* (2003) 226: 798-811
272. Yeh, T.C., Zhang, W., Ildstad, S.T., Ho, C.:  
Intracellular labeling of T-cells with superparamagnetic contrast agents.  
*Magn Reson Med* (1993) 30: 617-625
273. Yeh, T.C., Zhang, W., Ildstad, S.T., Ho, C.:  
In vivo dynamic MRI tracking of rat T-cells labeled with superparamagnetic iron-oxide  
particles. *Magn Reson Med* (1995) 33: 200-208
274. Zhang, S., Cai, J., Luo, Y., Han, C., Polissar, N.L., Hatsukami, T.S., Yuan, C.:  
Measurement of carotid wall volume and maximum area with contrast-enhanced 3D MR  
imaging: initial observations. *Radiology* (2003) 228: 200-205
275. Zhang, Y., Kohler, N., Zhang, M.:  
Intracellular uptake of poly(ethylene glycol) and folic acid modified magnetite nanoparticles.  
*Mater Res Soc Symp Proc* (2002) 676: Y981-Y985
276. Zheng, J., Bae, K.T., Woodard, P.K., Haacke, E.M., Li, D.:  
Efficacy of slow infusion of gadolinium contrast agent in three- dimensional MR coronary  
artery imaging. *J Magn Reson Imaging* (1999) 10: 800-805
277. Zheng, J., Li, D., Bae, K.T., Woodard, P.K., Haacke, E.M.:  
Three-dimensional gadolinium-enhanced coronary magnetic resonance angiography: initial  
experience. *J Cardiovasc Magn Reson* (1999) 1: 33-42
278. Zheng, J., Li, D., Cavagna, F.M., Harris, K., Klocke, F.J., Maggioni, F., Carr, J.,  
Simonetti, O., Laub, G., Finn, J.P.:  
Contrast-enhanced coronary MR angiography: relationship between coronary artery  
delineation and blood T1. *J Magn Reson Imaging* (2001) 14: 348-354