

7 LITERATURHINWEISE

- Achazi, R.K. (1989): Das Biotransformationssystem der Fische als Schadstoffindikator. Z. Angew. Zool., 76, 1-25.
- Achazi, R.K. (1995): The inhibition of microsomal EROD activity by a high-molecular protein. Z. Angew. Zool., 1, 141-153.
- Addison, R.F., J.F. Payne (1986): Assessment of hepatic mixed function oxidase induction in winter flounder (*Pseudopleuronectes americanus*) as a marine petroleum pollution monitoring technique, with an appendix describing practical field measurements of MFO activity. Can. Tech. Rept. Fish. Aquat. Sci., 1505, 22-37.
- Addison, R.F., A.J. Edwards (1988): Hepatic microsomal mono-oxygenase activity in flounder *Platichthys flesus* from polluted sites in Langesundfjord and from mesocosms experimentally dosed with diesel oil and copper. Mar. Ecol. Prog. Ser. 46, 51-54.
- Allen, Y., P. Matthiessen, A.P. Scott, S. Haworth, S. Feist, J.E. Thain (1999): The extent of oestrogenic contamination in the UK estuarine and marine environments – further surveys of flounder. Sci. Tot. Environ., 233, 5-20.
- AMAP (1999): Report of the workshop on combined effects of contaminants in the marine environment. AMAP Report 99:1, AMAP Oslo.
- Anders, K., H. Möller (1991): Epidemiologische Untersuchungen im Wattenmeer. Ber. Inst. Meeresk., 207.
- Anderson, M.J., M.R. Miller, D.E. Hinton (1996): In vitro modulation of 17-β-estradiol-induced vitellogenin synthesis: Effects of cytochrome P4501A1 inducing compounds on rainbow trout (*Oncorhynchus mykiss*) liver cells. Aquat. Toxicol., 34, 327-350.
- Andersson, T., U. Koivussari (1985): Influence of environmental temperature on the induction of xenobiotic metabolism by β-naphthoflavone in rainbow trout, *Salmo gairdneri*. Toxicol. Appl. Pharmacol., 80, 43-50.
- Ankley, G.T., R.E. Reinert, A.E. Wade, R.A. White (1985): Temperature compensation in the hepatic mixed function system of bluegill. Comp. Biochem. Physiol., 81C, 125-129.
- Ankley, G.T., V.S. Blazer, R.E. Reinert, M. Agosin (1986): Effects of Aroclor 1254 on cytochrome P-450-dependent monooxygenase, glutathione S-transferase, and UDP-glucuronyltransferase activities in channel catfish liver. Aquat. Toxicol., 9, 91-103.
- Anonymous (1993a): Monitoring and surveillance of biological contaminants and diseases in the aquatic environment 1990. Aquat. Environ. Monit. Rep., MAFF Direct. Fish. Res., Lowestoft, 35.
- Anonymous (1993b): Analysis and interpretation of benthic community data at sewage-sludge disposal sites. Aquat. Environ. Monit. Rep., MAFF Direct. Fish. Res., Lowestoft, 37.
- ARGE ELBE (1990): Nährstoffstudie der Elbe von Schnackenburg bis zur See. Arbeitsgemeinschaft für die Reinhalterung der Elbe (Hrsg.), Hamburg.
- ARGE ELBE (1992): Wassergütedaten der Elbe von Schnackenburg bis zur See – Zahlentafel 1991. Arbeitsgemeinschaft für die Reinhalterung der Elbe, Wassergütestelle Elbe (Hrsg.), Hamburg.
- ARGE ELBE (1993): Wassergütedaten der Elbe von Schmilka bis zur See – Zahlentafel 1993. Arbeitsgemeinschaft für die Reinhalterung der Elbe (Hrsg.), Hamburg.
- Arukwe, A., L. Förlin, A. Goksøyr (1997): Xenobiotic and steroid biotransformation enzymes in Atlantic salmon (*Salmo salar*) treated with an estrogenic compound, 4-nonylphenol. Environm. Toxicol. Chem., 16, 2576-2583.

- Becker, G.A. (1990): Die Nordsee als physikalisches System. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. von Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 11-27.
- Besselink, H.T., S. van Beusekom, E. Roex, A.D. Vethaak, J.H. Koeman, A. Brouwer (1996): Low hepatic 7-ethoxyresorufin-O-deethylase (EROD) activity and minor alterations in retinoid and thyroid hormone levels in flounder (*Platichthys flesus*) exposed to the polychlorinated biphenyl (PCB) mixture, clophen A50. Environ. Pollut., 92, 267-274.
- Besselink, H.T., M.S. Denison, M.E. Hahn, S.I. Karchner, A.D. Vethaak, J.H. Koeman, A. Brouwer (1998): Low inducibility of CYP1A activity by polychlorinated biphenyls (PCBs) in flounder (*Platichthys flesus*): Characterization of the Ah receptor and the role of CYP1A inhibition. Toxicological Sciences, 43, 161-171.
- Beyer, J., M. Sandvik, K. Hylland, E. Fjeld, E. Egaas, E. Aas, J.U. Skare, A. Goksøyr (1996): Contaminant accumulation and biomarker responses in flounder (*Platichthys flesus* L.) and Atlantic cod (*Gadus morhua* L.) exposed by caging to polluted sediments in Sørforden, Norway. Aquat. Toxicol., 36, 75-98.
- BgVV (2001): Tätigkeitsbericht 2000/2001, Fachbereich 8. Bundesinstitut für gesundheitlichen Verbraucherschutz und Veterinärmedizin (Hrsg.). Bericht nur als online-Version verfügbar: www.bgvv.de
- BLMP (2002): Meeresumwelt 1997 – 1998. Bundesamt für Seeschifffahrt und Hydrographie (BSH) (Hrsg.), Hamburg und Rostock, Online-Publikation: www.bsh.de/Meeresumwelt/BLMP.
- Bohl, H. (1957): Die Biologie der Kliesche (*Limanda limanda* L.) in der Nordsee. Ber. dt. wiss. Kommn. Meeresforsch., 15, 1-57.
- Boon, J.P., M.B. van Zantvoort, M.J.M.A. Govaert (1985): Organochlorines in benthic polychaetes (*Nephtys* spp.) and sediments from the southern North Sea. Identification of individual PCB components. Neth. J. Sea Res., 19, 93-109.
- Borchardt, T., S. Burchert, H. Hablitzel, L. Karbe, R. Zeitner (1988): Trace metal concentrations in mussels: comparison between estuarine, coastal and offshore regions in the southeastern North Sea from 1983 to 1986. Mar. Ecol. Prog. Ser., 42, 17-31.
- Bortz, J., G.A. Lienert, K. Boehnke (1990): Verteilungsfreie Methoden in der Biostatistik. Springer Verlag Berlin, Heidelberg, New York.
- Bradford, M.M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem., 72, 248-254.
- Brauer, F., H.-J. Pluta, E. Trölsch (1994): Untersuchung zur Biotransformation in der Fischleber (Entgiftungsaktivität der mischfunktionellen Oxidasen, MFO). Abschlussbericht Teilvorhaben 3 im FuE-Vorhaben „Fischkrankheiten in der Nordsee“, UFOPLAN-Nr. 10603900/03.
- Broeg, K., S. Zander, A. Diamant, W. Körting, G. Krüner, I. Paperna, H. von Westernhagen (1999): The use of fish metabolic, pathological and parasitological indices in pollution monitoring. I. North Sea. Helgol. Mar. Res., 53, 171-194.
- Brown, C.A., S. Black (1989): Membrane topology of mammalian cytochromes P-450 from liver endoplasmatic reticulum. J. Biol. Chem., 15, 4442-4449.
- Brüschiweiler, B.J., F.E. Würgler, K. Fent (1996): Inhibitory effects of heavy metals on cytochrome P4501A induction in permanent fish hepatoma cells. Arch. Environ. Contam. Toxicol., 31, 475-482.
- BSH (2002): Karten der Meeresoberflächentemperatur. Bundesamt für Seeschifffahrt und Hydrographie; regelmäßig aktualisierte Online-Darstellungen auf der homepage: www.BSH.de.
- Bucheli, T.D., K. Fent (1995): Induction of cytochrome P450 as a biomarker for environmental contamination in aquatic ecosystems. Critical Reviews in Environmental Science and Technology, 25, 201-268.

- Bückmann, A. (1929): Die Methodik fischereibiologischer Untersuchungen an Meeresfischen. In: E. Abderhalden (Hrsg.): Handbuch der biologischen Arbeitsmethoden, Abteilung IX, Teil 6, Verlag Urban u. Schwarzenberg Berlin, Wien.
- Buhler, D.R., J.-L. Wang-Buhler (1998): Rainbow trout cytochrome P450s: purification, molecular aspects, metabolic activity, induction and role in environmental monitoring. Comp. Biochem. Physiol., 121C, 107-137.
- Burke, M.D., R.T. Mayer (1974): Ethoxyresorufin: direct fluorimetric assay of a microsomal-O-dealkylation which is preferentially inducible by 3-methylcholanthrene. Drug Metab. Dispos., 2, 583-588.
- Büther, H. (1988): Distribution of chlorinated organic compounds in livers of dab (*Limanda limanda*) of the southern and central North Sea. Mitt. Geol.-Paläont. Inst. Univ. Hamburg, Heft 65, 497-541.
- Büther, H. (1990): Chlorierte organische Verbindungen in Fischen. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 274-281.
- Cameron, P., J. Berg, V. Dethlefsen, H. von Westernhagen (1992): Developmental defects in pelagic embryos of several flatfish species in the southern North Sea. Neth. J. Sea Res., 29, 239-256.
- Campos, W.L., M. Kloppmann, H. von Westernhagen (1994): Inferences from the horizontal distribution of dab *Limanda limanda* (L.) and flounder *Platichthys flesus* (L.) larvae in the southeastern North Sea. Neth. J. Sea Res., 32, 277-286.
- Carr, R.S., J.M. Neff (1988): Influence of prior exposure to xenobiotics on the metabolism and distribution of polychlorinated biphenyls and phenanthrene in winter flounder, *Pseudopleuronectes americanus*. Mar. Environ. Res., 24, 73-77.
- Claussen, T. (1988): Levels and spatial distribution of trace metals in dabs (*Limanda limanda*) of the southern North Sea. Mitt. Geol.-Paläont. Inst. Univ. Hamburg, Heft 65, 467-496.
- Collier, T.K., B.F. Anulacion, J.E. Stein, A. Goksøyr (1995): A field evaluation of cytochrome P4501A as a biomarker of contaminant exposure in three species of flatfish. Environm. Toxicol. Chem., Vol. 14, 143-152.
- Collier, T.K., B.F. Anulacion, B.D. Bill (1998): Hepatic CYP1A in winter flounder (*Pleuronectes americanus*) along the northeast coast: Results from the national benthic surveillance project. Mar. Poll. Bull., 37, 86-91.
- Cooreman, K. (2001): The influence of natural factors on EROD in dab. In: ICES (ed.): Report of the Working group on biological effects of contaminants. ICES CM 2001/E:03. Copenhagen, Denmark, 31-32.
- Daan, N., P.J. Bromley, J.R.G. Hislop, N.A. Nielsen (1990): Ecology of North Sea fishes. Neth. J. Sea Res., 26, 343-386.
- Daan, R., M. Mulder (1996): On the short-term and long-term impact of drilling activities in Dutch sector of the North Sea. ICES Journal of Marine Science, 53, 1036-1044.
- Damm, U., T. Lang, A. Rijnsdorp (1991): Movements of dab in the German Bight: Results of german and dutch tagging experiments in 1988, 1989. ICES CM/E:22, Copenhagen, Denmark.
- Davies, J.M., R. Hardy, A.D. McIntyre (1981): Environmental effects of North Sea oil operations. Mar. Poll. Bull., 12, No 12, 412-416.
- Davies, J.M., J.M. Addy, R.A. Blackman, J.R. Blanchard, J.E. Ferbrache, P.C. Moore, H.J. Somerville, A. Whitehead, T. Wilkinson (1984a): Environmental effects of the use of oil-based drilling muds in the North Sea. Mar. Poll. Bull., 15, No 10, 363-370.
- Davies, J.M., J.S. Bell (1984b): A comparison of the levels of hepatic aryl hydrocarbons hydroxylase in fish caught close to and distant from North Sea oil fields. Mar. Environ. Res., 14, 23-45.

- De Bruijn, J., W. Seinen, J. Hermens (1993): Biotransformation of organophosphorus compounds by rainbow trout (*Oncorhynchus mykiss*) liver in relation to bioconcentration. Environm. Toxicol. Chem., 12, 1041-1050.
- Dethlefsen, V. (1991): Probleme der Gewässerverschmutzung – Hohe Befallsraten der Kliesche (*Limanda limanda*) mit Flossenfäule vor der Humber-Mündung. Infn. Fischw., 38 (2), 6771.
- Dethlefsen, V., E. Huschenbeth (1986): Regional differences in organochlorine residues in livers of dab (*Limanda limanda*) and plaice (*Pleuronectes platessa*) of the southern North Sea. Arch. FischWiss., 37, 25-42.
- Dethlefsen, V., B. Watermann, M. Hoppenheit (1987): Diseases of North Sea dab (*Limanda limanda* L.) in relation to biological and chemical parameters. Arch. FischWiss., 37, 107-237.
- Dethlefsen, V., K. Söffker (1992): FuE-Vorhaben „Fischkrankheiten in der Nordsee“, Projektbericht Teilvorhaben 5 “Epidemiologie der Kliesche und Untersuchungen zur Organochlorkontamination“. UFOPLAN-Nr. 10603900/05.
- Diercking, R., L. Wehrmann (1991): Artenschutzprogramm Fische und Rundmäuler in Hamburg. Naturschutz und Landschaftspflege in Hamburg – Schriftenreihe der Umweltbehörde Nr. 38.
- Dornheim, H., G. Wegner (1992): Zur Lage der Jungfischbestände in der Nordsee. Infn. Fischw., 39, 60-70.
- Duncker, G. (1960): Die Fische der Nordmark. Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg N.F. Bd. III, Supplement (Hrsg.: E.M. Thiel), Kommisionsverlag Cram, De Gruyter und Co. Hamburg.
- Edwards, A.J., R.F. Addison, D.E. Willis, K.W. Renton (1988): Seasonal variation of hepatic mixed function oxidases in winter flounder (*Pseudopleuronectes americanus*). Mar. Environ. Res., 26, 299-309.
- Eggens, M.L., F. Galgani, J. Klungsøyr, J. Everts (1992): Hepatic EROD activity in dab, *Limanda limanda*, in the German Bight using an improved plate-reader method. In: Stebbing, A.R.D., V. Dethlefsen, M. Carr (eds.): Biological effects of contaminants in the North Sea - Results of the ICES/IOC Bremerhaven Workshop. Inter-Research, Amelinghausen. Mar. Ecol. Prog. Ser., 91, 71-75.
- Eggens, M.L., A. Bergman, D. Vethaak (1995a): Seasonal variation of hepatic EROD activity in flounder (*Platichthys flesus*) in the dutch Wadden Sea. Mar. Environ. Res., 39, 231-234.
- Eggens, M.L., A. Bergman, D. Vethaak, M. van der Weiden, M. Celander, J.P. Boon (1995b): Cytochrome P4501A indices as biomarkers of contaminant exposure: results of a field study with plaice (*Pleuronectes platessa*) and flounder (*Platichthys flesus*) from the southern North Sea. Aquat. Toxicol., 32, 211-225.
- Eggens, M.L., A. Opperhuizen, J.P. Boon (1996a): Temporal variation of CYP1A indices, PCB and 1-OH pyrene concentration in flounder, *Platichthys flesus*, from the Dutch Wadden Sea. Chemosphere, 33, 1579-1596.
- Eggens, M.L., A.D. Vethaak, M.J. Leaver, G.J.M.J. Horbach, J.P. Boon, W. Seinen (1996b): Differences in CYP1A response between flounder (*Platichthys flesus*) and plaice (*Pleuronectes platessa*) after long-term exposure to harbour dredged spoil in a mesocosm study. Chemosphere, 32, 1357-1380.
- Elcombe, C. R., R.B Franklin, J.J. Lech (1979): Induction of hepatic microsomal enzymes in rainbow trout. In: Khan, M.A.Q., J.J. Lech (eds.): Pesticide and Xenobiotic Metabolism in Aquatic Organisms. American Chemical Society, 319-337.
- Elskus, A.A., R. Pruell, J.J. Stegeman (1992): Endogenously-mediated, pretranslational suppression of cytochrome P4501A in PCB-contaminated flounder. Mar. Environ. Res., 34, 97-101.
- Elskus, A.A., J.J. Stegeman, J.W. Gooch (1994): Polychlorinated biphenyl congener distributions in winter flounder as related to gender, spawning site, and congener metabolism. Environ. Sci. Technol., 28, 401-407.

- EU-Kommission (2001): Mitteilung der Kommission zum sechsten Aktionsprogramm der Europäischen Gemeinschaft für die Umwelt – Umwelt 2010: Unsere Zukunft in unserer Hand. Kommission der Europäischen Gemeinschaften 31, 2001/0029 (COD).
- Fent, K. (1998): Ökotoxikologie. Thieme Verlag Stuttgart, New York.
- Fent, K., B.R. Woodin, J.J. Stegeman (1998): Effects of triphenyltin and other organotins on hepatic monooxygenase system in fish. *Comp. Biochem. Physiol.*, 121C, 277-288.
- Fiedler, M. (1991): Die Bedeutung von Makrozoobenthos und Zooplankton der Unterelbe als Fischnahrung. *Ber. Inst. Meeresk.* Kiel Nr. 204.
- Fock, H., K. Ricklefs (1996): Die Eider – Veränderungen seit dem Mittelalter. In: Lozán, J.L., H. Kausch (Hrsg.): Warnsignale aus Flüssen und Ästuaren. Parey Verlag Berlin, 39-43.
- Förlin, L. (1980): Effects of Clophen A50, 3-Methylcholanthrene, Pregnanolone-16 α -carbonitrile, and phenobarbital on the hepatic microsomal cytochrome P-450-dependent monooxygenase system in rainbow trout, *Salmo gairdneri*, of different age and sex. *Toxicol. Appl. Pharmacol.*, 54, 420-430.
- Förlin, L., T. Andersson, U. Koivusaari, T. Hansson (1984): Influence of biological and environmental factors on hepatic steroid and xenobiotic metabolism in fish: interaction with PCB and β -naphthoflavone. *Mar. Environ. Res.*, 14, 47-58.
- Förstner, U., W. Ahlf (1992): Metalle und ihr chemisch-physikalisches Verhalten in Ästuarien. In: Kausch, H. (Hrsg.): Die Unterelbe – natürlicher Zustand und Veränderungen durch den Menschen. *Ber. Zentr. Meeres- u. Klimaforsch.* Hamburg, 19, 61-72.
- Gaumert, T., (1992): Entwicklung der Schadstoffgehalte in frischen, schwebstoffbürtigen Sedimenten der Elbe bei Schnackenburg, 1984 – 1991. Arbeitgemeinschaft für die Reinhaltung der Elbe (Hrsg.), Hamburg.
- George, S., P. Young, M. Leaver, D. Clarke (1990): Activities of pollutant metabolising and detoxication systems in the liver of the plaice, *Pleuronectes platessa*: Sex and seasonal variations in non-induced fish. *Comp. Biochem. Physiol.*, 96C, No.1, 185-192.
- Goedecke, E., J. Smed, G. Tomczak (1967): Monatskarten des Salzgehaltes der Nordsee, dargestellt für verschiedene Tiefenhorizonte. *Dt. hydrogr. Z., Erg.-H. B*, Nr. 9.
- Goerke, H., K. Weber (1998): The bioaccumulation pattern of organochlorine residues in *Lanice conchilega* (Polychaeta) and its geographical variation between the English Channel and the German Bight. *Chemosphere*, 37, 1283-1298.
- Goksøyr, A., A.-M. Husøy, H.E. Larsen, J. Klungsøyr, S. Wilhelmsen, A. Maage, E. Brevik, T. Andersson, M. Celander, M. Pesonen, L. Förlin (1991): Environmental contaminants and biochemical responses in flatfish from the Hvaler Archipelago in Norway. *Arch. Environ. Contam. Toxicol.*, 21, 486-496.
- Goksøyr, A., L. Förlin (1992a): The cytochrome P-450 system in fish, aquatic toxicology and environmental monitoring. *Aquat. Toxicol.*, 22, 287-312.
- Goksøyr, A., H.E. Larsen, S. Blom, L. Förlin (1992b): Detection of Cytochrome P450 1A1 in North Sea dab liver and kidney. In: Stebbing, A.R.D., V. Dethlefsen, M. Carr (eds.): Biological effects of contaminants in the North Sea – Results of the ICES/IOC Bremerhaven Workshop. Inter-Research, Amelinghausen. *Mar. Ecol. Prog. Ser.*, 91, 83-88.
- Goksøyr, A., J. Beyer, A.-M. Husøy, H.E. Larsen, K. Westrheim, S. Wilhelmsen, J. Klungsøyr (1994): Accumulation and effects of aromatic and chlorinated hydrocarbons in juvenile Atlantic cod (*Gadus morhua*) caged in a polluted fjord (Sørfjorden, Norway). *Aquat. Toxicol.*, 29, 21-35.
- Goksøyr, A. (1995): Use of cytochrome P450 1A (CYP1A) in fish as a biomarker of aquatic pollution. *Archives in Toxicology*, 17, 80-95.

- Goksøyr, A., J. Beyer, E. Egaas, B.E. Grøsvik, K. Hylland, M. Sandvik, J.U. Skare (1996): Biomarker responses in flounder (*Platichthys flesus*) and their use in pollution monitoring. Mar. Poll. Bull., 33, 36-45.
- Greenpeace (2002): Die Nordsee, das Erdöl und die Offshore-Industrie am Beispiel von Esso. Greenpeace e. V., Hamburg. Online-Publikation: www.greenpeace.org
- Grinwis, G.C.M., H.T. Besselink, E.J. van den Brandhof, A.S. Bulder, M.Y. Engelsma, R.V. Kuiper, P.W. Wester, M.A. Vaal, A.D. Vethaak, J.G. Vos (2000a): Toxicity of TCDD in European flounder (*Platichthys flesus*) with emphasis on histopathology and cytochrome P4501A induction in several organ systems. Aquat. Toxicol., 50, 387-401.
- Grinwis, G.C.M., A.D. Vethaak, P.W. Wester, J.G. Vos (2000b): Toxicology of environmental chemicals in the flounder (*Platichthys flesus*) with emphasis on the immune system: field, semi-field (mesocosm) and laboratory studies. Toxicology letters, 112/113, 289-301.
- Haarich, M., V. Dethlefsen, T. Lang (1998): Langzeitstudien zur Belastung von Klieschen in der Nordsee mit chlorierten Kohlenwasserstoffen. In: Jahresbericht 1998. Inst. f. Fischereiökologie (Hrsg.), 42-43.
- Haasch, M.L., R. Prince, P.J. Wejksnora, K.R. Cooper, J.L. Lech (1993): Caged and wild fish: Induction of hepatic cytochrome P-450 (CYP1A1) as an environmental biomonitor. Environm. Toxicol. Chem., 12, 885-895.
- Hall, J.A., C.L.J. Frid, M.E. Gill (1997): The response of estuarine fish and benthos to an increasing discharge of sewage effluent. Mar. Poll. Bull., 34, 527-535.
- Hansen, P.D., R.F. Addison, D.E. Willis (1983): Hepatic microsomal O-de-ethylases in cod (*Gadus morhua*): their induction by arochlor 1254 but not by arochlor 1016. Comp. Biochem. Physiol., 74C, No. 1, 173-175.
- Hansen, P.D., R.F. Addison (1990): The use of mixed function oxidases (MFO) to support biological effects monitoring in the sea. ICES CM/E:33, Copenhagen, Denmark.
- Haux, C., L. Förlin (1988): Biochemical methods for detecting effects of contaminants on fish. Ambio, 17, 376-380.
- Henderson, P.A. (1998): On the variation in dab *Limanda limanda* recruitment: a zoogeographic study. J. Sea Res., 40, 131-142.
- Hodson, P.V., P.J. Klopper-Sams, K.R. Munkittrick, W.L. Lockhart, D.A. Metner, P.L. Luxon, I.R. Smith, M.M. Gagnon, M. Servos, J.F. Payne (1991): Protocols for measuring mixed function oxygenases of fish liver. Can. Tech. Rep. Fish. Aquat. Sci., 1829.
- Hoogweg, P.H.A. (1992): Das Aktionsprogramm Nordsee und seine Auswirkungen auf den Schutz der großen Flüsse. Vortrag, 13. Arbeitstagung der internationalen Arbeitsgemeinschaft der Wasserwerke im Rheineinzugsgebiet (IAWR), Scheveningen 1991.
- Hühnerfuss, H., B. Pfaffenberger, B. Gehrcke, L. Karbe, W.A. König, O. Landgraff (1995): Stereochemical effects of PCBs in the marine environment: Seasonal variation of coplanar and atropisomeric PCBs in blue mussels (*Mytilus edulis* L.) of the German Bight. Mar. Poll. Bull., 30, 332-340.
- Huntenburg, S., A. Zahrte, K. Ricklefs, D. Stüben (1995): Untersuchungen zur Schwermetallkonzentration im Eider-Ästuar. Meyniana, 47, 21-43.
- Hupkes, R. (1990): Pollution of the North Sea imposed by West European rivers (1984-1987). International Centre of Water Studies, Report 90.03.
- Hylland, K., T. Nissen-Lie, P. G. Christensen, M. Sandvik (1998): Natural modulation of hepatic metallothionein and cytochrome P4501A in flounder, *Platichthys flesus* L., Mar. Environ. Res., 46, 51-55.
- ICES (1993): Report of the ICES Advisory Committee on marine pollution, 1992, ICES Coop. Res. Rep. 190 JMG 18/9/1-E, Copenhagen, Denmark.

- ICES (1999): Report of the Working group on biological effects of contaminants. ICES CM 1999/E:03, Copenhagen, Denmark.
- ICES (2000): Report of the Working group on biological effects of contaminants. ICES CM 2000/E:04, Copenhagen, Denmark.
- ICES (2001): Report of the Working group on biological effects of contaminants. ICES CM 2001/E:03, Copenhagen, Denmark.
- James, M.O., M.A.Q. Khan, J.R. Bend (1979): Hepatic microsomal mixed-function oxidase activities in several marine species common to coastal Florida. *Comp. Biochem. Physiol.*, 62C, 155-164.
- Jarvie, H. P., C. Neal, A. D. Tappin (1997): European land-based pollutant loads to the North Sea: an analysis of the Paris Commission data and review of monitoring strategies. *Sci. Total Environ.*, 194/195, 39-58.
- Jiminez, B.D., L.S. Burtis, G.H. Ezell, B.Z. Egan, N.E. Lee, J.J. Beauchamp, J.F. McCarthy (1988): The mixed function oxydase system of bluegill sunfish, *Lepomis macrochirus*: correlation of activities in experimental and wild fish. *Environm. Toxicol. Chem.*, 7, 623-634.
- Kammann, U., O. Landgraff, H. Steinhart (1993): Distribution of aromatic organochlorines in livers and reproductive organs of male and female dabs from the German Bight. *Mar. Poll. Bull.*, 26, 629-635.
- Kannan, N., S. Tanabe, R. Tatsukawa (1988): Toxic potential of non-ortho and mono-ortho coplanar PCBs in commercial PCB preparations 2,3,7,8-T₄CDD toxicity equivalence factors approach. *Bull. Environ. Contam. Toxicol.*, 41, 267-276.
- Karbe, L. (1990): Kontamination mit Schwermetallen. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 188-198.
- Kausch, H. (1996): Die Elbe – ein immer wieder veränderter Fluss. In: Lozán, J.L., H. Kausch (Hrsg.): Warnsignale aus Flüssen und Ästuaren. Parey Verlag Berlin, 43-52.
- Kellermann, H.-J., M. Vobach (1997): Entgiftungsstoffwechsel der Nordsee-Kliesche (*Limanda limanda*). *Inf. Fischwirtsch.*, 44, 13-17.
- Kelly, A.G., D. Campbell (1994): Persistent organochlorine contaminants in fish and shellfish from scottish waters. *Scottish Fisheries Research Report*, 54
- Kerstan, M. (1991): The importance of rivers as nursery grounds for 0- and 1-Group flounder (*Platichthys flesus* L.) in comparison to the wadden sea. *Neth. J. Sea Res.*, 27, 353-366.
- Khan, R.A., D. Barker, R. Hooper, E.M. Lee (1992): Effect of pulp and paper effluent on a marine fish, *Pseudopleuronectes americanus*. *Bull. Environ. Contam. Tox.*, 48, 449-456.
- Kirby, M.F., P. Matthiessen, P. Neall, T. Tylor, C.R. Allchin, C.A. Kelly, D.L. Maxwell, J.E. Thain (1999): Hepatic EROD activity in flounder (*Platichthys flesus* L.) as an indicator of contaminant exposure in English estuaries. *Mar. Poll. Bull.*, 38, 676-686.
- Kirby, M.F., M.A. Blackburn, J.E. Thain, M.J. Waldock (1998): Assessment of water quality in estuarine and coastal waters of England and Wales using a contaminant concentration technique. *Mar. Poll. Bull.*, 36, 631-642.
- Knickmeyer, R. (1990a): Zooplankton – Kontaminationen mit chlorierten organischen Verbindungen. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 148-153.
- Knickmeyer, R. (1990b): Kontamination mit chlorierten organischen Verbindungen. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 198-202.

- Köhler, A. (1993): Untersuchungen zur Entgiftungsfunktion (Lysosomensystem) und Histopathologie der Fischleber. Abschlussbericht Teilvorhaben 2 im FuE-Vorhaben „Fischkrankheiten in der Nordsee“, UFOPLAN-Nr. 10603900/02.
- Koivusaari, U. (1983): Thermal acclimatization of hepatic polysubstrate monooxygenase and UDP-glucuronosyl-transferase of mature rainbow trout (*Salmo gairdneri*). J. Exp. Zool., 227, 35-42.
- Koivussari, U. (1984): Xenobiotic biotransformation in rainbow trout, *Salmo gairdneri*, with special reference to seasons and reproduction. Publications of the university of Kuopio. Natural sciences series original reports, 5/1984.
- Koivusaari, U., M. Harri, O. Hanninen (1981): Seasonal variation of hepatic in female and male rainbow trout (*Salmo gairdneri*). Comp. Biochem. Physiol., 70C, 149-157.
- Koivussari, U., T. Andersson (1984). Partial temperature compensation of hepatic biotransformation enzymes in juvenile rainbow trout (*Salmo gairdneri*) during the warming of water in spring. Comp. Biochem. Physiol., 78B, 223-226.
- Koopmann, G., H. Heinrich , D. Voppel, N.P. Rühl (1993): Transport, Umsatz und Variabilität von Schad- und Nährstoffen (TUVAS) in der Deutschen Bucht 1990 – 1992, Teile I und II; Bundesamt für Seeschifffahrt und Hydrographie (Hrsg.), Hamburg.
- Kröncke, I., R. Knust (1995): The Dogger Bank: a special ecological region in central North Sea. Helgoländer Meeresunters., 49, 335-353.
- Krüner, G., D. Janssen, H. v. Westernhagen (1996): Wissenschaftliche Grundlagen sowie Beschaffung und Bewertung von Daten für das Biologische Effektmonitoring der Nordsee. Ber. Biol. Anst. Helgoland Nr. 11.
- Krüner, G., H. v. Westernhagen (1999): Sources of measurement error in assays of EROD activity of fish for biological effects monitoring. Helgol. Mar. Res., 53, 250-256.
- Landwüst, C. von, K. Anders, U. Ballin, F. Brauer, P. Cameron, J. Cordt, V. Dethlefsen, S. Holst, A. Köhler-Günther, R. Kruse, K. Lee, H. Möller, M. Momme, N. von Neuhoff, H.-J. Pluta, W. Scharenberg, K. Söffker, N. Weis (1996): Fischkrankheiten in der Nordsee. Umweltbundesamt (Hrsg.), Texte 57/96.
- Lange, U. (1996): Natürliche Variabilität des schadstoffinduzierbaren Cytochroms P450 1A in der Leber von Klieschen (*Limanda limanda* [L.]) aus der Nordsee. Dissertation 1996. Berichte Zentr. Meeres- u. Klimaforsch. Hamburg Reihe E, Nr. 10.
- Lange, U., A. Goksøyr, D. Siebers, L. Karbe (1999): Cytochrome P450 1A-dependent enzyme activities in the liver of dab (*Limanda limanda*): kinetics, seasonal changes and detection limits. Comp. Biochem. Physiol., 123B, 361-371.
- Larsen, H.E., M. Celander, A. Goksøyr (1992): The cytochrome P450 system of Atlantic salmon (*Salmo salar*): II. Variations in hepatic catalytic activities and isozyme patterns during an annual reproductive cycle. Fish Physiol. Biochem., 10 (4), 291-301.
- Law, R.J., V.J. Dawes, R.J. Woodhead, P. Matthiessen (1997): Polycyclic aromatic hydrocarbons (PAH) in seawater around England and Wales. Mar. Poll. Bull., 34, 306-322.
- Leatherland, T.M. (1987): The estuary and Firth of Forth, Scotland: uses and aims. Proc. Royal Society Edinburgh, 93B, 285-297.
- Lech, J.J., J.R. Bend (1980): Relationship between biotransformation and the toxicity and fate of xenobiotic chemicals in fish. Environmental Health Perspectives, 34, 115-131.
- Lee, C.K.C. (1972): The biology and population dynamics of the common dab *Limanda limanda* (L.) in the North Sea. Dissertation, University of East Anglia.
- Lee, K., W. Scharenberg, N. Weis (1993): FuE-Vorhaben „Fischkrankheiten in der Nordsee“, Abschlussbericht Teilvorhaben 1b “Untersuchungen zur Organochlorkontamination“. UFOPLAN-Nr. 10603900/1b.

- Lemaire-Gony, S., P. Lemaire (1992): Interactive effects of cadmium and benzo(a)pyrene on cellular structure and biotransformation enzymes of the liver of the European eel *Anguilla anguilla*. Aquat. Toxicol., 22, 145-160.
- Levine, S.L., J.T. Oris, T.E. Wissing (1994): Comparison of P-4501A1 monooxygenase induction in gizzard shad (*Dorosoma cepedianum*) following intraperitoneal injection or continuous waterborne-exposure with benzo[a]pyrene: Temporal and dose-dependent studies. Aquat. Toxicol., 30, 61-75.
- Lienert, G.A. (1973): Verteilungsfreie Methoden der Biostatistik – Band I. Verlag Anton Hain Meisenheim am Glan.
- Lindström-Seppä, P., O. Hanninen, R. Hudd (1985): Biomonitoring of the petroleum spill in Vassa Archipelago, Finland, by biotransformation in fish. In: Vereczkey, L., K. Magyar (eds.): Cytochrome P-450: Biochemistry, Biophysics and Induction. Akadmiai, Kiado, Budapest, 357-360.
- Lindström-Seppä, P., A. Oikari (1990): Biotransformation and other toxicological and physiological responses in rainbow trout (*Salmo gairdneri* Richardson) caged in a lake receiving effluents of pulp and paper industry. Aquat. Toxicol., 16, 187-204.
- Lindström-Seppä, P., P.J. Korytko, M.E. Hahn, J.J. Stegeman (1994): Uptake of waterborn 3,3',4,4'-tetrachlorobiphenyl and organ and cell-specific induction of cytochrome P4501A in adult and larval fathead minnow *Pimephales promelas*. Aquat. Toxicol., 28, 147-167.
- Lindström-Seppä, P., J.J. Stegeman (1995): Sex differences in cytochrome P4501A induction by environmental exposure and β-Naphthoflavone in liver and extrahepatic organs of recrudescent Winter flounder. Mar. Environ. Res., 39, 219-223.
- Lohse, J. (1990): Chlorierte organische Verbindungen in Wasser und Sediment. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 75-85.
- Lohse, J., S. Winteler, K. Schweer, R. Leidecker (1995): Die Nordsee – Das Endlager der Chlorchemie. Greenpeace e. V. (Hrsg.), Hamburg.
- Loizeau, V., A. Menesguen (1993): A steady-state model of PCB accumulation in dab food web. Oceanologica Acta, 16, 633-639.
- Lowry, O. H., N.J. Rosebrough, A.L. Farr, R.J. Randall (1951): Protein measurement with folin phenol reagent. J. Biol. Chem., 193, 265-275.
- Lozán, J.L. (1989): Autökologische Untersuchung über die Kliesche (*Limanda limanda*, Linnaeus 1758) der Nordsee. Veröff. Inst. Küst.- u. Binnenfisch., Hamburg, 101, 1-199.
- Lozán, J.L. (1992): Sexual differences in food intake, digestive tract size, and growth performance of the dab *Limanda limanda* L. Neth. J. Sea Res., 29, 223-227.
- Lozán, J.L., W. Hickel, K. Reise, K. Ricklefs (1996): Wechselwirkungen zwischen Fluss und Meer. In: Lozán, J.L., H. Kausch (Hrsg.): Warnsignale aus Flüssen und Ästuaren. Parey Verlag Berlin, 75-85.
- Lye, C.M., C.L.J. Frid, M.E. Gill, D. McCormick (1997): Abnormalities in the reproductive health of flounder *Platichthys flesus* exposed to effluent from a sewage treatment works. Mar. Poll. Bull., 34, 34-41.
- Lyons, B.P., C. Stewart, M.F. Kirby (1999): The detection of biomarkers of genotoxin exposure in the european flounder *Platichthys flesus* collected from the river Tyne estuary. Mutat. Res., 446, 111-119.
- Mansuy, D. (1998): The great diversity of reactions catalyzed by cytochromes P450. Comp. Biochem. Physiol., 121C, 5-14.

- Matthiessen, P., Y.T. Allen, C.R. Allchin, S.W. Feist, M.F. Kirby, R.J. Law, A.P. Scott, J.E. Thain, K.V. Thomas (1998a): Oestrogenic endocrine disruption in flounder (*Platichthys flesus* L.) from United Kingdom estuarine and marine waters. Centre for Environment, Fisheries and Aquaculture Sciense, Lowestoft, UK, Sci. Ser., Tech. Rep., 107.
- Matthiessen, P., S. Bifield, F. Jarrett, M.F. Kirby, R.J. Law, W.R. McMinn, D.A. Sheahan, J.E. Thain, G.F. Whale (1998b): An assessment of sediment toxicity in the river Tyne estuary, UK by means of Bioassays. Mar. Environ. Res., 45, 1-15.
- Mehrtens, G., F. Latus (1999): Mixed function oxidase dependent biotransformation of polychlorinated biphenyls by different species of fish from the North Sea. Chemosphere, 38, 2995-3002.
- Millward, G.E., H.J. Kitts, L. Ebdon, J.I. Allen, A.W. Morris (1997): Arsenic in the Thames Plume, UK. Mar. Environ. Res., 44, 51-67.
- Möller, H. (1984): Daten zur Biologie der Elbfische. Verlag Möller, Kiel.
- Möller, H., K. Anders, M. Fiedler, G. Costa, B. Dieckwisch, S. Klatt, H. Lüchtenberg, C. Pohl, K. Rönnau, S. Schröder, M. Voigt (1988): Fischbestände und Fischkrankheiten in der Unterelbe 1984-1986. Verlag Möller, Kiel.
- Monosson, E., J.J. Stegeman (1991): Cytochrome P450E (P450IA) induction and inhibition in winter flounder by 3,3',4,4'-tetrachlorobiphenyl: comparison of response in fish from Georges bank and Narragansett bay. Environm. Toxicol. Chem., 10, 765-774.
- Morris, A.W. (1988): The estuaries of the Humber and the Thames. In: W. Salomons, B.L. Bayne, E.K. Duursma, U. Förstner (Hrsg.): The pollution of the North Sea – an assessment. Springer Verlag Berlin, 213-224.
- Müller, G. (1996): Schwermetalle und organische Schadstoffe in den Flusssedimenten. In: Lozán, J.L., H. Kausch (Hrsg.): Warnsignale aus Flüssen und Ästuaren. Parey Verlag Berlin, 113-124.
- Müller-Navarra, S.H., E. Mittelstaedt (1987): Schadstoffausbreitung und Schadstoffbelastung in der Nordsee – eine Modellstudie. Dt. hydrogr. Z., Erg.-H. B, Nr. 18.
- Muus, B.J., P. Dahlström (1991): Meeresfische, BLV Verlag München, Wien, Zürich.
- Neal, C., W.A. House, B.A. Whitton, G.J.L. Leeks (1998): Foreword to special issue: Water quality and biology of UK rivers entering in the North Sea: The land ocean interaction study LOIS and associated work. Sci. Tot. Environ., 210/211, 1-4.
- Neunaber, R., R.K. Achazi (1999): A novel, endogenous inhibitor of 7-ethoxyresorufin-O-deethylase activity isolated from liver cytosolic fractions of bream (*Abramis brama* L.). Comp. Biochem. Physiol., Part B 124, 89-100.
- Newstedt, J.L., J.P. Giesy, G.T. Ankley, D.E. Tillitt, R.A. Crawford, J.W. Gooch, P.D. Jones, M.S. Denison (1995): Development of toxic equivalency factors for PCB congeners and the assessment of TCDD and PCB mixtures in rainbow trout. Environm. Toxicol. Chem., 14, 861-871.
- Nicolas, J.-M. (1999): Vitellogenesis in fish and the effects of polycyclic aromatic hydrocarbon contaminants. Aquat. Toxicol., 45, 77-90.
- Nies, H. (1990): Radioaktive Substanzen. In: Lozán, J.L., W. Lenz, E. Rachor, B. Watermann und H. v. Westernhagen (Hrsg.): Warnsignale aus der Nordsee. Parey Verlag Berlin, Hamburg, 87-101.
- NRC (1987): Committee on biological markers. Environ. Health Perspect., 74, 3-9.
- NSTF (1993): North Sea Quality Status Report 1993. Oslo and Paris Commissions, London. Olsen and Olsen, Fredensborg, Denmark.
- O'Hare, D.B., R. Siddall, P.W.J. Robotham, R.A. Gill (1995): Influence of oestradiol on induction of EROD activity in roach (*Rutilus rutilus* L.). Chemosphere, 30, 1423-1428.

- Oikari, A., B. Jiminez (1992): Effects of hepatotoxicants on the induction of microsomal monooxygenase activity in sunfish liver by β -naphthoflavone and benzo[a]pyrene. Ecotox. Environm. Safety, 23, 89-102.
- OSPAR (1998): JAMP (Joint Assessment and Monitoring Programme) guidelines for contaminant-specific biological effects monitoring. OSPAR Commission, London.
- OSPAR (2000): Quality Status Report 2000. OSPAR Commission, London.
- Payne, J.F. (1976): Field evaluation of benzopyrene hydroxylase induction as a monitor for marine pollution. Science, 191, 945-946.
- Payne, J.F. (1984): Mixed-function oxygenases in biological monitoring programs review of potential usage in different phyla of aquatic animals. In: G. Persoone, E. Jaspers, C. Claus (eds.): Ecotoxicological testing for the marine environment. State Univ. Ghent and Inst. Mar. Scient. Res., Bredene, Belgium. Vol. 1, 625-655.
- Payne, J.F., W.R. Penrose (1975): Induction of aryl hydrocarbon benzo(a)pyrene hydroxylase in fish by petroleum. Bull. Environ. Contam. Tox., 14, 112-116.
- Payne, J.F., L.L. Fancy, A.D. Rahimtula, E.L. Porter (1987): Review and perspective on the use of mixed-function oxygenase enzymes in biological monitoring. Comp. Biochem. Physiol., 86C, 233-245.
- Perkins, E.J., D. Schlenk (1998): Immunochemical characterization of hepatic cytochrome P450 isozymes in the channel catfish: assessment of sexual, developmental and treatment-related effects. Comp. Biochem. Physiol., 121C, 305-310.
- Pfaffenberger, B., H. Hühnerfuss, O. Landgraff, L. Karbe (1994): Seasonal variation of the toxic potential of coplanar polychlorinated biphenyls in blue mussel (*Mytilus edulis* L.) from the German Bight. In: L. Karbe, K. Mädler, J. Westendorf (Hrsg.): Biologische Effekte von Schadstoffen und toxisches Potential von Wasser und Sediment in Elbe und Nordsee (BIO-TOX ELBE/BIOTOX NORDSEE). Berichte Zentr. Meeres- u. Klimaforsch., Hamburg, Reihe E. Nr. 7, 125-141.
- Pluta, H.-J., J. Hahn, P.-D. Hansen (1991): Fischkrankheiten im Wattenmeer, Teilvorhaben 3 – Untersuchungen zur Biotransformation („Entgiftungsaktivität“ der Mischfunktionellen Oxidasen, MFO) in der Fischleber. Abschlussbericht, UBA-Code Nr. 10204373.
- Power, M., M.J. Attrill, R.M. Thomas (1999): Trends in agricultural pesticide (atrazine, lindane, simazine) concentrations in the Thames estuary. Environ. Pollut., 104, 31-39.
- Reincke, H. (1992): Menschliche Eingriffe in das Elbesystem bis heute und notwendige Sanierungsmaßnahmen. In: Kausch, H. (Hrsg.): Die Unterelbe – Natürlicher Zustand und Veränderungen durch den Menschen. Berichte Zentr. Meeres- u. Klimaforsch., Hamburg, Nr. 19, 249-260.
- Renton, K.W., R.F. Addison (1992): Hepatic microsomal mono-oxygenase activity and P450Ia-mRNA in North Sea dab *Limanda limanda* from contaminated sites. In: Stebbing, A.R.D., V. Dethlefsen, M. Carr (eds.): Biological effects of contaminants in the North Sea - Results of the ICES/IOC Bremerhaven Workshop. Inter-Research, Amelinghausen. Mar. Ecol. Prog. Ser., 91, 65-69.
- Rice, D.W., C.P. Seltenrich, M.L. Keller, R.B. Spies, J.S. Felton (1994): Mixed-function oxidase-specific activity in wild and caged speckled sanddabs *Citharichthys stigmaeus* in Elkhorn Slough, Moss Landing Harbor and nearshore Monterey Bay, California. Environ. Pollut., 84, 179-188.
- Riedel-Lorjé, J. C., N. Möller-Lindenholz, B. Vaessen (1992): Salzgehalts- und Trübstoffverhältnisse in dem oberen Brackwassergebiet der Elbe. Arbeitsgemeinschaft für die Reinhalzung der Elbe (Hrsg.), Hamburg.

- Rijnsdorp, A.D., A.D. Vethaak, P.I. van Leeuwen (1992): Population biology of dab *Limanda limanda* in the southern North Sea. In: Stebbing, A.R.D., V. Dethlefsen, M. Carr (eds.): Biological effects of contaminants in the North Sea - Results of the ICES/IOC Bremerhaven Workshop. Inter-Research, Amelinghausen. Mar. Ecol. Prog. Ser., 91, 19-35.
- Robson, A.J., C. Neal (1997): A summary of regional water quality for eastern UK rivers. Sci. Tot. Environ., 194/195, 15-37.
- Saborowski, R. (1996): Zur Ökophysiologie der Kliesche, *Limanda limanda* (L.) – Einfluss saisonaler Zyklen auf das hepatische Entgiftungssystem. Dissertation Universität Hamburg 1996.
- Sachs, L. (1969): Angewandte Statistik. Springer Verlag Berlin.
- Safe, S. (1984): Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): Biochemistry, toxicology and mechanism of action. CRC Crit. Rev. Toxicol., 13, 319-393.
- Santillo, D. P. Johnston, A. Singhofen, J. Landmann, M. Krautter (1999): EU-Chemiepolitik – Der Ausweg. Greenpeace e. V. (Hrsg.), Hamburg.
- Sarasquete, C., H. Segner (2000): Cytochrome P4501A (CYP1A) in teleostean fishes. A review of immunohistochemical studies. Sci. Tot. Environ., 247, 313-332.
- Schirmer, M., (1996): Die Weser – eine Zustandsbeschreibung. In: Lozán, J.L., H. Kausch (Hrsg.): Warnsignale aus Flüssen und Ästuaren. Parey Verlag Berlin, 75-83.
- Schlenk, D. (1999): Necessity of defining biomarkers for use in ecological risk assessments. Mar. Poll. Bull., 39, 48-53.
- Schlenk, D., L.D. Peters, D.R. Livingstone (1996): Correction of Salinity with Flavin-containing monooxygenase activity but not cytochrome P450 activity in the euryhaline fish (*Platichthys flesus*). Biochem. Pharmacol., 52, 815-818.
- Sijm, D.T.H.M., W. Seinen, A. Opperhuizen (1992): Life-cycle biomagnification study in fish. Environ. Sci. Technol., 26, 2162-2174.
- Sijm, D.T.H.M., H. Weaver, A. Opperhuizen (1993): Congener-specific biotransformation and bioaccumulation of PCDDs and PCDFs from fly ash in fish. Environm. Toxicol. Chem., 12, 1895-1907.
- Sleiderink, H.M., J.P. Boon (1995): Cytochrome P450 1A response in North Sea dab, *Limanda limanda*, from offshore and coastal sites. Mar. Poll. Bull., 30, 660-666.
- Sleiderink, H.M., J. P. Boon (1996): Temporal induction pattern of hepatic cytochrome P450 1A in thermally acclimated dab (*Limanda limanda*) treated with 3,3',4,4'-tetrachlorobiphenyl (CB77). Chemosphere, 32, 2335-2344.
- Sleiderink, H.M., J. Beyer, J.M. Everaarts, J.P. Boon (1995a): Influence of temperature on cytochrome P450 1A in dab (*Limanda limanda*) from the southern North Sea: results from field surveys and a laboratory study. Mar. Environ. Res., 39, 67-71.
- Sleiderink, H.M., J. Beyer, E. Scholtens, A. Goksøyr, J. Nieuwenhuize, J.M. van Liere, J. M. Everaarts, J.P. Boon (1995b): Influence of temperature and polyaromatic contaminants on CYP1A levels in North Sea dab (*Limanda limanda*). Aquat. Toxicol., 32, 189-209.
- Sleiderink, H.M., J.M. Everaarts, A. Goksøyr, J.P. Boon (1995c): Hepatic cytochrome P-4501A induction in dab (*Limanda limanda*) after oral dosis with the polychlorinated biphenyl mixture clophen A40. Environm. Toxicol. Chem., 14, 679-687.
- Snowberger-Gray, E., B.R. Woodin, J.J. Stegemann, J.J. (1991): Sex differences in hepatic monooxygenases in winter flounder (*Pseudopleuronectes americanus*) and scup (*Stenotomus chrysops*) and regulation of P450 forms by estradiol. J. Exp. Zool., 259, 330-342.
- Spannhof, L. (1995): Einführung in die Fischphysiologie. Kovac Verlag Hamburg.

- Spies, R.B., D.W. Rice, Jr., J. Felton (1988): Effects of organic contaminants on reproduction of the starry flounder *Platichthys stellatus* in San Francisco Bay – I. Hepatic contamination and mixed-function oxidase (MFO) activity during the reproductive season. Mar. Biol., 98, 181-189.
- Stagg, R.M. (1998): The development of an international programme for monitoring the biological effects of contaminants in the OSPAR convention area. Mar. Environ. Res., 46, 307-313.
- Stagg, R.M., R.F. Addison (1995a): An inter-laboratory comparison of measurements of ethoxresorufin O-de-ethylase activity in dab (*Limanda limanda*) liver. Mar. Environ. Res., 40, 93-108.
- Stagg, R.M., A. McIntosh, P. Mackie (1995b): Elevation of hepatic monooxygenase activity in the dab (*Limanda limanda* L.) in relation to environmental contamination with petroleum hydrocarbons in the northern North Sea. Aquat. Toxicol., 33, 245-264.
- Stagg, R.M., C. Robinson, A.M. McIntosh, C.F. Moffat, D.W. Bruno (1998): The effects of the „Brear“ oil spill, Shetland isles, Scotland, on P4501A in farmed atlantic salmon (*Salmo salar*) and the common dab (*Limanda limanda*). Mar. Environ. Res., 46, 301-306.
- Stebbing, A.R.D., V. Dethlefsen (1992): Introduction to the Bremerhaven Workshop on biological effects monitoring. In: Stebbing, A.R.D., V. Dethlefsen, M. Carr (eds.): Biological effects of contaminants in the North Sea. Results of the ICES/IOC Bremerhaven Workshop. Inter-Research, Amelinghausen. Mar. Ecol. Prog. Ser., 91, 1-8.
- Stegeman, J.J. (1991): Nomenclature for hydrocarbon-inducible cytochrome P450 in fish. Mar. Environ. Res., 34, 133-138.
- Stegeman, J.J., A.M. Pajor, P. Thomas (1982): Influence of oestradiol and testosterone on cytochrome P-450 and monooxygenase activity in immature brook trout *Savelinus fontanilis*. Biochem. Pharmacol. 31 (24), 3979-3989.
- Stegeman, J.J., P.J. Klopper-Sams (1987): Cytochrome P-450 isozymes and monooxygenase activity in aquatic animals. Environ. Health Perspect., 71, 87-95.
- Stegeman, J.J., B.R. Woodin, A. Goksøyr (1988): Apparent cytochrome P-450 induction as an indication of exposure to environmental chemicals in the flounder *Platichthys flesus*. Mar. Ecol. Prog. Ser., 46, 55-60.
- Stein, J.E., W.L. Reichert, M. Nishimoto, U. Varanasi (1990): Overview of studies on liver carcinogenesis in english sole from puget sound; Evidence for a xenobiotic chemical etiology II: biochemical studies. Sci. Tot. Environ., 94, 51-69.
- Stevenson, C., N.G. Betty (1999): Distribution of copper, nickel and zinc in the Thames estuary. Mar. Poll. Bull., 38, 328-331.
- Sturm, R., J. Gandrass (1988): Verhalten von schwerflüchtigen Chlorkohlenwasserstoffen an Schwebstoffen des Elbe-Ästuars. Vom Wasser, 70, 271-279.
- Sulaiman, N., S. George, M.D. Burke (1991): Assessment of sublethal pollutant impact on flounders in an industrialised estuary using hepatic biochemical indices. Mar. Ecol. Prog. Ser., 68, 207-212.
- Summers, R.W. (1979): Life cycle and population ecology of the flounder *Platichthys flesus* (L.) in the Ythan estuary, Scotland. J. Nat. Hist., 13, 703-723.
- Sumpter, J.P. (1997): Environmental control of fish reproduction: A Different perspective. Fish Physiol. Biochem., 17, 25-31.
- Tarlebø, J., J.E. Solbakken, K.H. Palmkork (1985): Variation in hepatic arylhydrocarbon hydroxylase activity in flounder, *Platichthys flesus*: a baseline study. Helgoländer Meeresunters., 39, 187-199.
- Temming, A. (1989): Biologie und Populationsdynamik der Kliesche (*Limanda limanda*) (L.) in der Ostsee. Ber. Inst. Meeresk. Kiel Nr. 189.

- Thomas, K.V., J.E. Thain, M.J. Waldock (1999): Identification of toxic substances in United Kingdom estuaries. Environm. Toxicol. Chem., Vol. 18, 401-411.
- Tomczak, G., E. Goedecke (1962): Monatskarten der Temperatur der Nordsee, dargestellt für verschiedene Tiefenhorizonte. Dt. hydrogr. Z., Erg.-H. B, Nr. 7.
- UBA (1994): Daten zur Umwelt 1992/1993. Umweltbundesamt (Hrsg.). Erich Schmidt Verlag Berlin.
- UBA (1997): Daten zur Umwelt – Der Zustand der Umwelt in Deutschland - Ausgabe 1997. Umweltbundesamt (Hrsg.), Erich Schmidt Verlag, Berlin.
- Ueng, T.H., Y.-F. Ueng, S.S. Park (1992): Comparative induction of cytochrome P-450-dependent monooxygenases in the livers and gills of tilapia and carp. Aquat. Toxicol., 23, 49-64.
- Van der Land, M.A. (1991): Distribution of flatfish eggs in the 1989 egg surveys in the southeastern North Sea, and mortality of plaice and sole eggs. Neth. J. Sea Res., 27, 277-286.
- Van Schanke, A., J.P. Boon, Y. Aardoom, A. van Leest, F.J. van Schooten, L. Maas, M. van den Berg, J.M. Everaarts (2000): Effect of a dioxin-like PCB (126) on the biotransformation and genotoxicity of benzo[a]pyrene in the marine flatfish dab (*Limanda limanda*). Aquat. Toxicol., 50, 403-415.
- Vethaak, A.D., T. ap Rheinallt (1992): Fish deseases as a monitor for marine pollution: the case of the North Sea. Rev. Fish Biol. Fish., 2, 1-32.
- Wahl, E., P. Cameron, A. Köhler-Günther, H. v. Westernhagen, K. Anders, H. Möller, H. Büther, V. Dethlefsen, K. Söffker, P.-D. Hansen, H.-J. Pluta, U. Harms (1995): Fischkrankheiten im Wattenmeer. Umweltbundesamt (Hrsg.), Texte 51/95.
- Westernhagen, H. von, P. Cameron, D. Janssen, M. Kerstan (1995): Age and size dependent chlorinated hydrocarbon concentrations in marine teleosts. Mar. Poll. Bull., 30, 655-659.
- Westernhagen, H. von, G. Krüner, K. Broeg (1999): Ethoxyresorufin O-deethylase (EROD) activity in the liver of dab (*Limanda limanda* L.) and flounder (*Platichthys flesus* L.) from the German Bight. EROD expression and tissue contamination. Helgol. Mar. Res., 53, 244-249.
- Woggon, W.-D. (1988): Modelle für Cytochrom P450. Nachr. Chem. Tech. Lab., 36, 890-895.
- Woodhead, R.J., R.J. Law, P. Matthiessen (1999): Polycyclic aromatic hydrocarbons in surface sediments around England and Wales, and their possible biological significance. Mar. Poll. Bull., 38, 773-790.
- Zhou, J.L., T.W. Fileman, S. Evans, P. Donkin, R.F.C. Mantoura (1996): Seasonal distribution of dissolved pesticides and polynuclear aromatic hydrocarbons in the Humber estuary and Humber coastal zone. Mar. Poll. Bull., 32, 599-608.
- Zimmer, M., W. Ahlf (1994): Erarbeitung von Kriterien zur Ableitung von Qualitätszielen für Sedimente und Schwebstoffe. Umweltbundesamt (Hrsg.), Texte 69/94.