

6 LITERATURVERZEICHNIS

1. Ahn, J. and R.M. Johnstone, Origin of a soluble truncated transferrin receptor. *Blood*, 1993. **81**(9): p. 2442-51.
2. Aisen, P., Transferrin receptor I. *Int J Biochem Cell Biol*, 2004. **36**(11): p. 2137-43.
3. Akesson, A., P. Bjellerup and M. Vahter, Evaluation of kits for measurement of the soluble transferrin receptor. *Scand J Clin Lab Invest*, 1999. **59**(2): p. 77-81.
4. Alfalah, M., E.T. Parkin, R. Jacob, E.D. Sturrock, R. Mentele, et al., A point mutation in the juxtamembrane stalk of human angiotensin I-converting enzyme invokes the action of a distinct secretase. *J Biol Chem*, 2001. **276**(24): p. 21105-9.
5. Allen, J., K.R. Backstrom, J.A. Cooper, M.C. Cooper, T.C. Detwiler, et al., Measurement of soluble transferrin receptor in serum of healthy adults. *Clin Chem*, 1998. **44**(1): p. 35-9.
6. Althoff, K., J. Mullberg, D. Aasland, N. Voltz, K. Kallen, et al., Recognition sequences and structural elements contribute to shedding susceptibility of membrane proteins. *Biochem J*, 2001. **353**(Pt 3): p. 663-72.
7. Althoff, K., P. Reddy, N. Voltz, S. Rose-John and J. Mullberg, Shedding of interleukin-6 receptor and tumor necrosis factor alpha. Contribution of the stalk sequence to the cleavage pattern of transmembrane proteins. *Eur J Biochem*, 2000. **267**(9): p. 2624-31.
8. Altin, J.G. and E.B. Pagler, A one-step procedure for biotinylation and chemical cross-linking of lymphocyte surface and intracellular membrane-associated molecules. *Anal Biochem*, 1995. **224**(1): p. 382-9.
9. Alvarez, E., N. Girones and R.J. Davis, Intermolecular disulfide bonds are not required for the expression of the dimeric state and functional activity of the transferrin receptor. *Embo J*, 1989. **8**(8): p. 2231-40.
10. Alvarez, E., N. Girones and R.J. Davis, Inhibition of the receptor-mediated endocytosis of diferric transferrin is associated with the covalent modification of the transferrin receptor with palmitic acid. *J Biol Chem*, 1990. **265**(27): p. 16644-55.
11. Amour, A., P.M. Slocombe, A. Webster, M. Butler, C.G. Knight, et al., TNF-alpha converting enzyme (TACE) is inhibited by TIMP-3. *FEBS Lett*, 1998. **435**(1): p. 39-44.
12. Anderson, B.F., H.M. Baker, E.J. Dodson, G.E. Norris, S.V. Rumball, et al., Structure of human lactoferrin at 3.2-A resolution. *Proc Natl Acad Sci U S A*, 1987. **84**(7): p. 1769-73.
13. Arribas, J., L. Coodly, P. Vollmer, T.K. Kishimoto, S. Rose-John, et al., Diverse cell surface protein ectodomains are shed by a system sensitive to metalloprotease inhibitors. *J Biol Chem*, 1996. **271**(19): p. 11376-82.
14. Barton, J.C. and L.F. Bertoli, Hemochromatosis: the genetic disorder of the twenty-first century. *Nat Med*, 1996. **2**(4): p. 394-5.
15. Bastin, J.M., M. Jones, C.A. O'Callaghan, L. Schimanski, D.Y. Mason, et al., Kupffer cell staining by an HFE-specific monoclonal antibody: implications for hereditary haemochromatosis. *Br J Haematol*, 1998. **103**(4): p. 931-41.
16. Baynes, R.D., Transferrin reduces the production of soluble transferrin receptor. *Proc Soc Exp Biol Med*, 1995. **209**(3): p. 286-94.
17. Baynes, R.D., J.D. Cook, T.H. Bothwell, B.M. Friedman and T.E. Meyer, Serum transferrin receptor in hereditary hemochromatosis and African siderosis. *Am J Hematol*, 1994. **45**(4): p. 288-92.
18. Baynes, R.D., Y.J. Shih and J.D. Cook, Mechanism of production of the serum transferrin receptor. *Adv Exp Med Biol*, 1994. **356**: p. 61-8.
19. Baynes, R.D., Y.J. Shih, B.G. Hudson and J.D. Cook, Characterization of transferrin receptor released by K562 erythroleukemia cells. *Proc Soc Exp Biol Med*, 1991. **197**(4): p. 416-23.
20. Baynes, R.D., Y.J. Shih, B.G. Hudson and J.D. Cook, Production of the serum form of the transferrin receptor by a cell membrane-associated serine protease. *Proc Soc Exp Biol Med*, 1993. **204**(1): p. 65-9.
21. Baynes, R.D., Y.J. Shih, B.G. Hudson and J.D. Cook, Identification of the membrane remnants of transferrin receptor with domain-specific antibodies. *J Lab Clin Med*, 1994. **123**(3): p. 407-14.
22. Beard, P., E. Offord, N. Paduwat and H. Bruggmann, SV40 activates transcription from the transferrin receptor promoter by inducing a factor which binds to the CRE/AP-1 recognition sequence. *Nucleic Acids Res*, 1991. **19**(25): p. 7117-23.

23. Beauchamp, J.R. and P.G. Woodman, Regulation of transferrin receptor recycling by protein phosphorylation. *Biochem J*, 1994. **303** (Pt 2): p. 647-55.
24. Beguin, Y., Soluble transferrin receptor for the evaluation of erythropoiesis and iron status. *Clin Chim Acta*, 2003. **329**(1-2): p. 9-22.
25. Beguin, Y., H.A. Huebers, B. Josephson and C.A. Finch, Transferrin receptors in rat plasma. *Proc Natl Acad Sci U S A*, 1988. **85**(2): p. 637-40.
26. Bennett, M.J., J.A. Lebron and P.J. Bjorkman, Crystal structure of the hereditary haemochromatosis protein HFE complexed with transferrin receptor. *Nature*, 2000. **403**(6765): p. 46-53.
27. Birnboim, H.C. and J. Doly, A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res*, 1979. **7**(6): p. 1513-23.
28. Black, R.A., C.T. Rauch, C.J. Kozlosky, J.J. Peschon, J.L. Slack, et al., A metalloproteinase disintegrin that releases tumour-necrosis factor-alpha from cells. *Nature*, 1997. **385**(6618): p. 729-33.
29. Black, R.A. and J.M. White, ADAMs: focus on the protease domain. *Curr Opin Cell Biol*, 1998. **10**(5): p. 654-9.
30. Blobel, C.P., Metalloprotease-disintegrins: links to cell adhesion and cleavage of TNF alpha and Notch. *Cell*, 1997. **90**(4): p. 589-92.
31. Boumpas, D.T., E.D. Anastassiou, G.C. Tsokos, G. Thyphronitis and J.E. Balow, Gene transcription during in vitro activation of human B lymphocytes with *Staphylococcus aureus* Cowan I strain.. *J Immunol*, 1990. **145**(8): p. 2701-5.
32. Brakebusch, C., E.E. Varfolomeev, M. Batkin and D. Wallach, Structural requirements for inducible shedding of the p55 tumor necrosis factor receptor. *J Biol Chem*, 1994. **269**(51): p. 32488-96.
33. Brinkmann, U., Recombinant antibody fragments and immunotoxin fusions for cancer therapy. *In Vivo*, 2000. **14**(1): p. 21-7.
34. Brou, C., F. Logeat, N. Gupta, C. Bessia, O. LeBail, et al., A novel proteolytic cleavage involved in Notch signaling: the role of the disintegrin-metalloprotease TACE. *Mol Cell*, 2000. **5**(2): p. 207-16.
35. Buchegger, F., I.S. Trowbridge, L.F. Liu, S. White and J.F. Collawn, Functional analysis of human/chicken transferrin receptor chimeras indicates that the carboxy-terminal region is important for ligand binding. *Eur J Biochem*, 1996. **235**(1-2): p. 9-17.
36. Busfield, S.J., P.A. Tilbrook, B.A. Callus, A. Spadaccini, L. Kuhn, et al., Complex regulation of transferrin receptors during erythropoietin-induced differentiation of J2E erythroid cells--elevated transcription and mRNA stabilisation produce only a modest rise in protein content. *Eur J Biochem*, 1997. **249**(1): p. 77-84.
37. Butler, G.S., H. Will, S.J. Atkinson and G. Murphy, Membrane-type-2 matrix metalloproteinase can initiate the processing of progelatinase A and is regulated by the tissue inhibitors of metalloproteinases. *Eur J Biochem*, 1997. **244**(2): p. 653-7.
38. Cazzola, M., Y. Beguin, G. Bergamaschi, R. Guarnone, P. Cerani, et al., Soluble transferrin receptor as a potential determinant of iron loading in congenital anaemias due to ineffective erythropoiesis. *Br J Haematol*, 1999. **106**(3): p. 752-5.
39. Chan, L.N. and E.M. Gerhardt, Transferrin receptor gene is hyperepressed and transcriptionally regulated in differentiating erythroid cells. *J Biol Chem*, 1992. **267**(12): p. 8254-9.
40. Chan, R.Y., C. Seiser, H.M. Schulman, L.C. Kuhn and P. Ponka, Regulation of transferrin receptor mRNA expression. Distinct regulatory features in erythroid cells. *Eur J Biochem*, 1994. **220**(3): p. 683-92.
41. Cheng, Y., O. Zak, P. Aisen, S.C. Harrison and T. Walz, Structure of the human transferrin receptor-transferrin complex. *Cell*, 2004. **116**(4): p. 565-76.
42. Chitambar, C.R., A.L. Loebel and N.A. Noble, Shedding of transferrin receptor from rat reticulocytes during maturation in vitro: soluble transferrin receptor is derived from receptor shed in vesicles. *Blood*, 1991. **78**(9): p. 2444-50.
43. Chitambar, C.R. and Z. Zivkovic, Release of soluble transferrin receptor from the surface of human leukemic HL60 cells. *Blood*, 1989. **74**(2): p. 602-8.
44. Chitambar, C.R. and Z. Zivkovic-Gilgenbach, Influence of cellular iron status on the release of soluble transferrin receptor from human promyelocytic leukemic HL60 cells. *J Lab Clin Med*, 1990. **116**(3): p. 345-53.
45. Codony-Servat, J., J. Albanell, J.C. Lopez-Talavera, J. Arribas and J. Baselga, Cleavage of the HER2 ectodomain is a pervanadate-activable process that is inhibited by the tissue inhibitor of metalloproteinases-1 in breast cancer cells. *Cancer Res*, 1999. **59**(6): p. 1196-201.
46. Cook, J.D., S. Dassenko and B.S. Skikne, Serum transferrin receptor as an index of iron absorption. *Br J Haematol*, 1990. **75**(4): p. 603-9.
47. Costa, C., N.K. Bell, T.J. Stabel and W.L. Fodor, Use of porcine tumor necrosis factor receptor 1-Ig fusion protein to prolong xenograft survival. *Xenotransplantation*, 2004. **11**(6): p. 491-502.
48. Cowley, J.M., D.E. Janney, R.C. Gerkin and P.R. Buseck, The structure of ferritin cores determined by electron nanodiffraction. *J Struct Biol*, 2000. **131**(3): p. 210-6.
49. Cox, T.M., M.W. O'Donnell, P. Aisen and I.M. London, Hemin inhibits internalization of transferrin by reticulocytes and promotes phosphorylation of the membrane transferrin receptor. *Proc Natl Acad Sci U S A*, 1985. **82**(15): p. 5170-4.
50. Cuthbert, J.A., Iron, HFE, and hemochromatosis update. *J Investig Med*, 1997. **45**(9): p. 518-29.
51. Dassler, K., Der Sheddingprozeß des Transferrinrezeptors - Entwicklung von Detektionsverfahren und Messung der Sheddingaktivität. Diplomarbeit. 2000, Berlin: Freie Universität Berlin.
52. Dassler, K., K. Wandzik, R. Tauber, M. Kaup and H. Fuchs, Release of Soluble Transferrin Receptor is regulated by ferri-Transferrin. *Clin. Chem. Lab. Med.*, 2004. **42**(12).
53. Davies, P.S. and C.A. Enns, Expression of the hereditary hemochromatosis protein, HFE, increases ferritin levels by inhibiting iron export in HT29 cells. *J Biol Chem*, 2004.

54. Davis, R.J. and H. Meisner, Regulation of transferrin receptor cycling by protein kinase C is independent of receptor phosphorylation at serine 24 in Swiss 3T3 fibroblasts. *J Biol Chem*, 1987. **262**(33): p. 16041-7.
55. de Lima, G.A. and H.Z. Grotto, Soluble transferrin receptor and immature reticulocytes are not useful for distinguishing iron-deficiency anemia from heterozygous beta-thalassemia. *Sao Paulo Med J*, 2003. **121**(2): p. 90-1.
56. Dello Sbarba, P. and E. Rovida, Transmodulation of cell surface regulatory molecules via ectodomain shedding. *Biol Chem*, 2002. **383**(1): p. 69-83.
57. Do, S.I. and R.D. Cummings, Presence of O-linked oligosaccharide on a threonine residue in the human transferrin receptor. *Glycobiology*, 1992. **2**(4): p. 345-53.
58. Donovan, S.K., Peers under review. *Nature*, 2001. **413**(6857): p. 669.
59. Drakesmith, H., E. Sweetland, L. Schimanski, J. Edwards, D. Cowley, et al., The hemochromatosis protein HFE inhibits iron export from macrophages. *Proc Natl Acad Sci U S A*, 2002. **99**(24): p. 15602-7.
60. Dubljevic, V., A. Sali and J.W. Goding, A conserved RGD (Arg-Gly-Asp) motif in the transferrin receptor is required for binding to transferrin. *Biochem J*, 1999. **341** (Pt 1): p. 11-4.
61. Eichholtz, T., P. Vosseveld, M. van Overveld and H. Ploegh, Activation of protein kinase C accelerates internalization of transferrin receptor but not of major histocompatibility complex class I, independent of their phosphorylation status. *J Biol Chem*, 1992. **267**(31): p. 22490-5.
62. Elenius, K., G. Corfas, S. Paul, C.J. Choi, C. Rio, et al., A novel juxtamembrane domain isoform of HER4/ErbB4. Isoform-specific tissue distribution and differential processing in response to phorbol ester. *J Biol Chem*, 1997. **272**(42): p. 26761-8.
63. English, W.R., X.S. Puente, J.M. Freije, V. Knauper, A. Amour, et al., Membrane type 4 matrix metalloproteinase (MMP17) has tumor necrosis factor-alpha convertase activity but does not activate pro-MMP2. *J Biol Chem*, 2000. **275**(19): p. 14046-55.
64. English, W.R., G. Velasco, J.O. Stracke, V. Knauper and G. Murphy, Catalytic activities of membrane-type 6 matrix metalloproteinase (MMP25). *FEBS Lett*, 2001. **491**(1-2): p. 137-42.
65. Eyries, M., A. Michaud, J. Deinum, M. Agrapart, J. Chomilier, et al., Increased shedding of angiotensin-converting enzyme by a mutation identified in the stalk region. *J Biol Chem*, 2001. **276**(8): p. 5525-32.
66. Feder, J.N., D.M. Penny, A. Irrinki, V.K. Lee, J.A. Lebron, et al., The hemochromatosis gene product complexes with the transferrin receptor and lowers its affinity for ligand binding. *Proc Natl Acad Sci U S A*, 1998. **95**(4): p. 1472-7.
67. Feder, J.N., Z. Tsuchihashi, A. Irrinki, V.K. Lee, F.A. Mapa, et al., The hemochromatosis founder mutation in HLA-H disrupts beta2-microglobulin interaction and cell surface expression. *J Biol Chem*, 1997. **272**(22): p. 14025-8.
68. Feehan, C., K. Darlak, J. Kahn, B. Walcheck, A.F. Spatola, et al., Shedding of the lymphocyte L-selectin adhesion molecule is inhibited by a hydroxamic acid-based protease inhibitor. Identification with an L-selectin-alkaline phosphatase reporter. *J Biol Chem*, 1996. **271**(12): p. 7019-24.
69. Finch, C., Regulators of iron balance in humans. *Blood*, 1994. **84**(6): p. 1697-702.
70. Fitzgerald, M.L., Z. Wang, P.W. Park, G. Murphy and M. Bernfield, Shedding of syndecan-1 and -4 ectodomains is regulated by multiple signaling pathways and mediated by a TIMP-3-sensitive metalloproteinase. *J Cell Biol*, 2000. **148**(4): p. 811-24.
71. Fleming, M.D., M.A. Romano, M.A. Su, L.M. Garrick, M.D. Garrick, et al., Nramp2 is mutated in the anemic Belgrade (b) rat: evidence of a role for Nramp2 in endosomal iron transport. *Proc Natl Acad Sci U S A*, 1998. **95**(3): p. 1148-53.
72. Fleming, M.D., C.C. Trenor, 3rd, M.A. Su, D. Foernzler, D.R. Beier, et al., Microcytic anaemia mice have a mutation in Nramp2, a candidate iron transporter gene. *Nat Genet*, 1997. **16**(4): p. 383-6.
73. Flowers, C.H., B.S. Skikne, A.M. Covell and J.D. Cook, The clinical measurement of serum transferrin receptor. *J Lab Clin Med*, 1989. **114**(4): p. 368-77.
74. Fors, B.P., K. Goodarzi and U.H. von Andrian, L-selectin shedding is independent of its subsurface structures and topographic distribution. *J Immunol*, 2001. **167**(7): p. 3642-51.
75. Frazer, D.M. and G.J. Anderson, The orchestration of body iron intake: how and where do enterocytes receive their cues? *Blood Cells Mol Dis*, 2003. **30**(3): p. 288-97.
76. Fuchs, H. and R. Gessner, Iodination significantly influences the binding of human transferrin to the transferrin receptor. *Biochim Biophys Acta*, 2002. **1570**(1): p. 19-26.
77. Fuchs, H., U. Lucken, R. Tauber, A. Engel and R. Gessner, Structural model of phospholipid-reconstituted human transferrin receptor derived by electron microscopy. *Structure*, 1998. **6**(10): p. 1235-43.
78. Gatter, K.C., G. Brown, I.S. Trowbridge, R.E. Woolston and D.Y. Mason, Transferrin receptors in human tissues: their distribution and possible clinical relevance. *J Clin Pathol*, 1983. **36**(5): p. 539-45.
79. Gehrke, S.G., H. Kulaksiz, T. Herrmann, H.D. Riedel, K. Bents, et al., Expression of hepcidin in hereditary hemochromatosis: evidence for a regulation in response to the serum transferrin saturation and to non-transferrin-bound iron. *Blood*, 2003. **102**(1): p. 371-6.
80. Gerstein, M., B.F. Anderson, G.E. Norris, E.N. Baker, A.M. Lesk, et al., Domain closure in lactoferrin. Two hinges produce a see-saw motion between alternative close-packed interfaces. *J Mol Biol*, 1993. **234**(2): p. 357-72.
81. Giannetti, A.M., P.M. Snow, O. Zak and P.J. Bjorkman, Mechanism for multiple ligand recognition by the human transferrin receptor. *PLoS Biol*, 2003. **1**(3): p. E51.
82. Gross, C.N., A. Irrinki, J.N. Feder and C.A. Enns, Co-trafficking of HFE, a nonclassical major histocompatibility complex class I protein, with the transferrin receptor implies a role in intracellular iron regulation. *J Biol Chem*, 1998. **273**(34): p. 22068-74.

83. Gruenheid, S., M. Cellier, S. Vidal and P. Gros, Identification and characterization of a second mouse Nramp gene. *Genomics*, 1995. **25**(2): p. 514-25.
84. Gruner, S., M. Prostedna, M. Koch, Y. Miura, V. Schulte, et al., Relative antithrombotic effect of soluble GPVI dimer versus anti-GPVI antibodies in mice. *Blood*, 2004.
85. Gunshin, H., B. Mackenzie, U.V. Berger, Y. Gunshin, M.F. Romero, et al., Cloning and characterization of a mammalian proton-coupled metal-ion transporter. *Nature*, 1997. **388**(6641): p. 482-8.
86. Gutwein, P., S. Mechtshheimer, S. Riedle, A. Stoeck, D. Gast, et al., ADAM10-mediated cleavage of L1 adhesion molecule at the cell surface and in released membrane vesicles. *Faseb J*, 2003. **17**(2): p. 292-4.
87. Hall, D.R., J.M. Hadden, G.A. Leonard, S. Bailey, M. Neu, et al., The crystal and molecular structures of diferric porcine and rabbit serum transferrins at resolutions of 2.15 and 2.60 Å, respectively. *Acta Crystallogr D Biol Crystallogr*, 2002. **58**(Pt 1): p. 70-80.
88. Haro, H., H.C. Crawford, B. Fingleton, K. Shinomiya, D.M. Spengler, et al., Matrix metalloproteinase-7-dependent release of tumor necrosis factor- α in a model of herniated disc resorption. *J Clin Invest*, 2000. **105**(2): p. 143-50.
89. Hayes, G.R., C.A. Enns and J.J. Lucas, Identification of the O-linked glycosylation site of the human transferrin receptor. *Glycobiology*, 1992. **2**(4): p. 355-9.
90. Hayes, G.R., A. Williams, C.E. Costello, C.A. Enns and J.J. Lucas, The critical glycosylation site of human transferrin receptor contains a high-mannose oligosaccharide. *Glycobiology*, 1995. **5**(2): p. 227-32.
91. Hayes, G.R., A.M. Williams, J.J. Lucas and C.A. Enns, Structure of human transferrin receptor oligosaccharides: conservation of site-specific processing. *Biochemistry*, 1997. **36**(17): p. 5276-84.
92. Hinkle, C.L., S.W. Sunnarborg, D. Loisel, C.E. Parker, M. Stevenson, et al., Selective roles for tumor necrosis factor α -converting enzyme/ADAM17 in the shedding of the epidermal growth factor receptor ligand family: the juxtamembrane stalk determines cleavage efficiency. *J Biol Chem*, 2004. **279**(23): p. 24179-88.
93. Hirsch, S., R. Miskimins and W.K. Miskimins, Mitogenic activation of the transferrin receptor gene promoter is modulated by inhibitors of tyrosine kinases and tyrosine phosphatases. *Recept Signal Transduct*, 1996. **6**(3-4): p. 121-9.
94. Howard, L., Y. Zheng, M. Horrocks, R.A. Maciewicz and C. Blobel, Catalytic activity of ADAM28. *FEBS Lett*, 2001. **498**(1): p. 82-6.
95. Huebers, H.A., Y. Beguin, P. Pootrakul, D. Einspahr and C.A. Finch, Intact transferrin receptors in human plasma and their relation to erythropoiesis. *Blood*, 1990. **75**(1): p. 102-7.
96. Ito, N., S. Nomura, A. Iwase, T. Ito, F. Kikkawa, et al., ADAMs, a disintegrin and metalloproteinases, mediate shedding of oxytocinase. *Biochem Biophys Res Commun*, 2004. **314**(4): p. 1008-13.
97. Izumi, Y., M. Hirata, H. Hasuwa, R. Iwamoto, T. Umata, et al., A metalloprotease-disintegrin, MDC9/meltrin- γ /ADAM9 and PKC δ are involved in TPA-induced ectodomain shedding of membrane-anchored heparin-binding EGF-like growth factor. *Embo J*, 1998. **17**(24): p. 7260-72.
98. Johnstone, R.M., Cleavage of the transferrin receptor by human granulocytes: differential proteolysis of the exosome-bound TFR. *J Cell Physiol*, 1996. **168**(2): p. 333-45.
99. Johnstone, R.M., A. Mathew, A.B. Mason and K. Teng, Exosome formation during maturation of mammalian and avian reticulocytes: evidence that exosome release is a major route for externalization of obsolete membrane proteins. *J Cell Physiol*, 1991. **147**(1): p. 27-36.
100. Kaup, M., Identifizierung und funktionelle Charakterisierung von Transferrinrezeptor-Sheddingproteasen. Doktorarbeit. 2002, Berlin: Freie Universität Berlin.
101. Kaup, M., K. Dassler, U. Reineke, C. Weise, R. Tauber, et al., Processing of the human transferrin receptor at distinct positions within the stalk region by neutrophil elastase and cathepsin G. *Biol Chem*, 2002. **383**(6): p. 1011-20.
102. Kaup, M., K. Dassler, C. Weise and H. Fuchs, Shedding of the transferrin receptor is mediated constitutively by an integral membrane metalloprotease sensitive to tumor necrosis factor α protease inhibitor-2. *J Biol Chem*, 2002. **277**(41): p. 38494-502.
103. Kawabata, H., R. Yang, T. Hiram, P.T. Vuong, S. Kawano, et al., Molecular cloning of transferrin receptor 2. A new member of the transferrin receptor-like family. *J Biol Chem*, 1999. **274**(30): p. 20826-32.
104. Kinoshita, T., H. Sato, T. Takino, M. Itoh, T. Akizawa, et al., Processing of a precursor of 72-kilodalton type IV collagenase/gelatinase A by a recombinant membrane-type 1 matrix metalloproteinase. *Cancer Res*, 1996. **56**(11): p. 2535-8.
105. Klausner, R.D., J. van Renswoude, C. Kempf, K. Rao, J.L. Bateman, et al., Failure to release iron from transferrin in a Chinese hamster ovary cell mutant pleiotropically defective in endocytosis. *J Cell Biol*, 1984. **98**(3): p. 1098-101.
106. Kohgo, Y., T. Nishisato, H. Kondo, N. Tsushima, Y. Niitsu, et al., Circulating transferrin receptor in human serum. *Br J Haematol*, 1986. **64**(2): p. 277-81.
107. Kolkenbrock, H., L. Essers, N. Ulbrich and H. Will, Biochemical characterization of the catalytic domain of membrane-type 4 matrix metalloproteinase. *Biol Chem*, 1999. **380**(9): p. 1103-8.
108. Krause, A., S. Neitz, H.J. Magert, A. Schulz, W.G. Forssmann, et al., LEAP-1, a novel highly disulfide-bonded human peptide, exhibits antimicrobial activity. *FEBS Lett*, 2000. **480**(2-3): p. 147-50.
109. Kronke, M., W.J. Leonard, J.M. Depper and W.C. Greene, Sequential expression of genes involved in human T lymphocyte growth and differentiation. *J Exp Med*, 1985. **161**(6): p. 1593-8.
110. Kulaksiz, H., S.G. Gehrke, A. Janetzko, D. Rost, T. Bruckner, et al., Pro-hepcidin: expression and cell specific localisation in the liver and its regulation in hereditary haemochromatosis, chronic renal insufficiency, and renal anaemia. *Gut*, 2004. **53**(5): p. 735-43.
111. Laemmli, U.K., Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 1970. **227**(5259): p. 680-5.

112. Laftah, A.H., B. Ramesh, R.J. Simpson, N. Solanky, S. Bahram, et al., Effect of hepcidin on intestinal iron absorption in mice. *Blood*, 2004. **103**(10): p. 3940-4.
113. Lammich, S., E. Kojro, R. Postina, S. Gilbert, R. Pfeiffer, et al., Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc Natl Acad Sci U S A*, 1999. **96**(7): p. 3922-7.
114. Latif, R., T. Ando and T.F. Davies, Monomerization as a prerequisite for intramolecular cleavage and shedding of the thyrotropin receptor. *Endocrinology*, 2004. **145**(12): p. 5580-8.
115. Lawrence, C.M., S. Ray, M. Babyonyshev, R. Galluser, D.W. Borhani, et al., Crystal structure of the ectodomain of human transferrin receptor. *Science*, 1999. **286**(5440): p. 779-82.
116. Lebron, J.A., M.J. Bennett, D.E. Vaughn, A.J. Chirino, P.M. Snow, et al., Crystal structure of the hemochromatosis protein HFE and characterization of its interaction with transferrin receptor. *Cell*, 1998. **93**(1): p. 111-23.
117. Lebron, J.A., A.P. West, Jr. and P.J. Bjorkman, The hemochromatosis protein HFE competes with transferrin for binding to the transferrin receptor. *J Mol Biol*, 1999. **294**(1): p. 239-45.
118. Lee, G.R., S. Nacht, J.N. Lukens and G.E. Cartwright, Iron metabolism in copper-deficient swine. *J Clin Invest*, 1968. **47**(9): p. 2058-69.
119. Lee, K.Y. and S.I. Do, Differential and cell-type specific microheterogeneity of high mannose-type Asn-linked oligosaccharides of human transferrin receptor. *Mol Cells*, 2001. **12**(2): p. 239-43.
120. Leung, Y.M., W.H. Zhu and T.T. Loh, Apotransferrin can elevate intracellular free calcium ion and stimulate mitogenesis in human leukemic HL60 cells. *Biol Signals*, 1993. **2**(3): p. 117-25.
121. Levy, J.E., O. Jin, Y. Fujiwara, F. Kuo and N.C. Andrews, Transferrin receptor is necessary for development of erythrocytes and the nervous system. *Nat Genet*, 1999. **21**(4): p. 396-9.
122. Liliom, K., F. Orosz, L. Horvath and J. Ovadi, Quantitative evaluation of indirect ELISA. Effect of calmodulin antagonists on antibody binding to calmodulin. *J Immunol Methods*, 1991. **143**(1): p. 119-25.
123. Liu, R., J.Q. Guan, O. Zak, P. Aisen and M.R. Chance, Structural reorganization of the transferrin C-lobe and transferrin receptor upon complex formation: the C-lobe binds to the receptor helical domain. *Biochemistry*, 2003. **42**(43): p. 12447-54.
124. Lok, C.N., K.F. Chan and T.T. Loh, Transcriptional regulation of transferrin receptor expression during phorbol-ester-induced HL-60 cell differentiation. Evidence for a negative regulatory role of the phorbol-ester-responsive element-like sequence. *Eur J Biochem*, 1996. **236**(2): p. 614-9.
125. Lombard, M., A. Bomford, M. Hynes, N.V. Naoumov, S. Roberts, et al., Regulation of the hepatic transferrin receptor in hereditary hemochromatosis. *Hepatology*, 1989. **9**(1): p. 1-5.
126. Lottspeich, F. and H. Zorbas, *Bioanalytik*. 1998, Heidelberg Berlin: Spektrum Akademischer Verlag.
127. Ludeman, M.J., Y.W. Zheng, K. Ishii and S.R. Coughlin, Regulated shedding of PAR1 N-terminal exodomain from endothelial cells. *J Biol Chem*, 2004. **279**(18): p. 18592-9.
128. Lunn, C.A., X. Fan, B. Dalie, K. Miller, P.J. Zavodny, et al., Purification of ADAM 10 from bovine spleen as a TNFalpha convertase. *FEBS Lett*, 1997. **400**(3): p. 333-5.
129. MacGillivray, R.T., S.A. Moore, J. Chen, B.F. Anderson, H. Baker, et al., Two high-resolution crystal structures of the recombinant N-lobe of human transferrin reveal a structural change implicated in iron release. *Biochemistry*, 1998. **37**(22): p. 7919-28.
130. May, W.S. and G. Tyler, Phosphorylation of the surface transferrin receptor stimulates receptor internalization in HL60 leukemic cells. *J Biol Chem*, 1987. **262**(34): p. 16710-8.
131. McGraw, T.E., K.W. Dunn and F.R. Maxfield, Phorbol ester treatment increases the exocytic rate of the transferrin receptor recycling pathway independent of serine-24 phosphorylation. *J Cell Biol*, 1988. **106**(4): p. 1061-6.
132. McGraw, T.E., L. Greenfield and F.R. Maxfield, Functional expression of the human transferrin receptor cDNA in Chinese hamster ovary cells deficient in endogenous transferrin receptor. *J Cell Biol*, 1987. **105**(1): p. 207-14.
133. McKie, A.T., D. Barrow, G.O. Latunde-Dada, A. Rolfs, G. Sager, et al., An iron-regulated ferric reductase associated with the absorption of dietary iron. *Science*, 2001. **291**(5509): p. 1755-9.
134. McKie, A.T., P. Marciani, A. Rolfs, K. Brennan, K. Wehr, et al., A novel duodenal iron-regulated transporter, IREG1, implicated in the basolateral transfer of iron to the circulation. *Mol Cell*, 2000. **5**(2): p. 299-309.
135. Michelini, E., M. Mirasoli, M. Karp, M. Virta and A. Roda, Development of a Bioluminescence Resonance Energy-Transfer Assay for Estrogen-Like Compound in Vivo Monitoring. *Anal Chem*, 2004. **76**(23): p. 7069-7076.
136. Miskimins, W.K., A. McClelland, M.P. Roberts and F.H. Ruddle, Cell proliferation and expression of the transferrin receptor gene: promoter sequence homologies and protein interactions. *J Cell Biol*, 1986. **103**(5): p. 1781-8.
137. Mohler, K.M., P.R. Sleath, J.N. Fitzner, D.P. Cerretti, M. Alderson, et al., Protection against a lethal dose of endotoxin by an inhibitor of tumour necrosis factor processing. *Nature*, 1994. **370**(6486): p. 218-20.
138. Montosi, G., P. Paglia, C. Garuti, C.A. Guzman, J.M. Bastin, et al., Wild-type HFE protein normalizes transferrin iron accumulation in macrophages from subjects with hereditary hemochromatosis. *Blood*, 2000. **96**(3): p. 1125-9.
139. Moss, M.L., S.L. Jin, M.E. Milla, D.M. Bickett, W. Burkhart, et al., Cloning of a disintegrin metalloproteinase that processes precursor tumour-necrosis factor-alpha. *Nature*, 1997. **385**(6618): p. 733-6.
140. Mullberg, J., W. Oberthur, F. Lottspeich, E. Mehl, E. Dittrich, et al., The soluble human IL-6 receptor. Mutational characterization of the proteolytic cleavage site. *J Immunol*, 1994. **152**(10): p. 4958-68.
141. Nemeth, E., M.S. Tuttle, J. Powelson, M.B. Vaughn, A. Donovan, et al., Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science*, 2004. **306**(5704): p. 2090-3.

142. Nemeth, E., E.V. Valore, M. Territo, G. Schiller, A. Lichtenstein, et al., Hepcidin, a putative mediator of anemia of inflammation, is a type II acute-phase protein. *Blood*, 2003. **101**(7): p. 2461-3.
143. Nicolas, G., M. Bennoun, I. Devaux, C. Beaumont, B. Grandchamp, et al., Lack of hepcidin gene expression and severe tissue iron overload in upstream stimulatory factor 2 (USF2) knockout mice. *Proc Natl Acad Sci U S A*, 2001. **98**(15): p. 8780-5.
144. Nicolas, G., M. Bennoun, A. Porteu, S. Mativet, C. Beaumont, et al., Severe iron deficiency anemia in transgenic mice expressing liver hepcidin. *Proc Natl Acad Sci U S A*, 2002. **99**(7): p. 4596-601.
145. Nicolas, G., L. Viatte, D.Q. Lou, M. Bennoun, C. Beaumont, et al., Constitutive hepcidin expression prevents iron overload in a mouse model of hemochromatosis. *Nat Genet*, 2003. **34**(1): p. 97-101.
146. Olivares, M., T. Walter, J.D. Cook, E. Hertrampf and F. Pizarro, Usefulness of serum transferrin receptor and serum ferritin in diagnosis of iron deficiency in infancy. *Am J Clin Nutr*, 2000. **72**(5): p. 1191-5.
147. Omary, M.B. and I.S. Trowbridge, Covalent binding of fatty acid to the transferrin receptor in cultured human cells. *J Biol Chem*, 1981. **256**(10): p. 4715-8.
148. Osaki, S., D.A. Johnson and E. Frieden, The mobilization of iron from the perfused mammalian liver by a serum copper enzyme, ferroxidase I. *J Biol Chem*, 1971. **246**(9): p. 3018-23.
149. Paez, P.M., C.I. Garcia, C. Davio, A.T. Campagnoni, E.F. Soto, et al., Apotransferrin promotes the differentiation of two oligodendroglial cell lines. *Glia*, 2004. **46**(2): p. 207-17.
150. Pan, D. and G.M. Rubin, Kuzbanian controls proteolytic processing of Notch and mediates lateral inhibition during *Drosophila* and vertebrate neurogenesis. *Cell*, 1997. **90**(2): p. 271-80.
151. Pang, S., A.J. Chubb, S.L. Schwager, M.R. Ehlers, E.D. Sturrock, et al., Roles of the juxtamembrane and extracellular domains of angiotensin-converting enzyme in ectodomain shedding. *Biochem J*, 2001. **358**(Pt 1): p. 185-92.
152. Papanikolaou, G., M.E. Samuels, E.H. Ludwig, M.L. MacDonald, P.L. Franchini, et al., Mutations in HFE2 cause iron overload in chromosome 1q-linked juvenile hemochromatosis. *Nat Genet*, 2004. **36**(1): p. 77-82.
153. Park, C.H., E.V. Valore, A.J. Waring and T. Ganz, Hepcidin, a urinary antimicrobial peptide synthesized in the liver. *J Biol Chem*, 2001. **276**(11): p. 7806-10.
154. Parkin, E.T., F. Tan, R.A. Skidgel, A.J. Turner and N.M. Hooper, The ectodomain shedding of angiotensin-converting enzyme is independent of its localisation in lipid rafts. *J Cell Sci*, 2003. **116**(Pt 15): p. 3079-87.
155. Parkkila, S., A. Waheed, R.S. Britton, B.R. Bacon, X.Y. Zhou, et al., Association of the transferrin receptor in human placenta with HFE, the protein defective in hereditary hemochromatosis. *Proc Natl Acad Sci U S A*, 1997. **94**(24): p. 13198-202.
156. Pei, D., Identification and characterization of the fifth membrane-type matrix metalloproteinase MT5-MMP. *J Biol Chem*, 1999. **274**(13): p. 8925-32.
157. Peschon, J.J., D.S. Torrance, K.L. Stocking, M.B. Glaccum, C. Otten, et al., TNF receptor-deficient mice reveal divergent roles for p55 and p75 in several models of inflammation. *J Immunol*, 1998. **160**(2): p. 943-52.
158. Pigeon, C., G. Ilyin, B. Courselaud, P. Leroyer, B. Turlin, et al., A new mouse liver-specific gene, encoding a protein homologous to human antimicrobial peptide hepcidin, is overexpressed during iron overload. *J Biol Chem*, 2001. **276**(11): p. 7811-9.
159. Pikul, S., K.L. McDow Dunham, N.G. Almstead, B. De, M.G. Natchus, et al., Discovery of potent, achiral matrix metalloproteinase inhibitors. *J Med Chem*, 1998. **41**(19): p. 3568-71.
160. Powell, W.C., B. Fingleton, C.L. Wilson, M. Boothby and L.M. Matrisian, The metalloproteinase matrilysin proteolytically generates active soluble Fas ligand and potentiates epithelial cell apoptosis. *Curr Biol*, 1999. **9**(24): p. 1441-7.
161. Punnonen, K., K. Irjala and A. Rajamaki, Serum transferrin receptor and its ratio to serum ferritin in the diagnosis of iron deficiency. *Blood*, 1997. **89**(3): p. 1052-7.
162. Raja, K.B., D.J. Pountney, R.J. Simpson and T.J. Peters, Importance of anemia and transferrin levels in the regulation of intestinal iron absorption in hypotransferrinemic mice. *Blood*, 1999. **94**(9): p. 3185-92.
163. Ramalingam, T.S., A.P. West, Jr., J.A. Lebron, J.S. Nangiana, T.H. Hogan, et al., Binding to the transferrin receptor is required for endocytosis of HFE and regulation of iron homeostasis. *Nat Cell Biol*, 2000. **2**(12): p. 953-7.
164. Richter, G.W., Studies of iron overload. Lysosomal proteolysis of rat liver ferritin. *Pathol Res Pract*, 1986. **181**(2): p. 159-67.
165. Robache-Gallea, S., V. Morand, J.M. Bruneau, B. Schoot, E. Tagat, et al., In vitro processing of human tumor necrosis factor-alpha. *J Biol Chem*, 1995. **270**(40): p. 23688-92.
166. Roberts, M.R., Y. Han, A. Fienberg, L. Hunihan and F.H. Ruddle, A DNA-binding activity, TRAC, specific for the TRA element of the transferrin receptor gene copurifies with the Ku autoantigen. *Proc Natl Acad Sci U S A*, 1994. **91**(14): p. 6354-8.
167. Roetto, A., A. Totaro, A. Piperno, A. Piga, F. Longo, et al., New mutations inactivating transferrin receptor 2 in hemochromatosis type 3. *Blood*, 2001. **97**(9): p. 2555-60.
168. Rothenberger, S., B.J. Iacopetta and L.C. Kuhn, Endocytosis of the transferrin receptor requires the cytoplasmic domain but not its phosphorylation site. *Cell*, 1987. **49**(3): p. 423-31.
169. Roy, C.N., D.M. Penny, J.N. Feder and C.A. Enns, The hereditary hemochromatosis protein, HFE, specifically regulates transferrin-mediated iron uptake in HeLa cells. *J Biol Chem*, 1999. **274**(13): p. 9022-8.
170. Rutledge, E.A. and C.A. Enns, Cleavage of the transferrin receptor is influenced by the composition of the O-linked carbohydrate at position 104. *J Cell Physiol*, 1996. **168**(2): p. 284-93.
171. Rutledge, E.A., I. Gaston, B.J. Root, T.E. McGraw and C.A. Enns, The transferrin receptor cytoplasmic domain determines its rate of transport through the biosynthetic pathway and its susceptibility to cleavage early in the pathway. *J Biol Chem*, 1998. **273**(20): p. 12169-75.

172. Rutledge, E.A., F.A. Green and C.A. Enns, Generation of the soluble transferrin receptor requires cycling through an endosomal compartment. *J Biol Chem*, 1994. **269**(50): p. 31864-8.
173. Rutledge, E.A., B.J. Root, J.J. Lucas and C.A. Enns, Elimination of the O-linked glycosylation site at Thr 104 results in the generation of a soluble human-transferrin receptor. *Blood*, 1994. **83**(2): p. 580-6.
174. R'Zik, S. and Y. Beguin, Serum soluble transferrin receptor concentration is an accurate estimate of the mass of tissue receptors. *Exp Hematol*, 2001. **29**(6): p. 677-85.
175. Sadhukhan, R., G.C. Sen, R. Ramchandran and I. Sen, The distal ectodomain of angiotensin-converting enzyme regulates its cleavage-secretion from the cell surface. *Proc Natl Acad Sci U S A*, 1998. **95**(1): p. 138-43.
176. Salinas, M., J. Wang, M. Rosa de Sagarra, D. Martin, A.I. Rojo, et al., Protein kinase Akt/PKB phosphorylates heme oxygenase-1 in vitro and in vivo. *FEBS Lett*, 2004. **578**(1-2): p. 90-4.
177. Salter-Cid, L., A. Brunmark, P.A. Peterson and Y. Yang, The major histocompatibility complex-encoded class I-like HFE abrogates endocytosis of transferrin receptor by inducing receptor phosphorylation. *Genes Immun*, 2000. **1**(7): p. 409-17.
178. Sanger, F., S. Nicklen and A.R. Coulson, DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A*, 1977. **74**(12): p. 5463-7.
179. Schlondorff, J., L. Lum and C.P. Blobel, Biochemical and pharmacological criteria define two shedding activities for TRANCE/OPGL that are distinct from the tumor necrosis factor alpha convertase. *J Biol Chem*, 2001. **276**(18): p. 14665-74.
180. Schwager, S.L., A.J. Chubb, R.R. Scholle, W.F. Brandt, R. Mentele, et al., Modulation of juxtamembrane cleavage ("shedding") of angiotensin-converting enzyme by stalk glycosylation: evidence for an alternative shedding protease. *Biochemistry*, 1999. **38**(32): p. 10388-97.
181. Seiser, C., S. Teixeira and L.C. Kuhn, Interleukin-2-dependent transcriptional and post-transcriptional regulation of transferrin receptor mRNA. *J Biol Chem*, 1993. **268**(18): p. 13074-80.
182. Shih, Y.J., R.D. Baynes, B.G. Hudson, C.H. Flowers, B.S. Skikne, et al., Serum transferrin receptor is a truncated form of tissue receptor. *J Biol Chem*, 1990. **265**(31): p. 19077-81.
183. Shimada, T., H. Nakamura, E. Ohuchi, Y. Fujii, Y. Murakami, et al., Characterization of a truncated recombinant form of human membrane type 3 matrix metalloproteinase. *Eur J Biochem*, 1999. **262**(3): p. 907-14.
184. Shintani, N., Y. Kohgo, J. Kato, H. Kondo, K. Fujikawa, et al., Expression and extracellular release of transferrin receptors during peripheral erythroid progenitor cell differentiation in liquid culture. *Blood*, 1994. **83**(5): p. 1209-15.
185. Shirakabe, K., S. Wakatsuki, T. Kurisaki and A. Fujisawa-Sehara, Roles of Meltrin beta /ADAM19 in the processing of neuregulin. *J Biol Chem*, 2001. **276**(12): p. 9352-8.
186. Skikne, B.S., C.H. Flowers and J.D. Cook, Serum transferrin receptor: a quantitative measure of tissue iron deficiency. *Blood*, 1990. **75**(9): p. 1870-6.
187. Slack, B.E., L.K. Ma and C.C. Seah, Constitutive shedding of the amyloid precursor protein ectodomain is up-regulated by tumour necrosis factor-alpha converting enzyme. *Biochem J*, 2001. **357**(Pt 3): p. 787-94.
188. Sorensen, K., One reagent simultaneously identifies the salt and the protein peaks on desalting column. *Biotechniques*, 1992. **12**(2): p. 235-6.
189. Suominen, P., A. Virtanen, M. Lehtonen-Veromaa, O.J. Heinonen, T.T. Salmi, et al., Regression-based reference limits for serum transferrin receptor in children 6 months to 16 years of age. *Clin Chem*, 2001. **47**(5): p. 935-7.
190. Suzuki, M., G. Raab, M.A. Moses, C.A. Fernandez and M. Klagsbrun, Matrix metalloproteinase-3 releases active heparin-binding EGF-like growth factor by cleavage at a specific juxtamembrane site. *J Biol Chem*, 1997. **272**(50): p. 31730-7.
191. Taetle, R., S. Ralph, S. Smedsrud and I. Trowbridge, Regulation of transferrin receptor expression in myeloid leukemia cells. *Blood*, 1987. **70**(3): p. 852-9.
192. Tanaka, M., D. Nanba, S. Mori, F. Shiba, H. Ishiguro, et al., ADAM binding protein Eve-1 is required for ectodomain shedding of epidermal growth factor receptor ligands. *J Biol Chem*, 2004. **279**(40): p. 41950-9.
193. Tang, P., M.C. Hung and J. Klostergaard, Length of the linking domain of human pro-tumor necrosis factor determines the cleavage processing. *Biochemistry*, 1996. **35**(25): p. 8226-33.
194. Tanida, S., T. Joh, K. Itoh, H. Kataoka, M. Sasaki, et al., The mechanism of cleavage of EGFR ligands induced by inflammatory cytokines in gastric cancer cells. *Gastroenterology*, 2004. **127**(2): p. 559-69.
195. Taylor, R.M., J.B. Burritt, D. Baniulis, T.R. Foubert, C.I. Lord, et al., Site-Specific Inhibitors of NADPH Oxidase Activity and Structural Probes of Flavocytochrome b: Characterization of Six Monoclonal Antibodies to the p22phox Subunit. *J Immunol*, 2004. **173**(12): p. 7349-57.
196. Thomas, L., Labor und Diagnose. 5 ed. Vol. 1. 1998, Frankfurt: TH-Books Verlagsgesellschaft. 1567.
197. Thorstensen, K. and I. Romslo, Measurement of serum transferrin receptors in screening for hemochromatosis. *Clin Chem*, 1992. **38**(8 Pt 1): p. 1510.
198. Toki, F., D. Nanba, N. Matsuura and S. Higashiyama, Ectodomain shedding of membrane-anchored heparin-binding EGF like growth factor and subcellular localization of the C-terminal fragment in the cell cycle. *J Cell Physiol*, 2004.
199. Tong, X., H. Kawabata and H.P. Koefler, Iron deficiency can upregulate expression of transferrin receptor at both the mRNA and protein level. *Br J Haematol*, 2002. **116**(2): p. 458-64.
200. Towe, K.M., Structural distinction between ferritin and iron-dextran (imferon). An electron diffraction comparison. *J Biol Chem*, 1981. **256**(18): p. 9377-8.
201. Townsend, A. and H. Drakesmith, Role of HFE in iron metabolism, hereditary haemochromatosis, anaemia of chronic disease, and secondary iron overload. *Lancet*, 2002. **359**(9308): p. 786-90.
202. Trinder, D. and E. Baker, Transferrin receptor 2: a new molecule in iron metabolism. *Int J Biochem Cell Biol*, 2003. **35**(3): p. 292-6.

203. Turkewitz, A.P., J.F. Amatruda, D. Borhani, S.C. Harrison and A.L. Schwartz, A high yield purification of the human transferrin receptor and properties of its major extracellular fragment. *J Biol Chem*, 1988. **263**(17): p. 8318-25.
204. Uchiyama-Tanaka, Y., H. Matsubara, Y. Nozawa, S. Murasawa, Y. Mori, et al., Angiotensin II signaling and HB-EGF shedding via metalloproteinase in glomerular mesangial cells. *Kidney Int*, 2001. **60**(6): p. 2153-63.
205. Urban, S. and M. Freeman, Intramembrane proteolysis controls diverse signalling pathways throughout evolution. *Curr Opin Genet Dev*, 2002. **12**(5): p. 512-8.
206. Vey, M., W. Schafer, S. Berghofer, H.D. Klenk and W. Garten, Maturation of the trans-Golgi network protease furin: compartmentalization of propeptide removal, substrate cleavage, and COOH-terminal truncation. *J Cell Biol*, 1994. **127**(6 Pt 2): p. 1829-42.
207. Vincent, B., E. Paitel, P. Saftig, Y. Frobert, D. Hartmann, et al., The disintegrins ADAM10 and TACE contribute to the constitutive and phorbol ester-regulated normal cleavage of the cellular prion protein. *J Biol Chem*, 2001. **276**(41): p. 37743-6.
208. Virtanen, M.A., L.U. Viinikka, M.K. Virtanen, J.C. Svahn, R.M. Anttila, et al., Higher concentrations of serum transferrin receptor in children than in adults. *Am J Clin Nutr*, 1999. **69**(2): p. 256-60.
209. Vulpe, C.D., Y.M. Kuo, T.L. Murphy, L. Cowley, C. Askwith, et al., Hephhaestin, a ceruloplasmin homologue implicated in intestinal iron transport, is defective in the sla mouse. *Nat Genet*, 1999. **21**(2): p. 195-9.
210. Waheed, A., S. Parkkila, J. Saarnio, R.E. Fleming, X.Y. Zhou, et al., Association of HFE protein with transferrin receptor in crypt enterocytes of human duodenum. *Proc Natl Acad Sci U S A*, 1999. **96**(4): p. 1579-84.
211. Waheed, A., S. Parkkila, X.Y. Zhou, S. Tomatsu, Z. Tsuchihashi, et al., Hereditary hemochromatosis: effects of C282Y and H63D mutations on association with beta2-microglobulin, intracellular processing, and cell surface expression of the HFE protein in COS-7 cells. *Proc Natl Acad Sci U S A*, 1997. **94**(23): p. 12384-9.
212. Walev, I., P. Vollmer, M. Palmer, S. Bhakdi and S. Rose-John, Pore-forming toxins trigger shedding of receptors for interleukin 6 and lipopolysaccharide. *Proc Natl Acad Sci U S A*, 1996. **93**(15): p. 7882-7.
213. Wandzik, K., Untersuchungen zum Shedding des Transferrinrezeptors und anderer Membranproteine - Einfluß von TIMPs und ADAMs. Diplomarbeit. 2003, Berlin: Freie Universität Berlin.
214. Ward, R.J., R. Legssyer, C. Henry and R.R. Crichton, Does the haemosiderin iron core determine its potential for chelation and the development of iron-induced tissue damage? *J Inorg Biochem*, 2000. **79**(1-4): p. 311-7.
215. Weskamp, G., J. Schlondorff, L. Lum, J.D. Becherer, T.W. Kim, et al., Evidence for a critical role of the tumor necrosis factor alpha convertase (TACE) in ectodomain shedding of the p75 neurotrophin receptor (p75NTR). *J Biol Chem*, 2004. **279**(6): p. 4241-9.
216. West, A.P., Jr., M.J. Bennett, V.M. Sellers, N.C. Andrews, C.A. Enns, et al., Comparison of the interactions of transferrin receptor and transferrin receptor 2 with transferrin and the hereditary hemochromatosis protein HFE. *J Biol Chem*, 2000. **275**(49): p. 38135-8.
217. West, A.P., Jr., A.M. Giannetti, A.B. Herr, M.J. Bennett, J.S. Nangiana, et al., Mutational analysis of the transferrin receptor reveals overlapping HFE and transferrin binding sites. *J Mol Biol*, 2001. **313**(2): p. 385-97.
218. Will, H., S.J. Atkinson, G.S. Butler, B. Smith and G. Murphy, The soluble catalytic domain of membrane type 1 matrix metalloproteinase cleaves the propeptide of progelatinase A and initiates autoproteolytic activation. Regulation by TIMP-2 and TIMP-3. *J Biol Chem*, 1996. **271**(29): p. 17119-23.
219. Yamaji, S., P. Sharp, B. Ramesh and S.K. Srari, Inhibition of iron transport across human intestinal epithelial cells by hepcidin. *Blood*, 2004.
220. Yamashiro, D.J., B. Tycko, S.R. Fluss and F.R. Maxfield, Segregation of transferrin to a mildly acidic (pH 6.5) para-Golgi compartment in the recycling pathway. *Cell*, 1984. **37**(3): p. 789-800.
221. Yang, B., M.H. Hoe, P. Black and R.C. Hunt, Role of oligosaccharides in the processing and function of human transferrin receptors. Effect of the loss of the three N-glycosyl oligosaccharides individually or together. *J Biol Chem*, 1993. **268**(10): p. 7435-41.
222. Yefimova, M.G., I.S. Shcherbakova and N.D. Shushakova, Transferrin and ferritin modulate the activity of brain calcium-calmodulin-dependent phosphodiesterase. *Biochemistry (Mosc)*, 1997. **62**(2): p. 165-70.
223. Zerial, M., M. Suomalainen, M. Zanetti-Schneider, C. Schneider and H. Garoff, Phosphorylation of the human transferrin receptor by protein kinase C is not required for endocytosis and recycling in mouse 3T3 cells. *Embo J*, 1987. **6**(9): p. 2661-7.
224. Zhang, A.S., P.S. Davies, H.L. Carlson and C.A. Enns, Mechanisms of HFE-induced regulation of iron homeostasis: Insights from the W81A HFE mutation. *Proc Natl Acad Sci U S A*, 2003. **100**(16): p. 9500-5.
225. Zheng, Y., P. Saftig, D. Hartmann and C. Blobel, Evaluation of the contribution of different ADAMs to tumor necrosis factor alpha (TNFalpha) shedding and of the function of the TNFalpha ectodomain in ensuring selective stimulated shedding by the TNFalpha convertase (TACE/ADAM17). *J Biol Chem*, 2004. **279**(41): p. 42898-906.
226. Zhou, X.Y., S. Tomatsu, R.E. Fleming, S. Parkkila, A. Waheed, et al., HFE gene knockout produces mouse model of hereditary hemochromatosis. *Proc Natl Acad Sci U S A*, 1998. **95**(5): p. 2492-7.