

6. LITERATURVERZEICHNIS

Areosa SA, McShane R, Sherriff F. Memantine for dementia. Cochrane Database Syst Rev 2004;4:CD003154.

Bähr M, Vanselow J, Thanos S. In vitro regeneration of adult rat ganglion cell axons from retinal explants. Exp Brain Res 1988;73:393-401.

Bähr M, Bunge RP. Growth of adult retinal ganglion cell neurites on astrocytes. Glia 1990;3:293-300.

Berkelaar M, Clarke DB, Wang YC, Bray GM, Aguayo AJ. Axotomy results in delayed death and apoptosis of retinal ganglion cells in adult rats. J Neurosci 1994;14(7): 4368-4374.

Bernhardt R, Matus A. Light and electron microscopic studies of the distribution of microtubule-assiated protein 2 in rat brain: a difference between dendritic and axonal cytoskeletons. J Comp Neurol 1984;226(2):203-221.

Bickler PE, Buck LT. Adaptations of vertebrate neurons to hypoxia and anoxia: maintaining critical Ca^{2+} concentrations. J Exp Biol 1998;201:1141-1152.

Bignami A, Dahl D. Specificity of the glial fibrillary acidic protein for astroglia. J Histochem Cytochem 1977;25(6):466-469.

Blauw GJ, Lagaay AM, Smelt AHM, Westendorp RGJ. Stroke, statins, and cholesterol. A meta-analysis of randomized, placebo-controlled, double-blind trials with HMG-CoA reductase inhibitors. Stroke 1997;28:946-950.

Bösel J, Gandor F, Harms C, et al. Neuroprotective effects of atorvastatin against glutamate-induced excitotoxicity in primary cortical neurones. J Neurochem 2005;92(6):1386-1398.

Bolshakov KV, Kim KH, Potapjeva NN, et al. Design of antagonists for NMDA and AMPA receptors. *Neuropharm* 2005;49:144-155.

Boycott BB, Hopkins JM. Microglia in the retina of monkey and other mammals; its distinction from other types of glia and horizontal cells. *Neurosci* 1981;6:679-688.

Braestrup C, Nielsen M, Olsen CE. Urinary and brain beta-carboline-3-carboxylates as potent inhibitors of brain benzodiazepine receptors. *Proc Natl Acad Sci USA* 1980;77(4):2288-2292.

Brandstätter JH. Glutamate receptor in the rat retina: the molecular substrate for visual signal processing. *Curr Eye Res* 2002;25:327-331.

Caceres A, Binder LI, Payne MR, Bender P, Rebhun L, Steward O. Differential subcellular localization of tubulin and the microtubule-associated protein MAP2 in brain tissue as revealed by immunocytochemistry with monoclonal hybridoma antibodies. *J Neurosci* 1984;4(2):394-410.

Caffé AR, Ahuja P, Holmqvist B, et al. Mouse retina explants after long-term culture in serum free medium. *J Chem Neuroanat* 2001;22:263-273.

Caprioli J, Ishii Y, Kwong JMK. Retinal ganglion cell protection with geranylgeranyl-acetone, a heat shock protein inducer, in a rat glaucoma model. *Trans Am Ophthalmol Soc* 2003;101:39-50.

Carri NG, Ebendal T. Organotypic cultures of neural retina: Neurite outgrowth stimulated by brain extracts. *Dev Brain Res* 1983;6:219-229.

Choi DW. Calcium: still center-stage in hypoxic-ischemic neuronal death. *TINS* 1995; 18:58-60.

Crooks J, Kolb H. Localization of GABA, glycine, glutamate and tyrosine hydroxylase in the human retina. *J Comp Neurol* 1992;315:287-302.

Davanger S, Ottersen OP, Storm-Mathisen J. Glutamate, GABA and glycine in the human retina: An immunocytochemical investigation. *J Comp Neurol* 1991;311: 483-494.

Dirnagl U, Iadecola C, Moskowitz MA. Pathobiology of ischaemic stroke: an integrated view. *TINS* 1999;22:391-397.

Eccles JC, Schmidt RF, Willis WD. Pharmacological studies on presynaptic inhibition. *J Physiol* 1963; 185:298-322.

Eder M, Rammes G, Zieglgansberger W, Dodt HU. GABA_A and GABA_B receptors on neocortical neurons are differentially distributed. *Eur J Neurosci* 2001; 13:1065-1069.

Eichler W, Yafai Y, Keller T, Wiedemann P, Reichenbach A. PEDF derived from glial Müller cells: a possible regulator of retinal angiogenesis. *Exp Cell Res* 2004;299: 68-78.

Endres M, Laufs U, Huang Z, et al. Stroke protection by 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase inhibitors mediated by endothelial nitric oxide synthase. *Proc Natl Acad Sci USA* 1998;95:8880-8885.

Engelsberg K, Ehinger B, Wassélius J, Johansson K. Apoptotic cell death and microglial cell responses in cultured rat retina. *Graefe's Arch Clin Exp Ophthalmol* 2004; 242:229-239.

Engelsberg K, Johansson K, Ghosh F. Development of the embryonic porcine neuroretina in vitro. *Ophthalmic Res* 2005;37:104-111.

Ferreira IL, Duarte CB, Carvalho AP. Ca²⁺ influx through glutamate receptor-associated channels in retina cells correlates with neuronal cell death. *Eur J Pharmacol* 1996;302(1-3):153-162.

Garner CC, Tucker RP, Matus A. Selective localization of messenger RNA for cytoskeletal protein MAP2 in dendrites. *Nature* 1988;336:674-677.

Gerson RJ, MacDonald JS, Alberts AW, et al. Animal safety and toxicology of simvastatin and related hydroxy-methylglutaryl-coenzyme A reductase inhibitors: *Am J Med* 1989;87:28-38.

Garcia M, Forster V, Hicks D, Vecino E. Effects of Müller glia on cell survival and neuritogenesis in adult porcine retina in vitro. *Invest Ophthalmol Vis Sci* 2002;43: 3735-3743.

Garcia-Valenzuela E, Gorczyca W, Darzynkiewicz, Sharma SC. Apoptosis in adult retinal ganglion cells after axotomy. *J Neurobiol* 1994;25(4):431-438.

Gavrieli Y, Sherman Y, Ben-Sasson SA. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol* 1992; 119(3):493-501.

Goldblum D, Mittag T. Prospects for relevant glaucoma models with retinal ganglion cell damage in the rodent eye. *Vision Res* 2002;42:471-478.

Gralnick HR, Coller BS. Studies of the human factor VIII / von Willebrand's factor protein. II. Identification and characterization of the von Willebrand protein. *Blood* 1975;46(3):417-430.

Guerin CJ, Anderson DH, Fisher SK. Changes in intermediate filament immunolabeling occur in response to retinal detachment and reattachment in primates. *Invest Ophthalmol Vis Sci* 1990;31(8):1474-1482.

Haefely W, Kulcsar A, Mohler H, Pieri L, Polc P, Schaffner R. Possible involvement of GABA in the central actions of benzodiazepines. *Adv Biochem Psychopharmacol* 1975; 14:131-151.

- Hess DC, Demchuk AM, Brass LM, Yatsu FM. HMG-CoA reductase inhibitors (statins). A promising approach to stroke prevention. *Neurology* 2000;54:790-796.
- Hild W, Callas G. The behavior of retinal tissue in-vitro, light and electron microscopic observations. *Z Zellforsch* 1967;80:1-21.
- Hoff A, Hämerle H, Schlosshauer B. Organotypic culture system of chicken retina. *Brain Res Protoc* 1999;4:237-248.
- Horsburgh GM, Sefton AJ. Cellular degeneration and synaptogenesis in the developing retina of the rat. *J Comp Neurol* 1987;263:553-566.
- Ikonomovic MD, Mizukami K, Warde D, et al. Distribution of glutamate receptor subunit NMDAR1 in the hippocampus of normal elderly and patients with Alzheimer's disease. *Exp Neurol* 1999;160(1):194-204.
- Izumi Y, Shimamoto K, Benz AM, Hammerman SB, Olney JW, Zarumski CF. Glutamate transporters and retinal excitotoxicity. *Glia* 2002;39:58-68.
- Jacoby RA, Wu SM. AMPA-preferring receptors mediate excitatory non-NMDA responses of primate retinal ganglion cells. *Vis Neurosci* 2001;18:703-710.
- Kapuscinski J, Skoczylas B. Fluorescent complexes of DNA with DAPI 4,6-diamidine-2-phenyl indole.2HCl or DCI 4,6-dicarboxyamide-2-phenyl indole. *Nucl Acids Res* 1978;5:3775-3799.
- Kashiwagi K, Iizuka Y, Araie M, Suzuki Y, Tsukahara S. Effects of retinal glial cells on isolated rat retinal ganglion cells. *Invest Ophthalmol Vis Sci* 2001;42:2686-2694.
- Keinanen K, Wisden W, Sommer B, et al. A family of AMPA-selective glutamate receptors. *Science* 1990;249:556-560.

Kermer P, Klocker N, Weishaupt JH, Bahr M. Transsection of the optic nerve in rats: studying neuronal death and survival in vivo. *Brain Res Protoc* 2001;7:255-260.

Kerr JF, Wyllie AH, Currie AR. Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer* 1972;26(4):239-257.

Kim SU, Takahashi H. Tissue culture study of adult human retina neurons. *Invest Ophthalmol Vis Sci* 1988;29:1372-1379.

Knoflach F, Drescher U, Scheurer L, Malherbe P, Mohler H. Full and partial agonism displayed by benzodiazepine receptor ligands at recombinant γ -aminobutyric acid_A receptor subtypes. *J Pharmacol Exp Ther* 1993;266:385-391.

Knott RM, Robertson M, Muckersie E, et al. A model system for the study of human retinal angiogenesis: activation of monocytes and endothelial cells and the association with the expression of the monocarboxylate transporter type 1 (MCT-1). *Diabetologia* 1999;42:870-877.

Kolb H. How the retina works. *Am Sci* 2003;91:28-35.

Kretz A, Hermening SH, Isenmann S. A novel primary technique for adult retina allows for evaluation of CNS axon regeneration in rodents: *J Neurosci Methods* 2004; 136:207-219.

Kretz A, Schmeer C, Tausch S, Isenmann S. Simvastatin promotes heat shock protein 27 expression and Akt activation in the rat retina and protects axotomized retinal ganglion cells in vivo. *Neurobiol Dis* 2006;21:421-430.

Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *TINS* 1996;19: 312-318.

Kumpulainen T, Dahl D, Korhonen K, Nyström SHM. Immunolabeling of carbionic anhydrase isoenzyme C and glial fibrillary acidic protein in paraffin-embedded

tissue sections of human brain and retina. J Histochem Cytochem 1983;31(7): 879-886.

Lam TT, Abler AS, Kwong JMK, Tso MOM. N-methyl-D-aspartate (NMDA)-induced apoptosis in rat retina. Invest Ophthalmol Vis Sci 1999;40:2391-2397.

Laufs U, Gertz K, Huang P, et al. Atorvastatin upregulates type III nitric oxide synthase in thrombocytes, decreases platelet activation, and protects from cerebral ischemia in normocholesterolemic mice. Stroke 2000;31:2437-2449.

LaVail MM, Hild W. Histotypic organization of the rat retina in vitro. Z Zellforsch 1971;114:557-559.

Levin LA. Animal and culture models of glaucoma for studying neuroprotection. Eur J Ophthalmol 2001;11(2):S23-S29.

Liao JK. Beyond lipid lowering: the role of statins in vascular protection. Int J Cardiol 2002;86:15-18.

Linden R, Esbérard CEL. Displaced amacrine cells in the ganglion cell layer of the hamster retina. Vision Res 1987;27(7):1071-1076.

Lipton, S. A., 2003. Possible role for memantine in protecting retinal ganglion cells from glaucomatous damage. Surv Ophthalmol 2003;48:S38-46.

Luo X, Heidinger V, Picaud S, et al. Selective excitotoxic degeneration of adult pig retinal ganglion cells in vitro. Invest Ophthalmol Vis Sci 2001;42:1096-1106.

Luo X, Baba A, Matsuda T, Romano C. Susceptibilities to and mechanisms of excitotoxic cell death of adult mouse inner retinal neurons in dissociated culture. Invest ophthalmol Vis Sci 2004;45:4576-4582.

Massey SC. Cell types using glutamate as a neurotransmitter in the vertebrate retina.
Prog Ret Res 1990;9:399-425.

Meldrum BS. Glutamate as a neurotransmitter in the brain: review of physiology and pathology. J Nutr 2000; 130:S1007-1015.

Monyer H, Sprengel R, Schoepfer R, et al. Heteromeric NMDA receptors: molecular and functional distinction of subtypes. Science 1992;256:1217-1221.

Mosinger Ogilvie J, Speck JD, Lett JM, Fleming TT. A reliable method for organ culture of neonatal mouse retina with long-term survival. J Neurosci Methods 1999;87: 57-65.

Muir KW, Lees KR. Excitatory amino acid antagonists for acute stroke. Cochrane Database Syst Rev 2003;3:CD001244.

Neufeld G, Cohen T, Gengrinovitch S, Poltorak Z. Vascular endothelial growth factor (VEGF) and its receptors. FASEB J 1999;13:9-22.

Newman E, Reichenbach A. The Müller cell: a functional element of the retina. TINS 1996; 19:307-312.

Osborne NN. Benzodiazepine binding to bovine retina: Neurosci Lett 1980; 16(2):167-170.

Osborne NN, Wood JPM, Chidlow G, Bae J-H, Melena J, Nash MS. Ganglion cell death in glaucoma: what do we really know? Br J Ophthalmol 1999;83:980-986.

Osborne NN, Chidlow G, Layton CJ, Wood JP, Casson RJ, Melena J. Optic nerve and neuroprotection strategies. Eye 2004; 18:1075-1084.

- Pérez-Léon J, Frech MJ, Schröder JE, et al. Spontaneous synaptic activity in an organotypic culture of the mouse retina. *Invest Ophthalmol Vis Sci* 2003;44:1376-87.
- Perry VH. Evidence for an amacrine cell system in the ganglion cell layer of the rat retina. *Neurosci* 1981;6(5):931-944.
- Perry VH, Henderson Z, Linden R. Postnatal changes in retinal ganglion cell and optic axon populations in the pigmented rat. *J Comp Neurol* 1983;219:356-368.
- Pinzón-Duarte G, Kohler K, Arango-Gonzalez B, Guenther E. Cell differentiation, synaptogenesis, and influence of the retinal pigment epithelium in a rat neonatal organotypic retina culture. *Vision Res* 2000;40:3455–3465.
- Pollack MH, Worthington JJ, Manfro GG, Otto MW, Zucker BG. Abecarnil for the treatment of generalized anxiety disorder: a placebo-controlled comparison of two dosage ranges of abecarnil and buspirone. *J Clin Psychiatry* 1997;58:S19-S23.
- Quigley HA, Nickells RW, Kerrigan LA, Pease ME, Thibault DJ, Zack DJ. Retinal ganglion cell death in experimental glaucoma and after axotomy occurs by apoptosis. *Invest Ophthalmol Vis Sci* 1995;36:774-786.
- Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *Br J Ophthalmol* 2006;90:262-267.
- Rabacchi SA, Bonfani L, Liu X-H, Maffei L. Apoptotic cell death induced by optic nerve lesion in the neonatal rat. *J Neurosci* 1994;14:5292-5301.
- Rickels K, DeMartinis N, Aufdembrinke B, 2000. A double-blind, placebo-controlled trial of abecarnil and diazepam in the treatment of patients with generalized anxiety disorder. *J Clin Psychopharmacol* 2000;20:12-18.

Ruscher K, Rzeczinski S, Huang W, Freyer D, Victorov IV, Dirnagl U. Neuroprotective effects of the β -carboline abecarnil studied in cultured cortical neurons and organotypic retinal cultures. *Neuropharm* 2006; in Revision.

Sacks FM, Pfeffer MA, Moye LA, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med* 1996;335:1001-1009.

Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994;344:1383-9.

Schäbitz WR, Li F, Fisher M. The N-methyl-D-aspartate antagonist CNS 1102 protects cerebral gray and white matter from ischemic injury following temporary focal ischemia in rats. *Stroke* 2000;31(7):1709-1714.

Schmeer C, Kretz A, Isenmann S. Statin-mediated protective effects in the central nervous system: General mechanisms and putative role of stress proteins. *Restor Neurol Neurosci* 2006;24:79-95.

Schnitzer J. Astrocytes in the guinea pig, horse, and monkey retina: their occurrence coincides with the presence of blood vessels. *Glia* 1988; 1(1):74-89.

Semkova I, Peters S, Welsandt G, Janicki H, Jordan J, Schraermeyer U. Investigation of laser-induced choroidal neovascularization in the rat. *Invest Ophthalmol Vis Sci* 2003;44:5349-5354.

Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med* 1995;333(20):1301-1307.

Small GW, Bystritsky A. Double-blind, placebo-controlled trial of two doses of abecarnil for geriatric anxiety. *J Clin Psychiatry* 1997;58(11):24-29.

Stein A. Management of dyslipidemia. Am Heart J 2002; 144:S43-50.

Stephens DN, Schneider HH, Kehr W, et al. Abecarnil, a metabolically stable, anxioreactive beta-carboline acting at benzodiazepine receptors. J Pharmacol Exp Ther 1990;253(1):334-343.

Stolc S, Vlkolinsky R, Pavlasek J. Neuroprotection by the pyridoindole stobadine: a minireview. Brain Res Bull 1997;42:335-340.

Stone J, Itin A, Alon T, et al. Development of retinal vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression by neuroglia. J Neurosci 1995; 15(7):4738-4747.

Takano M, Horie H, Iijima Y, Dezava M, Sawada H, Ishikawa Y. Brain derived neurotrophic factor enhances neurite regeneration from retinal ganglion cells in aged human retina in vitro. Exp Eye Res 2002;74:319-323.

Thanos S, Vanselow J. Adult retinal ganglion cells retain the ability to regenerate their axons up to several weeks after axotomy. J Neurosci Res 1989;22(2):144-149.

Vallazza-Deschamps G, Fuchs C, Cia D, et al. Diltiazem-induced neuroprotection in glutamate excitotoxicity and ischemic insult of retinal neurons. Doc Ophthalmol 2005; 110:25-35.

Victorov IV, Lyjin AA, Aleksandrova OP. A modified roller method for organotypic brain cultures: free-floating slices of postnatal rat hippocampus. Brain Res Protoc 2001;7:30-37.

Victorov IV, Lyjin AA, Aleksandrova OP, Freier D, Dirnagl U. Roller organotypic cultures of postnatal retina. Bull Exp Biol Med 2004; 137:419-422.

- Villegas-Pérez MP, Vidal-Sanz M, Rasminksy M, Bray GM, Aguayo AJ. Rapid and protracted phases of retinal ganglion cell loss follow axotomy in the optic nerve of adult rats. *J Neurobiol* 1993;24(1):23-36.
- Vogel J, Möbius C, Kuschinsky W. Early Delineation of ischemic tissue in rat brain cryosections by high-contrast staining. *Stroke* 1999;30:1134-1141.
- Vorobjev VS, Sharanova IN, Skrebitsky VG, Schneider HH, Stephens DN. Abecarnil enhances GABA-induced currents in acutely isolated cerebellar Purkinje cells. *Neuropharm* 1995;34:157-163.
- Wein FB, Levin LA. Current understanding of neuroprotection in glaucoma. *Curr Opin Ophthalmol* 2002;13:61-67.
- Wong EHF, Kemp JA, Priestley T, Knight AR, Woodruff GN, Iversen LL. The anticonvulsant MK-801 is a potent N-methyl-D-aspartate antagonist. *Proc Natl Acad Sci USA* 1986;83:7104-7108.
- Yang XL. Characterization of receptors for glutamate and GABA in retinal neurons. *Progress Neurobiol* 2004;73:127-150.
- Yoneda S, Tanaka E, Goto W, Ota T, Hara H. Topiramate reduces excitotoxic and ischemic injury in the rat retina. *Brain Res* 2003;967:257-266.
- Young RW. Cell death during differentiation of the retina in the mouse. *J Comp Neurol* 1984;229:362-373.
- Zhang Y, Caffe AR, Azadi S, van Veen T, Ehinger B, Perez M-TR. Neuronal integration in an abutting-retinas culture system. *Invest Ophthalmol Vis Sci* 2003;44(11): 4936-4946.