

## V. Literaturverzeichnis

- Abu-Elteen, K.H., Elkarmi, A.Z., and Hamad, M. (2001) Characterization of phenotype-based pathogenic determinants of various *Candida albicans* strains in Jordan. *Jpn J Infect Dis* **54**: 229-236.
- Alarco, A.M., and Raymond, M. (1999) The bZip transcription factor Cap1p is involved in multidrug resistance and oxidative stress response in *Candida albicans*. *J Bacteriol* **181**: 700-708.
- Arnaud, M.B., Costanzo, M.C., Skrzypek, M.S., Binkley, G., Lane, C., Miyasato, S.R., and Sherlock, G. (2005) The *Candida* Genome Database (CGD), a community resource for *Candida albicans* gene and protein information. *Nucleic Acids Res* **33**: D358-363.
- Bain, J.M., Stubberfield, C., and Gow, N.A. (2001) Ura-status-dependent adhesion of *Candida albicans* mutants. *FEMS Microbiol Lett* **204**: 323-328.
- Barelle, C.J., Manson, C.L., MacCallum, D.M., Odds, F.C., Gow, N.A., and Brown, A.J. (2004) GFP as a quantitative reporter of gene regulation in *Candida albicans*. *Yeast* **21**: 333-340.
- Beausejour, A., Grenier, D., Goulet, J.P., and Deslauriers, N. (1998) Proteolytic activation of the interleukin-1beta precursor by *Candida albicans*. *Infect Immun* **66**: 676-681.
- Bektic, J., Lell, C.P., Fuchs, A., Stoiber, H., Speth, C., Lass-Flörl, C., Borg-von Zepelin, M., Dierich, M.P., and Wurzner, R. (2001) HIV protease inhibitors attenuate adherence of *Candida albicans* to epithelial cells in vitro. *FEMS Immunol Med Microbiol* **31**: 65-71.
- Bendtsen, J.D., Nielsen, H., von Heijne, G., and Brunak, S. (2004) Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* **340**: 783-795.
- Berg, F.T. (1846) Om Torsk Hos Barn. *L.J.Hjerta, Stockholm, Sweden*.
- Berkhout, C.M. (1923) De schimmelgeslachten *Monilia*, *Oidium*, *Oospora*, en *Torula*. *Dissertation. University of Utrecht, The Netherlands*.
- Borg-von Zepelin, M., Eiffert, H., Kann, M., and Ruchel, R. (1993) Changes in the spectrum of fungal isolates: results from clinical specimens gathered in 1987/88 compared with those in 1991/92 in the University Hospital Göttingen, Germany. *Mycoses* **36**: 247-253.
- Borg-von Zepelin, M., Beggah, S., Boggian, K., Sanglard, D., and Monod, M. (1998) The expression of the secreted aspartyl proteinases Sap4 to Sap6 from *Candida albicans* in murine macrophages. *Mol Microbiol* **28**: 543-554.
- Borg-von Zepelin, M., Meyer, I., Thomssen, R., Wurzner, R., Sanglard, D., Telenti, A., and Monod, M. (1999) HIV-Protease inhibitors reduce cell adherence of *Candida albicans* strains by inhibition of yeast secreted aspartic proteases. *J Invest Dermatol* **113**: 747-751.
- Borg, M., and Ruchel, R. (1988) Expression of extracellular acid proteinase by proteolytic *Candida* spp. during experimental infection of oral mucosa. *Infect Immun* **56**: 626-631.

- Brand, A., MacCallum, D.M., Brown, A.J., Gow, N.A., and Odds, F.C. (2004) Ectopic expression of URA3 can influence the virulence phenotypes and proteome of *Candida albicans* but can be overcome by targeted reintegration of URA3 at the RPS10 locus. *Eukaryot Cell* **3**: 900-909.
- Braun, B.R., and Johnson, A.D. (1997) Control of filament formation in *Candida albicans* by the transcriptional repressor TUP1. *Science* **277**: 105-109.
- Buurman, E.T., Westwater, C., Hube, B., Brown, A.J., Odds, F.C., and Gow, N.A. (1998) Molecular analysis of CaMnt1p, a mannosyl transferase important for adhesion and virulence of *Candida albicans*. *Proc Natl Acad Sci U S A* **95**: 7670-7675.
- Calderone, R.A. (1989) Host-parasite relationships in candidosis. *Mycoses* **32 Suppl 2**: 12-17.
- Calderone, R.A. (2002) *Candida and Candidiasis*: ASM Press.
- Cannon, R.D., Jenkinson, H.F., and Shepherd, M.G. (1992) Cloning and expression of *Candida albicans* ADE2 and proteinase genes on a replicative plasmid in *C. albicans* and in *Saccharomyces cerevisiae*. *Mol Gen Genet* **235**: 453-457.
- Cappellaro, C., Baldermann, C., Rachel, R., and Tanner, W. (1994) Mating type-specific cell-cell recognition of *Saccharomyces cerevisiae*: cell wall attachment and active sites of  $\alpha$ - and  $\alpha$ -agglutinin. *Embo J* **13**: 4737-4744.
- Caro, L.H., Tettelin, H., Vossen, J.H., Ram, A.F., van den Ende, H., and Klis, F.M. (1997) In silico identification of glycosyl-phosphatidylinositol-anchored plasma-membrane and cell wall proteins of *Saccharomyces cerevisiae*. *Yeast* **13**: 1477-1489.
- Carruthers, V.B., and Blackman, M.J. (2005) A new release on life: emerging concepts in proteolysis and parasite invasion. *Mol Microbiol* **55**: 1617-1630.
- Castellani, A. (1920) The etiology of thrush. *J.Trop.Med.Hyg.* **23**: 17-22.
- Cawley, N.X., Chen, H.C., Beinfeld, M.C., and Loh, Y.P. (1996) Specificity and kinetic studies on the cleavage of various prohormone mono- and paired-basic residue sites by yeast aspartic protease 3. *J Biol Chem* **271**: 4168-4176.
- Cawley, N.X., Olsen, V., Zhang, C.F., Chen, H.C., Tan, M., and Loh, Y.P. (1998) Activation and processing of non-anchored yapsin 1 (Yap3p). *J Biol Chem* **273**: 584-591.
- Cawley, N.X., Chino, M., Maldonado, A., Rodriguez, Y.M., Loh, Y.P., and Ellman, J.A. (2003) Synthesis and characterization of the first potent inhibitor of yapsin 1. Implications for the study of yapsin-like enzymes. *J Biol Chem* **278**: 5523-5530.
- Chandra, J., Kuhn, D.M., Mukherjee, P.K., Hoyer, L.L., McCormick, T., and Ghannoum, M.A. (2001a) Biofilm formation by the fungal pathogen *Candida albicans*: development, architecture, and drug resistance. *J Bacteriol* **183**: 5385-5394.
- Chandra, J., Mukherjee, P.K., Leidich, S.D., Faddoul, F.F., Hoyer, L.L., Douglas, L.J., and Ghannoum, M.A. (2001b) Antifungal resistance of

- candidal biofilms formed on denture acrylic in vitro. *J Dent Res* **80**: 903-908.
- Chen, G., Hata, N., and Zhang, M.Q. (2004) Transcription factor binding element detection using functional clustering of mutant expression data. *Nucleic Acids Res* **32**: 2362-2371.
- Chomczynski, P., and Sacchi, N. (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* **162**: 156-159.
- Colina, A.R., Aumont, F., Deslauriers, N., Belhumeur, P., and de Repentigny, L. (1996) Evidence for degradation of gastrointestinal mucin by *Candida albicans* secretory aspartyl proteinase. *Infect Immun* **64**: 4514-4519.
- Copping, V.M., Barelle, C.J., Hube, B., Gow, N.A., Brown, A.J., and Odds, F.C. (2005) Exposure of *Candida albicans* to antifungal agents affects expression of SAP2 and SAP9 secreted proteinase genes. *J Antimicrob Chemother* **55**: 645-654.
- Csank, C., Makris, C., Meloche, S., Schroppel, K., Rollinghoff, M., Dignard, D., Thomas, D.Y., and Whiteway, M. (1997) Derepressed hyphal growth and reduced virulence in a VH1 family-related protein phosphatase mutant of the human pathogen *Candida albicans*. *Mol Biol Cell* **8**: 2539-2551.
- Cutler, J.E., Friedman, L., and Milner, K.C. (1972) Biological and chemical characterization of toxic substances from *Candida albicans*. *Infect Immun* **6**: 616-627.
- d'Enfert, C., Goyard, S., Rodriguez-Arnaveilhe, S., Frangeul, L., Jones, L., Tekaia, F., Bader, O., Albrecht, A., Castillo, L., Dominguez, A., Ernst, J.F., Fradin, C., Gaillardin, C., Garcia-Sanchez, S., de Groot, P., Hube, B., Klis, F.M., Krishnamurthy, S., Kunze, D., Lopez, M.C., Mavor, A., Martin, N., Moszer, I., Onesime, D., Perez Martin, J., Sentandreu, R., Valentin, E., and Brown, A.J. (2005) CandidaDB: a genome database for *Candida albicans* pathogenomics. *Nucleic Acids Res* **33**: D353-357.
- Daniels, K.J., Srikantha, T., Lockhart, S.R., Pujol, C., and Soll, D.R. (2006) Opaque cells signal white cells to form biofilms in *Candida albicans*. *Embo J* **25**: 2240-2252.
- Davis, D., Wilson, R.B., and Mitchell, A.P. (2000) RIM101-dependent and-independent pathways govern pH responses in *Candida albicans*. *Mol Cell Biol* **20**: 971-978.
- De Bernardis, F., Cassone, A., Sturtevant, J., and Calderone, R. (1995) Expression of *Candida albicans* SAP1 and SAP2 in experimental vaginitis. *Infect Immun* **63**: 1887-1892.
- De Bernardis, F., Muhlschlegel, F.A., Cassone, A., and Fonzi, W.A. (1998) The pH of the host niche controls gene expression in and virulence of *Candida albicans*. *Infect Immun* **66**: 3317-3325.
- De Bernardis, F., Arancia, S., Morelli, L., Hube, B., Sanglard, D., Schafer, W., and Cassone, A. (1999) Evidence that members of the secretory aspartyl proteinase gene family, in particular SAP2, are virulence factors for *Candida vaginitis*. *J Infect Dis* **179**: 201-208.

- De Groot, P.W., Hellingwerf, K.J., and Klis, F.M. (2003) Genome-wide identification of fungal GPI proteins. *Yeast* **20**: 781-796.
- de Groot, P.W., de Boer, A.D., Cunningham, J., Dekker, H.L., de Jong, L., Hellingwerf, K.J., de Koster, C., and Klis, F.M. (2004) Proteomic analysis of *Candida albicans* cell walls reveals covalently bound carbohydrate-active enzymes and adhesins. *Eukaryot Cell* **3**: 955-965.
- De Groot, P.W., Ram, A.F., and Klis, F.M. (2005) Features and functions of covalently linked proteins in fungal cell walls. *Fungal Genet Biol* **42**: 657-675.
- de Nobel, J.G., Klis, F.M., Priem, J., Munnik, T., and van den Ende, H. (1990) The glucanase-soluble mannoproteins limit cell wall porosity in *Saccharomyces cerevisiae*. *Yeast* **6**: 491-499.
- Domergue, R., and Cormack, B.P. (2006) Transcriptional regulation of a *Candida glabrata* adhesin by environmental signals and promoter specific elements. *ASM conference*: abstract B20.
- Dostal, J., Merkerova, M., Hradilek, M., Pichova, I., and Hruskova-Heidingsfeldova, O. (2006) Secreted aspartic proteinases of *Candida parapsilosis*: regulation of transcription of *SAPP1* and *SAPP2* genes and activation of respective proteinase precursors. *ASM conference*: abstract A7.
- Douglas, L.J. (1987) Adhesion of *Candida* species to epithelial surfaces. *Crit Rev Microbiol* **15**: 27-43.
- Dujon, B., Sherman, D., Fischer, G., Durrens, P., Casaregola, S., Lafontaine, I., De Montigny, J., Marck, C., Neuveglise, C., Talla, E., Goffard, N., Frangeul, L., Aigle, M., Anthouard, V., Babour, A., Barbe, V., Barnay, S., Blanchin, S., Beckerich, J.M., Beyne, E., Bleykasten, C., Boisrame, A., Boyer, J., Cattolico, L., Confanioleri, F., De Daruvar, A., Despons, L., Fabre, E., Fairhead, C., Ferry-Dumazet, H., Groppi, A., Hantraye, F., Hennequin, C., Jauniaux, N., Joyet, P., Kachouri, R., Kerrest, A., Koszul, R., Lemaire, M., Lesur, I., Ma, L., Muller, H., Nicaud, J.M., Nikolski, M., Oztas, S., Ozier-Kalogeropoulos, O., Pellenz, S., Potier, S., Richard, G.F., Straub, M.L., Suleau, A., Swennen, D., Tekaia, F., Wesolowski-Louvel, M., Westhof, E., Wirth, B., Zeniou-Meyer, M., Zivanovic, I., Bolotin-Fukuhara, M., Thierry, A., Bouchier, C., Caudron, B., Scarpelli, C., Gaillardin, C., Weissenbach, J., Wincker, P., and Souciet, J.L. (2004) Genome evolution in yeasts. *Nature* **430**: 35-44.
- Egel-Mitani, M., Flygenring, H.P., and Hansen, M.T. (1990) A novel aspartyl protease allowing KEX2-independent MF alpha propheromone processing in yeast. *Yeast* **6**: 127-137.
- Eisenhaber, B., Schneider, G., Wildpaner, M., and Eisenhaber, F. (2004) A sensitive predictor for potential GPI lipid modification sites in fungal protein sequences and its application to genome-wide studies for *Aspergillus nidulans*, *Candida albicans*, *Neurospora crassa*, *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*. *J Mol Biol* **337**: 243-253.
- Ellepola, A.N., and Samaranyake, L.P. (1998) The effect of limited exposure to antimycotics on the relative cell-surface hydrophobicity and the

- adhesion of oral *Candida albicans* to buccal epithelial cells. *Arch Oral Biol* **43**: 879-887.
- Ernst, J.F. (2000) Transcription factors in *Candida albicans* - environmental control of morphogenesis. *Microbiology* **146 (Pt 8)**: 1763-1774.
- Falkow, S. (1988) Molecular Koch's postulates applied to microbial pathogenicity. *Rev Infect Dis* **10 Suppl 2**: S274-276.
- Felk, A. (2002) Sekretorische Aspartatproteasen von *Candida albicans* und ihre Bedeutung als Virulenzfaktoren. *Inauguraldissertation*: Freie Universität Berlin.
- Felk, A., Kretschmar, M., Albrecht, A., Schaller, M., Beinhauer, S., Nichterlein, T., Sanglard, D., Korting, H.C., Schafer, W., and Hube, B. (2002) *Candida albicans* hyphal formation and the expression of the Efg1-regulated proteinases Sap4 to Sap6 are required for the invasion of parenchymal organs. *Infect Immun* **70**: 3689-3700.
- Fernandez-Martinez, J., Brown, C.V., Diez, E., Tilburn, J., Arst, H.N., Jr., Penalva, M.A., and Espeso, E.A. (2003) Overlap of nuclear localisation signal and specific DNA-binding residues within the zinc finger domain of PacC. *J Mol Biol* **334**: 667-684.
- Fidel, P.L., Jr., Vazquez, J.A., and Sobel, J.D. (1999) *Candida glabrata*: review of epidemiology, pathogenesis, and clinical disease with comparison to *C. albicans*. *Clin Microbiol Rev* **12**: 80-96.
- Fonzi, W.A., and Irwin, M.Y. (1993) Isogenic strain construction and gene mapping in *Candida albicans*. *Genetics* **134**: 717-728.
- Fradin, C., De Groot, P., MacCallum, D., Schaller, M., Klis, F., Odds, F.C., and Hube, B. (2005) Granulocytes govern the transcriptional response, morphology and proliferation of *Candida albicans* in human blood. *Mol Microbiol* **56**: 397-415.
- Fraering, P., Imhof, I., Meyer, U., Strub, J.M., van Dorsselaer, A., Vionnet, C., and Conzelmann, A. (2001) The GPI transamidase complex of *Saccharomyces cerevisiae* contains Gaa1p, Gpi8p, and Gpi16p. *Mol Biol Cell* **12**: 3295-3306.
- Frieman, M.B., and Cormack, B.P. (2003) The omega-site sequence of glycosylphosphatidylinositol-anchored proteins in *Saccharomyces cerevisiae* can determine distribution between the membrane and the cell wall. *Mol Microbiol* **50**: 883-896.
- Gagnon-Arsenault, I., Parisé, L., Tardif, G., Malaisson, D., Boudrault, C., and Bourbonnais, Y. (2005) The yeast yapsin-1 endoprotease acts as a sheddase for the Gas1 glycanosyltransferase and its misregulation leads to cell wall defects. Poster presentation at the IPS Conference, Quebec, Canada.
- Gale, C.A., Bendel, C.M., McClellan, M., Hauser, M., Becker, J.M., Berman, J., and Hostetter, M.K. (1998) Linkage of adhesion, filamentous growth, and virulence in *Candida albicans* to a single gene, INT1. *Science* **279**: 1355-1358.
- Garcia, R., Bermejo, C., Grau, C., Perez, R., Rodriguez-Pena, J.M., Francois, J., Nombela, C., and Arroyo, J. (2004) The global transcriptional response to transient cell wall damage in

- Saccharomyces cerevisiae and its regulation by the cell integrity signaling pathway. *J Biol Chem* **279**: 15183-15195.
- Gerami-Nejad, M., Berman, J., and Gale, C.A. (2001) Cassettes for PCR-mediated construction of green, yellow, and cyan fluorescent protein fusions in *Candida albicans*. *Yeast* **18**: 859-864.
- Ghannoum, M., and Abu Elteen, K. (1986) Correlative relationship between proteinase production, adherence and pathogenicity of various strains of *Candida albicans*. *J Med Vet Mycol* **24**: 407-413.
- Ghannoum, M.A., Spellberg, B., Saporito-Irwin, S.M., and Fonzi, W.A. (1995) Reduced virulence of *Candida albicans* PHR1 mutants. *Infect Immun* **63**: 4528-4530.
- Ghannoum, M.A. (2000) Potential role of phospholipases in virulence and fungal pathogenesis. *Clin Microbiol Rev* **13**: 122-143, table of contents.
- Gillum, A.M., Tsay, E.Y., and Kirsch, D.R. (1984) Isolation of the *Candida albicans* gene for orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae* *ura3* and *E. coli* *pyrF* mutations. *Mol Gen Genet* **198**: 179-182.
- Giusani, A.D., Vinces, M., and Kumamoto, C.A. (2002) Invasive filamentous growth of *Candida albicans* is promoted by Czf1p-dependent relief of Efg1p-mediated repression. *Genetics* **160**: 1749-1753.
- Gola, S., Martin, R., Walther, A., Dunkler, A., and Wendland, J. (2003) New modules for PCR-based gene targeting in *Candida albicans*: rapid and efficient gene targeting using 100 bp of flanking homology region. *Yeast* **20**: 1339-1347.
- Granger, B.L., Flenniken, M.L., Davis, D.A., Mitchell, A.P., and Cutler, J.E. (2005) Yeast wall protein 1 of *Candida albicans*. *Microbiology* **151**: 1631-1644.
- Habermann, E., and Hardt, K.L. (1972) A sensitive and specific plate test for the quantitation of phospholipases. *Anal Biochem* **50**: 163-173.
- Hacker, J., and Heesemann, J. (2002) Molecular Infection Biology. Interactions Between Microorganisms and Cells. *John Wiley & Sons*: ISBN 0471178462.
- Hanahan, D. (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* **166**: 557-580.
- Hawser, S.P., and Douglas, L.J. (1994) Biofilm formation by *Candida* species on the surface of catheter materials in vitro. *Infect Immun* **62**: 915-921.
- Hazen, K.C. (1995) New and emerging yeast pathogens. *Clin Microbiol Rev* **8**: 462-478.
- Heidenreich, F., and Dierich, M.P. (1985) *Candida albicans* and *Candida stellatoidea*, in contrast to other *Candida* species, bind iC3b and C3d but not C3b. *Infect Immun* **50**: 598-600.
- Herrero, A.B., Magnelli, P., Mansour, M.K., Levitz, S.M., Bussey, H., and Abeijon, C. (2004) KRE5 gene null mutant strains of *Candida albicans* are avirulent and have altered cell wall composition and hypha formation properties. *Eukaryot Cell* **3**: 1423-1432.

- Hinnebusch, A.G., and Fink, G.R. (1983) Positive regulation in the general amino acid control of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* **80**: 5374-5378.
- Hoffman, C.S., and Winston, F. (1987) A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* **57**: 267-272.
- Hoover, C.I., Jantapour, M.J., Newport, G., Agabian, N., and Fisher, S.J. (1998) Cloning and regulated expression of the *Candida albicans* phospholipase B (PLB1) gene. *FEMS Microbiol Lett* **167**: 163-169.
- Hoyer, L.L. (2001) The ALS gene family of *Candida albicans*. *Trends Microbiol* **9**: 176-180.
- Hubbard, M.J., Markie, D., and Poulter, R.T. (1986) Isolation and morphological characterization of a mycelial mutant of *Candida albicans*. *J Bacteriol* **165**: 61-65.
- Hube, B., Turver, C.J., Odds, F.C., Eiffert, H., Boulnois, G.J., Kochel, H., and Ruchel, R. (1991) [Identification, cloning and characterization of the gene for the secretory aspartate protease of *Candida albicans*]. *Mycoses* **34 Suppl 1**: 59-61.
- Hube, B., Monod, M., Schofield, D.A., Brown, A.J., and Gow, N.A. (1994) Expression of seven members of the gene family encoding secretory aspartyl proteinases in *Candida albicans*. *Mol Microbiol* **14**: 87-99.
- Hube, B. (1996) *Candida albicans* secreted aspartyl proteinases. *Curr Top Med Mycol* **7**: 55-69.
- Hube, B., Sanglard, D., Odds, F.C., Hess, D., Monod, M., Schafer, W., Brown, A.J., and Gow, N.A. (1997) Disruption of each of the secreted aspartyl proteinase genes SAP1, SAP2, and SAP3 of *Candida albicans* attenuates virulence. *Infect Immun* **65**: 3529-3538.
- Hube, B., Stehr, F., Bossenz, M., Mazur, A., Kretschmar, M., and Schafer, W. (2000) Secreted lipases of *Candida albicans*: cloning, characterisation and expression analysis of a new gene family with at least ten members. *Arch Microbiol* **174**: 362-374.
- Hube, B., and Naglik, J. (2001) *Candida albicans* proteinases: resolving the mystery of a gene family. *Microbiology* **147**: 1997-2005.
- Jong, A.Y., Stins, M.F., Huang, S.H., Chen, S.H., and Kim, K.S. (2001) Traversal of *Candida albicans* across human blood-brain barrier in vitro. *Infect Immun* **69**: 4536-4544.
- Jung, U.S., and Levin, D.E. (1999) Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway. *Mol Microbiol* **34**: 1049-1057.
- Kamai, Y., Kubota, M., Kamai, Y., Hosokawa, T., Fukuoka, T., and Filler, S.G. (2002) Contribution of *Candida albicans* ALS1 to the pathogenesis of experimental oropharyngeal candidiasis. *Infect Immun* **70**: 5256-5258.
- Kapteyn, J.C., Van Den Ende, H., and Klis, F.M. (1999) The contribution of cell wall proteins to the organization of the yeast cell wall. *Biochim Biophys Acta* **1426**: 373-383.

- Kelly, M.T., MacCallum, D.M., Clancy, S.D., Odds, F.C., Brown, A.J., and Butler, G. (2004) The *Candida albicans* CaACE2 gene affects morphogenesis, adherence and virulence. *Mol Microbiol* **53**: 969-983.
- Kennedy, M.J., Rogers, A.L., Hanselmen, L.R., Soll, D.R., and Yancey, R.J., Jr. (1988) Variation in adherence and cell surface hydrophobicity in *Candida albicans* white and opaque phenotypes. *Mycopathologia* **102**: 149-156.
- Khalaf, R.A., and Zitomer, R.S. (2001) The DNA binding protein Rfg1 is a repressor of filamentation in *Candida albicans*. *Genetics* **157**: 1503-1512.
- King, R.D., Lee, J.C., and Morris, A.L. (1980) Adherence of *Candida albicans* and other *Candida* species to mucosal epithelial cells. *Infect Immun* **27**: 667-674.
- Kinneberg, K.M., Bendel, C.M., Jechorek, R.P., Cebelinski, E.A., Gale, C.A., Berman, J.G., Erlandsen, S.L., Hostetter, M.K., and Wells, C.L. (1999) Effect of INT1 gene on *Candida albicans* murine intestinal colonization. *J Surg Res* **87**: 245-251.
- Klis, F.M., Caro, L.H., Vossen, J.H., Kapteyn, J.C., Ram, A.F., Montijn, R.C., Van Berkel, M.A., and Van den Ende, H. (1997) Identification and characterization of a major building block in the cell wall of *Saccharomyces cerevisiae*. *Biochem Soc Trans* **25**: 856-860.
- Klis, F.M., de Groot, P., and Hellingwerf, K. (2001) Molecular organization of the cell wall of *Candida albicans*. *Med Mycol* **39 Suppl 1**: 1-8.
- Koelsch, G., Tang, J., Loy, J.A., Monod, M., Jackson, K., Foundling, S.I., and Lin, X. (2000) Enzymic characteristics of secreted aspartic proteases of *Candida albicans*. *Biochim Biophys Acta* **1480**: 117-131.
- Komano, H., and Fuller, R.S. (1995) Shared functions in vivo of a glycosylphosphatidylinositol-linked aspartyl protease, Mkc7, and the proprotein processing protease Kex2 in yeast. *Proc Natl Acad Sci U S A* **92**: 10752-10756.
- Komano, H., Rockwell, N., Wang, G.T., Krafft, G.A., and Fuller, R.S. (1999) Purification and characterization of the yeast glycosylphosphatidylinositol-anchored, monobasic-specific aspartyl protease yapsin 2 (Mkc7p). *J Biol Chem* **274**: 24431-24437.
- Kretschmar, M., Hube, B., Bertsch, T., Sanglard, D., Merker, R., Schroder, M., Hof, H., and Nichterlein, T. (1999) Germ tubes and proteinase activity contribute to virulence of *Candida albicans* in murine peritonitis. *Infect Immun* **67**: 6637-6642.
- Krysan, D.J., Ting, E.L., Abeijon, C., Kroos, L., and Fuller, R.S. (2005) Yapsins are a family of aspartyl proteases required for cell wall integrity in *Saccharomyces cerevisiae*. *Eukaryot Cell* **4**: 1364-1374.
- Kvaal, C., Lachke, S.A., Srikantha, T., Daniels, K., McCoy, J., and Soll, D.R. (1999) Misexpression of the opaque-phase-specific gene PEP1 (SAP1) in the white phase of *Candida albicans* confers increased virulence in a mouse model of cutaneous infection. *Infect Immun* **67**: 6652-6662.
- Leidich, S.D., Ibrahim, A.S., Fu, Y., Koul, A., Jessup, C., Vitullo, J., Fonzi, W., Mirbod, F., Nakashima, S., Nozawa, Y., and Ghannoum, M.A. (1998)

- Cloning and disruption of caPLB1, a phospholipase B gene involved in the pathogenicity of *Candida albicans*. *J Biol Chem* **273**: 26078-26086.
- Leng, P., Lee, P.R., Wu, H., and Brown, A.J. (2001) Efg1, a morphogenetic regulator in *Candida albicans*, is a sequence-specific DNA binding protein. *J Bacteriol* **183**: 4090-4093.
- Liu, H., Kohler, J., and Fink, G.R. (1994) Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. *Science* **266**: 1723-1726.
- Liu, H. (2002) Co-regulation of pathogenesis with dimorphism and phenotypic switching in *Candida albicans*, a commensal and a pathogen. *Int J Med Microbiol* **292**: 299-311.
- Lo, H.J., Kohler, J.R., DiDomenico, B., Loebenberg, D., Cacciapuoti, A., and Fink, G.R. (1997) Nonfilamentous *C. albicans* mutants are avirulent. *Cell* **90**: 939-949.
- Lockhart, S.R., Zhao, R., Daniels, K.J., and Soll, D.R. (2003) Alpha-pheromone-induced "shmooing" and gene regulation require white-opaque switching during *Candida albicans* mating. *Eukaryot Cell* **2**: 847-855.
- Lu, C.F., Kurjan, J., and Lipke, P.N. (1994) A pathway for cell wall anchorage of *Saccharomyces cerevisiae* alpha-agglutinin. *Mol Cell Biol* **14**: 4825-4833.
- Magee, B.B., Hube, B., Wright, R.J., Sullivan, P.J., and Magee, P.T. (1993) The genes encoding the secreted aspartyl proteinases of *Candida albicans* constitute a family with at least three members. *Infect Immun* **61**: 3240-3243.
- Masuoka, J., and Hazen, K.C. (1997) Cell wall protein mannosylation determines *Candida albicans* cell surface hydrophobicity. *Microbiology* **143 (Pt 9)**: 3015-3021.
- Masuoka, J., and Hazen, K.C. (1999) Differences in the acid-labile component of *Candida albicans* mannan from hydrophobic and hydrophilic yeast cells. *Glycobiology* **9**: 1281-1286.
- Mayor, S., and Riezman, H. (2004) Sorting GPI-anchored proteins. *Nat Rev Mol Cell Biol* **5**: 110-120.
- McLain, N., and Dolan, J.W. (1997) Phospholipase D activity is required for dimorphic transition in *Candida albicans*. *Microbiology* **143 (Pt 11)**: 3521-3526.
- Miller, M.G., and Johnson, A.D. (2002) White-opaque switching in *Candida albicans* is controlled by mating-type locus homeodomain proteins and allows efficient mating. *Cell* **110**: 293-302.
- Mitchell, A.P. (1998) Dimorphism and virulence in *Candida albicans*. *Curr Opin Microbiol* **1**: 687-692.
- Miyasaki, S.H., White, T.C., and Agabian, N. (1994) A fourth secreted aspartyl proteinase gene (SAP4) and a CARE2 repetitive element are located upstream of the SAP1 gene in *Candida albicans*. *J Bacteriol* **176**: 1702-1710.
- Monod, M., Togni, G., Hube, B., and Sanglard, D. (1994) Multiplicity of genes encoding secreted aspartic proteinases in *Candida* species. *Mol Microbiol* **13**: 357-368.

- Monod, M., Hube, B., Hess, D., and Sanglard, D. (1998) Differential regulation of SAP8 and SAP9, which encode two new members of the secreted aspartic proteinase family in *Candida albicans*. *Microbiology* **144 (Pt 10)**: 2731-2737.
- Moran, G., Stokes, C., Thewes, S., Hube, B., Coleman, D.C., and Sullivan, D. (2004) Comparative genomics using *Candida albicans* DNA microarrays reveals absence and divergence of virulence-associated genes in *Candida dubliniensis*. *Microbiology* **150**: 3363-3382.
- Morrow, B., Srikantha, T., and Soll, D.R. (1992) Transcription of the gene for a pepsinogen, PEP1, is regulated by white-opaque switching in *Candida albicans*. *Mol Cell Biol* **12**: 2997-3005.
- Morrow, B., Srikantha, T., Anderson, J., and Soll, D.R. (1993) Coordinate regulation of two opaque-phase-specific genes during white-opaque switching in *Candida albicans*. *Infect Immun* **61**: 1823-1828.
- Morschhauser, J., Virkola, R., Korhonen, T.K., and Hacker, J. (1997) Degradation of human subendothelial extracellular matrix by proteinase-secreting *Candida albicans*. *FEMS Microbiol Lett* **153**: 349-355.
- Muniz, M., Morsomme, P., and Riezman, H. (2001) Protein sorting upon exit from the endoplasmic reticulum. *Cell* **104**: 313-320.
- Munro, C.A., Winter, K., Buchan, A., Henry, K., Becker, J.M., Brown, A.J., Bulawa, C.E., and Gow, N.A. (2001) Chs1 of *Candida albicans* is an essential chitin synthase required for synthesis of the septum and for cell integrity. *Mol Microbiol* **39**: 1414-1426.
- Murad, A.M., Lee, P.R., Broadbent, I.D., Barelle, C.J., and Brown, A.J. (2000) Clp10, an efficient and convenient integrating vector for *Candida albicans*. *Yeast* **16**: 325-327.
- Murad, A.M., d'Enfert, C., Gaillardin, C., Tournu, H., Tekaia, F., Talibi, D., Marechal, D., Marchais, V., Cottin, J., and Brown, A.J. (2001a) Transcript profiling in *Candida albicans* reveals new cellular functions for the transcriptional repressors CaTup1, CaMig1 and CaNrg1. *Mol Microbiol* **42**: 981-993.
- Murad, A.M., Leng, P., Straffon, M., Wishart, J., Macaskill, S., MacCallum, D., Schnell, N., Talibi, D., Marechal, D., Tekaia, F., d'Enfert, C., Gaillardin, C., Odds, F.C., and Brown, A.J. (2001b) NRG1 represses yeast-hypha morphogenesis and hypha-specific gene expression in *Candida albicans*. *Embo J* **20**: 4742-4752.
- Naglik, J., Albrecht, A., Bader, O., and Hube, B. (2004) *Candida albicans* proteinases and host/pathogen interactions. *Cell Microbiol* **6**: 915-926.
- Naglik, J.R., Newport, G., White, T.C., Fernandes-Naglik, L.L., Greenspan, J.S., Greenspan, D., Sweet, S.P., Challacombe, S.J., and Agabian, N. (1999) In vivo analysis of secreted aspartyl proteinase expression in human oral candidiasis. *Infect Immun* **67**: 2482-2490.
- Naglik, J.R., Challacombe, S.J., and Hube, B. (2003) *Candida albicans* secreted aspartyl proteinases in virulence and pathogenesis. *Microbiol Mol Biol Rev* **67**: 400-428, table of contents.

- Negi, M., Tsuboi, R., Matsui, T., and Ogawa, H. (1984) Isolation and characterization of proteinase from *Candida albicans*: substrate specificity. *J Invest Dermatol* **83**: 32-36.
- Newport, G., and Agabian, N. (1997) KEX2 influences *Candida albicans* proteinase secretion and hyphal formation. *J Biol Chem* **272**: 28954-28961.
- Nielsen, H., Engelbrecht, J., Brunak, S., and von Heijne, G. (1997) Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng* **10**: 1-6.
- Niimi, M. (1996) Dibutyryl cyclic AMP-enhanced germ tube formation in exponentially growing *Candida albicans* cells. *Fungal Genet Biol* **20**: 79-83.
- Noble, S.M., and Johnson, A.D. (2005) Strains and strategies for large-scale gene deletion studies of the diploid human fungal pathogen *Candida albicans*. *Eukaryot Cell* **4**: 298-309.
- Odds, F.C. (1988) *Candida and Candidiasis*. London: Bailliere Tindal.
- Ollert, M.W., Sohnchen, R., Korting, H.C., Ollert, U., Brautigam, S., and Brautigam, W. (1993) Mechanisms of adherence of *Candida albicans* to cultured human epidermal keratinocytes. *Infect Immun* **61**: 4560-4568.
- Olsen, V., Cawley, N.X., Brandt, J., Egel-Mitani, M., and Loh, Y.P. (1999) Identification and characterization of *Saccharomyces cerevisiae* yapsin 3, a new member of the yapsin family of aspartic proteases encoded by the YPS3 gene. *Biochem J* **339 (Pt 2)**: 407-411.
- Olsen, V., and Loh, Y.P. (2000) In vivo processing of nonanchored Yapsin 1 (Yap3p). *Arch Biochem Biophys* **375**: 315-321.
- Pardo, M., Monteoliva, L., Vazquez, P., Martinez, R., Molero, G., Nombela, C., and Gil, C. (2004) PST1 and ECM33 encode two yeast cell surface GPI proteins important for cell wall integrity. *Microbiology* **150**: 4157-4170.
- Pichova, I., Pavlickova, L., Dostal, J., Dolejsi, E., Hruskova-Heidingsfeldova, O., Weber, J., Ruml, T., and Soucek, M. (2001) Secreted aspartic proteases of *Candida albicans*, *Candida tropicalis*, *Candida parapsilosis* and *Candida lusitanae*. Inhibition with peptidomimetic inhibitors. *Eur J Biochem* **268**: 2669-2677.
- Prasad, R., De Wergifosse, P., Goffeau, A., and Balzi, E. (1995) Molecular cloning and characterization of a novel gene of *Candida albicans*, CDR1, conferring multiple resistance to drugs and antifungals. *Curr Genet* **27**: 320-329.
- Ray, T.L., and Payne, C.D. (1988) Scanning electron microscopy of epidermal adherence and cavitation in murine candidiasis: a role for *Candida* acid proteinase. *Infect Immun* **56**: 1942-1949.
- Remold, H., Fasold, H., and Staib, F. (1968) Purification and characterization of a proteolytic enzyme from *Candida albicans*. *Biochim Biophys Acta* **167**: 399-406.
- Rex, J.H., Walsh, T.J., Sobel, J.D., Filler, S.G., Pappas, P.G., Dismukes, W.E., and Edwards, J.E. (2000) Practice guidelines for the treatment

- of candidiasis. Infectious Diseases Society of America. *Clin Infect Dis* **30**: 662-678.
- Richard, M., De Groot, P., Courtin, O., Poulain, D., Klis, F., and Gaillardin, C. (2002) GPI7 affects cell-wall protein anchorage in *Saccharomyces cerevisiae* and *Candida albicans*. *Microbiology* **148**: 2125-2133.
- Robin, C.P. (1853) Histoire naturelle des vegetaux. Parasites qui croissent sur l'homme et sur les animaux vivants. *Balliere, Paris, France*.
- Rodrigues, A.G., Mardh, P.A., Pina-Vaz, C., Martinez-de-Oliveira, J., and Fonseca, A.F. (1999) Germ tube formation changes surface hydrophobicity of *Candida* cells. *Infect Dis Obstet Gynecol* **7**: 222-226.
- Rüchel, R., Tegeler, R., and Trost, M. (1982) A comparison of secretory proteinases from different strains of *Candida albicans*. *Sabouraudia* **20**: 233-244.
- Rüchel, R. (1986) Cleavage of immunoglobulins by pathogenic yeasts of the genus *Candida*. *Microbiol Sci* **3**: 316-319.
- Rüchel, R. (1992) Proteinase. In *New Strategies in Fungal Disease*. Bennett, J.E., Hay, R.J., and Peterson, P.K. (ed). Edinburgh, pp. 17-31.
- Runeberg-Roos, P., Tormakangas, K., and Ostman, A. (1991) Primary structure of a barley-grain aspartic proteinase. A plant aspartic proteinase resembling mammalian cathepsin D. *Eur J Biochem* **202**: 1021-1027.
- Sanglard, D., Hube, B., Monod, M., Odds, F.C., and Gow, N.A. (1997a) A triple deletion of the secreted aspartyl proteinase genes SAP4, SAP5, and SAP6 of *Candida albicans* causes attenuated virulence. *Infect Immun* **65**: 3539-3546.
- Sanglard, D., Ischer, F., Monod, M., and Bille, J. (1997b) Cloning of *Candida albicans* genes conferring resistance to azole antifungal agents: characterization of CDR2, a new multidrug ABC transporter gene. *Microbiology* **143 (Pt 2)**: 405-416.
- Schaller, M., Korting, H.C., Schafer, W., Sanglard, D., and Hube, B. (1998a) [Investigations on the regulation of secreted aspartyl proteases in a model of oral candidiasis in vivo]. *Mycoses* **41 Suppl 2**: 69-73.
- Schaller, M., Schafer, W., Korting, H.C., and Hube, B. (1998b) Differential expression of secreted aspartyl proteinases in a model of human oral candidosis and in patient samples from the oral cavity. *Mol Microbiol* **29**: 605-615.
- Schaller, M., Korting, H.C., Schafer, W., Bastert, J., Chen, W., and Hube, B. (1999) Secreted aspartic proteinase (Sap) activity contributes to tissue damage in a model of human oral candidosis. *Mol Microbiol* **34**: 169-180.
- Schaller, M., Schackert, C., Korting, H.C., Januschke, E., and Hube, B. (2000) Invasion of *Candida albicans* correlates with expression of secreted aspartic proteinases during experimental infection of human epidermis. *J Invest Dermatol* **114**: 712-717.
- Schaller, M., Bein, M., Korting, H.C., Baur, S., Hamm, G., Monod, M., Beinhauer, S., and Hube, B. (2003) The secreted aspartyl proteinases Sap1 and Sap2 cause tissue damage in an in vitro model of vaginal

- candidiasis based on reconstituted human vaginal epithelium. *Infect Immun* **71**: 3227-3234.
- Schauer, F., and Hanschke, R. (1999) [Taxonomy and ecology of the genus *Candida*]. *Mycoses* **42 Suppl 1**: 12-21.
- Schweizer, A., Rupp, S., Taylor, B.N., Rollinghoff, M., and Schroppel, K. (2000) The TEA/ATTS transcription factor CaTec1p regulates hyphal development and virulence in *Candida albicans*. *Mol Microbiol* **38**: 435-445.
- Sinha, S., and Tompa, M. (2002) Discovery of novel transcription factor binding sites by statistical overrepresentation. *Nucleic Acids Res* **30**: 5549-5560.
- Slutsky, B., Buffo, J., and Soll, D.R. (1985) High-frequency switching of colony morphology in *Candida albicans*. *Science* **230**: 666-669.
- Slutsky, B., Staebell, M., Anderson, J., Risen, L., Pfaller, M., and Soll, D.R. (1987) "White-opaque transition": a second high-frequency switching system in *Candida albicans*. *J Bacteriol* **169**: 189-197.
- Smolenski, G., Sullivan, P.A., Cutfield, S.M., and Cutfield, J.F. (1997) Analysis of secreted aspartic proteinases from *Candida albicans*: purification and characterization of individual Sap1, Sap2 and Sap3 isoenzymes. *Microbiology* **143 (Pt 2)**: 349-356.
- Soll, D.R. (1997) Gene regulation during high-frequency switching in *Candida albicans*. *Microbiology* **143 (Pt 2)**: 279-288.
- Soll, D.R. (2002) *Candida* commensalism and virulence: the evolution of phenotypic plasticity. *Acta Trop* **81**: 101-110.
- Sonneborn, A., Tebarth, B., and Ernst, J.F. (1999) Control of white-opaque phenotypic switching in *Candida albicans* by the Efg1p morphogenetic regulator. *Infect Immun* **67**: 4655-4660.
- Staab, J.F., Ferrer, C.A., and Sundstrom, P. (1996) Developmental expression of a tandemly repeated, proline- and glutamine-rich amino acid motif on hyphal surfaces on *Candida albicans*. *J Biol Chem* **271**: 6298-6305.
- Staab, J.F., Bradway, S.D., Fidel, P.L., and Sundstrom, P. (1999) Adhesive and mammalian transglutaminase substrate properties of *Candida albicans* Hwp1. *Science* **283**: 1535-1538.
- Staib, F. (1965) Serum-proteins as nitrogen source for yeastlike fungi. *Sabouraudia* **4**: 187-193.
- Staib, P., Kretschmar, M., Nichterlein, T., Hof, H., and Morschhauser, J. (2000) Differential activation of a *Candida albicans* virulence gene family during infection. *Proc Natl Acad Sci U S A* **97**: 6102-6107.
- Staib, P., Binder, A., Kretschmar, M., Nichterlein, T., Schroppel, K., and Morschhauser, J. (2004) Tec1p-independent activation of a hypha-associated *Candida albicans* virulence gene during infection. *Infect Immun* **72**: 2386-2389.
- Stehr, F., Felk, A., Kretschmar, M., Schaller, M., Schafer, W., and Hube, B. (2000) [Extracellular hydrolytic enzymes and their relevance during *Candida albicans* infections.]. *Mycoses* **43 Suppl 2**: 17-21.
- Stoldt, V.R., Sonneborn, A., Leuker, C.E., and Ernst, J.F. (1997) Efg1p, an essential regulator of morphogenesis of the human pathogen *Candida*

- albicans, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. *Embo J* **16**: 1982-1991.
- Sugiyama, Y., Nakashima, S., Mirbod, F., Kanoh, H., Kitajima, Y., Ghannoum, M.A., and Nozawa, Y. (1999) Molecular cloning of a second phospholipase B gene, caPLB2 from *Candida albicans*. *Med Mycol* **37**: 61-67.
- Sullivan, D.J., Westerneng, T.J., Haynes, K.A., Bennett, D.E., and Coleman, D.C. (1995) *Candida dubliniensis* sp. nov.: phenotypic and molecular characterization of a novel species associated with oral candidosis in HIV-infected individuals. *Microbiology* **141 (Pt 7)**: 1507-1521.
- Sundstrom, P. (2002) Adhesion in *Candida* spp. *Cell Microbiol* **4**: 461-469.
- Takahashi, K., Inoue, H., Sakai, K., Kohama, T., Kitahara, S., Takishima, K., Tanji, M., Athauda, S.B., Takahashi, T., Akanuma, H., and et al. (1991) The primary structure of *Aspergillus niger* acid proteinase A. *J Biol Chem* **266**: 19480-19483.
- Timpel, C., Strahl-Bolsinger, S., Ziegelbauer, K., and Ernst, J.F. (1998) Multiple functions of Pmt1p-mediated protein O-mannosylation in the fungal pathogen *Candida albicans*. *J Biol Chem* **273**: 20837-20846.
- Timpel, C., Zink, S., Strahl-Bolsinger, S., Schroppel, K., and Ernst, J. (2000) Morphogenesis, adhesive properties, and antifungal resistance depend on the Pmt6 protein mannosyltransferase in the fungal pathogen *Candida albicans*. *J Bacteriol* **182**: 3063-3071.
- Togni, G., Sanglard, D., Quadroni, M., Foundling, S.I., and Monod, M. (1996) Acid proteinase secreted by *Candida tropicalis*: functional analysis of preproregion cleavages in *C. tropicalis* and *Saccharomyces cerevisiae*. *Microbiology* **142 (Pt 3)**: 493-503.
- Tokunaga, M., Kusamichi, M., and Koike, H. (1986) Ultrastructure of outermost layer of cell wall in *Candida albicans* observed by rapid-freezing technique. *J Electron Microsc (Tokyo)* **35**: 237-246.
- Tournu, H., Tripathi, G., Bertram, G., Macaskill, S., Mavor, A., Walker, L., Odds, F.C., Gow, N.A., and Brown, A.J. (2005) Global role of the protein kinase Gcn2 in the human pathogen *Candida albicans*. *Eukaryot Cell* **4**: 1687-1696.
- Tripathi, G., Wiltshire, C., Macaskill, S., Tournu, H., Budge, S., and Brown, A.J. (2002) Gcn4 co-ordinates morphogenetic and metabolic responses to amino acid starvation in *Candida albicans*. *Embo J* **21**: 5448-5456.
- Tsuboi, R., Ogawa, H., Bramono, K., Richardson, M.D., Shankland, G.S., Crozier, W.J., Sei, Y., Ninomiya, J., Nakabayashi, A., Takaiuchi, I., and et al. (1994) Pathogenesis of superficial mycoses. *J Med Vet Mycol* **32 Suppl 1**: 91-104.
- Van Der Vaart, J.M., te Biesebeke, R., Chapman, J.W., Klis, F.M., and Verrips, C.T. (1996) The beta-1, 6-glucan containing side-chain of cell wall proteins of *Saccharomyces cerevisiae* is bound to the glycan core of the GPI moiety. *FEMS Microbiol Lett* **145**: 401-407.
- von Heijne, G. (1983) Patterns of amino acids near signal-sequence cleavage sites. *Eur J Biochem* **133**: 17-21.

- von Heijne, G. (1984) How signal sequences maintain cleavage specificity. *J Mol Biol* **173**: 243-251.
- Walther, A., and Wendland, J. (2003) An improved transformation protocol for the human fungal pathogen *Candida albicans*. *Curr Genet* **42**: 339-343.
- Wang, W., Cherry, J.M., Botstein, D., and Li, H. (2002) A systematic approach to reconstructing transcription networks in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* **99**: 16893-16898.
- Watts, H.J., Cheah, F.S., Hube, B., Sanglard, D., and Gow, N.A. (1998) Altered adherence in strains of *Candida albicans* harbouring null mutations in secreted aspartic proteinase genes. *FEMS Microbiol Lett* **159**: 129-135.
- White, T.C., Miyasaki, S.H., and Agabian, N. (1993) Three distinct secreted aspartyl proteinases in *Candida albicans*. *J Bacteriol* **175**: 6126-6133.
- White, T.C., and Agabian, N. (1995) *Candida albicans* secreted aspartyl proteinases: isoenzyme pattern is determined by cell type, and levels are determined by environmental factors. *J Bacteriol* **177**: 5215-5221.
- Wingard, J.R. (1995) Importance of *Candida* species other than *C. albicans* as pathogens in oncology patients. *Clin Infect Dis* **20**: 115-125.
- Wright, R.J., Carne, A., Hieber, A.D., Lamont, I.L., Emerson, G.W., and Sullivan, P.A. (1992) A second gene for a secreted aspartate proteinase in *Candida albicans*. *J Bacteriol* **174**: 7848-7853.
- Wu, T., and Samaranayake, L.P. (1999) The expression of secreted aspartyl proteinases of *Candida* species in human whole saliva. *J Med Microbiol* **48**: 711-720.
- Yonezawa, S., Takahashi, T., Wang, X.J., Wong, R.N., Hartsuck, J.A., and Tang, J. (1988) Structures at the proteolytic processing region of cathepsin D. *J Biol Chem* **263**: 16504-16511.
- Zakrzewska, A., Boorsma, A., Brul, S., Hellingwerf, K.J., and Klis, F.M. (2005) Transcriptional response of *Saccharomyces cerevisiae* to the plasma membrane-perturbing compound chitosan. *Eukaryot Cell* **4**: 703-715.
- Zaragoza, O., Rodriguez, C., and Gancedo, C. (2000) Isolation of the MIG1 gene from *Candida albicans* and effects of its disruption on catabolite repression. *J Bacteriol* **182**: 320-326.
- Zhao, X., Oh, S.H., Cheng, G., Green, C.B., Nuessen, J.A., Yeater, K., Leng, R.P., Brown, A.J., and Hoyer, L.L. (2004) ALS3 and ALS8 represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p. *Microbiology* **150**: 2415-2428.