

6

Literaturverzeichnis

1. Akerstrom G, Malmaeus J, Grimelius L, Ljunghall S, Bergstrom R. Histological changes in parathyroid glands in subclinical and clinical renal disease. An autopsy investigation. *Scand J Urol Nephrol.* 1984;18(1):75-84.
2. Anthony LB, Martin W, Delbeke D, Sandler M. Somatostatin receptor imaging: predictive and prognostic considerations. *Digestion.* 1996;57 Suppl 1:50-3.
3. Amthauer H, Ruf J, Bohmig M, Lopez-Hanninen E, Rohlfing T, Wernecke KD, Plockinger U, Gutberlet M, Lemke AJ, Steinmuller T, Wiedenmann B, Felix R. Diagnosis of neuroendocrine tumours by retrospective image fusion: is there a benefit? *Eur J Nucl Med Mol Imaging.* 2004;31:342-8.
4. Amthauer H, Ruf J, Rösch T, Ricke J. Bildgebung bei neuroendokrinen Tumoren. *Onkologe.* 2004;10:588-97.
5. Amthauer H, Denecke T, Rohlfing T, Ruf J, Bohmig M, Gutberlet M, Plockinger U, Felix R, Lemke AJ. Value of image fusion using single photon emission computed tomography with integrated low dose computed tomography in comparison with a retrospective voxel-based method in neuroendocrine tumours. *Eur Radiol.* 2005;15:1456-62.
6. Amthauer H, Wurm R, Kuczer D, Ruf J, Michel R, Eisenacher J, Stockhammer F, Denecke T, Felix R, Plotkin M. Relevance of image fusion with MRI for the interpretation of I-123 iodo-methyl-tyrosine scans in patients with suspected recurrent or residual brain tumor. *Clin Nucl Med.* 2006 (im Druck).
7. Baum U, Anders K, Steinbichler G, Lell M, Greess H, Riedel T, Kachelriess M, Kalender WA, Bautz WA. Improvement of image quality of multislice spiral CT scans of the head and neck region using a raw data-based multidimensional adaptive filtering (MAF) technique. *Eur Radiol.* 2004;14:1873-81.

8. Beil W, Rohr K, Stiehl S. Investigation of Approaches for the Localization of Anatomical Landmarks in 3D Medical Images. In: Lemke HU, Vannier MW und Inamura K, eds. Computer Assisted Radiology and Surgery, No. 1134 in Excerpta Medica, International Congress Series, Amsterdam: Elsevier, 1997:265-70.
9. Biersack HJ, Coenen HH, Stocklin G, Reichmann K, Bockisch A, Oehr P, Kashab M, Rollmann O. Imaging of brain tumors with L-3-[123I]iodo-alpha-methyl tyrosine and SPECT. J Nucl Med. 1989;30:110-2.
10. Bocher M, Balan A, Krausz Y, Shrem Y, Lonn A, Wilk M, Chisin R. Gamma camera-mounted anatomical X-ray tomography: technology, system characteristics and first images. Eur J Nucl Med. 2000;27:619-27.
11. Bock JC, Sander B, Schedel H, Felix R. Magnetic resonance tomographic characterization of intracranial tumors by evaluating the blood-brain barrier and regional cerebral circulation. Aktuelle Radiol. 1993;3:346-50.
12. Braun V, Dempf S, Tomczak R, Wunderlich A, Weller R, Richter HP. Functional cranial neuronavigation. Direct integration of fMRI and PET data. J Neuroradiol. 2000;27:157-63.
13. Bronstein AD, Nyberg DA, Schwartz AN, Shuman WP, Griffin BR. Soft-tissue changes after head and neck radiation: CT findings. AJNR Am J Neuroradiol. 1989;10:171-5.
14. Caixas A, Berna L, Hernandez A, Tebar FJ, Madariaga P, Vegazo O, Bittini AL, Moreno B, Faure E, Abos D, Piera J, Rodriguez JM, Farrerons J, Puig-Domingo M. Efficacy of preoperative diagnostic imaging localization of technetium 99m-sestamibi scintigraphy in hyperparathyroidism. Surgery. 1997;121:535-41.
15. Chisin R. Nuclear medicine in head and neck oncology: reality and perspectives. J Nucl Med. 1999;40:91-5.

16. Collignon A, Maes F, Delaere D, Vandermeulen D, Suetens P, Marchal G. Automated multi-modality image using information theory. In: Bizais Y and Barillot C, eds. Information Processing in Medical Imaging. Dordrecht, The Netherlands: Kluwer Academic Publishers, 1995:263-74.
17. Cook GJ, Ott RJ. Dual-modality imaging. *Eur Radiol*. 2001;11:1857-58.
18. Davis JP, Maisey MN, Chevretton EB. Positron emission tomography--a useful imaging technique for otolaryngology, head and neck surgery? *J Laryngol Otol*. 1998;112:125-7.
19. Delbeke D, Meyerowitz C, Lapidus RL, Maciunas RJ, Jennings MT, Moots PL, Kessler RM. Optimal cutoff levels of F-18 fluorodeoxyglucose uptake in the differentiation of low-grade from high-grade brain tumors with PET. *Radiology*. 1995;195:47-52.
20. Denecke T, Hildebrandt B, Lehmkuhl L, Peters N, Nicolaou A, Pech M, Riess H, Ricke J, Felix R, Amthauer H. Fusion imaging using a hybrid SPECT-CT camera improves port perfusion scintigraphy for control of hepatic arterial infusion of chemotherapy in colorectal cancer patients. *Eur J Nucl Med Mol Imaging*. 2005;32:1003-10.
21. De Salles AA, Brekhus SD, De Souza EC, Behnke EJ, Farahani K, Anzai Y, Lufkin R. Early postoperative appearance of radiofrequency lesions on magnetic resonance imaging. *Neurosurgery*. 1995;36:932-6.
22. Di Chiro G, DeLaPaz RL, Brooks RA, Sokoloff L, Kornblith PL, Smith BH, Patronas NJ, Kufta CV, Kessler RM, Johnston GS, Manning RG, Wolf AP. Glucose utilization of cerebral gliomas measured by [18F] fluorodeoxyglucose and positron emission tomography. *Neurology*. 1982;32:1323-9.
23. Dierickx LO, Lahoutte T, Deron P, Caveliers V, Vanhove C, Everaert H, Bossuyt A. Diagnosis of recurrent head and neck squamous cell carcinoma with 3-[123I]iodo-L-alpha-methyltyrosine SPET. *Eur J Nucl Med*. 2001;28:282-7.

24. Dierickx LO, Everaert H, Deron P, Voordeckers M, Lahoutte T, Bossuyt A. Evaluation of the response to therapy of head and neck squamous cell carcinoma by using 3-[123I]iodo-L-alpha-methyl tyrosine and single photon emission tomography. *Nucl Med Commun.* 2003;24:633-41.
25. Even-Sapir E, Keidar Z, Sachs J, Engel A, Bettman L, Gaitini D, Guralnik L, Werbin N, Iosilevsky G, Israel O. The new technology of combined transmission and emission tomography in evaluation of endocrine neoplasms. *J Nucl Med.* 2001;42:998-1004.
26. Flamen P, Bernheim N, Deron P, Caveliers V, Chavatte K, Franken PR, Bossuyt A. Iodine-123 alpha-methyl-l-tyrosine single-photon emission tomography for the visualization of head and neck squamous cell carcinomas. *Eur J Nucl Med.* 1998;25:177-81.
27. Forster GJ, Laumann C, Nickel O, Kann P, Rieker O, Bartenstein P. SPET/CT image co-registration in the abdomen with a simple and cost-effective tool. *Eur J Nucl Med Mol Imaging.* 2003;30:32-9.
28. Freudenberg LS, Antoch G, Jentzen W, Pink R, Knust J, Gorges R, Muller SP, Bockisch A, Debatin JF, Brandau W. Value of (124)I-PET/CT in staging of patients with differentiated thyroid cancer. *Eur Radiol.* 2004;14:2092-8.
29. Gabriel M, Hausler F, Bale R, Moncayo R, Decristoforo C, Kovacs P, Virgolini I. Image fusion analysis of (99m)Tc-HYNIC-Tyr(3)-octreotide SPECT and diagnostic CT using an immobilisation device with external markers in patients with endocrine tumours. *Eur J Nucl Med Mol Imaging.* 2005;32:1440-51.
30. Giordano A, Rubello D, Casara D. New trends in parathyroid scintigraphy. *Eur J Nucl Med.* 2001;28:1409-20.

31. Goerres GW, Kamel E, Heidelberg TN, Schwitter MR, Burger C, von Schulthess GK. PET-CT image co-registration in the thorax: influence of respiration. *Eur J Nucl Med Mol Imaging*. 2002;29:351-60.
32. Goldman S, Levivier M, Pirotte B, Brucher JM, Wikler D, Damhaut P, Dethy S, Brotchi J, Hildebrand J. Regional methionine and glucose uptake in high-grade gliomas: a comparative study on PET-guided stereotactic biopsy. *J Nucl Med*. 1997;38:1459-62.
33. Gotway MB, Reddy GP, Webb WR, Morita ET, Clark OH, Higgins CB. Comparison between MR imaging and 99mTc MIBI scintigraphy in the evaluation of recurrent or persistent hyperparathyroidism. *Radiology*. 2001;218:783-90.
34. Grosu AL, Feldmann H, Dick S, Dzewas B, Nieder C, Gumprecht H, Frank A, Schwaiger M, Molls M, Weber WA. Implications of IMT-SPECT for postoperative radiotherapy planning in patients with gliomas. *Int J Radiat Oncol Biol Phys*. 2002;54:842-54.
35. Grosu AL, Lachner R, Wiedenmann N, Stark S, Thamm R, Kneschaurek P, Schwaiger M, Molls M, Weber WA. Validation of a method for automatic image fusion (BrainLAB System) of CT data and 11C-methionine-PET data for stereotactic radiotherapy using a LINAC: first clinical experience. *Int J Radiat Oncol Biol Phys*. 2003;56:1450-63.
36. Grosu AL, Weber WA, Franz M, Stark S, Piert M, Thamm R, Gumprecht H, Schwaiger M, Molls M, Nieder C. Reirradiation of recurrent high-grade gliomas using amino acid PET (SPECT)/CT/MRI image fusion to determine gross tumor volume for stereotactic fractionated radiotherapy. *Int J Radiat Oncol Biol Phys*. 2005;63:511-9.
37. Guth-Tougelidis B, Muller S, Mehdorn MM, Knust EJ, Dutschka K, Reiners C. Uptake of DL-3-123I-iodo-alpha-methyltyrosine in recurrent brain tumors. *Nuklearmedizin*. 1995;34:71-5.

38. Hasegawa BH, Stebler B, Rutt BK, Martinez A, Gingold EL, Barker CS, Faulkner KG, Cann CE, Boyd DP. A prototype high-purity germanium detector system with fast photon-counting circuitry for medical imaging. *Med Phys.* 1991;18:900-9.
39. Herrmann KA, Waggershauser T, Sittek H, Reiser MF. Liver intraarterial chemotherapy: use of the femoral artery for percutaneous implantation of catheter-port systems. *Radiology.* 2000;215:294-9.
40. Hill DL, Hawkes DJ, Crossman JE, Gleeson MJ, Cox TC, Bracey EE, Strong AJ, Graves P. Registration of MR and CT images for skull base surgery using point-like anatomical features. *Br J Radiol.* 1991;64:1030-5.
41. Hofler H, Stier A, Schusdziarra V, Siewert JR. Classification of neuroendocrine tumors of the gastrointestinal tract and pancreas and its therapeutic relevance. *Chirurg.* 1997;68:107-15.
42. Hosten N, Kreissig R, Puls R, Amthauer H, Beier J, Rohlfing T, Stroszczynski C, Herbel A, Lemke AJ, Felix R. Fusion of CT and PET data: methods and clinical relevance for planning laser-induced thermotherapy of liver metastases. *Rofo.* 2000;172:630-5.
43. Hutton BF, Braun M, Thurfjell L, Lau DY. Image registration: an essential tool for nuclear medicine. *Eur J Nucl Med Mol Imaging.* 2002;29:559-77.
44. Hutton BF, Braun M. Software for image registration: algorithms, accuracy, efficacy. *Semin Nucl Med.* 2003;33:180-92.
45. Ishiwata K, Kubota K, Murakami M, Kubota R, Sasaki T, Ishii S, Senda M. Re-evaluation of amino acid PET studies: can the protein synthesis rates in brain and tumor tissues be measured in vivo? *J Nucl Med.* 1993;34:1936-43.
46. Israel O, Keidar Z, Iosilevsky G, Bettman L, Sachs J, Frenkel A. The fusion of anatomic and physiologic imaging in the management of patients with cancer. *Semin Nucl Med.* 2001;31:191-205.

47. Kaczirek K, Prager G, Kienast O, Dobrozemsky G, Dudczak R, Niederle B, Kurtaran A. Combined transmission and (99m)Tc-sestamibi emission tomography for localization of mediastinal parathyroid glands. *Nuklearmedizin*. 2003;42:220-3.
48. Kaschten B, Stevenaert A, Sadzot B, Deprez M, Degueldre C, Del Fiore G, Luxen A, Reznik M. Preoperative evaluation of 54 gliomas by PET with fluorine-18-fluorodeoxyglucose and/or carbon-11-methionine. *J Nucl Med*. 1998;39:778-85.
49. Krausz Y, Lebensart PD, Klein M, Weininger J, Blachar A, Chisin R, Shiloni E. Preoperative localization of parathyroid adenoma in patients with concomitant thyroid nodular disease. *World J Surg*. 2000;24:1573-8.
50. Krausz Y, Bettman L, Guralnik L, Yosilevsky G, Keidar Z, Bar-Shalom R, Even-Sapir E, Chisin R, Israel O. Technetium-99m-MIBI SPECT/CT in primary hyperparathyroidism. *World J Surg*. 2006;30:76-83.
51. Krenning EP, Kwekkeboom DJ, Bakker WH, Breeman WA, Kooij PP, Oei HY, van Hagen M, Postema PT, de Jong M, Reubi JC, et al. Somatostatin receptor scintigraphy with [111In-DTPA-D-Phe1]- and [123I-Tyr3]-octreotide: the Rotterdam experience with more than 1000 patients. *Eur J Nucl Med*. 1993;20:716-31.
52. Krummeich C, Holschbach M, Stocklin G. Direct n.c.a. electrophilic radioiodination of tyrosine analogues; their in vivo stability and brain-uptake in mice. *Appl Radiat Isot*. 1994;45:929-35.
53. Kulke MH, Mayer RJ. Carcinoid tumors. *N Engl J Med*. 1999;340:858-68.
54. Kuwert T, Woesler B, Morgenroth C, Lerch H, Schafers M, Palkovic S, Matheja P, Brandau W, Wassmann H, Schober O. Diagnosis of recurrent glioma with SPECT and iodine-123-alpha-methyl tyrosine. *J Nucl Med*. 1998;39:23-7.

55. Lang TF, Hasegawa BH, Liew SC, Brown JK, Blakespoor SC, Reilly SM, Gingold EL, Cann CE. Description of a prototype emission-transmission computed tomography imaging system. *J Nucl Med.* 1992;33:1881-7.
56. Langen KJ, Ziemons K, Kiwit JC, Herzog H, Kuwert T, Bock WJ, Stocklin G, Feinendegen LE, Muller-Gartner HW. 3-[123I]iodo-alpha-methyltyrosine and [methyl-11C]-L-methionine uptake in cerebral gliomas: a comparative study using SPECT and PET. *J Nucl Med.* 1997;38:517-22.
57. Lehner K, Reiser M, Gebhardt U, Heuck A, Schaff J. DSA--control of implanted devices for arterial hepatic perfusion. *Cardiovasc Intervent Radiol.* 1987;10:71-4.
58. Lemke AJ, Sander B, Balzer T, Geens V, Hosten N, Felix R. Safety and use of gadobutrol in patients with brain tumors (phase III trial). *Rofo.* 1997;167:591-8.
59. Lemke AJ, Niehues SM, Amthauer H, Rohlfing T, Hosten N, Felix R. Clinical use of digital retrospective image fusion of CT, MRI, FDG-PET and SPECT - fields of indications and results. *Rofo.* 2004;176:1811-8.
60. Lemke AJ, Niehues SM, Hosten N, Amthauer H, Boehmig M, Stroszczynski C, Rohlfing T, Rosewicz S, Felix R. Retrospective digital image fusion of multidetector CT and 18F-FDG PET: clinical value in pancreatic lesions--a prospective study with 104 patients. *J Nucl Med.* 2004;45:1279-86.
61. Lester H, Arridge SR. A survey of non-linear medical image registration. *Pattern recognition.* 1999; 32:129-49.
62. Lopez Hanninen E, Vogl TJ, Steinmuller T, Ricke J, Neuhaus P, Felix R. Preoperative contrast-enhanced MRI of the parathyroid glands in hyperparathyroidism. *Invest Radiol.* 2000;35:426-30.
63. Lubin E, Cyjon A, Neuman M. Technetium-99m macroaggregates for the study of the distribution of arterial infusion chemotherapy in the liver. *Clin Nucl Med.* 1987;12:385-8.

64. MACG (Meta Analysis Group in Cancer). Reappraisal of hepatic arterial infusion in the treatment of nonresectable liver metastases from colorectal cancer. Meta-Analysis Group in Cancer. *J Natl Cancer Inst.* 1996;88:252-8.
65. Maintz JB, Viergever MA. A survey of medical image registration. *Med Image Anal.* 1998;2:1-36.
66. Maisey MN, Hawkes DJ, Lukawieck-Vydelingum AM. Synergistic imaging. *Eur J Nucl Med.* 1992;19:1002-5.
67. Matheja P, Weckesser M, Rickert Ch, Franzius Ch, Palkovic S, Riemann B, Schober O. I-123-lodo-alpha-methyl tyrosine SPECT in non-parenchymal brain tumours. *Nuklearmedizin.* 2002;41:191-6.
68. Mazzeo S, Caramella D, Lencioni R, Molea N, De Liperi A, Marcocci C, Miccoli P, Iacconi P, Bossio GB, Viacava P, Lazzeri E, Bartolozzi C. Comparison among sonography, double-tracer subtraction scintigraphy, and double-phase scintigraphy in the detection of parathyroid lesions. *AJR Am J Roentgenol.* 1996;166:1465-70.
69. Modlin IM, Sandor A. An analysis of 8305 cases of carcinoid tumors. *Cancer.* 1997;79:813-29.
70. Modlin IM, Kidd M, Latich I, Zikusoka MN, Shapiro MD. Current status of gastrointestinal carcinoids. *Gastroenterology.* 2005;128:1717-51.
71. Moka D, Voth E, Larena-Avellaneda A, Schicha H. 99m-Tc-MIBI SPECT parathyroid gland scintigraphy for the preoperative localization of small parathyroid gland adenomas. *Nuklearmedizin.* 1997;36:240-4.
72. Moka D, Voth E, Dietlein M, Larena-Avellaneda A, Schicha H. Technetium 99m-MIBI-SPECT: A highly sensitive diagnostic tool for localization of parathyroid adenomas. *Surgery.* 2000;128:29-35.

73. Moka D, Voth E, Dietlein M, Larena-Avellaneda A, Schicha H. Preoperative localization of parathyroid adenomas using 99mTc-MIBI scintigraphy. *Am J Med.* 2000;108:733-6.
74. Morimoto M, Satake M, Sekiguchi R, Haruno M, Moriyama N. Optimal injection protocol for CT evaluation during hepatic arterial infusion chemotherapy. *Invest Radiol.* 1999;34:744-50.
75. Myers E, Leffall L. Head and neck oncology: diagnosis, treatment and rehabilitation. Boston, USA: Little, Brown and Company, 1991.
76. Oberg K, Eriksson B. Nuclear medicine in the detection, staging and treatment of gastrointestinal carcinoid tumours. *Best Pract Res Clin Endocrinol Metab.* 2005;19:265-76.
77. Oberg K. Neuroendocrine tumors of the gastrointestinal tract: recent advances in molecular genetics, diagnosis, and treatment. *Curr Opin Oncol.* 2005;17:386-91.
78. Ogawa T, Inugami A, Hatazawa J, Kanno I, Murakami M, Yasui N, Mineura K, Uemura K. Clinical positron emission tomography for brain tumors: comparison of fludeoxyglucose F 18 and L-methyl-11C-methionine. *AJNR Am J Neuroradiol.* 1996;17:345-53.
79. Paulino AC, Thorstad WL, Fox T. Role of fusion in radiotherapy treatment planning. *Semin Nucl Med.* 2003;33:238-43.
80. Pascher A, Steinmuller T, Radke C, Hosten N, Wiedenmann B, Neuhaus P, Bechstein WO. Primary and secondary hepatic manifestation of neuroendocrine tumors. *Langenbecks Arch Surg.* 2000;385:265-70.
81. Pelizzari CA, Chen GT, Spelbring DR, Weichselbaum RR, Chen CT. Accurate three-dimensional registration of CT, PET, and/or MR images of the brain. *J Comput Assist Tomogr.* 1989;13:20-6.

82. Perault C, Schvartz C, Wampach H, Liehn JC, Delisle MJ. Thoracic and abdominal SPECT-CT image fusion without external markers in endocrine carcinomas. The Group of Thyroid Tumoral Pathology of Champagne-Ardenne. *J Nucl Med*. 1997;38:1234-42.
83. Perrier ND, Ituarte PH, Morita E, Hamill T, Gielow R, Duh QY, Clark OH. Parathyroid surgery: separating promise from reality. *J Clin Endocrinol Metab*. 2002;87:1024-9.
84. Pfannenberg AC, Eschmann SM, Horger M, Lamberts R, Vonthein R, Claussen CD, Bares R. Benefit of anatomical-functional image fusion in the diagnostic work-up of neuroendocrine neoplasms. *Eur J Nucl Med Mol Imaging*. 2003;30:835-43.
85. Pietrzyk U, Herholz K, Fink G, Jacobs A, Mielke R, Slansky I, Wurker M, Heiss WD. An interactive technique for three-dimensional image registration: validation for PET, SPECT, MRI and CT brain studies. *J Nucl Med*. 1994;35:2011-8.
86. Plockinger U, Rindi G, Arnold R, Eriksson B, Krenning EP, de Herder WW, Goede A, Caplin M, Oberg K, Reubi JC, Nilsson O, Delle Fave G, Ruszniewski P, Ahlman H, Wiedenmann B; European Neuroendocrine Tumour Society. Guidelines for the diagnosis and treatment of neuroendocrine gastrointestinal tumours. A consensus statement on behalf of the European Neuroendocrine Tumour Society (ENETS). *Neuroendocrinology*. 2004;80:394-424.
87. Plotkin M, Eisenacher J, Bruhn H, Wurm R, Michel R, Stockhammer F, Feussner A, Dudeck O, Wust P, Felix R, Amthauer H. ¹²³I-IMT SPECT and ^{1H} MR-spectroscopy at 3.0 T in the differential diagnosis of recurrent or residual gliomas: a comparative study. *J Neurooncol*. 2004;70:49-58.
88. Plotkin M, Amthauer H, Eisenacher J, Wurm R, Michel R, Wust P, Stockhammer F, Rottgen R, Gutberlet M, Ruf J, Felix R. Value of ¹²³I-IMT SPECT for diagnosis of recurrent non-astrocytic intracranial tumours. *Neuroradiology*. 2005;47:18-26.

89. Plotkin M, Wurm R, Eisenacher J, Szerewicz K, Michel R, Schlenger L, Pech M, Denecke T, Kuczer D, Bischoff A, Felix R, Amthauer H. Combined SPECT/CT imaging using ¹²³I-IMT in the detection of recurrent or persistent head and neck cancer. *Eur Radiol.* 2006;16:503-11.
90. Pons F, Torregrosa JV, Fuster D. Biological factors influencing parathyroid localization. *Nucl Med Commun.* 2003;24:121-4.
91. Prager G, Czerny C, Kurtaran A, Passler C, Scheuba C, Niederle B. The value of preoperative localization studies in primary hyperparathyroidism. *Chirurg.* 1999;70:1082-8.
92. Reinhardt MJ, Wiethoelter N, Matthies A, Joe AY, Strunk H, Jaeger U, Biersack HJ. PET recognition of pulmonary metastases on PET/CT imaging: impact of attenuation-corrected and non-attenuation-corrected PET images. *Eur J Nucl Med Mol Imaging.* 2006;33:134-9.
93. Ricke J, Klose KJ, Mignon M, Oberg K, Wiedenmann B. Standardisation of imaging in neuroendocrine tumours: results of a European delphi process. *Eur J Radiol.* 2001;37:8-17.
94. Ricke J, Hildebrandt B, Miersch A, Nicolaou A, Warschewske G, Teichgraber U, Lopez Hanninen E, Riess H, Felix R. Hepatic arterial port systems for treatment of liver metastases: factors affecting patency and adverse events. *J Vasc Interv Radiol.* 2004;15:825-33.
95. Rohlfing T, Beier J, Graf R, Wolf M, Wust P, Felix R. Automated integration of CT, MRI and PET for Planning and Dose Distribution Computation in Radiation Therapy. In: Lemke HU, Vannier MW, Inamura K, Farman AG, eds, Computer Assisted Radiology and Surgery (CARS), No. 1191 in Excerpta Medica, International Congress Series, Amsterdam, The Netherlands: Elsevier, 1999:279-284.

96. Rohlfing T, West JB, Beier J, Liebig T, Taschner CA, Thomale UW. Registration of functional and anatomical MRI: accuracy assessment and application in navigated neurosurgery. *Comput Aided Surg.* 2000;5:414-25.
97. Rohlfing T, Beier J. Gesteigerte Registrierungsgenauigkeit verrauschter Bilddaten durch Conincidence Thresholding. In: Horsch A, Lehmann T, eds, *Bildverarbeitung für die Medizin 2000 – Algorithmen, Systeme, Anwendungen*. Informatik Aktuell, Berlin, Deutschland: Springer Verlag, 2000.
98. Rubello D, Casara D, Pagetta C, Piotto A, Pelizzo MR, Shapiro B. Determinant role of Tc-99m MIBI SPECT in the localization of a retrotracheal parathyroid adenoma successfully treated by radioguided surgery. *Clin Nucl Med.* 2002;27:711-5.
99. Ruf J, Lehmkuhl L, Bertram H, Sandrock D, Amthauer H, Humplik B, Munz DL, Felix R. Impact of SPECT and integrated low-dose CT after radioiodine therapy on the management of patients with thyroid carcinoma. *Nucl Med Commun.* 2004;25:1177-82.
100. Ruf J, Lopez Hanninen E, Steinmuller T, Rohlfing T, Bertram H, Gutberlet M, Lemke AJ, Felix R, Amthauer H. Preoperative localization of parathyroid glands. Use of MRI, scintigraphy, and image fusion. *Nuklearmedizin.* 2004;43:85-90.
101. Sadahiro S, Suzuki T, Ishikawa K, Yasuda S, Tajima T, Makuuchi H, Saitoh T, Murayama C. Prophylactic hepatic arterial infusion chemotherapy for the prevention of liver metastasis in patients with colon carcinoma: a randomized control trial. *Cancer.* 2004;100:590-7.
102. Samnick S, Bader JB, Hellwig D, Moringlane JR, Alexander C, Romeike BF, Feiden W, Kirsch CM. Clinical value of iodine-123-alpha-methyl-L-tyrosine single-photon emission tomography in the differential diagnosis of recurrent brain tumor in patients pretreated for glioma at follow-up. *J Clin Oncol.* 2002;20:396-404.

103. Sato N, Suzuki M, Kuwata N, Kuroda K, Wada T, Beppu T, Sera K, Sasaki T, Ogawa A. Evaluation of the malignancy of glioma using ¹¹C-methionine positron emission tomography and proliferating cell nuclear antigen staining. *Neurosurg Rev.* 1999;22:210-4.
104. Schillaci O, Simonetti G. Fusion imaging in nuclear medicine--applications of dual-modality systems in oncology. *Cancer Biother Radiopharm.* 2004;19:1-10.
105. Schillaci O. Functional-anatomical image fusion in neuroendocrine tumors. *Cancer Biother Radiopharm.* 2004;19:129-34.
106. Schillaci O, Danieli R, Manni C, Capoccetti F, Simonetti G. Technetium-99m-labelled red blood cell imaging in the diagnosis of hepatic haemangiomas: the role of SPECT/CT with a hybrid camera. *Eur J Nucl Med Mol Imaging.* 2004;31:1011-5.
107. Schmidbauer S, Ladurner R, Juckstock H, Trupka AW, Mussack T, Hallfeldt KK. Surgical and adjuvant therapy of neuroendocrine tumors of the gastrointestinal tract and their metastases. A retrospective analysis of personal patient group. *Chirurg.* 2001;72:945-52.
108. Seki H, Ozaki T, Takaki S, Ooi H, Oda J, Shiina M. Using slow-infusion MR arteriography and an implantable port system to assess drug distribution at hepatic arterial infusion chemotherapy. *AJR Am J Roentgenol.* 2003;180:681-6.
109. Shekhar R, Walimbe V, Raja S, Zagrodsky V, Kanvinde M, Wu G, Bybel B. Automated 3-dimensional elastic registration of whole-body PET and CT from separate or combined scanners. *J Nucl Med.* 2005 Sep;46:1488-96.
110. Shreve PD. Adding structure to function. *J Nucl Med.* 2000;41:1380-2.
111. Slooter GD, Mearadji A, Breeman WA, Marquet RL, de Jong M, Krenning EP, van Eijck CH. Somatostatin receptor imaging, therapy and new strategies in patients with neuroendocrine tumours. *Br J Surg.* 2001;88:31-40.

112. Som PM. Detection of metastasis in cervical lymph nodes: CT and MR criteria and differential diagnosis. *AJR Am J Roentgenol.* 1992;158:961-9.
113. Studholme C, Hill DL, Hawkes DJ. Automated three-dimensional registration of magnetic resonance and positron emission tomography brain images by multiresolution optimization of voxel similarity measures. *Med Phys.* 1997;24:25-35.
114. Studholme C, Hill DLG, Hawkes DJ. An Overlap Invariant Entropy Measure of 3D Medical Image Alignment. *Pattern Recognition.* 1999;33:71-86.
115. Sugimoto E, Lorelius LE, Eriksson B, Oberg K. Midgut carcinoid tumours. CT appearance. *Acta Radiol.* 1995;36:367-71.
116. Suzuki Y, Kobayashi S, Yasuda S. Application of single-photon emission computed tomography (SPECT) with ^{99m}Tc -MAA in evaluation of perfusion patterns during hepatic infusion chemotherapy. *Ann Nucl Med.* 1991;5:123-6.
117. Takebayashi S, Hidai H, Chiba T, Takagi Y, Nagatani Y, Matsubara S. Hyperfunctional parathyroid glands with ^{99m}Tc -MIBI scan: semiquantitative analysis correlated with histologic findings. *J Nucl Med.* 1999;40:1792-7.
118. Torregrosa JV, Fernandez-Cruz L, Canalejo A, Vidal S, Astudillo E, Almaden Y, Pons F, Rodriguez M. (^{99m}Tc)sestamibi scintigraphy and cell cycle in parathyroid glands of secondary hyperparathyroidism. *World J Surg.* 2000;24:1386-90.
119. Townsend DW, Cherry SR. Combining anatomy and function: the path to true image fusion. *Eur Radiol.* 2001;11:1968-74.
120. Udelsman R. Six hundred fifty-six consecutive explorations for primary hyperparathyroidism. *Ann Surg.* 2002;235:665-70.
121. Viergever MA, Maintz JB, Stokking R. Integration of functional and anatomical brain images. *Biophys Chem.* 1997;68:207-19.

122. Viergever MA, Maintz JB, Niessen WJ, Noordmans HJ, Pluim JP, Stokking R, Vincken KL. Registration, segmentation, and visualization of multimodal brain images. *Comput Med Imaging Graph*. 2001;25:147-51.
123. Viola P, Wells WM. Alignment by maximization of mutual information. *Int J Comput Vision*. 1997;24:137-154.
124. von Schulthess GK, Pelc NJ. Integrated-modality imaging: the best of both worlds. *Acad Radiol*. 2002;9:1241-4.
125. Wahl RL, Quint LE, Cieslak RD, Aisen AM, Koeppe RA, Meyer CR. "Anatometabolic" tumor imaging: fusion of FDG PET with CT or MRI to localize foci of increased activity. *J Nucl Med*. 1993;34:1190-7.
126. Wakamatsu H, Noguchi S, Yamashita H, Yamashita H, Tamura S, Jinnouchi S, Nagamachi S, Futami S. Technetium-99m tetrofosmin for parathyroid scintigraphy: a direct comparison with (99m)Tc-MIBI, (201)Tl, MRI and US. *Eur J Nucl Med*. 2001;28:1817-27.
127. Weber W, Bartenstein P, Gross MW, Kinzel D, Daschner H, Feldmann HJ, Reidel G, Ziegler SI, Lumenta C, Molls M, Schwaiger M. Fluorine-18-FDG PET and iodine-123-IMT SPECT in the evaluation of brain tumors. *J Nucl Med*. 1997;38:802-8.
128. Weiss M, Beneke F, Schmid R, Dresel S, Hahn K. Can supplemental imaging with SPECT technique improve the diagnostic value of preoperative Tc-99m-MIBI scintigraphy in primary hyperparathyroidism? *Med Klin (Munich)*. 2002;97:389-95.
129. West J, Fitzpatrick JM, Wang MY, Dawant BM, Maurer CR Jr, Kessler RM, Maciunas RJ, Barillot C, Lemoine D, Collignon A, Maes F, Suetens P, Vandermeulen D, van den Elsen PA, Napel S, Sumanaweera TS, Harkness B, Hemler PF, Hill DL, Hawkes DJ, Studholme C, Maintz JB, Viergever MA, Malandain G, Pennec X, Noz M, Maguire GQ, Pollack M, Pelizzari CA, Robb RA, Hanson D, Woods RP. Comparison and evaluation of retrospective intermodality brain image registration techniques. *J Comput Assist Tomogr*. 1997;21:554-66.

130. West J, Fitzpatrick JM, Wang MY, Dawant BM, Maurer CR Jr, Kessler RM, Maciunas RJ. Retrospective intermodality registration techniques for images of the head: surface-based versus volume-based. *IEEE Trans Med Imaging*. 1999;18:144-50.
131. Wiedenmann B, Jensen RT, Mignon M, Modlin CI, Skogseid B, Doherty G, Oberg K. Preoperative diagnosis and surgical management of neuroendocrine gastroenteropancreatic tumors: general recommendations by a consensus workshop. *World J Surg*. 1998;22:309-18.
132. Wong TZ, Turkington TG, Hawk TC, Coleman RE. PET and brain tumor image fusion. *Cancer J*. 2004;10:234-42.
133. Ziessman HA, Gyves JW, Juni JE, Wahl FL, Thrall JH, Ensminger WD, Goldstein HA, Dubiansky V. Atlas of hepatic arterial perfusion scintigraphy. *Clin Nucl Med*. 1985;10:675-81.
134. Ziessman HA, Wahl RL, Juni JE, Gyves JE, Ensminger WD, Thrall JH, Keyes JW Jr, Walker SC. The utility of SPECT for ^{99m}Tc -MAA hepatic arterial perfusion scintigraphy. *AJR Am J Roentgenol*. 1985;145:747-51.