

11. LITERATURVERZEICHNIS

1. Altmannsberger, S.H.G., Paneitz, A., Siegmund, W., Kroemer, H.K. (2001). Wirkung der AT₁-Rezeptor-antagonisten. *Pharmazie in unserer Zeit* 30. Jhg. (4):296-301.
2. Ast I. (2000). Normothermic hemoperfusion of isolated pig hearts-Characterization of a new model with consideration of the myocardial oxygen partial pressure in regular perfusion and with occlusion of a coronary artery branch. *Dissertation*. FU-Berlin
3. Bers, D.M. (2002). Cardiac excitation-contraction coupling. *Nature*, Vol. 415:198-205.
4. Boban, M., Stowe D.F., Kampine, J.P., Goldberg, A.H., Bosnjak, Z.J. (1993). Effects of 2, 3-butanedione monoxime in isolated hearts: protection during reperfusion after global ischemia. *J Thorac Cardiovasc Surg* 105(3):532-40.
5. Bruckschlegel, G., Holmer, S.R., Schunkert, H., Jandeleit, K., Grimm, D., Muders, F., Kromer, E.P., Rieger, G.A. (1995). Blockade of the renin-angiotensin system in cardiac pressure overload hypertrophy in rats. *Hypertension* 25(2):250-9.
6. Buckberg, G.D. (1987). Strategies and logic of cardioplegic delivery to prevent, avoid, and reverse ischemic and reperfusion damage. *J Thorac Cardiovasc Surg* 93(1):127-39.
7. Budrikis, A., Bolys, R., Liao, Q., Ingemansson, R., Sjoberg, T., Steen, S. (1998). Function of adult pig hearts after 2 and 12 hours of cold cardioplegic preservation. *Ann Thorac Surg* 66(1):73-8.
8. Budrikis, A., Liao, Q., Bolys, R., Westerlaken, B., Steen, S. (1999). Effects of cardioplegic flushing, storage, and reperfusion on coronary circulation in the pig. *Ann Thorac Surg* 67(5):1345-9.
9. Cleutjens, J.P.M., Blankesteijn, W.M., Daemen M.F., Smits, J.F. (1999). The infarcted myocardium: Simply dead tissue, or lively target for therapeutic interventions *Cardiovasc Res* 44(2):232-241.
10. Danser, A.H., Chowdury, S., de Lannoy, L.M., van der Giessen, W.J., Saxen, P.R., Schalekamp, M.A. (1995). Conversion and degradation of [125I] labelled angiotensin I in isolated perfused porcine coronary and carotid arteries. *Cardiovasc Res* 29(6):789-95.
11. Danser, A.H., van Kats, J.P., Admiraal, P.J., Derkx, F.H., Lamers, J.M., Verdouw, P.D., Saxena, P.R., Schalekamp, M.A. (1994). Cardiac rennin and angiotensins. Uptake from plasma versus in situ synthesis. *Hypertension* 24:37-48.
12. Danser, A.H., Saris, J.J., Schnijt M.P., van Kats, J.P. (1999). Is there a local renin-angiotensin system in the heart? *Cardiovasc Res* 44(2):252-65.
13. Danser, A.H. (2003). Local renin-angiotensin systems: the unanswered questions. *Int J Biochem Cell Biol* 35(6):759-68.

14. Deetjen, P., Speckmann E.-J. (1994). Physiologie, 2. Aufl., München/Wien/Baltimore, Verlag Urban & Schwarzenberg.
15. de Lannoy, L.M., Danser, A.H., van Kats, J.P., Schoemaker, R.G., Saxena, P.R., Schalekamp, M.A. (1997). Renin-angiotensin-system components in the interstitial fluid of the isolated perfused rat heart. Local production of angiotensin I. *Hypertension* 29(6):1240-51.
16. Döring, H.J. (1990). The isolated perfused heart according to Langendorff-technique-function-application. *Physiol Bohemoslov* 39(6):481-504.
17. Ensinger, H., Weichel, T., Lindner, K.H., Grunert, A., Ahnefeld, F.W. (1993). Effects of norepinephrine, epinephrine and dopamine infusion on oxygenconsumption in volunteers. *Crit Care Med* 21(10):1502-8.
18. Farquharson, C.J.A., Struthers A.D. (2002). Gradual Reactivation over time of vascular tissue angiotensin I to Angiotensin II conversion during chronic lisinopril therapy in chronic heart failure. *JACC* 39:767-775
19. Ferrario, C.M. and J.M. Flack (1996). Pathologic consequences of increased angiotensin II activity. *Cardiovasc Drugs Ther* 10(5): 511-8.
20. Friedman, B.J., Grinberg, O.Y., Gringberg, S.A., Swartz, H.M. (1997). Myocardial oxygen tension in isolated erythrocyte-perfused rat hearts and comparison with crystalloid media. *J Mol Cell Cardiol* 29(10):2855-8.
21. Fries, R. (1992). Fleischhygiene und Lebensmitteluntersuchung. Stuttgart, Verlag Eugen Ulmer.
22. Furuta, H., Guo, D.F., Inagami, T. (1992). Molecular cloning and sequencing of the gene encoding human angiotensin II type 1 receptor. *Biochem Biophys Res Commun* 183(1):8-13.
23. Gao, Z.P., Downey, H.F., Suu, J., He, M.X., Mallet, R.T. (1997). Adenosine receptor blockade enhance glycolysis in hypoperfused guinea-pig myocardium. *Cardiovasc Res* 33(1):31-44.
24. Garcia-Dorado, D., Theroux, P., Duran, J.M., Solares, J., Alonso, J., Sanz, E., Munoz, R., Elizag, J., Botas, J., Fernandez-Aviles, F. (1992). Selective inhibition of the contractile apparatus. A new approach to modification of infarct size, infarct composition, and infarct geometry during coronary artery occlusion and reperfusion. *Circulation* 85(3):1160-74.
25. Gibbs, C.L. and C.J. Barclay. (1995). Cardiac efficiency. *Cardiovasc Res* 30(5):627-34.
26. Gorelik, G., Carbini, L.A., Scicli, A.G. (1998). Angiotensin 1-7 induces bradykinin-mediated relaxation in porcine coronary artery. *J Pharmacol Exp Ther* 286(1):403-10.
27. Grimm, D., Cameron D., Griese, D.P., Riegger, G.A.J., Kromer, E.P. (1998). Differential effects of growth hormone on cardiomyocyte and extracellular matrix

- protein remodeling following experimental myocardial infarction. *Cardiovasc Res* 40:297-306.
28. Grynberg, A. and L. Demaison (1996). Fatty acid oxidation in the heart. *J Cardiovasc Pharmacol* 28 Suppl. 1:S11-7.
 29. Hellige, G., Paul, M., Schrader, J. (1998). Herz-Kreislauf-Forum. München, Arcis Verlag.
 30. Henrion, D., Kubis, N., Levy, B.I. (2001). Physiological and pathophysiological functions of the AT (2) subtype receptor of angiotensin II: from large arteries to the microcirculation. *Hypertension* 38(5):1150-7.
 31. Henze, E., Schoser, K., Lietzenmayer, R., Schnur, G., Sauer, M., Kunz, R., Claußer, M., Adam, W.E., (1989) Studies of myocardial metabolism using P-31-NMR spectroscopy and a simple heart perfusion model of slaughtered animals- an approach to avoiding in vivo animal experiments. *Z cardiol* 78(9):553-60.
 32. Horiuchi, M., Akishita, M., Dzau, V.J. (1999). Recent progress in angiotensin II type 2 receptor research in the cardiovascular system. *Hypertension* 33(2):613-21.
 33. Horiuchi, M. (2002). Recent progress in AT1 and AT2 receptor research. *Nippon Rinsho* 60(10):1873-80.
 34. Horstick, G., Heimann, A., Gotze, O., Hafner, G., Berg, O., Boehmer, P., Becker, P., Darius, H., Rupprecht, H.J., Loos, M., Bhakdi, S., Meyer, J., Kempski, O. (1997). Intracoronary application of C1 esterase inhibitor improves cardiac function and reduces myocardial necrosis in an experimental model of ischemia and reperfusion. *Circulation* 95:701-8.
 35. Hupf, H., Grimm, D., Riegger, G.A., Schunkert, H. (1999). Evidence for a vasopressin system in the rat heart. *Circ Res* 84(3):365-70.
 36. Imanaka-Yoshida, K., Hiroe, M., Nishikawa, T., Ishiyama, S., Shimojo, T., Otha, Y., Sakakura, T., Yoshida, T. (2001). Tenascin-C modulates adhesion of cardiomyocytes to extracellular matrix during tissue remodeling after myocardial infarction. *Lab Invest* 81:1015-24.
 37. Jöhren, O., Dendorfer, A., Dominiak, P. (2004). Cardiovascular and renal function of angiotensin II type-2 receptors. *Cardiovasc Res* 62(3):460-7.
 38. Jugdutt, B.I. und Amy, R.W.M. (1996). Healing after myocardial infarction in the dog: changes in infarct hydroxyproline and topography. *J Am Coll Cardiol* 7:91-102.
 39. Jugdutt, B.I., Joljart, M.J., Khan, M.I. (1996). Rate of collagen deposition during healing and ventricular remodelling after myocardial infarction in rat and dog models. *Circulation* 94:94-101.
 40. Kaschina, E. and T. Unger (2003). Angiotensin AT1/AT2 receptors: regulation, signalling and function. *Blood Press* 12(2):70-88.

41. Kimose, H.H., Helligso, P., Randsbaek, F., Kim, Y., Botker, H.E., Hansen, S.B., Thomassen, A.R., Nielsen, T.T. (1996). Improved recovery after cold crystalloid cardioplegia using low-dose glutamate enrichment during reperfusion after aortic unclamping: a study in isolated blood-perfused pig hearts. *Thorac Cardiovasc Surg* 44:118-25.
42. Klinke, R and S. Silbernagl (1996). Lehrbuch der Physiologie. 2. Aufl., Stuttgart/New York, Georg Thieme Verlag.
43. Klövekorn, W. P. (1990). Myocardial oxygen supply in hemodilution in patients with healthy hearts and heart surgery patients. *Infusionstherapie* 17(2):24-7.
44. Knickel, U.R., Wilczek, C. and K. Jöst (1996). MemoVet-Praxisleitfaden, Tiermedizin. Stuttgart; New York, Schattauer Verlag.
45. Kossmehl, P., Schönberger, J., Shakibaei, M., Faramarzi, S., Kurth, E., Habighorst, B., von Bauer, R., Wehland, M., Kreutz, R., Infanger, M., Schulze-Tanzil, G., Paul, M., Grimm, D. (2005). Increase of fibronectin and osteopontin in porcine hearts following ischemia and reperfusion. *J Mol Med* 16: 1432-40
46. Leung, P.S. (2004). The peptide hormone angiotensin II: its new functions in tissues and organs. *Curr Protein Pept Sci* 5(4):267-73.
47. Liebich, H.G. (2004). Funktionelle Histologie der Haussäugetiere. 4.Auflage, Schattauer GmbH, Stuttgart.
48. Lindner, K.H. und F.W. Ahnefeld (1989). Comparison of epinephrine and norepinephrine in the treatment of asphyxial or fibrillatory cardiac arrest in porcine model. *Crit Care Med* 17(5):437-41.
49. Lindner, K.H., Ahnefeld, F.W., Grunert, A. (1991). Epinephrine versus norepinephrine in prehospital ventricular fibrillation. *Am J Cardiol* 67(5):427-8.
50. Löscher, W., Ungemach, F.R. und Kroker, R. (1999) Pharmakotherapie bei Haus- und Nutztieren. 4. Aufl., Berlin, Verlag Paul Parey
51. Lazar, H.L., Bao, Y., Rivers, S., Colton, T., Bernard S.A. (2001). High Tissue affinity angiotensin-converting enzyme inhibitors improve endothelial function and reduce infarct size. *Ann Thorac surg* 72: 548-554.
52. Lopaschuk, G.D. (2004). Modulation of energy metabolism as an approach to treating heart failure. *Cardiovasc JS Afr* 15(4 Suppl 1):2.
53. Lucius, R., Gallinat, S., Busche, S., Rosenstiel, P., Unger, T. (1999). Beyond blood pressure: new roles for angiotensin II. *Cell Mol Life Sci* 56(11-12):1008-19.
54. Maisch, B. (1995). Extrazelluläre Matrix und Interstitium des Herzens: „Restriktion“ ist häufiger als gedacht. *Herz* 20: 75-80.
55. Martin, M.M., Su, B., Elton, T.S. (1994). Molecular cloning of the human angiotensin II type 2 receptor cDNA. *Biochem Biophys Res Commun* 205(1):645-51.

56. Mitrovic, V., Mudra, H., Bonzel, T., Schmidt, W., Schlepper, M. (1996). Hemodynamic and humoral effects of parenteral therapy with intravenously administered ACE inhibitor quinaprilat in patients with advanced heart failure. *Z Kardiol* 85(11):828-38.
57. Myers, P.R., Katwa, L.C., Tanner, M., Morrow, C., Guarda, E., Parker, J.L. (1994). Effects of angiotensin II on canine and porcine coronary epicardial and resistance arteries. *J Vasc Res* 31(6):338-46.
58. Modersohn, D., Grosse-Siestrup, C., Liu, C., Konertz, W. (1997). Evaluations of regional and global heart function and heart metabolism in hemoperfused swine hearts. *Biomed Tech (Berl)* 42 Suppl.:393-4.
59. Modersohn, D., S. Eddicks, Grosse-Siestrup, C., Ast, I., Holinski, S., Konertz, W. (2001). Isolated hemoperfused heart model of slaughterhouse pigs. *Int J Artif Organs* 24(4):215-21.
60. Mukoyama, M., Nakajima, M., Horiuchi, M., Sasamura, H., Pratt, R.E., Dzau, V.J. (1993). Expression cloning of type 2 angiotensin II receptor reveals a unique class of seven-transmembrane receptors. *J Biol Chem* 268:24539-24542.
61. Neely, J.R., Liebermeister, H., Battersby, E.J., Morgan, H.E. (1967). Effect of pressure development on oxygen consumption by isolated rat heart. *Am J Physiol* 212(4):804-14.
62. Nickel, R., Schummer, A. and E. Seiferle (1996). Lehrbuch der Anatomie der Haustiere, Band III, Kreislaufsystem, Haut und Hautorgane, Berlin, Paul Parey Verlag.
63. Nogueira, A.C., Wagner, S., Riebeling, I., Klug, S. (1999). The isolated normothermic hemoperfused porcine leg as model for pharmacological and toxicological investigations. *Altex* 16(2):90-94.
64. Nogueira, A.C., Ast, I., Patone, G., Perschel, F.H., Grimm, D., Paul, M. (2003). Functional effects of acute coronary occlusion and catecholnergic stimuli on the isolated normothermic hemoperfused porcine heart. *Clin Exp Hypertens* 25(4):235-55.
65. Plonait, H. and K. Bickhardt. (2004). Lehrbuch der Schweinekrankheiten, 4. Aufl., Berlin, Paul Parey Verlag.
66. Radiometer Referenzhandbuch ABLTM 505
67. Richey, J.M., Ader, M., Moore, D., Bergmann, D.N. (1999). Angiotensin II induces insulin resistance independent of changes in interstitial insulin. *Am J Physiol* 277(5 Pt 1):E920-6.
68. Rigler, R. (1932). Das Vorkammerkreislaufpräparat. *Arch exper Path Pharmakol* 163:295-310.
69. Rhode, E. (1910). Stoffwechseluntersuchungen am überlebenden Warmblüterherzen. I. Mitteilung. Zur Physiologie des Herzstoffwechsels. *Z. Physiol. Chem.* 68:181-235.

70. Rosenstrauch, D., Akay, H.M., Bolukoglu, H., Behrens, L., Bryant, L., Herrera, P., Eya, K., Tuzun, E., Clubb, F.J. Jr., Radovancevic, B., Frazier, H., Kadipasaoglu, A. (2003). Ex vivo resuscitation of adult pig hearts. *Tex Heart Inst J* 30(2):121-7.
71. Rothermund, L., Kreutz, R., Kossmehl, P., Fredersdorf, S., Shakibaei, M., Schulze-Tanzii, G., Paul, M., Grimm, D. (2002). Early onset of chondroitin sulfate and osteopontin expression in angiotensin II-dependent left ventricular hypertrophy. *Am J Hypertens* 15:644-52.
72. Sadoshima, J. (2000). Cytokine actions of angiotensin II. *Circ Res* 86(12):1187-9.
73. Sambraus, H.H. (1994). Atlas der Nutztierrassen. 4. Auflage, Verlag Eugen Ulmer.
74. Schauf, C.L., Moffett D.F. und S.B. Moffett. (1993). *Medizinische Physiologie*. Berlin und Hamburg, Paul Parey Verlag.
75. Scheunert, A. and A. Trautmann. (1987) Lehrbuch der Veterinärphysiologie. 7. Auflage, Verlag Paul Parey.
76. Schmermund, A., Lerman, L.O., Rumberger, J.A., Lund, P.E., Pfeifer, E.A., Sheedy, P.F. 2nd, Ritmann, E.L. (2000). Effects of acute and chronic angiotensin receptor blockade on myocardial vascular blood volume and perfusion in a pig model of coronary microembolization. *Am J Hypertens* 13(7):827-37.
77. Schölmerich, J. (2003). Medizinische Therapie in Klinik und Praxis. Berlin/Heidelberg/New York/Hongkong/London/Mailand/Paris/Tokio; Springer Verlag.
78. Shibutani, M., Uneyama, C., Miyazaki, K., Toyoda, K., Hirose, M. (2000). Methacarn fixation: a novel tool for analysis of gene expressions in paraffin-embedded tissue specimens. *Lab Invest* 80(2):199-208.
79. Spinale, F.G., Gunasinghe, H., Sprunger, P.D., Baskin, J.M., Bradham, W.C. (2002). Extracellular degradative pathway in myocardial remodeling and progression to heart failure. *J Card Fail* 8:332-8.
80. Stolte, M. (1981). Anatomie und Pathologie der Koronararterien. Erlangen, perimed-Verlagsgesellschaft
81. Stowe, D.F., Boban, M., Graf, B.M., Kampine, J.P., Bosnjak, Z.J. (1994). Contraction uncoupling with butanedione monoxime versus low calcium or high potassium solutions on flow and contractile functions of isolated hearts after prolonged hypothermic perfusion. *Circulation* 89(5):2412-20.
82. Stringham, J.C., Paulsen, K.L., Southard, J.H., Fields, B.L., Belzer, F.O. (1992). Improved myocardial ischemic tolerance by contractile inhibition with 2,3 butanedione monoxime. *Ann Thorac Surg* 54:852-860.
83. Strohmenger, H.U., Lindner, K.H., Wienen, W., Radermacher, P., (1998). Effects of an angiotensin II antagonist on organ perfusion during the post-resuscitation phase in pigs. *Crit Care (Lond)* 2(2):49-55.

84. Takayanagi, R., Ohnaka, K., Sakai, Y., Nakao, R., Yanase, T., Haji, M., Inagami, T., Furuta H., Gou, D.F., Nakamura, M. (1992). Molecular cloning, sequence analysis and expression of a cDNA encoding human type-1 angiotensin II receptor. *Biochem Biophys Res Commun* 182(2):910-6.
85. Texter, M., Lees, R.S., Pitt, B., Dinsmore, R.E., Uprichard, A.C. (1993). The QUinapril Ischemic Event Trial (QUIET) design and methods: evaluation of chronic ACE inhibitor therapy after coronary artery intervention. *Cardiovasc Drugs Ther* 7 (2):273-82.
86. Thielscher, H.H. (1985). Oxygen supply and lactate utilization in the porcine heart. *Berl Munch Tierärztl Wochenschr* 98(2):54-7.
87. Thomas, W.G. and F.A. Mendelsohn (2003). Angiotensin receptors: form and function and distribution. *Int J Biochem Cell Biol* 35(6):774-9.
88. Traquandi, C. and Riva, E. (1997). Cardiac effects of Angiotensin I and Angiotensin II: Dose-Response studies in the isolated perfused rat heart. *Pharmacol Res* 37(1):57-63.
89. Tsuzuki, S., Ichiki, T., Nakakub, H., Kitami, Y., Guo, D.F., Shirai, H., Inagami, T. (1994). Molecular cloning and expression of the gene encoding human angiotensin II type 2 receptor. *Biochem Biophys Res Commun* 200(3):1449-54.
90. Uneyama, C., Shibutani, M., (2002). Methacarn fixation for genomic DNA analysis in microdissected, paraffin-embedded tissue specimens. *J of Histochemistry and Cytochemistry* Vol 50:1237-1245.
91. Unger, T. (2002). The role of the renin-angiotensin system in the development of cardiovascular disease. *Am J Cardiol* 89(2A):3A-9A; discussion 10A.
92. Unger, T. (2003). Blood pressure lowering and renin-angiotensin system blockade. *J Hypertens* 21 Suppl 6:S3-7.
93. Urata, H., Kinoshita, A., Misono, K.S., Bumpus, F. M., Husain, A. (1990). Identification of a highly specific chymase as the major angiotensin II-forming enzyme in the human heart. *J Biol Chem* 265:22348-22357
94. van Kats, J.P., de Lannoy, L.M., Danser, A.H., van Meegen, J.R., Verdouw, P.D., Schalekamp, M.A. (1997). Angiotensin II type 1 (AT₁) receptor-mediated accumulation of angiotensin II in tissue and its intracellular half-life in vivo. *Hypertension* 30: 42-49.
95. van Kats, J.P., Dunker D.J., Haitsma, D.B., Schuijt, M.P., Niebuur, R., Stubenitsky, R., Moomsma, F., Schalekamp, M.A., Verdouw, P.D., Danser, A.H. (2000). Angiotensin-converting-enzyme inhibition and angiotensin II typ 1 receptor blockade prevent cardiac remodeling in pigs after myocardial infarction: role of tissue angiotensin II. *Circulation* 102:1556-1563.

96. Varagic, J. and E.D. Frohlich (2002). Local cardiac renin-angiotensin system: hypertension and cardiac failure. *J Mol Cell Cardiol* 34(11):1435-42.
97. Verdouw, P.D., Wolffenbuttel, B.H., van der Giessen, W.J. (1983). Domestic pigs in the study of myocardial ischemia. *Eur Heart J* 4 Suppl C:61-7.
98. Verdouw, P.D., van den Doel, M.A., de Zeeuw, S., Duncker, D.J. (1998). Animal models in the study of myocardial ischaemia and ischaemic syndromes. *Cardiovasc Res* 39(1):121-35.
99. von Baeyer, H., Stahl, K., Hausler, M., Meissler, M., Unger, V., Frank, J., Grosse-Siestrup, C., Kaczmarczyk, G., Affeld, K., Flraig, H.J., Steinbach, B. (1997). A new method of ex vivo whole blood perfusion of isolated mammalian organs, exemplified by the kidney of swine. *Biomed Tech* (Berlin), 43(3):61-8.
100. Wagner, S., Nogueira, A.C., Paul, M., Heydeck, D., Klug, S., Christ, B. (2003). The isolated normothermic hemoperfused porcine forelimb as a test system for transdermal absorption studies. *J Artif Organs* 6(3):183-91.
101. Wiggers, H., Klebe, T., Heickendorf, L., Host, N.B., Danielsen, C.C., Baandrup, U., Andersen H.R. (1997). Ischaemia and reperfusion of the porcine myocardium: effect on collagen. *J Mol Cell Cardiol* 29:289-99.
102. Xia, Q.G., Chung, O., Spitznagel, H., Illner, S., Janichen, G., Rossius, B., Gohlke, P., Unger, T. (2001). Significance of timing of angiotensin AT1 receptor blockade in rats with myocardial infarction-induced heart failure. *Cardiovasc Res* 49(1):110-7.