

## 8. Literaturverzeichnis

Ahlers D., Aurich J., Bleckmann E., Bienek A., Emmert M., Hoedemaker M., Lange J., Mansfeld S., Scheibl P., Zaremba W., Zerbe H. (2000a):

Verlauf des Puerperiums und das weitere Schicksal von Rindern ohne und mit Retentio secundinarium nach Abkalbung mit tierärztlicher Geburtshilfe und intrauteriner Applikation von Ampicillin-Cloxacillin- oder Tetracyclin-haltigen Uterusstäben, Teil 1.

Tierärztl. Umschau 9, 479-488

Ahlers D., Aurich J., Bleckmann E., Bienek A., Emmert M., Hoedemaker M., Lange J., Mansfeld S., Scheibl P., Zaremba W., Zerbe H. (2000b):

Verlauf des Puerperiums und das weitere Schicksal von Rindern ohne und mit Retentio secundinarium nach Abkalbung mit tierärztlicher Geburtshilfe und intrauteriner Applikation von Ampicillin-Cloxacillin- oder Tetracyclin-haltigen Uterusstäben, Teil 2.

Tierärztl. Umschau 10, 565-575

Archbald L.F., Tran T., Thomas P.G.A., Lyle S. (1990):

Apparent failure of Prostaglandin  $F_{2\alpha}$  to improve the reproductive efficiency of postpartum dairy cows that had experienced dystocia and/or retained fetal membranes.

Theriogenology 34, 1025-1034

Arthur G.H. (1979):

Retention of the afterbirth in cattle: A review and commentary.

Vet. Annual 18, 26-36

Arthur G.H., Noakes D.E., Pearson H. (1995):

Veterinary reproduction and obsterics, Bailliere and Tindall, 283.

Aslan S., Gupte Sh., Arbeiter K. (1989):

Das Puerperium von Kühen mit und ohne Retentio secundinarium und nach manipulierter Geburt.

Wien. Tierärztl. Mschr. 76, 386-388

Ball L., Olson J.R., Mortimer R.G. (1984):

Therapeutic considerations for the postpartum bovine uterus.

Soc. for Therio. Newsletter 7, 4-5

Barnouin J., Chassagne M. (1991):

An aetiological hypothesis for the nutrition-induced association between retained placenta and milk fever in the dairy cow.

Ann. Rech. Vet. 22, 331-343

Beetz O. (2002):

Studie zur Wirksamkeit von Ceftiofur bei fieberhaften Puerperalstörungen des Rindes.

Freie Universität Berlin, FB Veterinärmedizin, Diss.

Bekana M., Jonsson P., Ekman T., Kindahl H. (1994):

Intrauterine bacterial findings in postpartum cows with retained fetal membranes.

J. Vet. Med. A. 41, 663-670

Bolinder A., Seguin B., Kindahl H., Bouley D., Otterby D. (1988):

Retained fetal membranes in cows: Manual removal versus nonremoval and its effect on reproductive performance.

Theriogenology 30, 45-56

Borsberry S., Dobson H. (1989):

Periparturient diseases and their effect on reproductive performance in five dairy herds.

Vet. Rec. 124, 217-219

Bostedt H., Sobiraj A. (1985):

Zu Therapie und Nachsorge bei Kühen mit Nachgeburtshaltung.

Tierärztl. Umschau 40, 133-140

Breztlaff K. (1984):

Pharmacology of the uterus.

Proceedings, Xth Int. Cong. Anim. Reprod. and AI; Urbana-Campaign, 9, XI 39-43

Bretzlaff K. (1988):

Physiology and pharmacology of the postpartum cow and retained fetal membranes.

Proceedings, Annu. Conv. Am. Assoc. Bovine Practice. 20, 71

Bretzlaff K.N., Ott R.S., Koritz G.D., Bevill R.F., Gustafsson B.K., Davis L.E. (1983a):

Distribution of oxytetracycline in the healthy and diseased postpartum genital tract of cows.

Am. J. Vet. Res. 44, 760-763

Bretzlaff K.N., Ott R.S., Koritz G.D., Bevill R.F., Gustafsson B.K., Davis L.E. (1983b):

Distribution of oxytetracycline in genital tract tissues of postpartum cows given the drug by intravenous and intrauterine routes.

Am. J. Vet. Res. 44, 764-769

Brooks, G. (2001):

Comparison of two treatments after retained fetal membranes on clinical signs in cattle.

Vet. Rec. 148, 243-244

Chenault J.R., McAllister J.F., Chester S.T., Dame K.J., Kausche F.M., Robb E.J. (2004):

Efficacy of ceftiofur hydrochloride sterile suspension administered parenterally for the treatment of acute postpartum metritis in dairy cows.

J. Am. Vet. Med. Assoc. 224, 1634-1639

Chew, B.P., Keller H.F., Erb R.E., Malven P.V. (1977):

Periparturient concentrations of prolactin, progesterone and the estrogens in blood plasma of cows retaining and not retaining fetal membranes.

J. Anim. Sci. 44, 1055-1060

Clarke C.R., Brown S.A., Streeter R.N., Clarke J.M., Hamlow P.J., Callahan J.K., Hubbard V.L., Speedy A.K., Burrows G.E. (1996):

Penetration of parenterally administered Ceftiofur into sterile vs. *Pasteurella haemolytica*-infected tissue chambers in cattle.

J. Vet. Pharmacol. Therap. 19, 376-381

Cohen R.O., Colodner R., Ziv G., Keness J. (1996):

Isolation and antimicrobial susceptibility of obligate anaerobic bacteria recovered from the uteri of dairy cows with retained fetal membranes and postparturient endometritis.

J. Vet. Med. B 43, 193-199

Correa M.T., Erb H., Scarlett J. (1993):

Path analysis for seven postpartum disorders of Holstein cows.

J. Dairy Sci. 75, 1305-1312

Davies C.J., Hill J.R., Edwards J.L., Schrick F.N., Fisher P.J., Eldridge J.A., Schlafer D.H. (2004):

Major histocompatibility antigen expression on the bovine placenta: Its relationship to abnormal and retained placenta.

Anim. Reprod. Sci. 82-83, 267-280

De Bois C.H.W. (1982):

Some aspects of the therapy and prophylaxis of retained placenta and puerperal endometritis in the cow.

In.: Karg H. und Schallenberger E. (Hrsg.): Factors affecting fertility in the post partum cow.

Martinus Nijhoff Publishing, The Hague, 479-509

Dinsmore R.P., Stevens R.D., Cattell M.B., Salman M.D., Sundlof S.F. (1996):

Oxytetracycline residues in milk after intrauterine treatment of cows with retained fetal membranes.

JAVMA 209, 1753-1755

Dobson D.P., Noakes D.E. (1990):

Use of a uterine pessary to prevent infection of the uterus of the cow after parturition.

Vet. Rec. 127, 128-131

Dohmen M.J.W., Joop K., Sturk A., Bols P.E.J., Lohuis J.A.C.M. (2000):

Relationship between intra-uterine bacterial contamination, endotoxin levels and the development of endometritis in postpartum cows with dystocia or retained fetal membranes.

Theriogenology 54, 1019-1032

Dosogne H., Burvenich C., Lohuis J.A. (1999):

Acyloxyacyl hydrolase activity of neutrophil leukocytes in normal early postpartum dairy cows and in cows with retained placenta.

Theriogenology 51, 867-874

Drillich M. (2000):

Vergleich des strategischen Einsatzes von Prostaglandin F<sub>2</sub>alpha mit konventionellen Methoden des Fruchtbarkeitsmanagements in zwei Milchviehbetrieben.

Freie Universität Berlin, FB Veterinarmedizin, Diss.

Drillich M., Beetz O., Pfützner A., Sabin M., Sabin H.-J., Kutzer P., Nattermann H., Heuwieser W. (2001):

Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows

J. Dairy Sci. 84, 2010-2017

Drillich M., Pfützner A., Sabin H.-J., Sabin M., Heuwieser W. (2003):

Comparison of two protocols for the treatment of retained fetal membranes in dairy cattle.

Theriogenology 59, 951-960

Du Bois P.R., Williams W.F. (1980):

Increased incidence of retained placenta associated with heat stress in dairy cows.

Theriogenology 13, 115-121

Dyrendahl I., Mattson J., Pehrson B. (1977):

Retained placenta in cattle – incidence, clinical data and effects on fertility.

Zbl. Vet. Med. A. 24, 529-541

Echternkamp S.E., Gregory K.E. (1999):

Effects of twinning on gestation length, retained placenta and dystocia.

J. Anim. Sci. 77, 39-47

Eiler H. (1997):

Retained placenta.

In: Youngquist R.S. (Hrsg.): Current Therapy in Large Animal Theriogenology.

W.B. Saunders Company, Philadelphia, 2. Auflage, 340-348

Eiler H., Hopkins F.M. (1992):

Bovine retained placenta: Effects of collagenase and hyaluronidase on detachment of placenta.

Biology of Reprod. 46, 580-585

Eiler H., Hopkins F.M. (1993):

Successful treatment of retained placenta with umbilical cord injections of collagenase.

JAVMA 203, 436-443

Eiler H., Hopkins F.M., Armstrong-Backus C.S., Lyke W.A. (1984):

Uterotonic effect of Prostaglandine  $F_{2\alpha}$  and oxytocin on the postpartum cow.

Am. J. Vet. Res. 45, 1011-1014

Erb H.N., Guard C.L., Hillman R.B., Oltenacu P.A., Powers P.A., Smith M.C., Smith R.D., White M.E. (1985):

Path model of reproductive disorders and performance, milk fever, mastitis, milk yield and culling in Holstein Cows.

J. Dairy Sci. 68, 3337-3349

Erb H.N., Martin S.W., Ison N., Swaminathan S. (1981):

Interrelationships between production and reproductive diseases in Holstein cows.

Conditional relationships between production and disease.

J. Dairy Sci. 64, 272-281

Esslemont R.J., Peeler E.J. (1993):

The scope for raising margins in dairy herds by improving fertility and health.

Br. Vet. J. 149, 537-547

- Fleischer P., Metzner M., Beyerbach M., Hoedemaker M., Klee W. (2001):  
The relationship between milk yield and the incidence of some diseases in dairy cows.  
J. Dairy Sci. 84, 2025-2035
- Fürstenberg A., Busch W., Fürstenberg L., Münchow H. (1990):  
Untersuchung zur Ätiologie der Retentio secundinarium beim Rind.  
Mh. Vet.-Med. 45, 493-496
- Gaverick H.A. (1997):  
Ovarian follicular cysts in dairy cows.  
J. Dairy Sci. 80, 995-1004
- Geiser S., Siegenthaler H.P., Martig J. (1995):  
Untersuchungen zur Wirksamkeit der lokalen Behandlung mit einem PVP-Jodpräperat bei der  
Nachgeburtshaltung der Kuh.  
Berl. Münch. Tierärztl. Wschr. 108, 264-268
- Gilbert R.O., Gröhn Y.T., Guard C.L., Surman V. (1993a):  
Partum neutrophil function in cows which retain fetal membranes.  
Res. Vet. Sci. 55, 15-19
- Gilbert R.O., Gröhn Y.T., Miller P.M., Hoffman D.J. (1993b):  
Effect of parity on periparturient neutrophil function in dairy cows.  
Vet. Immunology and Immunopathology 36, 75-82
- Golub L.M., Goodson J.M., Lee H.M., Vidal A.M., McNamara T.F., Ramamurthy N.S.  
(1985):  
Tetracyclines inhibit tissue collagenase; effects of ingested low-dose and local delivery  
systems.  
J. Periodontol. 56, 93-97

Gröhn Y.T., Erb H.N., McCulloch C.E., Saloniemi H.S. (1990):

Epidemiology of reproductive disorders in dairy cattle: Associations among host characteristics, disease and production.

Prev. Vet. Med. 8, 25-39

Grunert E. (1983):

Ätiologie, Pathogenese und Therapie der Nachgeburtsverhaltung beim Rind

Wien. Tierärztl. Mschr. 70, 230-235

Grunert E. (1984):

Placental separation/retention in the bovine.

Proceedings, Xth Int. Cong. Anim. Reprod. and AI; Urbana-Campaign, 9, XI 25-32

Grunert E. (1990):

Weiblicher Geschlechtsapparat.

in: Dirksen G., Gründer H.-D. und Stöber M. (Hrsg.): Die klinische Untersuchung des Rindes.

Verlag Paul Parey, Berlin und Hamburg, 3. Auflage, 472-548

Grunert E. und Andresen P. (1996):

Geburtshilfe.

In: Grunert E. (Hrsg.): Buiatrik Band 1.

Verlag M.&H. Schaper Alfred, Hannover, 5. Auflage. 129-190.

Grunert E., Zaremba W. (1979):

Untersuchungen über negative Einflüsse von endogenen und exogenen Faktoren auf das Frühpuerperium des Rindes.

Dtsch. Tierärztl. Wschr. 86, 461-464

Gunnink J.W. (1984):

Retained placenta and leucocytic activity.

The Vet. Quarterly 6, 49-51



Gustafsson B. (1980):

Disposition of antimicrobial drugs in the female genital system.

Proceedings, XIth Int. Congr. Anim. Reprod. 2, 485-490

Gustafsson B.K. (1984):

Therapeutic strategies involving antimicrobial treatment of the uterus in large animals.

JAVMA 185, 1194-1198

Gustafson B.K., Ott R.S. (1981):

Current trends in the treatment of genital infections in large animals.

Comp. Cont. Ed. Pract. Vet. 3, 147-151

Haaland M.A., Manspeaker J.F., Moreland T.W. (1984):

Antibiotic residues in milk after intrauterine infusion.

Vet. Med. Sm. Anim. Clin. 79, 383-386

Halpern N.E., Erb H.N., Smith R.D. (1985):

Duration of retained fetal membranes and subsequent fertility in dairy cows.

Theriogenology 23, 807-813

Heinonen M., Heinonen K. (1989):

Retained placenta in cattle: The effect of treatment or nontreatment on puerperal diseases and subsequent fertility.

Acta Vet. Scand. 30, 425-429

Helmuth R. (1999):

Einsatz antimikrobiell wirksamer Substanzen in der Veterinärmedizin.

Bundesgesundheitsblatt 42, 26-34

Herschler R.C., Lawrence J.R. (1984):

A prostaglandin analogue for therapy of retained placenta.

Vet Medicine, 822-826

Heuwieser W., Grunert E. (1987):

Significance of chemotactic activity for placental expulsion in cattle.

Theriogenology 27, 907-912

Heuwieser W., Mansfeld R. (1992):

Beurteilung der Körperkondition bei Milchkühen. Teil 2.

Wie Sie ihre Herde durch die Beurteilung der Körperkondition besser in den Griff bekommen.

Milchpraxis 30, 10-14

Hindson J.C. (1976):

Retention of the fetal membranes in cattle.

Vet. Rec. 99, 49-50

Hoffmann B. (1993):

Endokrinologie der Hochträchtigkeit, während des Partus und im Puerperium.

In: Grunert E. und Arbeiter K. (Hrsg.): Richter J. und Götze R.: Tiergeburtshilfe.

Verlag Paul Parey, Berlin und Hamburg, 4. Auflage.

Holt L.C., Whittier W.D., Gwazdauskas F.C., Vinson W.E. (1989):

Early postpartum reproductive profiles in Holstein cows with retained placenta and uterine discharge.

J. Dairy Sci. 72, 533-539

Jeffrey J.J. (1991):

Collagen and Collagenase: Pregnancy and partuition.

Semin. Perinatol. 15, 118-126

Jeffrey J.J., Ehlich L.S., Roswit W.T. (1991):

Serotonin: An inducer of collagenase in myometrial smooth muscle cells.

J. Cell. Physiol. 146, 399-406

Joosten I., Stelwagen J., Dijkhuizen A.A. (1988):

Economic and reproductive consequences of retained placenta in dairy cattle.

Vet. Rec. 123, 53-57

Kaneko K., Kawakami S., Miyoshi M., Abukawa T., Yamanaka S., Mochizuki M., Yoshihara S. (1997):

Effect of retained placenta on subsequent bacteriological and cytological intrauterine environment and reproduction in Holstein dairy cows.

Theriogenology 48, 617-624

Kelton D.F., Lissemore K.D., Martin R.E. (1998):

Recommendations for recording and calculating the incidence of selected clinical diseases of dairy cattle.

J. Dairy Sci. 81, 2502-2509

Kimura K., Goff J.P., Kehrli M.E., Reinhardt T.A. (2002):

Decreased neutrophil function as a cause of retained placenta in dairy cattle.

J. Dairy Sci. 85, 544-550

Kossaibati M.A., Esslemont R.J. (1995):

Daisy Report number 4. Wastage in dairy herds.

Reading University, Reading.

Kudlac E. (1991):

Einige Fragen der Therapie und Prophylaxe der Nachgeburtshaltung bei Kühen und ihre weitere Fruchtbarkeit.

Tierärztl. Umschau 46, 407-411

Kurth A. (2001):

Feldstudie zur Überprüfung der klinischen Wirksamkeit von Prostaglandin  $F_{2\alpha}$  als Ergänzung zu einer konventionellen Therapie der Retentio secundinarium bei Milchkühen.

Freie Universität Berlin, FB Veterinärmedizin, Diss.

Königsson K., Gustafsson H., Gunnarsson A., Kindahl H. (2001):

Clinical and bacteriological aspects on the use of tetracycline and flunixin in primiparous cows with induced retained placenta and postpartal endometritis.

Reprod. Domest. Anim. 36, 247-256

Kroker R., Löscher W., Simunek J., Trolldenier H., Ungemach F.R. (1996):

Chemotherapie bakterieller Infektionen.

In: Frey H.-H., Löscher W.: Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin.

Ferdinand Enke Verlag, Stuttgart, 1. Auflage, 454-508

Lawrence K. (2001):

Developing responsible use of antibiotics protocols.

Proceedings, 34<sup>th</sup> Ann. AABP Conv., 10-13

Laven R.A., Peters A.R. (1996):

Bovine retained placenta: aetiology, pathogenesis and economic loss.

Vet. Rec. 139, 465-471

LeBlanc S.J., Duffield T.F., Leslie K.E., Bateman K.G., Keefe G.P., Walton J.S., Johnson W.H. (2002):

The effect of treatment of clinical endometritis on reproductive performance in dairy cows.

J. Dairy Sci. 85, 2237-2249

Leidl W., Hegner P., Rockel P. (1980a):

Investigations on the PGF<sub>2α</sub> concentration in maternal and foetal cotyledons of cows with and without retained foetal membranes.

Zbl. Vet. Med. A. 27, 691-696

Leidl W., Hegner P., Uhlig A., Schels H. (1980b):

Untersuchung zur Ätiologie der Retentio secundinarum beim Rind.

Proceedings, XIth. Int. Cong. on disease of Cattle, Tel Aviv, 973-978

Loeffler S.H., de Vries M.J., Schukken Y.H. (1999):

The effects of disease occurrence, milk yield and body condition on fertility of dairy cows.

J. Dairy Sci. 82, 2589-2604

Lüpschen C. (1997):

Aussagekraft des Body-Condition-Score Test im Hinblick auf das Partus-Syndrom.

Bonn, Hohe Landwirtschaftl. Fakultät der Rheinischen Friedrich-Wilhelms-Universität, Diss.

Luginbühl A., Küpfer U. (1980a):

Bakteriologische Befunde im Geschlechtsapparat von Kühen im Puerperium.

Mitteilung III.

Schweiz. Arch. Tierheilk. 123, 13-18

Luginbühl A., Küpfer U. (1980b):

Bakteriologische Befunde im Geschlechtsapparat von Kühen im Puerperium.

Mitteilung IV.

Schweiz. Arch. Tierheilk. 123, 629-637

Markusfeld O., Galon N., Ezra E. (1997):

Body condition score, health, yield and fertility in dairy cows.

Vet. Rec. 141, 67-72

Martin J.M., Wilcox C.J., Moya J., Klebanow E.W. (1986):

Effects of retained fetal membranes on milk yield and reproductive performance.

J. Dairy Sci. 69, 1166-1168

Metzner M., Heuwieser W., Klee W. (1993):

Die Beurteilung der Körperkondition (body condition scoring) im Herdenmanagement.

Prakt. Tierarzt 74, 991-998

Morrione T.G., Zairu M. (1964):

Injury influence on resorption of uterine collagen.

Arch. Pathol. 78, 591-600

Muller L.D., Owens M.J. (1974):

Factors associated with the incidence of retained placentas.

J. Dairy Sci. 57, 725-728

Neu H.C. (1982):

The new beta-lactamase-stable Cephalosporins.

Ann. Intern. Med. 97, 408-419

Noakes D.E. (1991):

Non-specific uterine infections in cattle and their treatment.

Proceedings of the British Cattle Veterinary Association, University of Reading, Reading, 47-62

Oettel M. (1996):

Endokrinopharmakologie.

In: Frey H.-H., Löscher W.: Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin.

Ferdinand Enke Verlag, Stuttgart, 1. Auflage, 370-423

Okano, A., Tomizuka, T. (1987):

Ultrasonic observation of postpartum uterine involution in the cow.

Theriogenology 27, 369-376.

Okker H., Schmitt E.J., Vos P.L.A.M., Scherpenisse P., Bergenwerff A.A., Jonker F.H.

(2002):

Pharmacokinetics of ceftiofur in plasma and uterine secretions and tissues after subcutaneous postpartum administration in lactating dairy cows.

J. vet. Pharmacol. Therap. 25, 33-38

Peter A.T., Bosu W.T.K. (1988):

Relationship of uterine infections and folliculogenesis in dairy cows during early puerperium.

Theriogenology 30, 1045-1051

Peters A.R., Laven R.A. (1996):

Treatment of bovine retained placenta and its effects.

Vet. Rec. 139, 535-539

Paisley L.G., Mickelsen W.D., Anderson P.B. (1986):

Mechanisms and therapy for retained fetal membranes and uterine infections of cows: A review.

Theriogenology 25, 353-381

Piper E.L., Combs K.J., Peterson H.P. (1978):

Influence of corticoid induced parturition on postpartum uterine activity.

Vet. Med. Sm. Anim. Clin. 73, 1447-1449

Rajala P.J., Gröhn Y.T. (1998):

Effects of dystocia, retained placenta and metritis on milk yield in dairy cows.

J. Dairy Sci. 81, 3172-3181

Rasmussen F.E., Wiltbank M.C., Christensen J.O., Grummer R.R. (1996):

Effects of fenprostalene and estradiol-17 benzoate on parturition and retained placenta in dairy cows and heifers.

J. Dairy Sci. 79, 227-234

Risco C.A., Archbald L.F., Elliott J.T., Chavatte P. (1994):

Effect of hormonal treatment on fertility in dairy cows with dystocia or retained fetal membranes at parturition.

J. Dairy Sci. 77, 2562-2569

Risco C.A., Hernandez J. (2003):

Comparison of ceftiofur hydrochloride and estradiol cypionate for metritis prevention and reproductive performance in dairy cows affected with retained fetal membranes.

Theriogenology 60, 47-58

Romaniuk J. (1985):

Nachgeburtverhalten bei den Milchkühen – Vorkommen und Einfluss auf die Fruchtbarkeit.

Tierärztl. Umschau 40, 130-133

Rüsse I. (1991):

Frühgravidität, Implantation und Plazentation.

In: Rüsse I., Sinowatz F. (Hrsg.): Lehrbuch der Embryologie der Haustiere.

Verlag Paul Parey, Berlin und Hamburg, 1. Auflage. 153-218

Russe M. (1982):

Myometrial activity postpartum.

In: Karg H., Schallenberger E. (Hrsg.): Factors influencing fertility in the postpartum cow.

Martinus Nijhoff Publishing, The Hague, 55-60

Sandals W.C.D., Curtis R.A., Cote J.F., Martin S.W. (1979):

The effect of retained placenta and metritis complex on reproductive performance in dairy cattle – a case control study.

Can. Vet. J. 20, 131-135

Sande M.A., Mandell G.I. (1980):

Antimicrobial agents – tetracyclines and chloramphenicol.

In: Gilman A.G., Goodman L.S., Gilman A. (Hrsg.): The pharmacological basis of therapeutics.

Macmillan Pub. Co., New York, 1080-1105

Sarges J., Heuwieser W., Schlüns J., Drewes B. (1998):

Immunohistological examination on the distribution of collagen types I, III, IV und V in bovine post partum placentomes.

J Vet. Med. A. 45, 1-10

Schnorr B. (1996):

Plazentation beim Säuger und Embryonalhüllen beim Vogel.

In Schnorr B. (Hrsg.): Embryologie der Haustiere.

Ferdinand Enke Verlag, Stuttgart, 1. Auflage, 66-99



Schulz L.-Cl., Merkt H. (1956):

Morphologische Befunde an exstirpierten Plazentomen, zugleich ein Beitrag zur Ätiologie der Retentio secundinarium beim Rind.

Mh. Vet. Med. 11, 712-714

Schukken Y.H., Erb H.E., Scarlett J.M. (1989):

A hospital based study of the relationship between retained placenta and mastitis in dairy cows.

Cornell Vet. 79, 319-326

Schukken Y.H., Erb H.N., Smith R.D (1988):

The relationship between mastitis and retained placenta in a commercial population of Holstein dairy cows.

Prev. Vet. Med. 5 188-190

Senger P.L. (2002):

Placentation, the endocrinology of gestation and parturition.

In: Pathways to Pregnancy and Parturition.

Pullman, Washington, 2. Auflage.

Seguin B.E., Morrow D.A., Lewis T.M. (1974):

Luteolysis, luteostasis and the effects of Prostaglandin  $F_{2\alpha}$  on cows after endometrial irritation.

Am. J. Vet. Res. 35, 57-61

Sharpe K.L., Eiler H., Hopkins F.M. (1990):

Changes in the proportion of type I and type III collagen in the developing and retained bovine placentome.

Biol. of Reprod. 43, 229-235

Sheldon I.M., Rycroft A.N., Zhou C. (2004):

Association between postpartum pyrexia and uterine bacterial infection in cattle.

Vet. Rec. 64, 289-293

Smith B.I., Donovan G.A., Risco C., Littell R., Young C., Stanker L.H., Elliott J. (1998):  
Comparison of various antibiotic treatments for cows diagnosed with toxic puerperal metritis.  
J. Dairy Sci. 81, 1555-1562

Sobiraj A., Hermülheim A., Herfen K., Schulz S. (1998):  
Einfluss verschiedener Uterotonika auf den Nachgeburtsabgang bei Rindern nach  
konservativen und operativen geburtshilflichen Eingriffen.  
Tierärztl. Umschau 53, 392-399

Stevens R.D., Dinsmore R.P., Cattell M.B. (1995):  
Evaluation of the use of intrauterine infusions of oxytetracycline, subcutaneous injections of  
fenprostalene, or a combination of both, for the treatment of retained fetal membranes in dairy  
cows.  
JAVMA 207, 1612-1615

Strahl H. (1906):  
Die Embryonalhüllen der Säuger und die Placenta.  
In: Hertwig O. (Hrsg.): Handbuch der vergleichenden und experimentellen  
Entwicklungslehre.

Tischer M. (1998):  
Vergleich von intrauterinen Arzneimittelapplikationen mit einem strategischen  
Prostaglandinprogramm zur Behandlung von chronischen Endometritiden in einer  
Milchviehherde.  
Freie Universität Berlin, FB Veterinärmedizin, Diss.

Ungemach F.R. (1999):  
Einsatz von Antibiotika in der Veterinärmedizin: Konsequenzen und rationaler Umgang.  
Tierärztl. Praxis 27(G), 335-340

Vandenplassche M. (1984):  
Stimulation and inhibition of phagozytosis in domestic animals.  
Proceedings, Xth. Int. Cong. Anim. Reprod. And AI, Urbana-Campaign, III: 475

Vandenplassche M., Bouters R. (1982):

The impact of gynaecological and obstetrical problems resulting out of pregnancy and parturition.

In: Karg H., Schallenberger E. (Hrsg.): Factors influencing fertility in the postpartum cow, Martinus Nijhoff Publishing, The Hague, 30-44

Van Werven T., Schukken Y.H., Lloyd J., Brand A., Heeringa H.Tj., Shea M. (1992):

The effects of duration of retained placenta on reproduction, milk production, postpartum diseases and culling rate.

Theriogenology 37, 1191-1203

Venable J.H., McDonald L.E. (1958):

Postparturient bovine uterine motility: Normal and after experimentally produced retention of the fetal membranes.

Am. J. Vet. Res. 19, 308-313

Werner E. (1996):

Desinfektionsmittel.

In: Frey H.-H., Löscher W.: Lehrbuch der Pharmakologie und Toxikologie für die Veterinärmedizin.

Ferdinand Enke Verlag, Stuttgart, 1. Auflage, 575-595

WHO. Recommendations (1997):

In: WHO-Proceedings: The medical impact of the use of antimicrobials in food animals.

WHO/EMC/ZOO/97.4, 11-16

Wischral A., Verreschi I.T.N., Lima S.B., Hayashi L.F., Barnabe R.C. (2001):

Pre-partum profile of steroids and prostaglandin in cows with or without foetal membrane retention.

Anim. Reprod. Sci. 67, 181-188

Zdunczyk S., Ahlers D., Grunert E. (1992):

Zur Frage der Beziehungen zwischen bovinen klinischen Mastitiden zum Zeitpunkt der Abkalbung und Retentio secundinarum.

Dtsch. Tierärztl. Wschr. 99, 353-392

Zhou C., Boucher J.F., Dame K.J. (2001a):

A multi-location clinical study of ceftiofur for the treatment of postpartum cows with elevated temperature and other complications

Proceedings, 34<sup>th</sup> Ann AABP Conv, 211-212

Zhou C., Boucher J.F., Dame K.J. (2001b):

Multilocation trial of ceftiofur for treatment of postpartum cows with fever.

JAVMA 219, 805-808

Ziv G. (1980):

Review of pharmacology of antimicrobial drugs employed in veterinary obstetrics.

Proceedings, IXth. Int Cong. Anim. Reprod. and AI, Madrid, 466-471

Ziv G. (1984):

Minimal inhibitory concentrations of antibiotics for pathogenic bacteria as determined aerobically and anaerobically in uterine secretions.

Proceedings, Xth. Int. Cong. Anim. Reprod. and AI, Urbana-Campaign, I: 420