

7. Literaturverzeichnis

Abramowski, D., Rigo, M., Duc, D., Hoyer, D., Staufenbiel, M.:
 Localization of the 5-hydroxytryptamine_{2C} receptor in human and rat brain using specific antisera.
Neuropharmacology 34: 1635-1645, 1995

Adham, N., Kao, H.-T., Schechter, L. E., Bard, J., Olsen, M., Urquhart, D., Durkin, M., Hartig, P. R., Winshank, R. L., Branchek, T. L.:
 Cloning of another human serotonin receptor (5-HT_{1F}): a fifth 5-HT₁ receptor sub-type coupled to the inhibition of adenylyl cyclase.
Proc. Natn. Acad. Sci. U. S. A. 89: 408-412, 1993

Amlaiky, N., Ramboz, S., Boschert, U., Plassat, J. L., Hen, R.:
 Isolation of a mouse 5-HT_{1E}-like serotonin receptor expressed predominantly in hippocampus.
J. Biol. Chem. 267: 19761-19764, 1992

Andrews, N., Hogg, S., Gonzalez, L. E., File, S. E.:
 5-HT_{1A} receptors in the median raphe nucleus and dorsal hippocampus may mediate anxiolytic and anxiogenic behaviours respectively.
Europ. J. Pharmacol. 264: 259-264, 1994

Azmitia, E. C., Buchan, A. M., Williams, J.H.:
 Structural and functional restoration by collateral sprouting of hippocampal 5-HT axons.
Nature 274: 374-376, 1978

Azmitia, E. C., Segal, M.:
 An autoradiographic analysis of the different ascending projections of the dorsal and median raphe nuclei in the rat.
J. Comp. Neurol. 179: 641-668, 1978

Baker, K. G., Halliday, G. M.:
 Ascending noradrenergic and serotonergic systems in the human brainstem.
In: Neurotransmitters in the human brain. D. J. Tracey, G. Paxinos, J. Stone (Hrsg.),
Adv. Behav. Biol. 43: 155-171, 1995

Baraban, J. M., Aghajanian, G. K.:
 Noradrenergic innervation of serotonergic neurons in the dorsal raphe: Demonstration by electron microscopic autoradiography.
Brain Res. 204: 1, 1981

Barnes, J. M., Barnes, N. M., Cooper, S. J.:
 Behavioural pharmacology of 5-HT₃ receptor ligands.
Neurosci. Biobehav. Rev. 16: 107-113, 1992

Baumgarten, H. G., Björklund, A., Lachenmayer, L., Nobin, A., Stenevi, U.:
 Long-lasting depletion of brain serotonin by 5,6-dihydroxytryptamine.
Acta physiol. Scand. Suppl. 373: 1-15, 1971

Baumgarten, H. G., Lachenmayer, L.:
 5,7-Dihydroxytryptamine: Improvement in chemical lesioning of indoleamine neurons in the mammalian brain.
Z. Zellforsch. 135: 399-414, 1972

- Baumgarten, H. G., Björklund, A.:
 Neurotoxic indolamines and monoamine neurons.
Annu. Rev. Pharmacol. 16: 101-111, 1976
- Beinfeld, M. C., Meyer, D. K., Eskay, R. L., Jensen, J. T., Brownstein, M. J.:
 The distribution of cholecystokinin-immunoreactivity in the central nervous system of the rat, as determined by radioimmunoassay.
Brain Res. 212: 51-57, 1981
- Bito, L., Davson, H., Levin, E., Murray, M., Snider, N.:
 The concentrations of free amino acids and other electrolytes in cerebrospinal fluid, in dialysate of brain, and blood plasma of the dog.
J. Neurochem. 13: 1057-1067, 1966
- Björklund, A., Nobin, A., Stenevi, U.:
 The use of neurotoxic dihydroxytryptamines as tools for morphological studies and localized lesioning of central indolamine neurons.
Z. Zellforsch. 145: 479-501, 1973
- Blakely, R. D., Berson, H. E., Fremeau, R. T., Caron, M. C., Peek, M. M., Prince, H. K., Bradley, C. C.:
 Cloning and expression of functional transporter from rat brain.
Nature 354: 66-70, 1991
- Boissier, J. R., Simon, P Lwoff, J. M.:
 L'utilisation d'une reaction particulière de la souris (methode de la planche a trous) pour l'étude des medicamentes psychotropes.
Therapie 19: 571-589, 1964
- Boosalis, M. G., Gemayel, N., Lee, A., Bray, G. A., Laine, L., Cohen, H.:
 Cholecystokinin and satiety: Effect of hypothalamic obesity and gastric bubble insertion.
Am. J. Physiol. 262: R241-R244, 1992
- Brewerton, T. D., Brandt, H. A., Lesem, D. T., Murphy, D. L., Jimerson, D. C.:
 Serotonin in eating disorders.
In: Serotonin in major psychiatric disorders, E. Coccaro, D. Murphy (Hrsg), pp. 153-184.
 American Psychiatric Press, Washington, 1990
- Brewerton, T. D., Mueller, E. A., Lesem, M. D., Brandt, H. A., Quearry, B., George, D. T., Murphy, D. L., Jimerson, D. C.:
 Neuroendocrine responses to m-chlorophenylpiperazine and L-tryptophan in bulimia.
Arch. Gen. Psychiatry 49: 852-861, 1992
- Briley, M.:
 Imipramine binding: its relationship with serotonin uptake and depression.
In: Neuropharmacology of serotonin, A. R. Green (Hrsg.), pp. 50-78.
 Oxford University Press, Oxford, 1985
- Briley, M., Chopin, P., Moret, C.:
 Effect of serotonergic lesion on „anxious“ behaviour measured in the elevated plus-maze test in the rat.
Psychopharmacology 101: 187-189, 1990

- Brodal, A., Taber, E., Walberg, F.:
 The raphe nuclei of the brain stem in the cat. II. Efferent connections.
J. Comp. Neurol. 114: 239-260, 1960
- Bruinvels, A.T., Palacios, J. M., Hoyer, D.:
 Autoradiographic characterization and localisation of 5-HT_{1D} compared to 5-HT_{1B} binding sites in rat brain.
Naunyn Schmiedebergs Arch. Pharmacol. 347: 569-582, 1993
- Brunner, E., Langer, F.:
 Nichtparametrische Analyse longitudinaler Daten.
 Oldenbourg Verlag, 1999
- Burhol, P. G., Jenssen, T. G., Jorde, R., Lygren, I., Johnson, J. A.:
 Plasma cholecystokinin (CCK) before and after a jeuno-ileal bypass operation in obese patients with reference to appetite regulation.
Int. J. Obes. 8: 233-236, 1984
- Cadogan, A. K., Kendall, D. A., Fink, H., Marsden, C. A.:
 Serotonin and behaviour: *In vivo* assessment of pre- and post-synaptic mechanisms.
In: Monitoring Molecules in Neuroscience, 6th International conference on in vivo methods, A. Louilot (Hrsg.), pp. 217-218.
 Talence, Univ. Bordeaux I., 1994
- Cajal, S., Ramon, Y.:
 Histologie du systeme nerveux de l'homme et des vertebres.
Maloine, Paris, 1911
- Carli, M., Samanin, R.:
 Potential anxiolytic properties of 8-hydroxy-2-(Di-N-propylamino)tetralin, a selective serotonin1A receptor agonist.
Psychopharmacology. 94: 94-91, 1988
- Carli, M., Prontera, C., Samanin, R.:
 Evidence that central 5-hydroxytyraminergic neurones are involved in the anxiolytic activity of buspirone.
Br. J. Pharmacol. 96: 829-836, 1989
- Carson, M. J., Thomas, E.A., Danielson, P.E., Sutcliffe, J. G.:
 The 5-HT5A serotoninreceptor is expressed predominantly in astrocytes in which it inhibits cAMP accumulation: a mechanism of reactive astrocytes.
Glia 17: 317-326, 1996
- Casper, R. C.:
 On the emergence of bulimia nervosa as a syndrome: a historical view.
Int. J. Eat. Dis. 2: 3-16, 1983
- Castelao, J. R., Ferreira, L., Gelders, Y. J., Heylen, S. L. E.:
 The efficacy of the D-2 and 5-HT₂ antagonist risperidone (R 64766) in psychotic patients.
Schizophrenia Res. 2: 411-415, 1989
- Choi, D. S., Maroteaux, L.:
 Immunohistochemical localisation of the serotonin 5-HT_{2B} in mouse gut, cardiovascular system and brain.
FEBS Lett. 391: 45-61, 1996

Chopin, P., Pellow, S., File, S. E.:

The effects of yohimbine on exploratory and locomotor behaviour are attributable to its effects at noradrenaline and not benzodiazepine receptors.

Neuropharmacology 25: 53-57, 1986

Chopin, P., Briley, M.:

Animal models of anxiety.

TIPS 8: 383-388, 1987

Commissaris, R. L., Lyness, W. H., Rech, R. H.:

The effects of *d*-lysergic acid diethylamide (LSD), 2,5-dimethoxy-4-methylamphetamine (DOM), pentobarbital and methaqualone on punished responding in control and 5,7-dihydroxytryptamine-treated rats.

Pharmacol. Biochem. Behav. 14: 617-623, 1981.

Cooper, S. J., Dourish, C. T.:

Multiple cholecystokinin (CCK) receptors and CCK-monoamine interactions are instrumental in the control of feeding.

Physiol. Behav. 48: 849-857, 1990

Corrodi, H., Fuxe, K., Lindbrink, P., Olson, L.:

Minor tranquilizers stress and central catecholamine neurones.

Brain Res. 29: 1-6, 1971

Costall, B., Naylor, R. J.:

Anxiolytic effects of 5-HT₃ antagonists in animals.

In: 5-HT1A agonists, 5-HT3 antagonists and benzodiazepines, their comparative behavioural pharmacology, R. J. Rodgers, S. J. Cooper (Hrsg.), pp. 133-157.

Wiley, Chichester, 1991

Costall, B., Naylor, R. J.:

The pharmacology of the 5-HT₄ receptor.

Int. Clin. Psychopharmacol. 8 [Suppl. 2]: 11-18, 1993

Costall, B., Naylor, R. J.:

5-HT₃ receptor antagonists in the treatment of cognitive disorders.

In: 5-hydroxytryptamine₃ receptor antagonists, F. D. King, B. J. Jones, G. J. Sanger (Hrsg.), pp. 203-219.

CRC Inc., Boca Raton, 1994

Cotton, R.G.H., McAdam, W., Jennings I. G., Morgan, F. J.:

A monoclonal antibody to aromatic amino acid hydroxylases. Identification of the epitope.

Biochem. J. 255: 193-196, 1988

Crawley, J. N., Hays, S. E., Paul, S. M.:

Vagotomy abolishes the inhibitory effects of cholecystokinin on rat exploratory behavior.

Eur. J. Pharmacol. 73: 379-380, 1981

Crawley, J. N.:

Comparative distribution of cholecystokinin and other neuropeptides.

Ann. NY Acad. Sci. 448: 1-8, 1985

Critchley, M. A. E., Njung'e, K., Handley, S. L.:
 Actions and some interactions of 5-HT_{1A} ligands in the elevated X-maze and effects of dorsal raphe lesions.
Psychopharmacology 106: 484-490, 1992

Curzon, G., Kennett, G. A.:
 m-CPP: a tool for studying behavioural responses associated with 5-HT_(1C) receptors.
Trends Pharmacol. Sci. 11:181-182, 1990

Dahlström, A., Fuxe, K.:
 Evidence for the existence of monoamine-containing neurons in the central nervous system.
 I. Demonstration of monoamines in the cell bodies of brain stem neurons.
Acta Physiol. Scand. 62 [Suppl 232]: 1-55, 1964

Danner, H., Pfister, C.:
 Untersuchungen zur Zyroarchitektonik des Nucleus Raphe dorsalis der Ratte.
J. Hirnforsch. 21: 655-664, 1980

Dawson, T. M., Wamsley, J. F.:
 Autoradiographic localization of [³H]imipramine binding sites: Association with serotonergic neurons.
Brain Res. Bull. 11: 325-334, 1983

Deakin, J. F. W., File, S. E., Hyde, J. R. G., MacLeod, N. K.:
 Ascending 5-HT pathways and behavioural habituation.
Pharm. Biochem. Behav. 10: 687-694, 1979

Delgado, J. M., DeFeudis, F. V., Roth, R. H., Ryugo, D. K., Mitruka, B. M.:
 Dialytrode for longterm intracerebral perfusion in awake monkeys.
Arch. Int. Pharmacodyn. Ther. 198: 9-21, 1972

D'Haenen, H., Bossuyt, A., Mertens, J., Bossuyt-Piron, C., Gijsemans, M., Kaufman, L.:
 SPECT imaging of serotonin₂ receptors in depression.
Psychiatry Res. 45: 227-237, 1992

Dolberg, O. T., Sasson, Y., Cohen, R., Zohar, J.:
 The relevance of behavioral probes in obsessive-compulsive disorder.
Eur. Neuropsychopharmacol. 5: 161-162, 1995

Dourish, C. T., Coughlan, J., Hawley, D., Clark, M., Iversen, S. D.:
 Blockade of CCK-induced hypophagia and prevention of morphine tolerance by the CCK antagonist
 L-364,718.
In: Cholecystokinin antagonists, R. Y. Wang, R. Schoenfeld (Hrsg.), pp. 307-326.
 New York, Alan R. Liss, Inc., 1988a

Dourish, C. T., Cooper, S. J., Gilbert, F., Coughlan, J., Iversen, S. D.:
 The 5-HT_{1A} agonist 8-OH-DPAT increases consumption of palatable wet mash and liquid diets in the rat.
Psychopharmacology 94: 58-63, 1988b

Dourish, C. T., Ruckert, A. C., Tattersall, F. D., Iversen, S. D.:
 Evidence that decreased feeding induced by systemic injection of cholecystokinin is mediated by CCK-A receptors.
Eur. J. Pharmacol. 173: 233-234, 1989

Drewnowski, A., Kurth, C., Holden-Wiltse, J., Saari, J.:
 Food preferences in human obesity: Carbohydrate versus fats.
Appetite 18: 207-221, 1992

Dumuis, A., Bouhelal, R., Sebben, M., Bockaert, J.:
 A 5-HT receptor in the central nervous system, positively coupled with adenylate cyclase, is antagonised by ICS 205930.
Eur. J. Pharmacol. 146: 187-188, 1988

Eglen, R. M., Bonhaus, D. W., Clark, R. D., Daniels, S., Leung, E., Wong, E. H. F., Fontana, D. J.:
 Effects of a selective and potent 5-HT₄ receptor agonist, RS-67333, and antagonist, RS-67532, in a rodent model of spatial learning and memory.
Br. J. Pharmacol. 116: 235p, 1995a

Eglen, R. M., Wong, E. H. F., Dumuis, A., Bockaert, J.:
 Central 5-HT₄ receptors.
Trends Pharmacol. Sci. 16: 391-398, 1995b

el-Hawary, M. B., Feldberg, W.:
 Effect of 5-hydroxytryptophan acting from the cerebral ventricles on 5-hydroxytryptamine output and body temperature.
J. Physiol. (Lond) 186(2): 401-415, 1966

Erlander, M. G., Lovenberg, T. W., Baron, B. M., Delecea, L., Danielson, P. E., Racke, M., Slone, A. L., Siegel, B. W., Foye, P. E., Cannon, K., Bruns, J. E., Sutcliffe, J. G.:
 Two members of a distinct subfamily of 5-hydroxytryptamine receptors differentially expressed in rat brain.
Proc. Natl. Acad. Sci. USA 90: 3452-3456, 1993

Falck, B., Hillarp, N. A., Thieme, G., Torp, A.:
 Fluorescence of catecholamines and related compounds condensed with formaldehyde.
J. Histchem. Cytochem. 10: 348-354, 1962

Fan, Z. W., Eng, J., Miedel, M., Hulmes, J. D., Pan, Y.-C. E., Yallow, R. S.:
 Cholecystokinin octapeptides purified from chinchilla and chicken brains.
Brain Res. Bull. 18: 757-760, 1987

Fernstrom, J. D., Wurtman, R. J.:
 Brain serotonin content: Increase following ingestion of carbohydrate diet.
Science 174: 1023-1025, 1971

File, S. E.:
 Effects of parachlorophenylalanine and amphetamine on habituation of exploration.
Pharmac. Biochem. Behav. 6: 151-156, 1977

File, S. E.:
 Chemical lesions of both dorsal and median raphe nuclei and changes in social and aggressive behaviour in rats.
Pharmacol. Biochem. Behav. 12: 855-859, 1980

File, S. E., Hyde, J. R. G.:
 The effects of *p*-chlorophenylalanine and ethanolamine-*O*-sulphate in an animal test of anxiety.
J. Pharm. Pharmacol. 20: 735-738, 1977

File, S. E., Hyde, J. R. G., McLeod, N. K.:
 5,7-dihydroxytryptamine lesions of dorsal and median raphé nuclei and performance in the social interaction test of anxiety and in a home-cage aggression test.
J. Affect. Dis. 1: 115-122, 1979

File, S. E., Lister, R. G.:
 β -CCE and chlordiazepoxide reduce exploratory head-dipping and rearing: no mutual antagonism.
Neuropharmacology 21: 1215-1218, 1982

File, S. E., Pellow, S., Braestrup, C.:
 Effects of the β -carboline, FG 7142, in the social interaction test of anxiety and the holeboard: correlations between behavior and plasma concentrations.
Pharmac. Biochem. Behav. 23: 33-36, 1985

File, S. E., Wardill, A. G.:
 Validity of head-dipping as a measure of exploration in a modified holeboard.
Psychopharmacology 44: 53-59, 1975

Fluoxetine Bulimia Nervosa Collaborative Group:
 Fluoxetine in the treatment of bulimia nervosa: a multicenter placebo-controlled, double-blind trial.
Arch. Gen. Psychiatry 49: 39-147, 1992

Foguet, M., Hoyer, D., Pardo, L. A., Kluxen, F. W., Kalkman, H. O., Stuhmer, W., Lübbert, H.:
 Cloning and functional characterization of the rat stomach fundus serotonin receptor.
EMBO J. 11: 3481-3487, 1992

Fozard, J. R.:
 Neuronal 5-HT receptor in the periphery.
Neuropharmacology 23: 1473-1486, 1984

Fozard, J. R.:
 Role of 5-HT₃ receptors in nociception.
In: 5-hydroxytryptamine₃ receptor antagonists, F. D. King, B. J. Jones, G. J. Sanger (Hrsg.), pp. 241-253.
 CRC Inc., Boca Raton, 1994

Fozard, J. R., Gray, J. A.:
 5-HT_{1C} receptor activation: a key step in the initiation of migraine?
Trends Pharmacol. Sci. 10: 307-309, 1989

Frankfurt, M., Azmitia, E.:
 Regeneration of serotonergic fibers in the rat hypothalamus following unilateral 5,7-dihydroxytryptamine injection.
Brain Res. 298: 273-282, 1984

Frey, P.:
 Cholecystokinin octapeptide (CCK 26-33) nonsulfated octapeptide and tetrapeptide (CCK 30-33) in rat brain: Analysis by high pressure liquid chromatography (HPLC) and radioimmunoassay (RIA).
Neurochem. Int. 5: 811-815, 1983

Gaddum, J. H.:
 Antagonism between LSD and 5-hydroxytryptamine.
J. Physiol. Lond.: 121, 1953

Gaddum, J. H., Picarelli, Z. P.:
 Two kinds of tryptamine receptors.
Br. J. Pharmac. 12: 323-328, 1957

Gaddum, J. H.:
 Push-pull cannulae.
J. Physiol. 49: 133-143, 1962

Garlow, S. J., Morilak, D. A., Dean, R. R., Roth, B. L., Ciaranello, R. D.:
 Production and characterization of a specific 5-HT₂ receptor antibody.
Brain Res. 615: 113-120, 1993

Geary, N., Kissileff, H. R., Pi-Sunyer, F. X., Hinton, V.:
 Individual, but not simultaneous, glucagon and cholecystokinin infusions inhibit feeding in men.
Am. J. Physiol. 262: R975-R980, 1992

Geller, I., Blum, K.:
 The effect of 5-HTP on para-Chlorophenylalanine (p-CPA) attenuation of „conflict“ behavior.
Eur. J. Pharmacol. 9: 319-324, 1970

Geraciotti, T. D., Jr., Liddle, R. A.:
 Impaired cholecystokinin secretion in bulimia nervosa.
N. Engl. J. Med. 319: 683-688, 1988

Geraciotti, T. D., Jr., Liddle, R. A., Altemus, M., Demitrack, M. A., Gold, P. W.:
 Regulation of appetite and cholecystokinin secretion in anorexia nervosa.
Am. J. Psychiatry 149: 958-961, 1992

Gershon, M., Tamir, R.:
 Serotonectin and the family of proteins that binds serotonin.
Biochem. Pharmacol. 33: 3115-3118, 1984

Geyer, M. A., Petersen, L. R., Rose, G. J.:
 Effects of serotonergic lesions on investigatory responding by rats in a holeboard.
Behav. Neural Biol. 30: 160-177, 1980

Ghelardini, C., Galeotti, N., Casamenti, F., Malmberg-Aiello, P., Pepeu, G., Gualtieri, F., Bartolini, A.:
 Central cholinergic antinociception induced by 5-HT₄ agonists: BIMU 1 and BIMU 8.
Life Sci. 58: 2297-2309, 1996

Gibbs, J., Young, R. C., Smith, G. P.:
 Cholecystokinin decreases food intake in rats.
J. Comp. Physiol. Psychol. 84: 488-495, 1973a

Gibbs, J., Young, R. C., Smith, G. P.:
 Cholecystokinin elicits satiety in rats with open gastric fistulas.
Nature 245: 323-325, 1973b

Glatt, C. E., Snowman, A. M., Sibley, D. R., Snyder, D. R.:
 Clozapine: selective labeling of sites resembling 5HT₆ serotonin receptors may reflect psychoactive profile.
Mol. Med. 1: 398-406, 1995

Göthert, M.:

5-HT receptor mediating pre-synaptic autoinhibition in central serotonergic nerve terminals.
In: Serotonin: Actions, Receptors, Pathophysiology, E. J. Mylecharane, J. A. Angus, I. S. de la Lande, P. P. A. Humphrey (Hrsg.), pp. 56-63.
 MacMillan Press, Hounds Mills, 1989

Goldbloom, D. S., Garfinkel, P. E.:

The serotonin hypothesis of bulimia nervosa: theory and evidence.
Can. J. Psychiatry 35(9): 741-744, 1990

Golden, R. N., Gilmore, J. H., Corrigan, M. H., Ekstrom, R. D., Knight, B. T., Garbutt, J. C.:
 Serotonin, suicide, and aggression: clinical studies.
J. Clin. Psychiatry 52(Suppl): 61-69, 1991

Goodwin, G. M., De Souza, R. J., Green, A.R., Heal, D. J.:

The pharmacology of the behavioural and hypothermic response of rats to 8-hydroxy-2-(di-n-propylamino)tetralin (8-OH-DPAT).
Psychopharmacology 91: 506-511, 1987

Goodwin, G. M., Fairburn, C. G., Cowen, P. J.:

Dieting changes serotonergic function in women, not men: Implications for the aetiology of anorexia nervosa?
Psychol. Med. 17: 839-842, 1987

Gorzalka, B. B., Mendelson, S. D., Watson, N. V.:

Serotonin receptor subtypes and sexual behavior.
Ann. NY Acad. Sci. 600: 435-444, 1990

Graeff, F. G., Silveira Filho, N. G.:

Behavioral inhibition induced by electrical stimulation of the median raphe nucleus of the rat.
Physiol. Behav. 21: 477-484, 1978

Graeff, F. G., Viana, M. B., Mora, P. O.:

Dual role of 5-HT in defense and anxiety.
Neurosci. Biobehav. Rev. 21: 791-799, 1997

Griebel, G.:

5-Hydroxytryptamine-interacting drugs in animal models of anxiety disorders: More than 30 years of research.

Pharmac. Ther. 65: 319-395, 1995

Grignaschi, G., Mantelli, B., Fracasso, C., Anelli, M., Caccia, S., Samanin, R.:

Reciprocal interaction of 5-hydroxytryptamine and cholecystokinin in the control of feeding patterns in rats.

Br. J. Pharmacol. 109: 491-494, 1993

Grossmann, C. J., Kilpatrick, G. J., Bunce, K. T.:

Development of a radioligand binding assay for 5-HT₄ receptors in guinea-pig and rat brain.

Br. J. Pharmacol. 109: 618-624, 1993

Gustafson, E. L., Durkin, M. M., Bard, J. A., Zgombick, J., Branchek, T. A.:

A receptor autoradiographic and in situ hybridization analysis of the distribution of the 5-HT₇ receptor in rat brain.

Br. J. Pharmacol. 117: 657-666, 1996

Haan, E. A.; Jennings, I. G., Cuello, A. C., Nakata, H., Chow, C. W., Kushinsky, R., Brittingham, J., Cotton, R. G. H.: Identification of serotonergic neurons in the human brain by a monoclonal antibody binding to all three aromatic amino acid hydroxylases.
Brain Res. 426: 19-27, 1987

Haefely, W., Pieri, L., Polc, P., Schaffner, R.: General pharmacology and neuropharmacology of benzodiazepine derivatives.
In: Handbook of Experimental Pharmacology, Vol. 55 Pt II, F. Hoffmeister, G. Stille (Hrsg.), pp. 13-262.
 Springer Verlag, Berlin, 1981

Halasy, K., Miettinen, R., Szabat, E., Freund, T. F.: GABAergic interneurons are the major postsynaptic targets of median raphe afferents in the rat dentate gyrus.
Eur. J. Neurosci. 4: 144-153, 1992

Haleem, D. J., Kennett, G. A., Curzon, G.: Hippocampal 5-HT synthesis is greater in female rats than in males and more decreased by 5-HT_{1A} agonist 8-OH-DPAT.
J. Transm. Gen. Sec. 79: 93-101, 1990

Halliday, G., Harding, A., Paxinos G.: Serotonin and tachykinin systems.
In: The rat nervous system, G. Paxinos (Hrsg.), pp. 929-974.
 Academic Press, San Diego, 1995

Halmi, K. A., Eckert, E., LaDu T. J., Cohen, J.: Anorexia nervosa treatment efficacy of cyproheptadine and amitryptyline.
Arch. Gen. Psychiatry 43: 177-181, 1986

Handley, S. L., Mithani, S.: Effects of alpha-adrenoceptor agonists and antagonists in a maze-exploration model of 'fear'-motivated behaviour.
Naunyn-Schmiedeberg's Arch. Pharmacol. 327: 1-5, 1984

Hanson, H., Strouse, J.: Effects of the CCK antagonist L-364,718 on food intake and on the blockade of feeding produced by exogenous CCK in the rat.
Fed. Proc. 46: 1480, 1987

Haring, J. H.: Reorganization of the area dentata serotonergic plexus after lesions of the median raphe nucleus.
J. Comp. Neurol. 306: 576-584, 1991

Harty, R. F., Pearson, P. H., Solomon, T. E., McGuigan, J. E.: Cholecystokinin, vasoactive intestinal peptide and peptide histidine methionine responses to feeding in anorexia nervosa.
Regul. Pept. 36: 141-150, 1991

Hays, S. E., Beinfeld, M. C., Jensen, R. T., Goodwin, F. K., Paul, S. M.: Demonstration of a putative receptor site for cholecystokinin in rat brain.
Neuropeptides 1: 53-62, 1980

Hertz, L., Tamir, H.:

Some properties of an astrocytic protein fraction that binds serotonin.

J. Neurochem. 37: 1331-1334, 1981

Heuring, R. E., Peroutka, S. J.:

Characterization of a novel [³H]5-hydroxytryptamine binding site subtype in bovine brain membranes.

J. Neurosci. 7: 894-903, 1987

Higgins, G. A., Joharchi, N., Nguyen P., Sellers, E. M.:

Effect of the 5-HT₃ receptor antagonists, MDL72222 and ondansetron on morphine place conditioning.

Psychopharmacology 106: 315-320, 1992

Hill, A. J., Blundell, J. E.:

Model system for investigating the actions of anorectic drugs: Effect of d-fenfluramine on food intake, nutrient selection, food preferences, meal patterns, hunger and satiety in healthy human subjects.

In: Advances in the Biosciences, Vol. 60, E. Ferrari, F. Brambilla (Hrsg.), pp. 377-389.

Pergamon Press, New York, 1986

Hölzel, B., Pfister, C.:

Zur Neurontypisierung des Nucleus centralis superior (Nucleus raphe medianus) der Ratte.

J. Hirnforsch. 24: 593-597, 1983

Hofmann, A.:

The discovery of LSD and subsequent investigations in naturally occurring hallucinogens.

In: Discoveries in biological psychiatry, F. J. Jr. Ayd, b. Blackwell (Hrsg.), pp. 91-106.

Lippincott, Philadelphia, Toronto, 1970

Hogg, S.:

A review of the validity and variability of the elevated plus-maze as an animal model of anxiety.

Pharmacol. Biochem. Behav. 54(1): 21-30, 1996

Hornung, J.-P., Fritschy, J.-M., Törk, I.:

Distribution of two morphologically distinct subsets of serotonergic axons in the cerebral cortex of the marmoset.

J. Comp. Neurol. 297: 165-181, 1990

Hoyer, D., Clarke, D. E., Fozard, J. R., Hartig, P. R., Martin, G. R., Mylecharane, E. J., Saxena, P. R., Humphrey, P. P. A.:

VII. International union of pharmacology classification of receptors for 5-hydroxytryptamine (serotonin).

Pharmacol. Rev. 46: 157-203, 1994

Hoyer, D., Martin, G.:

5-HT receptor classification and nomenclature: towards a harmonization with the human genome.

Pharmacology 36: 419-428, 1997

Innis, R. B., Synder, S. H.:

Cholecystokinin receptor binding in brain and pancreas: regulation of pancreatic binding by cyclic and acyclic guanine nucleotides.

Eur. J. Pharmacol. 65: 123-124, 1980

Inoue, T., Tsuchiya, K., Koyama, T.:
 Regional changes in dopamine and serotonin activation with various intensity of physical and psychological stress in the rat brain.
Pharmacol. Biochem. Behav. 49: 911-920, 1994

Ivy, A. C., Oldberg, E.:
 A hormone mechanism for gallbladder contraction and evacuation.
Am. J. Physiol. 86: 599-613, 1928

Jakeman, L. B., To, Z P., Eglen, R. M., Wong, E. H., Bonhaus, D. W.:
 Quantitative autoradiography of 5-HT₄ receptors in brains of three species using two structurally distinct radioligands, [³H]GR113808 and [³H]BIMU-1.
Neuropharmacology 33: 1027-1038, 1994

Jacobs, B. L.:
 Serotonin and Behavior: Emphasis on motor control.
J. Clin. Psychiatry 52(12 suppl): 17-23, 1991

Jenner, P., Chadwick, D., Reynolds, E. H., Marsden, C. D.:
 Altered 5HT metabolism with clonazepam, diazepam and diphenylhydantoin.
J. Pharm. Pharmac. 27: 707-710, 1975

Jimerson, D. C., Lesem, M. D., Kaye, W. H., Hegg, A. P., Brewerton, T. D.:
 Eating disorders and depression: is there a serotonin connection?
Biol. Psychiatry 28: 443-454, 1990

Jouvet, M.:
 Neurophysiology of the states of sleep.
Physiol. Reviews 47(2): 117-177, 1967

Kaneyuki, T., Morimasa, T., Shohmori, T.:
 Action of peripherally administered cholecystokinin on monoaminergic and GABAergic neurons in the rat brain.
Acta Med. Okayama 43: 153-159, 1989

Kaye, W. H., Weltzin, T. E.:
 Serotonin activity in anorexia and bulimia nervosa: relationship to the modulation of feeding and mood.
J. Clin. Psychiatry 52(Suppl.): 41-48, 1991

Kaye, W. H., Weltzin, T. E., McKee, M., McConaha, C., Hansen, D., Hsu, L. K. G.:
 Laboratory assessment of feeding behavior in bulimia nervosa and healthy women: methods for developing a human feeding laboratory.
Am. J. Clin. Nutri. 55: 372-380, 1992

Kennett, G. A., Curzon, G.:
 Evidence that hypophagia induced by mCPP and TFMPP requires 5-HT_{1C} and 5-HT_{1B} receptors; hypophagia induced by RU 24969 only requires 5-HT_{1B} receptors.
Psychopharmacology 96: 93-100, 1988

Kennett, G. A., Whitton, P., Shah, K., Curzon, G.:
 Anxiogenic-like effects of mCPP and TFMPP in animal models are opposed by 5-HT_{1C} receptor antagonists.
Eur. J. Pharmacol. 164: 445-454, 1989

Kia, H. K., Miquel, M.-C., Brisorgueil, M.-J., Daval, G. D., Riad, M., Mestikawy, S. E., Hamon, M., Vergé, D.:
Immunocytochemical localization of serotonin_{1A} receptors in the rat central nervous system.

J. Comp. Neurol. 365: 289-305, 1996

Kirby, L. G., Kreiss, D. S., Singh, A., Lucki, I.:

Effect of destruction of serotonin neurons on basal and fenfluramine-induced serotonin release in striatum.

Synapse 20: 99-105, 1995

Kiser, R. S., Brown, C. A., Sanghera, M. K., German, D. C.:

Dorsal raphe nucleus stimulation reduces centrally-elicited fearlike behavior.

Brain Res. 191: 265-272, 1980

Klein, D. F., Fink, M.:

Psychiatric reaction patterns to imipramine.

Am. J. Psychiatry 119: 432-438, 1962

Koelliker, A.:

Der feinere Bau des verlängerten Markes.

Anat. Anz. 6: 427-431, 1891

Koenig, J. F. R., Klipper, R. A., Lange, B., Grunwald, U.:

The rat brain. A stereotaxic atlas of forebrain and lower parts of the brain stem.

Williams & Wilkins, Baltimore, 1963

Kohen, R., Metcalf, M. A., Khan, N., Druck, T., Huebner, K., Lachowicz, J. E., Meltzer, H. Y., Sibley, D. R., Roth, B. L., Hamblin, M. W.:

Cloning, characterization, and chromosomal localization of a human 5-HT₆ serotonin receptor.

J. Neurochem. 66: 47-56, 1996

Kursar, J. D., Nelson, D. L., Wainscott, D. B., Cohen, M. L., Baez, M.:

Molecular cloning, functional expression, and pharmacology of a novel serotonin receptor (5-hydroxytryptamine_{2F}) from rat stomach fundus.

Mol. Pharmacol. 42: 227-234, 1992

Larsen, P. J., Hay-Schmidt, A., Vrang, N., Mikkelsen, J. D.:

Origin of projections from the midbrain raphe nuclei to the hypothalamic paraventricular nucleus in the rat: A combined retrograde and anterograde tracing study.

Neuroscience 70: 963-988, 1996

Lauder, J. M.:

Hormonal and humoral influences on brain development.

Psychoneuroendocrinology 8: 121-155, 1983

Leibowitz, S. F., Jhanwar-Uniyal, M.:

5-HT_{1A} and 5-HT_{1B} receptor binding sites in discrete hypothalamic nuclei: Relation to circadian rhythm and gender.

Soc. Neurosci. Abs. 16: 294, 1990

Leibowitz, S. F., Lucas, D. J., Leibowitz, K. L., Jhanwar, Y. S.:

Developmental patterns of macronutrient intake in female and male rats from weaning to maturity.

Physiol. Behav. 50: 1167-1174, 1991

Leibowitz, S. F., Weiss, G. F., Shor-Posner, G.:
 Hypothalamic serotonin: pharmacological, biochemical and behavioral analyses of its feeding-suppressive action.
Clin. Neuropharmacol. 11: S51-S71, 1988

Leonhardt, S., Herrick-Davies, K., Titeler, M.:
 Detection of a novel serotonin receptor subtype (5-HT_{1E}) in human brain: interaction with a GTP-binding protein.
J. Neurochem. 53: 465-471, 1989

Lister, R. G.:
 The effects of ethanol on exploration in DAB/2 and C57BL/6 mice.
Alcohol 4: 17-19, 1987a

Lister, R. G.:
 Interactions of Ro 15-4513 with diazepam, sodium pentobarbital and ethanol in a holeboard test.
Pharmac. Biochem. Behav. 28: 75-79, 1987b

Lister, R. G.:
 The benzodiazepine receptor inverse agonist FG 7142 and Ro 15-4513 both reverse some of the behavioral effects of ethanol in a holeboard test.
Life Sci. 41: 1481-1489, 1987c

Lister, R. G.:
 The use of a plus-maze to measure anxiety in the mouse.
Psychopharmacology 92: 180-185, 1987d

Lopez-Ibor, J. J. Jr., Fernandez-Cordoba, E.:
 La monoclormipramina en enfermos resistentes a otros tratamientos.
Actas Luso Esp. Neurol. Psiquiatr. 26: 119-147, 1967

Lopez-Rubalcava, C., Saldivar, A., Fernandez-Guasti, A.:
 Interaction of GABA and serotonin in the anxiolytic action of diazepam and serotonergic anxiolytics.
Pharmacol. Biochem. Behav. 43: 433-440, 1992

Lorens, S. A.:
 Some behavioral effects of serotonin depletion depend on method: A comparison of 5,7-dihydroxytryptamine, p-chlorophenylalanine, p-chloroamphetamine and electrolytic raphe lesions.
Ann. NY Acad. Sci. 305: 532-555, 1978

Lovenberg, T. W., Baron, B. M., De Lecea, L., Miller, J. D., Prosser, R. A., Rea, M. A., Foye, P. E., Racke, M., Slone, A. L., Siegel, B. W., Danielson, P. E., Sutcliffe, J. G., Erlander, M. G.:
 A novel adenylyl cyclase-activating serotonin receptor (5-HT₇) implicated in the regulation of mammalian circadian rhythms.
Neuron 11: 449-458, 1993a

Lovenberg, T. W., Erlander, M. G., Baron, M. G., Racke, M., Slone A. L., Siegel, B. W., Craft, C. M., Burns, J. E., Danielson, P. E., Sutcliffe, J. G.:
 Molecular cloning and functional expression of 5-HT_{1E}-like rat and human 5-hydroxytryptamine receptor genes.
Proc. Natl. Acad. Sci. USA 90: 2184-2188, 1993b

Lowther, S., De Paermentier, F., Rufus Crompton, M., Horton, R.:
 The distribution of 5-HT_(1D) and 5-HT_(1E) binding sites in human brain.
Eur. J. Pharmacol. 222: 137-142, 1992

Lydiard, R. B., Brewerton, T. D., Fossey, M. D.:
 CSF cholecystokinin octapeptide in bulimia nervosa and normal controls.
Am. J. Psychiatry 150: 1099-1101, 1993

MacIntosh, F. C., Oborin, P. E.:
 Release of acetylcholine from intact cerebral cortex.
Abstracts of the XIX. Int. Physiol. Congress: 580-581, 1953

Maroteaux, L., Saudou, F., Amlaiky, N., Boschert, U., Plassat, J. L., Hen, R.:
 Mouse 5HT_{1B} serotonin receptor: cloning, functional expression, and localization in motor control centers.
Neurobiology 89: 3020-3024, 1992

Marsden, C. A., Beckett, S. R. G., Wilson, W., Bickerdike, M., Fink, H., Rex, A., Cadogan, A.-K.:
 Serotonin involvement in animal models of anxiety and panic.
In: Serotonin in the central nervous system and Periphery. A. Takade, G. Curzon (Hrsg.),
 pp. 135-143.
 Elsevier, Amsterdam, 1995

Marsden, C. A., Sleight, A. J., Fone, K. C. F., Jonson, J. V., Crespi, F., Martin, K. F., Garrett, J. C.,
 Bennett, G. W.:
 Functional identification of 5HT receptor subtypes.
Comp. Biochem. Physiol. 93A: 107-114, 1989

Massari, V. J., Tizabi, Y., Jacobowitz, D. M.:
 Potential noradrenergic regulation of serotonergic neurones in the median raphe nucleus.
Exp. Brain Res. 34: 177, 1979

Matthes, H., Boschert, U., Amlaiky, N., Grailhe, R., Plassat, J. L., Muscatelli, G., Mattei, M. G.,
 Hen, R.:
 Mouse 5-hydroxytryptamine_{5A} and 5-hydroxytryptamine_{5B} receptors define a new family of serotonin receptors: cloning, functional expression and chromosomal localisation.
Mol. Pharmacol. 43: 313-319, 1993

McQuade, R., Sharp, T.:
 Release of cerebral 5-hydroxytryptamine evoked by electrical stimulation of the dorsal and median raphe nuclei: effect of a neurotoxic amphetamine.
Neuroscience 68: 1079-1088, 1995

Melton, P. M., Kissileff, H. R., Pi-Sunyer, F. X.:
 Cholecystokinin (CCK-8) affects gastric pressure and ratings of hunger and fullness in women.
Am. J. Physiol. 263: R452-R456, 1992

Mengod, G., Nguyen, H., Le, H., Waeber, C., Lubbert, H., Palacios, J. M.:
 The distribution and cellular localization of the serotonin 1C receptor messenger mRNA in the rodent brain examined by in situ hybridization histochemistry comparison receptor binding distribution.
Neurosci. 35: 577-591, 1990a

Mengod, G., Palacios, J. M., Wiederhold, K. H., Hoyer, D.:
 5-hydroxytryptamine receptor histochemistry: comparison of receptor mRNA distribution and radioligand autoradiography in the brain.
In: Serotonergic neurons and 5-HT receptors in the CNS, H. G. Baumgarten, M. Göthert (Hrsg.),
 pp.213-237. Springer, 1997

Mengod, G., Pompeiano, M., Martinez-Mir, M. I., Palacios, J.M.: Localization of the mRNA of the 5-HT₂ receptor by in situ hybridization histochemistry. Correlation with the distribution of receptor sites.
Brain Res. 524: 139-143, 1990b

Meyer-Bernstein, E. L., Blanchard, J. H., Morin, L. P.: The serotonergic projection from the median raphe nucleus to the suprachiasmatic nucleus modulates activity phase onset, but not other circadian rythm parameters.
Brain Res. 755: 112-120, 1997

Miczek, K. A.: Effects of scopolamine, amphetamine and benzodiazepines on conditioned suppression.
Pharmacol. Biochem. Behav. 1: 401-411, 1973

Middlemiss, D. N., Fozard, J. R.: 8-Hydroxy-2-(di-n-propylamino) tetralin discriminates between subtypes of the 5-HT₁-recognition site.
Eur. J. Pharmac. 90: 151-153, 1983

Millar, J., O'Connor, J. J., Trout, S. J., Kruk, Z. L.: Continous scan cyclic voltammetry (CSCV): A new highspeed electrochemical method for monitoring neuronal dopamine release.
J. Neurosci. Methods 43: 109-118, 1992

Miller, K. J., Teitler, M.: Quantitative autoradiography of 5-CT sensitive (5-HT_{1D}) and 5-CT insensitive (5-HT_{1E}) serotonin receptors in human brain.
Neurosci. Lett. 136: 223-226, 1992

Molliver, M. E., Berger, U. V., Mamounas, L. A., Molliver, D. C., O'Hearn, E., Wilson, M. A.: Neurotoxicity of MDMA and related compounds: anatomic studies.
Ann. N. Y. Acad. Sci. 600: 640-664, 1990

Montgomery, K. C.: The relation between fear induced by novel stimulation and exploratory behaviour.
J. comp. Physiol. Psychol. 48: 254-260, 1955

Monsma, F. J., JR., Shen, Y., Ward, R.P., Hamblin, M. W., Sibley, D. R.: Cloning and expression of a novel serotonin receptor with high affinity for tricyclic psychotropic drugs.
Mol. Pharmacol. 43: 320-327, 1993

Morley, J. E.: Species differences in the response to cholecystokinin.
Ann. NY Acad. Sci. 448: 413-416, 1985

Morrow, B. A., Roth, R. H.: Serotonergic lesions alter cocaine-induced locomotor behavior and stress-activation of the mesocorticolimbic dopamine system.
Synapse 23: 174-181, 1996

Muurahainen, N. E., Kissileff, H. R., Lachaussee, J., Pi-Sunyer, F. X.: Effect of a soup preload on reduction of food intake by cholecystokinin in humans.
Am. J. Physiol. 260: R672-R680, 1991

Nashold, B. S., Wilson, N. P., Slaughter, C. S.:
 Sensations evoked by stimulation in the midbrain of man.
J. Neurosurg. 30: 14-24, 1969

O'Connor, J. J., Kruk, Z. L.:
 Pharmacological characteristics of 5-hydroxytryptamine autoreceptors in rat brain slices incorporating the dorsal raphe or the suprachiasmatic nucleus.
Br. J. Pharmacol. 106: 524-532, 1992

O'Neill, M. F., Kitchener, S. J., Dourish, C. T., S. J., Iversen, S. D.:
 The CCK antagonist devazepide and L-365,260 promote weight gain in weanling rats.
Br. J. Pharmacol. 99: 137P, 1990

Palkovits, M., Saavedra, J. M., Jacobowitz, D., Kiser, J. S., Zaborsky, L., Brownstein, M. J.:
 Serotonergic innervation of the forebrain: Effect of lesions on serotonin and tryptophan hydroxylase levels.
Brain Res. 130: 121-134, 1977

Panocka, I., Ciccociopo, R., Plidori, C., Pompei, P., Massi, M.:
 The 5-HT₄ receptor antagonist, GR 113808, reduces ethanol intake in alcohol preferring rats.
Pharmacol. Biochem. Behavior 52: 255-259, 1995

Patel, T. D., Azmitia, E. C., Zhou, F. C.:
 Increased 5-HT_{1A} receptor immunoreactivity in the rat hippocampus following 5,7-dihydroxytryptamine lesions in the cingulum bundle and fimbria-fornix.
Behav. Brain Res. 73: 319-323, 1996

Paxinos, G., Watson, C.:
 The rat brain in stereotaxic coordinates.
 Academic Press, 1986

Pazos, A., Hoyer, D., Palacios, J. M.:
 The binding of serotonergic ligands to the porcine choroid plexus: characterization of a new type of serotonin recognition site.
Eur. J. Pharmacol. 106: 539-546, 1984

Pedigo, B. W., Yamamura, H. I., Nelson, D. L.:
 Discrimination of multiple serotonin binding sites by the neuroleptic spiperone in rat brain.
J. Neurochem. 36: 220-226, 1981

Pellow, S., Chopin, P., File, S. E., Briley, M.:
 Validation of open: closed arm entries in an elevated plus-maze as a measure of anxiety in the rat.
J. Neurosci. Meth. 14: 149-167, 1985

Peroutka, S. J., Snyder, S. H.:
 Multiple serotonin receptors: differential binding of [³H]-5-hydroxytryptamine, [³H]-lysergic acid diethylamide and [³H]-spiroperidol.
Molec. Pharmac. 108: 260P, 1979

Philipp, E., Pirke, K.-M., Keliner, M. B., Krieg, J.-C.:
 Disturbed cholecystokinin secretion in patients with eating disorders.
Life Sci. 48: 2442-2450, 1991

Pirke, K. M., Kellner, M. B., Friess, E., Krieg, J. C., Fichter, M. M.:
Satiety and cholecystokinin.
Int. J. Disord. 15: 63-69, 1994

Pi-Sunyer, X., Kissileff, H. R., Thornton, J., Smith, G. P.:
C-terminal octapeptide of cholecystokinin decreases food intake in obese men.
Physiol. Behav. 29: 627-630, 1982

Plassat, J. L., Amlaiky, N., Hen, R.:
Molecular cloning of a mammalian serotonin receptor that activates adenylyl cyclase.
Mol. Pharmacol. 44: 229-236, 1993

Plassat, J. L., Boschert, U., Amlaiky, N., Hen, R.:
The mouse 5-HT₅ receptor reveals a remarkable heterogeneity within the 5-HT_{1D} receptor family.
EMBO J. 11: 4779-4786, 1992

Poeschla, B., Gibbs, J., Simansky, K. J., Greenberg, D., Smith, G. P.:
Cholecystokinin-induced satiety depends on activation of 5-HT_{1C} receptors.
Am. J. Physiol. Regul. Integr. Comp. Physiol. 264 (33): R62-R64, 1993

Poeschla, B., Gibbs, J., Simansky, K. J., Smith, G. P.:
The 5-HT_{1A} agonist 8-OH-DPAT attenuates the satiating action of cholecystokinin.
Pharmacol. Biochem. Behav. 42: 541-543, 1992

Rapport, M. M., Green, A. A., Page, I. H.:
Crystalline serotonin.
Science 108: 329, 1948

Rehfeld, J. F.:
Immunochemical studies on cholecystokinin. II. Distribution and molecular heterogeneity in the central nervous system and small intestine of man and dog.
J. Biol. Chem. 253: 4022-4030, 1978

Rex, A., Marsden, C. A., Fink, H.:
Effect of diazepam on cortical 5-HT release and behavior in the guinea-pig on exposure to the elevated plus maze.
Psychopharmacology 110: 490-496, 1993

Rex, A., Fink, H., Marsden, C. A.:
Effects of BOC-CCK-4 and L 365.260 on cortical 5-HT release in guinea-pigs on exposure to the elevated plus maze.
Neuropharmacology 33: 559-565, 1994

Romero, L., Jernej, B., Bel, N., Cicin-Sain, L., Cortés, R., Artigas, F.:
Basal and stimulated extracellular serotonin concentration in the brain of rats with altered serotonin uptake.
Synapse 28: 313-321, 1998

Rotman, A., Daly, J. W., Creveling, C. R.:
Oxygen-dependent reaction of 6-hydroxydopamine, 5,6-dihydroxytryptamine, and related compounds with proteins in vitro: a model for cytotoxicity.
Mol. Pharmacol. 12: 877-899, 1976

- Ruat, M., Traiffort, E., Arrang, J. M., Tardivel-Lacombe, J., Diaz, J., Leurs, R., Schwartz, J. C.: A novel rat serotonin (5-HT₆) receptor: molecular cloning, localisation and stimulation of cAMP accumulation.
Biochem. Biophys. Res. Commun. 193: 268-276, 1993
- Saito, H. Matsumoto, M., Togashi, H., Yoshioka, M.: Functional interaction between serotonin and other neuronal systems: focus on in vivo microdialysis studies.
Jpn. J. Pharmacol. 70: 203-205, 1996
- Sakaguchi, T, Bray, G. A.: Effect of norepinephrine, serotonin and tryptophan on the firing rate of sympathetic nerves.
Brain Res. 492: 271-280, 1989
- Saner, A., Pletscher, A.: Effects of diazepam on cerebral 5-hydroxytryptamine synthesis.
Eur. J. Pharmacol. 55: 315-318, 1979
- Sankaran, H., Goldfine, I. D., Deveney, C. W., Wong, K. Y., Williams, J. A.: Binding of cholecystokinin to high affinity receptors on isolated rat pancreatic acini.
J. Biol. Chem. 255: 1849-1853, 1980
- Sarna, G.S., Hutson, P. H., Tricklbank, M. D., Curzon, G.: Determination of brain 5-hydroxytryptamine turnover in freely moving rats using repeated sampling of cerebrospinal fluid.
J. Neurochem. 40: 383-388, 1983
- Saxena P. R., Den Boer, M. O.: Pharmacology of antimigraine drugs.
J. Neurol. 238: S28-S35, 1991
- Schildkraut, J. J.: The catecholamine hypothesis of affective disorders: A review of supporting evidence.
Am. J. Psychiatry 122: 509-522, 1965
- Sheard, M. H.: The effect of p-chlorophenylalanine on behavior in rats: relation to brain serotonin and 5-hydroxyindolacetic acid.
Brain Res. 15: 524-528, 1969
- Shimizu, H., Tatsuno, T., Tanaka, H., Hirose, A., Araki, Y., Nakamura, M.: Serotonergic mechanisms in anxiolytic effect of tandospirone in the vogel conflict test.
Jpn. J. Pharmacol. 59: 105-112, 1992
- Shor-Posner, G., Azar, A. P., Insinga, S., Leibowitz, S. F.: Deficits in the control of food intake after hypothalamic paraventricular nucleus lesions.
Phys. Behav. 35: 883-890, 1985
- Silverman, J. A.: Anorexia nervosa clinical and metabolic observations.
Int. J. Eating Disorders 2: 159-166, 1983
- Silverstone, T., Goodall, E.: Serotonergic mechanisms in human feeding: The pharmacological evidence.
Appetite 7: 85-97, 1986

- Simon, P., Panissaud, C., Constanttin, J.:
 Anxiogenic-like effects induced by stimulation of dopamine receptors.
Pharmac. Biochem. Behav. 45: 685-690, 1993
- Sleight, A. J., Carolo, C., Petit, N., Zwingelstein, C., Bourson, A.:
 Identification of 5-hydroxytryptamine₇ receptor binding sites in rat hypothalamus: sensitivity to chronic antidepressant treatment.
Mol. Pharmacol. 47: 99-103, 1995
- Smith, G. P., Gibbs, J.:
 The development and proof of the CCK hypothesis of satiety.
In: Multiple cholecystokinin receptors in the CNS, C. T. Dourish, S. J. Cooper, S. D. Iversen, L. L. Iverson (Hrsg.), pp. 166-182.
 Oxford University Press, 1992
- Söderpalm, B., Engel, A. E.:
 The 5,7-DHT-induced anticonflict effect is dependent on intact adrenocortical function.
Life Sci. 51: 315-326, 1992
- Soubrié, P.:
 Reconciling the role of central serotonin neurons in human and animal behavior.
The behavioral and brain sciences 9: 319-364, 1986
- South, E. H., Ritter, R. C.:
 Capsaicin application to central or peripheral vagal fibers attenuates CCK satiety.
Peptides 9: 601-612, 1988
- Stallone, D., Nicolaïdis, S., Gibbs, J.:
 Cholecystokinin-induced anorexia depends on serotonergic function.
Am. J. Physiol. Regul. Integr. Comp. Physiol. 256: R1138-R1141, 1989
- Stanley, B. G., Schwartz, D. H., Hernandez, L., Leibowitz, S. F., Hoebel, B. G.:
 Patterns of extracellular 5-hydroxyindolacetic acid (5-HIAA) in the paraventricular hypothalamus (PVN): Relation to rhythm and deprivation-induced eating behavior.
Pharmacol. Biochem. Behav. 33: 257-260, 1989
- Stein, L., Wise, C. D., Berger, B. D.:
 Antianxiety action of benzodiazepines: Decrease in activity of serotonin neurons in the punishment system.
In: The Benzodiazepines, S. Garattini, E. Mussini, L. O. Randall (Hrsg.), pp. 299-326.
 Raven Press, New York, 1973
- Steinbusch, H. W. M., Verhofstad, A. A. J., Joosten, H. W. J.:
 Localization of serotonin in the central nervous system by immunohistochemistry: description of a specific and sensitive technique and some applications.
Neuroscience 3: 811-819, 1978
- Sumner, M. J., Humphrey, P. P. A.:
 5-HT_{1D} binding sites in porcine brain can be subdivided by GR43175.
Br. J. Pharmac. 98: 29-31, 1989
- Tabatabaie, T., Dryhurst, G.:
 Molecular mechanisms of action of 5,6- and 5,7-dihydroxytryptamine.
In: Highly selective neurotoxins, R. M. Kostrzewa (Hrsg.), pp. 269-291.
 Humana Press, Totowa, 1998

- Takao, K., Nagatani, T., Kasahara, K.-I., Hashimoto, S.:
 Role of the central serotonergic system in the anticonflict effects of d-AP159.
Pharmac. Biochem. Behav. 43: 503-508, 1992
- Tamai, H., Takemura, J., Kobayashi, N., Matsubayashi, S., Matsukara, S., Nakagawa, T.:
 Changes in plasma cholecystokinin concentration after oral glucose tolerance test in anorexia nervosa before and after therapy.
Metabolism 42: 581-584, 1993
- Tenen, S. S.:
 The effects of p-chlorophenylalanine, a serotonin depletor, on avoidance acquisition, pain sensitivity and related behavior in the rat.
Psychopharmacologia 10: 204-219, 1967
- Thiébot, M.-H., Hamon, M., Soubrié, P.:
 The involvement of nigral serotonin innervation in the control of punishment-induced behavioral inhibition in rats.
Pharmac. Biochem. Behav. 19: 225-229, 1983
- Thoenen, H., Tanzer, J. P.:
 Chemical sympathectomy by selective destruction of adrenergic nerve endings with 6-hydroxydopamine.
Arch. Pharmacol. Exp. Pathol. 261: 271-288, 1968
- Traber, J., Glaser, T.:
 5-HT_{1A} receptor related anxiolytics.
Trends Pharmacol. Sci. 8: 432-437, 1987
- Twarog, B. M., Page, I. H.:
 Serotonin content of some mammalian tissues and urine and a method for its determination.
Am. J. Physiol. 175: 157-161, 1953
- Ueda, S., Kawata, M.:
 Regeneration of serotonergic immunoreactive fibers in the brain of 5,6-dihydroxytryptamine treated rat.
J. Brain. Res. 35: 159-180, 1994
- Ungerstedt, U., Herrera-Marschitz, M., Jungnelius, U., Stahle, L., Tossman, U., Zetterstrom, T.:
 Dopamin synaptic mechanisms reflected in studies combining behavioural recordings and brain dialysis.
Adv. in Biosci. 37: 219-231, 1982
- Ungerstedt, U., Hallstrom, A.:
 In vivo microdialysis - a new approach to the analysis of neurotransmitters in the brain.
Life Sci. 41: 861-864, 1987
- Vanderhaeghen, J. J., Signeau, J. C., Gepts, W.:
 New peptide in the vertebrate CNS reacting with anti gastrin antibodies.
Nature 257: 604-605, 1975
- Van Praag, H. M., Korf, J., Puite, J.:
 5-Hydroxyindolacetic acid levels in the cerebrospinal fluid of depressive patients treated with probenecid.
Nature 25: 1259-1260, 1970

- Waeber, C., Dixon, K., Hoyer, D., Palacios, J. M.:
 Localisation by autoradiography of neuronal 5-HT₃ receptors in mouse CNS.
Eur. J. Pharmacol. 151: 351-352, 1988
- Waeber, C., Hoyer, D., Palacios, J. M.:
 5-HT₃ receptors in the human brain: autoradiographic visualisation using [³H]ICS 205-930.
Neuroscience 31: 393-400, 1989
- Waeber, C., Sebben, M., Nieoullon, A., Bockaert, J., Dumuis, A.:
 Regional distribution and ontogeny of 5-HT₄ binding sites in rodent brain.
Neuropharmacology 33: 527-541, 1994
- Weiss, G. F., Rogacki, N., Fueg, A., Buchen, D., Leibowitz, S. F.:
 Impact of hypothalamic d-norfenfluramine and peripheral d-fenfluramine injection on macronutrient intake in the rat.
Brain Res. Bull. 25: 849-859, 1990
- Weiss, G. F., Rogacki, N., Fueg, A., Buchen, D., Suh, J. S., Wong, D. T., Leibowitz, S. F.:
 Effect of hypothalamic and peripheral fluoxetine injection on natural patterns of macronutrient intake in the rat.
Psychopharmacology 105: 467-476, 1991
- Wieland, S., Lucki, I.:
 Antidepressant-like activity of 5-HT_{1A} agonists measured with the forced swim test.
Psychopharmacology 101: 497-504, 1990
- Wooley, D. W., Shaw, E.:
 Some neurophysiological aspects of serotonin.
Br. Med. J. 2: 122, 1954
- Wright, I. K.; Upton, N.; Marsden, C. A.:
 Effect of established and putative anxiolytics on extracellular 5-HT and 5-HIAA in the ventral hippocampus of rats during behavior on the elevated X-maze.
Psychopharmacology 109: 338-346, 1992
- Wurtmann, J. J., Wurtmann, R. J.:
 d-Fenfluramine selectively decreases carbohydrate but not protein intake in obese subjects.
Int. J. Obesity 8: 79-84, 1984
- Wurtmann, R. J., Wurtmann, J. J.:
 Carbohydrate and depression.
Sci. Am. 260: 68-75, 1989
- Yakel, J. L., Jackson, M. B.:
 5-HT₃ receptors mediate rapid responses in cultured hippocampus and a clonal cell line.
Neuron 1: 615-621, 1988
- Yen, C. Y., Stanger, R. L., Millman, N.:
 Ataractic suppression of isolation-induced aggressive behavior.
Arch. Int. Pharmacodyn. 123: 179-185, 1959
- Zhou, f. C., Azmitia, E. C.:
 Induced homotypic sprouting of serotonergic fibers in hippocampus. II. An immunocytochemistry study.
Brain Research 373: 337-348, 1986