

6. Zitierte Literatur

- Baer K, Essrich C, Balsiger S, Wick MJ, Harris RA, Fritschy JM, Luscher B (2000) Rescue of gamma2 subunit-deficient mice by transgenic overexpression of the GABAA receptor gamma2S or gamma2L subunit isoforms. *Eur J Neurosci* 12: 2639-2643.
- Baer K, Essrich C, Benson JA, Benke D, Bluethmann H, Fritschy JM, Luscher B (1999) Postsynaptic clustering of gamma-aminobutyric acid type A receptors by the gamma3 subunit in vivo. *Proc Natl Acad Sci U S A* 96: 12860-12865.
- Bamforth FJ, Johnson JL, Davidson AG, Wong LT, Lockitch G, Applegarth DA (1990) Biochemical investigation of a child with molybdenum cofactor deficiency. *Clin Biochem* 23: 537-542.
- Barnes EM, Jr. (1996) Use-dependent regulation of GABAA receptors. *Int Rev Neurobiol* 39: 53-76.
- Bedford FK, Kittler JT, Muller E, Thomas P, Uren JM, Merlo D, Wisden W, Triller A, Smart TG, Moss SJ (2001) GABAA receptor cell surface number and subunit stability are regulated by the ubiquitin-like protein Plic-1. *Nat Neurosci* 4: 908-916.
- Beemer FA, Duran M, Wadman SK, Cats BP (1985) Absence of hepatic molybdenum cofactor. An inborn error of metabolism associated with lens dislocation. *Ophthalmic Paediatr Genet* 5: 191-195.
- Bell RM (1986) Protein kinase C activation by diacylglycerol second messengers. *Cell* 45: 631-632.
- Bernard C (1856) Analyse physiologique des propriétés des systèmes musculaires et nerveux au moyen de curare. *Comptes rendus hebdomadaires de l'Académie des sciences, Paris* 43: 825-829.
- Bhalla T, Rosenthal JJ, Holmgren M, Reenan R (2004) Control of human potassium channel inactivation by editing of a small mRNA hairpin. *Nat Struct Mol Biol* 11: 950-956.
- Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, Südhof TC (2002) SynCAM, a Synaptic Adhesion Molecule That Drives Synapse Assembly. *Science* 297: 1525-1531.
- Borgdorff AJ, Choquet D (2002) Regulation of AMPA receptor lateral movements. *Nature* 417: 649-653.
- Bray D, Levin MD, Morton-Firth CJ (1998) Receptor clustering as a cellular mechanism to control sensitivity. *Nature* 393: 85-88.
- Breitinger HG, Becker CM (2002) The inhibitory glycine receptor-simple views of a complicated channel. *Chembiochem* 3: 1042-1052.
- Carroll RC, Zukin RS (2002) NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci* 25: 571.
- Charych EI, Yu W, Miralles CP, Serwanski DR, Li X, Rubio M, De Blas AL (2004) The brefeldin A-inhibited GDP/GTP exchange factor 2, a protein involved in vesicular trafficking, interacts with the beta subunits of the GABA receptors. *J Neurochem* 90: 173-189.

- Colledge M, Froehner SC (1998) Signals mediating ion channel clustering at the neuromuscular junction. *Curr Opin Neurobiol* 8: 357-363.
- Craig AM, Blackstone CD, Haganir RL, Banker G (1994) Selective clustering of glutamate and gamma-aminobutyric acid receptors opposite terminals releasing the corresponding neurotransmitters. *Proc Natl Acad Sci U S A* 91: 12373-12377.
- Dahan M, Levi S, Luccardini C, Rostaing P, Riveau B, Triller A (2003) Diffusion dynamics of glycine receptors revealed by single-quantum dot tracking. *Science* 302: 442-445.
- Danglot L, Triller A, Bessis A (2003) Association of gephyrin with synaptic and extrasynaptic GABA(A) receptors varies during development in cultured hippocampal neurons. *Mol Cell Neurosci* 23: 264-278.
- de Lima AD, Merten MD, Voigt T (1997) Neuritic differentiation and synaptogenesis in serum-free neuronal cultures of the rat cerebral cortex. *J Comp Neurol* 382: 230-246.
- Diez-Guerra FJ, Avila J (1993) MAP2 phosphorylation parallels dendrite arborization in hippocampal neurones in culture. *Neuroreport* 4: 419-422.
- Diez-Guerra FJ, Avila J (1995) An increase in phosphorylation of microtubule-associated protein 2 accompanies dendrite extension during the differentiation of cultured hippocampal neurones. *Eur J Biochem* 227: 68-77.
- Dresbach T, Qualmann B, Kessels MM, Garner CC, Gundelfinger ED (2001) The presynaptic cytomatrix of brain synapses. *Cell Mol Life Sci* 58: 94-116.
- Durkin JP, Tremblay R, Buchan A, Blosser J, Chakravarthy B, Mealing G, Morley P, Song D (1996) An early loss in membrane protein kinase C activity precedes the excitatory amino acid-induced death of primary cortical neurons. *J Neurochem* 66: 951-962.
- Durkin JP, Tremblay R, Chakravarthy B, Mealing G, Morley P, Small D, Song D (1997) Evidence that the early loss of membrane protein kinase C is a necessary step in the excitatory amino acid-induced death of primary cortical neurons. *J Neurochem* 68: 1400-1412.
- Ehlers MD (2001) Reinsertion or degradation of AMPA receptors determined by activity-dependent endocytic sorting. *Neuron* 28: 511-525.
- Essrich C, Lorez M, Benson JA, Fritschy JM, Luscher B (1998) Postsynaptic clustering of major GABAA receptor subtypes requires the gamma 2 subunit and gephyrin. *Nat Neurosci* 1: 563-571.
- Esteller M (2005) DNA methylation and cancer therapy: new developments and expectations. *Curr Opin Oncol* 17: 55-60.
- Faber DS, Korn H (1988) Synergism at central synapses due to lateral diffusion of transmitter. *Proc Natl Acad Sci U S A* 85: 8708-8712.
- Feng G, Steinbach JH, Sanes JR (1998) Rapsyn clusters neuronal acetylcholine receptors but is inessential for formation of an interneuronal cholinergic synapse. *J Neurosci* 18: 4166-4176.

- Fierro L, Llano I (1996) High endogenous calcium buffering in Purkinje cells from rat cerebellar slices. *J Physiol* 496: 617-625.
- Fritschy JM, Brunig I (2003) Formation and plasticity of GABAergic synapses: physiological mechanisms and pathophysiological implications. *Pharmacol Ther* 98: 299-323.
- Fu Z, Washbourne P, Ortinski P, Vicini S (2003) Functional excitatory synapses in HEK293 cells expressing neuroligin and glutamate receptors. *J Neurophysiol* 90: 3950-3957.
- Gallo G, Letourneau PC (2004) Regulation of growth cone actin filaments by guidance cues. *J Neurobiol* 58: 92-102.
- Gardiol A, Racca C, Triller A (1999) Dendritic and postsynaptic protein synthetic machinery. *J Neurosci* 19: 168-179.
- Garner CC, Zhai RG, Gundelfinger ED, Ziv NE (2002) Molecular mechanisms of CNS synaptogenesis. *Trends Neurosci* 25: 243-250.
- Gillen CM, Brill S, Payne JA, Forbush B (1996) Molecular cloning and functional expression of the K-Cl cotransporter from rabbit, rat, and human. A new member of the cation-chloride cotransporter family. *J Biol Chem* 271: 16237-16244.
- Golgi C (1873) *Gazetta Medica Italiana*.
- Graf ER, Zhang X, Jin SX, Linhoff MW, Craig AM (2004) Neurexins Induce Differentiation of GABA and Glutamate Postsynaptic Specializations via Neuroligins. *Cell* 119: 1013-1026.
- Grenningloh G, Pribilla I, Prior P, Multhaup G, Beyreuther K, Taleb O, Betz H (1990) Cloning and expression of the 58 kd beta subunit of the inhibitory glycine receptor. *Neuron* 4: 963-970.
- Grutzendler J, Kasthuri N, Gan WB (2002) Long-term dendritic spine stability in the adult cortex. *Nature* 420: 812-816.
- Guo C, Levine H (1999) A thermodynamic model for receptor clustering. *Biophys J* 77: 2358-2365.
- Guo Y, Sanchez C, Udin SB (2001) MAP2 phosphorylation and visual plasticity in *Xenopus*. *Brain Res* 905: 134-141.
- Gurevich I, Englander MT, Adlersberg M, Siegal NB, Schmauss C (2002) Modulation of serotonin 2C receptor editing by sustained changes in serotonergic neurotransmission. *J Neurosci* 22: 10529-10532.
- Harlan JE, Hajduk PJ, Yoon HS, Fesik SW (1994) Pleckstrin homology domains bind to phosphatidylinositol-4,5-bisphosphate. *Nature* 371: 168-170.
- Henneberger C, Juttner R, Schmidt SA, Walter J, Meier J, Rothe T, Grantyn R (2005) GluR- and TrkB-mediated maturation of GABA receptor function during the period of eye opening. *Eur J Neurosci* 21: 431-440.

- Herb A, Burnashev N, Werner P, Sakmann B, Wisden W, Seeburg PH (1992) The KA-2 subunit of excitatory amino acid receptors shows widespread expression in brain and forms ion channels with distantly related subunits. *Neuron* 8: 775-785.
- Hoshi M, Akiyama T, Shinohara Y, Miyata Y, Ogawara H, Nishida E, Sakai H (1988) Protein-kinase-C-catalyzed phosphorylation of the microtubule-binding domain of microtubule-associated protein 2 inhibits its ability to induce tubulin polymerization. *Eur J Biochem* 174: 225-230.
- Hübner CA, Stein V, Hermans-Borgmeyer I, Meyer T, Ballanyi K, Jentsch TJ (2001) Disruption of KCC2 reveals an essential role of K-Cl cotransport already in early synaptic inhibition. *Neuron* 30: 515-524.
- Huntsman MM, Tran BV, Potkin SG, Bunney WEJ, Jones EG (1998) Altered ratios of alternatively spliced long and short gamma2 subunit mRNAs of the gamma-amino butyrate type A receptor in prefrontal cortex of schizophrenics. *Proc Natl Acad Sci U S A* 95: 15066-15071.
- Husi H, Grant SG (2001) Proteomics of the nervous system. *Trends Neurosci* 24: 259-266.
- Husi H, Ward MA, Choudhary JS, Blackstock WP, Grant SG (2000) Proteomic analysis of NMDA receptor-adhesion protein signaling complexes. *Nat Neurosci* 3: 661-669.
- Jonas P, Bischofberger J, Sandkuhler J (1998) Corelease of two fast neurotransmitters at a central synapse. *Science* 281: 419-424.
- Jones MV, Westbrook GL (1996) The impact of receptor desensitization on fast synaptic transmission. *Trends Neurosci* 19: 96-101.
- Kabir N, Schaefer AW, Nakhost A, Sossin WS, Forscher P (2001) Protein kinase C activation promotes microtubule advance in neuronal growth cones by increasing average microtubule growth lifetimes. *J Cell Biol* 152: 1033-1044.
- Kaila K, Voipio J (1987) Postsynaptic fall in intracellular pH induced by GABA-activated bicarbonate conductance. *Nature* 330: 163-165.
- Kaplan DR, Miller FD (2000) Neurotrophin signal transduction in the nervous system. *Curr Opin Neurobiol* 10: 381-391.
- Keller CA, Yuan X, Panzanelli P, Martin ML, Alldred M, Sassoe-Pognetto M, Luscher B (2004) The gamma2 subunit of GABA(A) receptors is a substrate for palmitoylation by GODZ. *J Neurosci* 24: 5881-5891.
- Kennedy MB (1997) The postsynaptic density at glutamatergic synapses. *Trends Neurosci* 20: 264-268.
- Kennedy MB (2000) Signal-processing machines at the postsynaptic density. *Science* 290: 750-754.
- Kim E, Sheng M (2004) PDZ domain proteins of synapses. *Nat Rev Neurosci* 5: 771-781.
- Kim JH, Haganir RL (1999) Organization and regulation of proteins at synapses. *Curr Opin Cell Biol* 11: 248-254.

- Kins S, Betz H, Kirsch J (2000) Collybistin, a newly identified brain-specific GEF, induces sub-membrane clustering of gephyrin. *Nat Neurosci* 3: 22-29.
- Kirschuk S, Veselovsky N, Grantyn R (1999a) Relationship between presynaptic calcium transients and postsynaptic currents at single gamma-aminobutyric acid (GABA)ergic boutons. *Proc Natl Acad Sci U S A* 96: 7520-7525.
- Kirschuk S, Veselovsky N, Grantyn R (1999b) Single-bouton-mediated synaptic transmission: postsynaptic conductance changes in their relationship with presynaptic calcium signals. *Pflügers Arch* 438: 716-724.
- Kirov SA, Goddard CA, Harris KM (2004) Age-dependence in the homeostatic upregulation of hippocampal dendritic spine number during blocked synaptic transmission. *Neuropharmacology* 47: 640-648.
- Kirov SA, Sorra KE, Harris KM (1999) Slices have more synapses than perfusion-fixed hippocampus from both young and mature rats. *J Neurosci* 19: 2876-2886.
- Kirsch J, Betz H (1998) Glycine-receptor activation is required for receptor clustering in spinal neurons. *Nature* 392: 717-720.
- Kirsch J, Kuhse J, Betz H (1995) Targeting of glycine receptor subunits to gephyrin-rich domains in transfected human embryonic kidney cells. *Mol Cell Neurosci* 6: 450-461.
- Kirsch J, Langosch D, Prior P, Littauer UZ, Schmitt B, Betz H (1991) The 93-kDa glycine receptor-associated protein binds to tubulin. *J Biol Chem* 266: 22242-22245.
- Kirsch J, Wolters I, Triller A, Betz H (1993) Gephyrin antisense oligonucleotides prevent glycine receptor clustering in spinal neurons. *Nature* 366: 745-748.
- Kittler JT, Delmas P, Jovanovic JN, Brown DA, Smart TG, Moss SJ (2000) Constitutive endocytosis of GABA_A receptors by an association with the adaptin AP2 complex modulates inhibitory synaptic currents in hippocampal neurons. *J Neurosci* 20: 7972-7977.
- Kittler JT, Moss SJ (2003) Modulation of GABA(A) receptor activity by phosphorylation and receptor trafficking: implications for the efficacy of synaptic inhibition. *Curr Opin Neurobiol* 13: 341-347.
- Kneussel M, Haverkamp S, Fuhrmann JC, Wang H, Wassle H, Olsen RW, Betz H (2000) The gamma-aminobutyric acid type A receptor (GABAAR)-associated protein GABARAP interacts with gephyrin but is not involved in receptor anchoring at the synapse. *Proc Natl Acad Sci U S A* 97: 8594-8599.
- Kohler M, Burnashev N, Sakmann B, Seeburg PH (1993) Determinants of Ca²⁺ permeability in both TM1 and TM2 of high affinity kainate receptor channels: diversity by RNA editing. *Neuron* 10: 491-500.
- Kuhse J, Schmieden V, Betz H (1990) Identification and functional expression of a novel ligand binding subunit of the inhibitory glycine receptor. *J Biol Chem* 265: 22317-22320.
- Legendre P, Müller E, Badiu CI, Meier J, Vannier C, Triller A (2002) Desensitization of homomeric alpha1 glycine receptor increases with receptor density. *Mol Pharmacol* 62: 817-827.

- Luscher B, Keller CA (2004) Regulation of GABAA receptor trafficking, channel activity, and functional plasticity of inhibitory synapses. *Pharmacol Ther* 102: 195-221.
- Malinow R, Malenka RC (2002) AMPA receptor trafficking and synaptic plasticity. *Annu Rev Neurosci* 25: 103-126.
- Matlin KS, Simons K (1983) Reduced temperature prevents transfer of a membrane glycoprotein to the cell surface but does not prevent terminal glycosylation. *Cell* 34: 233-243.
- Meier J, Akyeli J, Kirischuk S, Grantyn R (2003) GABAA receptor activity and PKC control inhibitory synaptogenesis in CNS tissue slices. *Mol Cell Neurosci* 23: 600-613.
- Meier J, De Chaldee M, Triller A, Vannier C (2000a) Functional Heterogeneity of Gephyrins. *Mol Cell Neurosci* 16: 566-577.
- Meier J, Grantyn R (2004a) A gephyrin-related mechanism restraining glycine receptor anchoring at GABAergic synapses. *J Neurosci* 24: 1398-1405.
- Meier J, Grantyn R (2004b) Preferential accumulation of GABA(A) receptor γ 2L, not γ 2S, cytoplasmic loops at rat spinal cord inhibitory synapses. *J Physiol* 559: 355-365.
- Meier J, Henneberger C, Melnick I, Racca C, Harvey RJ, Heinemann U, Schmieden V, Grantyn R (2005) RNA editing produces P185L single amino acid polymorphism in glycine receptor α 3 resulting in high agonist sensitivity. *Nat Neurosci* in Revision.
- Meier J, Meunier-Durmort C, Forest C, Triller A, Vannier C (2000b) Formation of glycine receptor clusters and their accumulation at synapses. *J Cell Sci* 113: 2783-2795.
- Meier J, Vannier C, Serge A, Triller A, Choquet D (2001) Fast and reversible trapping of surface glycine receptors by gephyrin. *Nat Neurosci* 4: 253-260.
- Meyer G, Kirsch J, Betz H, Langosch D (1995) Identification of a gephyrin binding motif on the glycine receptor beta subunit. *Neuron* 15: 563-572.
- Moss SJ, Doherty CA, Huganir RL (1992) Identification of the cAMP-dependent protein kinase and protein kinase C phosphorylation sites within the major intracellular domains of the beta 1, gamma 2S, and gamma 2L subunits of the gamma-aminobutyric acid type A receptor. *J Biol Chem* 267: 14470-14476.
- Moss SJ, Smart TG (2001) Constructing inhibitory synapses. *Nat Rev Neurosci* 2: 240-250.
- Mumm JS, Kopan R (2000) Notch signaling: from the outside in. *Dev Biol* 228: 151-165.
- Niell CM, Meyer MP, Smith SJ (2004) In vivo imaging of synapse formation on a growing dendritic arbor. *Nat Neurosci* ..
- Nilsson T, Hoe MH, Slusarewicz P, Rabouille C, Watson R, Hunte F, Watzel G, Berger EG, Warren G (1994) Kin recognition between medial Golgi enzymes in HeLa cells. *EMBO J* 13: 562-574.
- Nusser Z, Lujan R, Laube G, Roberts JD, Molnar E, Somogyi P (1998) Cell type and pathway dependence of synaptic AMPA receptor number and variability in the hippocampus. *Neuron* 21: 545-559.

- Obrietan K, van den Pol AN (1995) GABA neurotransmission in the hypothalamus: developmental reversal from Ca²⁺ elevating to depressing. *J Neurosci* 15: 5065-5077.
- Palade GE, Palay SL (1954) Electron microscope observations of interneuronal and neuromuscular synapses. *Anat Rec* 118: 335-336.
- Payne JA, Stevenson TJ, Donaldson LF (1996) Molecular characterization of a putative K-Cl cotransporter in rat brain. A neuronal-specific isoform. *J Biol Chem* 271: 16245-16252.
- Pfeiffer F, Graham D, Betz H (1982) Purification by affinity chromatography of the glycine receptor of rat spinal cord. *J Biol Chem* 257: 9389-9393.
- Pisani A, Bonsi P, Calabresi P (2004) Calcium signaling and neuronal vulnerability to ischemia in the striatum. *Cell Calcium* 36: 277-284.
- Powell LM, Wallis SC, Pease RJ, Edwards YH, Knott TJ, Scott J (1987) A novel form of tissue-specific RNA processing produces apolipoprotein-B48 in intestine. *Cell* 50: 831-840.
- Prange O, Wong TP, Gerrow K, Wang YT, El Husseini A (2004) A balance between excitatory and inhibitory synapses is controlled by PSD-95 and neuroligin. *Proc Natl Acad Sci U S A* ..
- Prior P, Schmitt B, Grenningloh G, Pribilla I, Multhaup G, Beyreuther K, Maulet Y, Werner P, Langosch D, Kirsch J (1992) Primary structure and alternative splice variants of gephyrin, a putative glycine receptor-tubulin linker protein. *Neuron* 8: 1161-1170.
- Purkinje J (1837) Über die gangliösen Körperchen in Verschiedenen Teilen des Gehirns. *Berliner Versammlung deutscher Naturforscher* 179.
- Purpura DP (1975) Dendritic differentiation in human cerebral cortex: normal and aberrant developmental patterns. *Adv Neurol* 12: 91-134.
- Quinlan JJ, Firestone LL, Homanics GE (2000) Mice lacking the long splice variant of the gamma 2 subunit of the GABA(A) receptor are more sensitive to benzodiazepines. *Pharmacol Biochem Behav* 66: 371-374.
- Ramming M, Kins S, Werner N, Hermann A, Betz H, Kirsch J (2000) Diversity and phylogeny of gephyrin: Tissue-specific splice variants, gene structure, and sequence similarities to molybdenum cofactor-synthesizing and cytoskeleton-associated proteins. *Proc Natl Acad Sci U S A* 97: 10266-10271.
- Rao A, Cha EM, Craig AM (2000) Mismatched appositions of presynaptic and postsynaptic components in isolated hippocampal neurons. *J Neurosci* 20: 8344-8353.
- Rasmussen H, Rasmussen T, Triller A, Vannier C (2002) Strychnine-Blocked Glycine Receptor Is Removed from Synapses by a Shift in Insertion/Degradation Equilibrium. *Mol Cell Neurosci* 19: 201-215.
- Rees MI, Harvey K, Ward H, White JH, Evans LI, Duguid IC, Hsu CC, Coleman SL, Miller J, Baer K, Waldvogel HJ, Gibbon F, Smart TG, Owen MJ, Harvey RJ, Snell RG (2003) Isoform heterogeneity of the human gephyrin gene (GPHN), binding domains to the glycine receptor and mutation analysis in hyperekplexia. *J Biol Chem* 278: 24688-24696.

- Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K (1999) The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* 397: 251-255.
- Rivera C, Voipio J, Thomas-Crusells J, Li H, Emri Z, Sipila S, Payne JA, Minichiello L, Saarma M, Kaila K (2004) Mechanism of activity-dependent downregulation of the neuron-specific K-Cl cotransporter KCC2. *J Neurosci* 24: 4683-4691.
- Roberts AA, Kellogg CK (2000) Synchronous postnatal increase in alpha1 and gamma2L GABA(A) receptor mRNAs and high affinity zolpidem binding across three regions of rat brain. *Brain Res Dev Brain Res* 119: 21-32.
- Rosdahl JA, Mourton TL, Brady-Kalnay SM (2002) Protein kinase C delta (PKCdelta) is required for protein tyrosine phosphatase mu (PTPmu)-dependent neurite outgrowth. *Mol Cell Neurosci* 19: 292-306.
- Rosenberg M, Meier J, Triller A, Vannier C (2001) Dynamics of Glycine Receptor Insertion in the Neuronal Plasma Membrane. *J Neurosci* 21: 5036-5044.
- Sabatini DM, Barrow RK, Blackshaw S, Burnett PE, Lai MM, Field ME, Bahr BA, Kirsch J, Betz H, Snyder SH (1999) Interaction of RAFT1 with gephyrin required for rapamycin-sensitive signaling. *Science* 284: 1161-1164.
- Salama-Cohen P, Arevalo MA, Meier J, Grantyn R, Rodriguez-Tebar A (2005) NGF Controls Dendrite Development in Hippocampal Neurons by Binding to p75NTR and Modulating the Cellular Targets of Notch. *Mol Biol Cell* 16: 339-347.
- Sanes JR, Lichtman JW (2001) Induction, assembly, maturation and maintenance of a postsynaptic apparatus. *Nat Rev Neurosci* 2: 791-805.
- Scheiffele P (2003) Cell-Cell Signaling During Synapse Formation in the CNS. *Annu Rev Neurosci* 26: 485-508.
- Scheiffele P, Fan J, Choih J, Fetter R, Serafini T (2000) Neuroligin expressed in nonneuronal cells triggers presynaptic development in contacting axons. *Cell* 101: 657-669.
- Schrenk K, Kapfhammer JP, Metzger F (2002) Altered dendritic development of cerebellar Purkinje cells in slice cultures from protein kinase Cgamma-deficient mice. *Neurosci* 110: 675-689.
- Schwarz G, Schrader N, Mendel RR, Hecht HJ, Schindelin H (2001) Crystal structures of human gephyrin and plant cnx1 g domains: comparative analysis and functional implications. *J Mol Biol* 312: 405-418.
- Schwarz N, Kim EY, Winking J, Paulukat J, Schindelin H, Schwarz G (2004a) Biochemical characterization of the high affinity binding between the glycine receptor and gephyrin. *J Biol Chem* 279: 18733-18741.
- Schwarz N, Kim EY, Winking J, Paulukat J, Schindelin H, Schwarz G (2004b) Biochemical characterization of the high affinity binding between the glycine receptor and gephyrin. *J Biol Chem* 279: 18733-18741.

- Schweizer C, Balsiger S, Bluethmann H, Mansuy IM, Fritschy JM, Mohler H, Luscher B (2003) The gamma2 subunit of GABA(A) receptors is required for maintenance of receptors at mature synapses. *Mol Cell Neurosci* 24: 442-450.
- Seeburg PH, Hartner J (2003) Regulation of ion channel/neurotransmitter receptor function by RNA editing. *Curr Opin Neurobiol* 13: 279-283.
- Seitanidou T, Triller A, Korn H (1988) Distribution of glycine receptors on the membrane of a central neuron: an immunoelectron microscopy study. *J Neurosci* 8: 4319-4333.
- Serge A, Fourgeaud L, Hemar A, Choquet D (2002) Receptor activation and homer differentially control the lateral mobility of metabotropic glutamate receptor 5 in the neuronal membrane. *J Neurosci* 22: 3910-3920.
- Shepherd GM, Erulkar SD (1997) Centenary of the synapse: from Sherrington to the molecular biology of the synapse and beyond. *Trends Neurosci* 20: 385-392.
- Singh B, Henneberger C, Meier J, Rodriguez-Tebar A, Grantyn R (2005) Dendritic BDNF expression in BDNF deficient hippocampal neurons differentially influences glutamatergic and GABAergic synapse formation. in Vorbereitung.
- Skuse GR, Cappione AJ, Sowden M, Metheny LJ, Smith HC (1996) The neurofibromatosis type I messenger RNA undergoes base-modification RNA editing. *Nucleic Acids Res* 24: 478-485.
- Sola M, Bavro VN, Timmins J, Franz T, Ricard-Blum S, Schoehn G, Ruigrok RW, Paarmann I, Saiyed T, O'Sullivan GA, Schmitt B, Betz H, Weissenhorn W (2004) Structural basis of dynamic glycine receptor clustering by gephyrin. *EMBO J* 23: 2510-2519.
- Sola M, Kneussel M, Heck IS, Betz H, Weissenhorn W (2001) X-ray crystal structure of the trimeric N-terminal domain of gephyrin. *J Biol Chem* 276: 25294-25301.
- Stallmeyer B, Schwarz G, Schulze J, Nerlich A, Reiss J, Kirsch J, Mendel RR (1999) The neurotransmitter receptor-anchoring protein gephyrin reconstitutes molybdenum cofactor biosynthesis in bacteria, plants, and mammalian cells. *Proc Natl Acad Sci U S A* 96: 1333-1338.
- Tardin C, Cognet L, Bats C, Lounis B, Choquet D (2003) Direct imaging of lateral movements of AMPA receptors inside synapses. *EMBO J* 22: 4656-4665.
- Tovar KR, Westbrook GL (2002) Mobile NMDA Receptors at Hippocampal Synapses. *Neuron* 34: 255-264.
- Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K (2002) Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* 420: 788-794.
- Tremblay R, Chakravarthy B, Hewitt K, Tauskela J, Morley P, Atkinson T, Durkin JP (2000) Transient NMDA receptor inactivation provides long-term protection to cultured cortical neurons from a variety of death signals. *J Neurosci* 20: 7183-7192.
- Triller A, Choquet D (2003) Synaptic structure and diffusion dynamics of synaptic receptors. *Biol Cell* 95: 465-476.

- Triller A, Cluzeaud F, Korn H (1987) gamma-Aminobutyric acid-containing terminals can be apposed to glycine receptors at central synapses. *J Cell Biol* 104: 947-956.
- Triller A, Cluzeaud F, Pfeiffer F, Betz H, Korn H (1985) Distribution of glycine receptors at central synapses: an immunoelectron microscopy study. *J Cell Biol* 101: 683-688.
- Varoqueaux F, Jamain S, Brose N (2004) Neuroligin 2 is exclusively localized to inhibitory synapses. *Eur J Cell Biol* 83: 449-456.
- Volk DW, Lewis DA (2002) Impaired prefrontal inhibition in schizophrenia: relevance for cognitive dysfunction. *Physiol Behav* 77: 501-505.
- Walter J, Henneberger C, Meier J, Müller U, Grantyn R (2005) Antagonists to group I metabotropic glutamate receptors leads to an up-regulation of the number of inhibitory terminals in acute brain slices. in Vorbereitung.
- Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW (1999) GABA(A)-receptor-associated protein links GABA(A) receptors and the cytoskeleton. *Nature* 397: 69-72.
- Wang JB, Burt DR (1991) Differential expression of two forms of GABAA receptor gamma 2-subunit in mice. *Brain Res Bull* 27: 731-735.
- Whitford KL, Dijkhuizen P, Polleux F, Ghosh A (2002) Molecular control of cortical dendrite development. *Annu Rev Neurosci* 25:127-49. Epub;2002 Mar;19.: 127-149.
- Whiting P, McKernan RM, Iversen LL (1990) Another mechanism for creating diversity in gamma-aminobutyrate type A receptors: RNA splicing directs expression of two forms of gamma 2 phosphorylation site. *Proc Natl Acad Sci U S A* 87: 9966-9970.
- Whittingham TS, Lust WD, Christakis DA, Passonneau JV (1984) Metabolic stability of hippocampal slice preparations during prolonged incubation. *J Neurochem* 43: 689-696.
- Wong RO, Ghosh A (2002) Activity-dependent regulation of dendritic growth and patterning. *Nat Rev Neurosci* 3: 803-812.
- Xiang S, Nichols J, Rajagopalan KV, Schindelin H (2001) The crystal structure of Escherichia coli MoeA and its relationship to the multifunctional protein gephyrin. *Structure* 9: 299-310.
- Yeramian E, Trautmann A, Claverie P (1986) Acetylcholine receptors are not functionally independent. *Biophys J* 50: 253-263.
- Yoshimura Y, Yamauchi Y, Shinkawa T, Taoka M, Donai H, Takahashi N, Isobe T, Yamauchi T (2004) Molecular constituents of the postsynaptic density fraction revealed by proteomic analysis using multidimensional liquid chromatography-tandem mass spectrometry. *J Neurochem* 88: 759-768.
- Zhou FX, Cocco MJ, Russ WP, Brunger AT, Engelman DM (2000) Interhelical hydrogen bonding drives strong interactions in membrane proteins. *Nat Struct Biol* 7: 154-160.

6.1 Eidesstattlicher Erklärung gemäß Habilitationsordnung der Charité

EIDESSTATTLICHE VERSICHERUNG

gemäß Habilitationsordnung der Medizinischen Fakultät Charité

Hiermit erkläre ich, daß

- keine staatsanwaltschaftlichen Ermittlungsverfahren gegen mich anhängig sind,
- weder früher noch gleichzeitig ein Habilitationsverfahren durchgeführt oder angemeldet wurde bzw. welchen Ausgang ein durchgeführtes Habilitationsverfahren hatte;
- die vorgelegte Habilitationsschrift ohne fremde Hilfe verfaßt, die beschriebenen Ergebnisse selbst gewonnen wurden, sowie die verwendeten Hilfsmittel, die Zusammenarbeit mit anderen Wissenschaftlerinnen oder Wissenschaftlern und technischen Hilfskräften und die Literatur vollständig angegeben sind,
- mir die geltende Habilitationsordnung bekannt ist.

.....
Datum

.....
Unterschrift