

Bibliography

- [1] <http://lpdaac.usgs.gov/gtopo30/gtopo30.html>.
- [2] <http://www.mcst.ssai.biz/mcstweb/performance/terra/rsb-snr.html>.
- [3] Vaisala Inc., Upper Air Systems product information reference no. RO422-2, 1989.
- [4] P. Albert, R. Bennartz, and J. Fischer. Remote sensing of atmospheric water vapour from backscattered sunlight in cloudy atmospheres. *Journal of Atmospheric and Oceanic Technology*, 18:865–874, 2001.
- [5] B. Bartsch and J. Fischer. Passive remote sensing of columnar water vapour content over land surfaces. *MPI-report No.234, Hamburg, Germany, ISBN 0937-1060*, 1997.
- [6] D. Belmiloud, R. Schermaul, K. Smith, J. Brault, R. Learner, D. Newnham, and J. Tennyson. New studies of the visible and near-infrared absorption by water vapor and some problems with the database. *Geophysical Research Letters*, 27:3703–3706, 2000.
- [7] R. Bennartz and J. Fischer. A modified k-distribution approach applied to narrow band water vapour and oxygen absorption estimates in the near infrared. *J Quant. Spectrosc. Radiat. Transfer*, 1999.
- [8] R. Bennartz and J. Fischer. Retrieval of columnar water vapour over land from backscattered solar radiation using the medium resolution imaging spectrometer. *Remote Sensing of Environment*, 78:274–283, 2001.

- [9] J.-L. Bézy, S. Delwart, and M. Rast. Meris - a new generation of ocean-colour sensor onboard envisat. *esa bulletin*, 103:48–56, 2000.
- [10] S. Bouiffiés, F. M. Bréon, D. Tanré, and P. Dubuisson. Atmospheric water vapor estimate by a differential absorption technique with the polarization and directionality of the earth reflectances (polder) instrument. *Journal of Geophysical Research*, 102:3831–3841, 1997.
- [11] D. E. Bowker, R. E. Davis, D. L. Myrick, K. Stacy, and W. T. Jones. Spectral reflectances of natural targets for use in remote sensing studies. *NASA Ref. Publ. 1139*, 1985.
- [12] L. R. Brown, R. A. Toth, and M. Dulick. Empirical line parameters of (H₂O)-O-16 near 0.94 μm : Positions, intensities and air-broadening coefficients. *Journal of Molecular Spectroscopy*, 212:57–82, 2002.
- [13] I. Chernykh and R. Eskridge. Determination of cloud amount and level from radiosonde soundings. *Journal of Applied Meteorology*, 35:1362–1369, 1996.
- [14] A. Dudia. *Reference forward model version 3: software user's manual*, Eur. Space Technol. Cent. (ESTEC) Document PO-MA-OXF-GS-0003. European Space Agency (ESA), Paris, 1997.
- [15] F. Fell and J. Fischer. MOMO-The FU-Berlin Radiative Transfer Model based on the Matrix-Operator-Method. Model Description and User Manual. *MAST-II, Technical Progress Report, Programm of the European Commission, Contract Number MAS2-CT92-0020*, 1994.
- [16] F. Fell and J. Fischer. Numerical simulation of the light field in the atmosphere-ocean system using the matrix-operator method. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 1257, May 2000.
- [17] J. Fischer. High resolution spectroscopy for remote sensing of physical cloud properties and water vapour. In: *Current Problems in Atmospheric Radiation*, Ed. Lenoble and Geleyn, Deepak Publishing:151–156, 1988.

- [18] J. Fischer, R. Preusker, and L. Schüller. ATBD cloud top pressure. Algorithm Theoretical Basis Document PO-TN-MEL-GS-0006, European Space Agency, 1997.
- [19] R. Frouin, P. Y. Deschamps, and P. Lecomte. Determination from space of atmospheric total water vapour amounts by differential absorption near 940 nm: Theory and airborne verification. *Journal of Applied Meteorology*, 29:448–460, 1990.
- [20] B.-C. Gao, A. F. H. Goetz, E. R. Westwater, J. E. Conel, and R. O. Green. Possible near-ir channels for remote sensing precipitable water vapour from geostationary satellite platforms. *Journal of Applied Meteorology*, 32:1791–1801, 1993.
- [21] B.-C. Gao and J. Kaufman. The modis near-ir water vapour algorithm. Technical Report ATBD-MOD-03, 1998.
- [22] G. Gendt, C. Reigber, and G. Dick. Near real-time water vapor estimation in a german gps network-first results from the ground program of the hgf gasp project. *Physics and Chemistry of the Earth, Part A: Solid Earth and Geodesy*, 26:413–416, 2001.
- [23] L. P. Giver, J. C. Chackerian, and P. Varanasi. Visible and near-infrared H_2^{16}O line intensity corrections for HITRAN-96. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 66:101–105, 2000.
- [24] N. Gustafsson, L. Berre, S. Hörnquist, X.-Y. Huang, M. Lindskog, B. Navascués, K. S. Mogensen, and S. Thorsteinsson. Three-dimensional variational data assimilation for a limited area model - part i: General formulation and the background error constraint. *Tellus*, 53A(4):425–446, 2001.
- [25] Y. Han and E. R. Westwater. Remote sensing of tropospheric water vapor and cloud liquid water by integrated ground-based sensors. *Journal of Atmospheric and Oceanic Technology*, 12(5):1050–1059, 1995.

- [26] J. Koistinen and D. B. Michelson. Baltex weather radar-based precipitation products and their accuracies. *Boreal Environment Research*, (7):253–263, 2002.
- [27] R. L. Kurucz. The solar irradiance by computation. In G. P. Anderson, R. H. Picard, and J. H. Chetwyn, editors, *Proceedings of the 17th Annual Review Conference on Atmospheric Transmission models, Spec. Rep.*, volume 274, page 332. Phillips Lab. Geophys. Dir., Hanson A. F. B, MA, 1995.
- [28] J. C. Liljegren and B. M. Lesht. Measurements of integrated water vapor and cloud liquid water from microwave radiometers at the doe arm cloud and radiation testbed in the U.S. southern great plains. In *Proceedings of the Int. Geoscience and Remote Sensing Symp. (IGARSS)*, pages 1675–1677, Lincoln, NE, 1996.
- [29] M. Lindskog, N. Gustafsson, B. Navascués, K. S. Mogensen, X.-Y. Huang, x. Yang, U. Andrae, L. Berre, S. Thorsteinsson, and J. Rantakokko. Three-dimensional variational data assimilation for a limited area model - part ii: Observation handling and assimilation experiments. *Tellus*, 53A(4):447–468, 2001.
- [30] A. C. Lorenc. Atmospheric data assimilation. Meteorological Office Forecasting Research Div., Scientific paper 34, The Met Office, UK, 1995.
- [31] D. B. Michelson and J. Koistinen. Gauge-radar-network adjustment for the baltic sea experiment. *Phys. Chem. Earth (B)*, 25(10-12):915–920, 2000.
- [32] Jacquinet-Husson N., E. Arié, J. Ballard, A. Barbe, G. Bjoraker, B. Bonnet, L. R. Brown, C. Camy-Peyret, J. P. Champion, A. Chédin, A. Chusin, C. Clerbaux, G. Duxbury, J.-M. Flaud, N. Fourrié, A. Fayt, G. Graner, R. Gamache, A. Goldman, V. Golovko, G. Guelachvili, J. M. Hatmann, J. C. Hilico, J. Hillman, G. Lefèvre, E. Lellouch, S. N. Mikhaïlenko, O. V. Naumenko, V. Nemtchinov, D. A. Newnham, A. Nikitin, J. Orphal, A. Perrin, D. C. Reuter, C. P. Rinsland, L. Rosenmann, L. S. Rothamn, N. A. Scott, J. Selby, L. N. Sinita, J. M. Sirota, A. M. Smith, K. M. Smith, V. G. Tyuterev, R. H. Tipping,

- S. Urban, P. Varanasi, and M. Weber. The 1997 spectroscopic geisa data-bank. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 62:205–254, 1999.
- [33] C. Reigber, G. Gendt, G. Dick, and M. Tomassini. Near-real-time water vapor monitoring for weather forecasts. *GPS World*, 13:18–27, 2002.
- [34] L. S. Rothman, K. Chance, J. Schroeder, and A. Goldman. A new edition of HITRAN database. *Eleventh ARM Science Team Meeting Proceedings, Atlanta, GA, March 19-23, 2001*.
- [35] Rothman L. S., C. P. Rinsland, A. Goldman, S. T. Massie, D. P. Edwards, J. M. Flaud, A. Perrin, C. Camy-Peyret, V. Dana, J.-Y. Mandin, J. Schroeder, A. McCann, R. R. Gamache, R. B. Wattson, K. Yoshino, K. V. Chance, K. W. Jucks, L. R. Brown, V. Nemtchinov, P. Varanasi, C. Camy-Peyret, V. Dana, J. Y. Mandin, J. Schroeder, A. McCann, R. R. Gamache, R. B. Wattson, K. Yoshino, K. V. Chance, K. W. Jucks, L. R. Brown, V. Nemtchinov, and P. Varanasi. The HITRAN molecular spectroscopic database and HAWKS (HITRAN atmospheric workstation): 1996 edition. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 60:665–710, 1998.
- [36] R. Schermaul, R. C. M. Learner, D. A. Newnham, R. G. Williams, J. Ballard, N. F. Zobov, D. Belmiloud, and J. Tennyson. The water vapor spectrum in the region 8600-15,000 cm^{-1} : experimental and theoretical studies for a new spectral line database 1: laboratory measurements. *Journal of Molecular Spectroscopy*, 208:32–42, 2002.
- [37] K. M. Smith, D. A. Newnham, and R. G. Williams. Collision-induced absorption of solar radiation in the atmosphere by molecular oxygen at 1.27 μm ; field observations and model calculations. *Journal of Geophysical Research*, 106:7541–7552, 2001.
- [38] G. L. Stephens. *Remote sensing of the lower atmosphere : an introduction*. Oxford University Press, New York, 1994.

- [39] S. Tahl and M. v. Schoenermark. Determination of the column water vapour of the atmosphere using backscattered solar radiation measured by the Modular Optoelectronic Scanner (MOS). *Int. J. Remote Sensing*, 19(17):3223–3236, 1998.
- [40] D. D. Turner, B. M. Lesht, S. A. Clough, J. C. Liljegren, H. E. Revercomb, and D. C. Tobin. Dry bias and variability in vaisala rs80-h radiosondes: The arm experience. *Journal of Atmospheric and Oceanic Technology*, 20:117–128, 2003.
- [41] M. Vesperini, F.-M. Breon, and D. Tanre. Atmospheric water vapour content from spaceborne polder measurements. *IEEE Trans. Geosci. Remote Sensing*, 37:1613–1619, 1999.
- [42] G. Zimmermann and A. Neumann. MOS-a spaceborne imaging spectrometer for ocean remote sensing. *Backscatter*, 8(2):9–13, 1997.
- [43] G. Zimmermann, A. Neumann, H. Sümlich, and H. Schwarzer. MOS/PRIRODA - an imaging VIS/NIR spectrometer for ocean remote sensing. In *SPIE*, volume 1937, pages 201–206, 1993.