

6. Literaturverzeichnis

- 1 Aasvang E., Kehlet H. Surgical management of chronic pain after inguinal hernia repair. *British Journal of Surgery* 2005; 92: 795-801
- 2 Amid P.K. Groin hernia repair: open techniques. *World Journal of Surgery* 2005 (Accessed October 3, 2005, at www.springerlink.com/media/988T6MYGQGDQN6XHD)
- 3 Amid P.K. Lichtenstein tension-free hernioplasty: its inception, evolution, and principles. *Hernia* 2004; 8: 1-7
- 4 Amid P.K., Shulman A.G., Lichtenstein I.L. A five-step technique for local anesthesia in inguinal hernia repair. *Chirurg* 1994; 65: 388-390
- 5 Anadol A.Z., Ersoy E., Taneri F., Tekin E. Outcome and cost comparison of laparoscopic transabdominal preperitoneal hernia repair versus open Lichtenstein technique. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2004; 14(3): 159-163
- 6 Anderson B., Hallén M., Leveau P., Bergenfelz A., Westerdahl J. Laparoscopic extraperitoneal repair versus open mesh repair: a prospective randomised controlled trial. *Surgery* 2003; 133: 464-472
- 7 Arlt G., Schumpelick V. Die Leistenhernienoperation nach Shouldice. *Zentralblatt Chirurgie* 2002; 127: 565-569
- 8 Arvidsson D., Berndsen F.H., Larsson L.G. et al. (SMIL-group) Randomized clinical trial comparing 5-year recurrence rate after laparoscopic versus Shouldice repair of primary inguinal hernia. *British Journal of Surgery* 2005; 92: 1085-1091
- 9 Aufenacker T.J., van Geldere D., van Mesdag T. et al. The role of antibiotic prophylaxis in prevention of wound infection after Lichtenstein open mesh repair of primary inguinal hernia. A multicenter double-blind randomized controlled trial. *Annals of Surgery* 2004; 240: 955-961
- 10 Bagley F.H., Louras J.C., Conway M.A. Pain management after hernia repair *Journal American College of Surgeons* 2005; 201(1) Letters: 154-155
- 11 Balen E.M., Ferrer J.V., Blasquez L., Herrera J., Lera J.M. Recurrences after prosthetic repair of inguinal hernias by Lichtenstein technique. *Hernia* 2000; 4: 13-36
- 12 Barrat C., Surlin V., Bordea A., Champault G. Management of recurrent inguinal hernias: A prospective study of 163 cases. *Hernia* 2003; 7: 125-129
- 13 Bay-Nielsen M., Nilsson E., Nordin P., Kehlet H. Chronic pain after open mesh and sutured repair of indirect inguinal hernia in young males. *British Journal of Surgery* 2004; 91: 1372-1376

- 14 Bay-Nielsen M., Nordin P., Nilsson E., Kehlet H. Operative findings in recurrent hernia after a Lichtenstein procedure. *The American Journal of Surgery* 2001; 182: 134-136
- 15 Bay-Nielsen M., F. M. Perkins, Kehlet H. Pain and functional impairment 1 year after inguinal herniorrhaphy: a national questionnaire study. *Annals of Surgery* 2001; 233: 1-7
- 16 Bay-Nielsen M., Kehlet H., Strand L. et al. Quality assessment of 26 304 herniorrhaphies in Denmark: a prospective nationwide study. *Lancet* 2001; 358: 1124-1128
- 17 Berndsen F.H., Bjursten L.-M., Simanaitis M., Montgomery A. Does mesh implantation affect the spermatic cord structures after inguinal hernia surgery?. *European Surgical Research* 2004; 36: 318-322
- 18 Berndsen F., Petersson U., Montgomery A. Endoscopic repair of bilateral inguinal hernias – short and late outcome. *Hernia* 2002; 5: 192-195
- 19 Berndsen F., Arvidsson D., Enander L.-K. et al. Postoperative convalescence after inguinal hernia surgery: prospective randomized multicenter study of laparoscopic versus Shouldice inguinal hernia repair in 1042 patients. *Hernia* 2002; 6: 56-61
- 20 Berndsen F., Sevonius D. Changing the path of inguinal hernia surgery decreased the recurrence rate ten-fold. Report from a county hospital. *European Journal of Surgery* 2002;168:592-596
- 21 Bittner R., Sauerland S., Schmedt C.-G. Comparison of endoscopic techniques versus Shouldice and other open non mesh techniques for inguinal hernia repair. *Surgical Endoscopy* 2005; 19: 605-615
- 22 Bittner R. Ergebnisse der Magdeburger Konsensuskonferenz zur Hernienchirurgie. *Zentralblatt Chirurgie* 2003; 128: 549-550
- 23 Bittner R., Schmidt C.-G., Schwarz J., Kraft K., Leibl B. J. Laparoscopic transperitoneal procedure for routine repair of groin hernia. *British Journal of Surgery* 2002; 89: 1062-1066
- 24 Bloor K., Freemantle N., Khadjesari Z., Maynard A. Impact of NICE guidance on laparoscopic surgery for inguinal hernias: analysis of interrupted time series. *British Medical Journal* 2003; 326: 578
- 25 Bobrzynski A., Budzynski A., Biesiada Z., Kowalczyk M., Lubikowski J., Sienko J. Experience – the key factor in successful laparoscopic total extraperitoneal and transabdominal preperitoneal hernia repair. *Hernia* 2001; 5: 80-83
- 26 Bozuk M., Schuster R., Stewart D., Hicks K., Greaney G., Waxman K. Disability and chronic pain after open mesh and laparoscopic inguinal hernia repair *American Surgeon* 2003; 69: 839-841

- 27 Bringman S., Ramel S., Heikkinen T.J., Engmund T., Westman B., Anderberg B. Tension-free inguinal hernia repair: TEP versus mesh-plug versus Lichtenstein. A prospective randomized controlled trial. *Annals of Surgery* 2003; 237: 142-147
- 28 Bringman S., Heikkinen T.J., Wollert S. et al. Early results of a single-blinded, randomized, controlled, Internet-based multicenter trial comparing Prolene and Vypro II mesh in Lichtenstein hernioplasty. *Hernia* 2003 (Accessed March 13, 2006, at <http://springerlink.metapress.com/media/56WE0CGMRG6TVJE>)
- 29 Bringman S., Ramel S., Nyberg B., Anderberg B. Introduction of herniorraphy with mesh plug and patch. *European Journal of Surgery* 2000; 166:310-312
- 30 Brisinda G. Randomized trial comparing the Prolene® hernia system, mesh plug repair and Lichtenstein method for open inguinal hernia repair. *British Journal of Surgery* 2005 (Letter); 92: 493-494
- 31 Callesen T., Bech K., Kehlet H. One-thousand consecutive inguinal hernia repairs under unmonitored local anesthesia. *Anesthesia & Analgesia* 2001; 93: 1373-1376
- 32 Celdran A., Frieyro O., de la Pinta J.C. et al. The role of antibiotic prophylaxis on wound infection after mesh hernia repair under local anesthesia on an ambulatory basis. *Hernia* 2004; 8: 20-22
- 33 Champault G., Barrat C. Inguinal hernia repair with beta glucan-coated mesh: results of two-year follow up. *Hernia* 2005; 9: 125-130
- 34 Chiofalo R., Holzinger F., Klaiber C. Total endoskopische präperitoneale Netzplastik bei primären und Rezidivleistenhernien. Gibt es Unterschiede?. *Chirurg* 2001; 72: 1485-1491
- 35 Chuback J.A., Singh R. S., Sills C., Dick L. S. Small bowel obstruction resulting from mesh plug migration after open inguinal hernia repair. *Surgery* 2000; 127: 475-476
- 36 Cochrane Collaboration: Scott N., Go P.M.N.Y.H., McCormack K., Ross S.J., Grant A.M. Open mesh versus non-mesh for groin hernia repair (Review), In: *The Cochrane Library* 2006, Issue 1; John Wiley & Sons Ltd. eds., Oxford, England
- 37 Colak T., Akca T., Kanik A., Aydin S. Randomized clinical trial comparing laparoscopic totally extraperitoneal approach with open mesh repair in inguinal hernia. *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques Tech* 2003; 13: 191-195
- 38 Coskun F., Ozmen M.M., Moran M., Ozozan O. New technique for inguinal hernia repair. *Hernia* 2005; 9: 32-36
- 39 Courtney C.A., Duffy K., Serpell M. G., O'Dwyer P. J. Outcome of patients with severe chronic pain following repair of groin hernia. *British Journal of Surgery* 2002; 89: 1310-1314

- 40 Czechowski A., Schafmayer A. TAPP versus TEP. Eine retrospektive Analyse fünf Jahre postoperative nach transperitonealer und endoskopisch-extraperitonealer Leisten-und Schenkelhernienversorgung. *Chirurg* 2003; 74: 1143-1148
- 41 Deysine M. Post mesh herniorrhapy infection control: Are we doing all we can?. *Hernia* 2004; 8: 90-91
- 42 Dieudonné G. Plug repair of groin hernias: a 10-year experience. *Hernia* 2002; 5: 189-191
- 43 Dittrick G.W., Ridl K., Kuhn J. A., McCarty T. M. Routine ilioinguinal nerve excision in inguinal hernia repairs. *American Journal of Surgery* 2004; 188: 736-740
- 44 DiVita G., Milano S., Frazzetta M. et al. Tension-free hernia repair is associated with an increase in inflammatory response markers against the mesh. *American Journal of Surgery* 2000; 180: 203-207
- 45 Douek M., Smith G., Oshowo A., Stoker D.L., Wellwood J.M. Prospective randomised controlled trial of laparoscopic versus open inguinal hernia mesh repair: five year follow up. *British Medical Journal* 2003; 326: 1012-1013
- 46 EU Hernia Trialists Collaboration: Repair of groin hernia with synthetic mesh: meta-analysis of randomised controlled trials. *Annals of Surgery* 2002; 235: 322-332
- 47 EU Hernia Trialists Collaboration: Mesh compared with non-mesh methods of open groin hernia repair: systematic review of randomised controlled trials. *British Journal of Surgery* 2000; 87: 854-859
- 48 EU Hernia Trialists Collaboration: Laparoscopic compared with open methods of groin hernia repair: systematic review of randomised controlled trials. *British Journal of Surgery* 2000; 87: 860-867
- 49 Fasih T., Mahapatra T.K., Waddington R.T. Early results of inguinal hernia repair by the 'mesh plug' technique-first 200 cases. *Annals of Royal College of Surgeons England* 2000; 82: 396-400
- 50 Feliu X., Jaurrieta E., Vinas X., Macarulla E., Abad J.M., Fernandez-Sallent E. Recurrent inguinal hernia: a ten-year review. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2004; 14(6): 362-367
- 51 Feliu X., Torres G., Martinez-Rodenas F., Fernandez-Sallent E., Pie J. Preperitoneal repair for recurrent inguinal hernia: laparoscopic and open approach. *Hernia* 2003; (Accessed May 22, 2003, at <http://springerlink.metapress.com/media/9CX>)
- 52 Fingerhut A., Millat B., Bataille N. et al. Laparoscopic hernia repair in 2000: Update of the European Association for Endoscopic Surgery (E.A.E.S.) Consensus Conference in Madrid, June 1994. *Surgical Endoscopy* 2000; 15: 1061-1065
- 53 Fischer S., Troidl H., MacLean A.A., Koehler L., Paul A. Prospective double-blind randomised study of a new regimen of pre-emptive analgesia for inguinal hernia repair:

evaluation of postoperative pain course. *European Journal of Surgery* 2000; 166: 545-551

54 Fleming W.R., Elliott T.B., Jones R.McL., Hardy K.J. Randomized clinical trial comparing totally extraperitoneal inguinal hernia repair with the Shouldice technique. *British Journal of Surgery* 2001; 88: 1183-1188

55 Forte A., D'Urso A., Palumbo P. et al. Inguinal hernioplasty: the gold standard of hernia repair. *Hernia* 2003; 7: 35-38

56 Friemert B., Faoual J., Hölldobler G., Becker H.P., Lampl L., Gerngroß H. Eine prospektiv randomisierte Studie zur Leistenhernienoperation nach Shouldice. *Chirurg* 2000; 71: 52-57

57 Gerber S., Hämmerli P.A., Glättli A. Laparoskopische transabdominale präperitoneale Hernienplastik. Evaluation der zugangsbedingten Komplikationen. *Chirurg* 2000; 71: 824-828

58 Ghandimi B.M., Langer C., Becker H. Zum kanzerogenen Potential von Biomaterialien in der Hernienchirurgie. *Chirurg* 2002; 73: 833-838

59 Gianetta E., Cuneo S., Vitale B., Camerini G., Marini P., Stella M. Anterior tension-free repair of recurrent inguinal hernia under local anesthesia. *Annals of Surgery* 2001; 231: 132-136

60 Gönüllü N.N., Cubucu A., Alponat A. Comparison of local and general anesthesia in tension-free (Lichtenstein) hernioplasty: a prospective randomized trial. *Hernia* 2002; 6: 29-32

61 Goldstein H.S., Rabaza J.R., Gonzalez A.M., Verdeja J.C. Evaluation of pain and disability in plug repair with the aid of a personal digital assistant. *Hernia* 2003; 7: 25-28

62 Goldstein H.S. A university experience using mesh in inguinal hernia repair. *Hernia* 2002; 5: 182-185

63 Goyal S., Abbasakoor F., Stephenson B.M. Experience with the preperitoneal "plug and patch" inguinal hernia repair. *British Journal of Surgery* 1999; 86: 1284-1285

64 Grant A.M., Scott N.W., O'Dwyer P.J. Five-year follow-up of a randomized trial to assess pain and numbness after laparoscopic or open repair of groin hernia. *British Journal of Surgery* 2004; 91: 1570-1574

65 Grant A.M. Laparoscopic versus open groin hernia repair: meta-analysis of randomised trials based on individual patient data. *Hernia* 2002; 6: 2-10

66 Grant A.M. Open mesh versus non-mesh repair of groin hernia meta-analysis of randomized trials based on individual patient data. *Hernia* 2002; 6: 130-136

- 67 Haapaniemi S., Nilsson E. Recurrence and pain three years after groin hernia repair. Validation of postal questionnaire and selective physical examination as a method of follow-up. *European Journal of Surgery* 2002; 168: 22-28
- 68 Haapaniemi S., Gunnarsson U., Nordin P., Nilsson E. Reoperation after recurrent groin hernia repair. *Annals of Surgery* 2001; 234: 122-126
- 69 Hair A., Duffy K., McLean J. et al. Groin hernia repair in Scotland. *British Journal of Surgery* 2000; 87: 1722-1726
- 70 Hildebrandt J., Levantin O. Spannungsfreie Methoden der Operation primärer Leistenbrüche. Vergleich der endoskopischen, total extraperitonealen Hernioplastik mit der Operation nach Lichtenstein. *Chirurg* 2003; 74: 915-921
- 71 Hindmarsh A.C. , Cheong E., Lewis M. P. N., Rhodes M. Attendance at a pain clinic with severe chronic pain after open and laparoscopic hernia repairs. *British Journal of Surgery* 2003; 90: 1152-1154
- 72 Huang C.S., Huang C.C., Lien H.H. Prolene hernia system compared with mesh plug technique: a prospective study of short- to mid-term outcomes in primary groin hernia repair. *Hernia* 2005;9:167-171
- 73 Ismail W., Agrawal A., Zia M.I. Fate of chronically infected onlay mesh in groin wound. *Hernia* 2002; 6: 79-81
- 74 Johansson B., Hallerbäck B., Glise H., Anesten B., Smedberg S., Román J. Laparoscopic mesh versus open preperitoneal mesh versus conventional technique for inguinal hernia repair: a randomized multicenter trial (SCUR Hernia Repair Study). *Annals of Surgery* 1999; 230: 225-231
- 75 Jones K.R., Burney R. E., Peterson M., Christy B. Return to work after inguinal hernia repair. *Surgery* 2001; 129:128
- 76 Junge K., Peiper C., Rosch R., Lynen P., Schumpelick V. Effect of tension induced by Shouldice repair on post-operative course and long-term outcome. *European Journal of Surgery* 2002; 168: 329-333
- 77 Kapisir S.A., Brough W.A., Royston C.M.S., O'Boyle C., Sedman P.C. Laparoscopic transabdominal preperitoneal (TAPP) hernia repair. A 7-year two-center experience in 3017 patients. *Surgical Endoscopy* 2001; 15: 972-975
- 78 Kark A.E., Kurzer M.N., Belsham P.A. Three thousand one hundred seventy-five primary inguinal hernia repairs: advantages of ambulatory open mesh repair using local anesthesia. *Journal American College of Surgeons* 1998; 86: 447-456
- 79 Kehlet H., Bay-Nielsen M., Kingsnorth A. Chronic postherniorrhapy pain – a call for uniform assessment. *Hernia* 2002; 6: 178-181
- 80 Kingsnorth A.N. Treating inguinal hernias. Open mesh Lichtenstein operation is preferred over laparoscopy. *British Medical Journal* 2004; 328: 59-60

- 81 Kingsnorth A.N., Bowley D.M.G., Porter C. A prospective study of 1000 hernias: results of the Plymouth Hernia Service. *Annals of the Royal College of Surgeons of England* 2003; 85: 18-22
- 82 Kingsnorth A.N., LeBlanc K.A. In: *Management of abdominal hernias*. Edward Arnold Verlag; 3. Auflage, 2003: 40-47
- 83 Kingsnorth A.N., Wright D., Porter C. S., Robertson G. Prolene hernia system compared with Lichtenstein patch: a randomised double-blind study of short-term and medium-term outcomes in primary inguinal hernia repair. *Hernia* 2002; 6: 113-119
- 84 Kingsnorth A.N., Hyland M.E., Porter C. A., Sodergren S. Prospective double-blind randomized study comparing Perfix plug and patch with Lichtenstein patch in inguinal hernia repair: one year quality of life. *Hernia* 2000; 4: 255-258
- 85 Kingsnorth A.N., Porter C.S., Bennett D.H., Walker A.J., Hyland M.E., Sodergren S. Lichtenstein patch or Perfix plug-and-patch in inguinal hernia: a prospective double-blind randomized controlled trial of short term outcome. *Surgery* 2000; 127: 276-283
- 86 Klinge U., Junge K., Strumpf M., Öttinger A.P., Klosterhalfen B. Functional and morphological evaluation of a low-weight, monofilament polypropylene mesh for hernia repair. *Applied Biomaterials* 2002; 63: 129-136
- 87 Koch A., Edwards A., Haapaniemi S., Nordin P., Kald A. Prospective evaluation of 6895 groin hernia repairs in women. *British Journal of Surgery* 2005; 92:1553-1558
- 88 Köninger J., J. Redecke, M. Butters Chronic pain after hernia repair: a randomized trial comparing Shouldice, Lichtenstein and TAPP. *Langenbecks Archives of Surgery* 2004; 389: 361-365
- 89 Koukourou A., Lyon W., Rice J., Wattchow D.A. Prospective randomised trial of polypropylene mesh compared with nylon darn in inguinal hernia repair *British Journal of Surgery* 2001, 88, 931-934
- 90 Kumar S., Wilson R. G., Nixon S. J., Macintyre I. M. C. Chronic pain after laparoscopic and open mesh repair of groin hernia. *British Journal of Surgery* 2002; 89: 1476-1479
- 91 Kurzer M., Belsham P.A., Kark A.E. Prospective study of open preperitoneal mesh repair for recurrent inguinal hernia. *British Journal of Surgery* 2002; 89: 90-93
- 92 Lau H., Patil N. G., Yuen W. K., Lee F. Prevalence and severity of chronic groin pain after endoscopic extraperitoneal hernioplasty. *Surgical Endoscopy* 2003; 17: 1620-1623
- 93 Lau H., Lee F. Determinant factors of pain after ambulatory inguinal herniorrhaphy: a multi-variate analysis. *Hernia* 2001; 5: 17-20
- 94 LeBlanc K.A. Complications associated with the plug-and-patch method of inguinal herniorrhaphy. *Hernia* 2001; 5: 135-138

- 95 Leibl B.J., Bittner R. Alloplastische Materialien in der Hernienchirurgie. *Viszeralchirurgie* 2004; 39: 393-397
- 96 Leibl B.J., Schmedt C. G., Ulrich M., Kraft K., Bittner R. Recurrence after endoscopic transperitoneal hernia repair (TAPP): causes, reparative techniques, and results of reoperation. *Journal of American College of Surgeons* 2000; 190: 651-655
- 97 Leibl B.J., Schmedt C. G., Ulrich M., Kraft K., Bittner R. Laparoskopische Leistenhernientherapie (TAPP) als Ausbildungsoperation. *Chirurg* 2000; 71(8): 939-942
- 98 Leibl B.J., Däubler P., Schmedt C. G., Kraft K., Bittner R. Long-term results of a randomized clinical trial between laparoscopic hernioplasty and Shouldice repair. *British Journal of Surgery* 2000; 87: 780-783
- 99 Liem M.S., van Duyn E. B., van der Graaf Y., van Vroonhoven T. J. M. V. Recurrences after conventional anterior and laparoscopic inguinal hernia repair: a randomized comparison. *Annals of Surgery* 2003; 237: 136-141
- 100 Lorenz D., Stark E., Oestreich K., Richter A. Laparoscopic hernioplasty versus conventional hernioplasty (Shouldice): results of a randomized trial. *World Journal of Surgery* 2000; 24: 739-746
- 101 Macintyre I.M.C. Inguinal hernia repair – How I do it. *Journal of Royal College of Surgeons Edinburgh* 2001; 46: 349-353
- 102 Mahmud S. Randomized clinical trial comparing 5-year recurrence rate after laparoscopic versus Shouldice repair of primary inguinal hernia. *British Journal of Surgery (Letter)* 2006; 93(1):122-3
- 103 Mahon D., Decadt B., Rhodes M. Prospective randomized trial of laparoscopic (transabdominal preperitoneal) versus open (mesh) repair for bilateral and recurrent inguinal hernia. *Surgical Endoscopy* 2003; 17: 1386-1390
- 104 Mayagoitia J.C. Inguinal hernioplasty with the Prolene Hernia System. *Hernia* 2004; 8: 64-66
- 105 McCloud J.M., Evans D.S. Day-case laparoscopic hernia repair in a single unit. *Surgical Endoscopy* 2003; 17: 491-493
- 106 McCormack K., Wake B., Perez J. et al. Laparoscopic surgery for inguinal hernia repair: systematic review of effectiveness and economic evaluation. *Health Technology Assessment* 2005; 9: 14
- 107 McCormack K., Grant A., Scott N. for the EU Hernia Trialists Collaboration: Value of updating a systemic review in surgery using individual patient data. *British Journal of Surgery* 2004; 91: 495-499
- 108 Medical Research Council Laparoscopic Groin Hernia Trial Group Cost-utility analysis of open versus laparoscopic groin hernia repair: results from a multi-centre randomized clinical trial. *British Journal of Surgery* 2001; 88: 653-661

- 109 Melis P., van der Drift D.G.F., Sybrandy R., Go P.M.N.Y.H. High recurrence rate 12 years after primary inguinal hernia repair. *European Journal of Surgery* 2000; 166: 313-314
- 110 Memon M.A., Cooper N.J., Memon B., Memon M.I., Abrams K.R. Meta-analysis of randomized clinical trials comparing open and laparoscopic inguinal hernia repair. *British Journal of Surgery* 2003; 90: 1479-1492
- 111 Millikan K.W., Cummings B., Doolas A. A prospective study of the mesh-plug hernioplasty. *The American Surgeon* 2001; 67(3): 285-289
- 112 Miedema B.W., Ibrahim S.M., Davis B.D., Koivunen D.G. A prospective trial of primary inguinal hernia repair by surgical trainees. *Hernia* 2004;8:28-32
- 113 Müller J.M. (Hrsg.) *Manual der Klinik für Allgemein-, Viszeral-, Gefäß-, Thoraxchirurgie Universitätsklinikums Charité*, Blackwell Wissenschaftsverlag GmbH Berlin 1997
- 114 Muldoon R.L., Marchant K., Johnson D.D., Yoder G.G., Read R.C., Hauer-Jensen M. Lichtenstein versus anterior preperitoneal prosthetic mesh placement in open inguinal hernia repair: a prospective randomized trial. *Hernia* 2004; 8: 98-103
- 115 Murata N., Ishida H., Makita Y. et al. Muscle strength and walking ability after laparoscopic hernia repair versus conventional repair. *Surgery Today* 2003; 33: 259-263
- 116 National Health Service National Institute for Clinical Excellence (NICE) Guidance on the use of laparoscopic surgery for inguinal hernia. In: *Technology Appraisal Guidance No.18*, 2001: 1-103 (Accessed October 22, 2006, at www.nice.org.uk/page.aspx?o=20663)
- 117 Neumayer L., Giobbie-Hurder A., Jonasson O. et al. Open mesh versus laparoscopic mesh repair of inguinal hernia. *New England Journal of Medicine* 2004; 350: 1819-1827
- 118 Nienhuijs S.W., Boelens O.B.A., Strobbe L.J.A. Pain after anterior mesh repair. *Journal of American College of Surgeons* 2005; 200: 885-889
- 119 Nienhuijs S.W., van Oort I., Keemers-Gels M.E., Strobbe L.J.A., Rosmann C. Randomized trial comparing the Prolene® hernia system, mesh plug repair and Lichtenstein method for open inguinal hernia repair. *British Journal of Surgery* 2005; 92: 33-38
- 120 Nordin P., Haapaniemi S., van Der Linden W., Nilsson E. Choice of anesthesia and risk of reoperation for recurrence in groin hernia repair. *Annals of Surgery* 2004; 240: 187-192
- 121 Nordin P., Bartelmess P., Jansson C., Svensson C., Edmund G. Randomized trial of Lichtenstein versus Shouldice hernia repair in general surgical practice. *British Journal of Surgery* 2002; 89: 45-49

- 122 Nyhus L.M. Evolution of hernia repair: a salute to Professor Piero Pietri. *Hernia* 2002; 5: 196-199
- 123 Nyhus L.M. Ubiquitous use of prosthetic mesh in inguinal hernia repair: the dilemma. *Hernia* 2000; 4: 184-186
- 124 O'Dwyer P. J., Kingsnorth A. N., Molloy R. G., Small P. K., Lammers B., Horeysek G. Randomized clinical trial assessing impact of a lightweight or heavyweight mesh on chronic pain after inguinal hernia repair. *British Journal of Surgery* 2005; 92: 166-170
- 125 Page B., Paterson C., Young D., O'Dwyer P.J. Pain from primary inguinal hernia and the effect of the repair on pain. *British Journal of Surgery* 2002; 89: 1315-1318
- 126 Paajanen H. Do absorbable mesh sutures cause less chronic pain than nonabsorbable sutures after Lichtenstein inguinal herniorrhaphy?. *Hernia* 2002; 6: 26-28
- 127 Pavlidis T.E. Comparison between modern mesh and conventional mesh methods of inguinal hernia repair. *Minerva Chirurgica* 2002; 57: 7-12
- 128 Pawanindra L., Kajla R.K., Chander J., Saha R., Ramteke V.K. Randomized controlled study of laparoscopic total extraperitoneal versus open Lichtenstein hernia repair. *Surgical Endoscopy* 2003; 17: 850-856
- 129 Peiper C. Netze in der Leistenhernienchirurgie. *Zentralblatt Chirurgie* 2002; 127: 573-577
- 130 Pélissier E. P. Inguinal hernia: the size of the mesh. *Hernia* 2000; 5: 169-171
- 131 Perkins F.M., Kehlet H. Chronic pain as an outcome of surgery: a review of predictive factors. *Anesthesiology* 2004; 93: 1123-1133
- 132 Poobalan A. S., Bruce J., Cairns W. et al. A review of chronic pain after inguinal herniorrhaphy. *Clinical Journal of Pain* 2003; 19: 48-54
- 133 Poobalan A. S., Bruce J., King P.M., Chambers W.A., Krukowski Z.H., Smith W.C.S. Chronic pain and quality of life following open inguinal hernia repair. *British Journal of Surgery* 2001; 88: 1122-1126
- 134 Post S., Weiss B., Willer M., Neufang T., Lorenz D. Randomized clinical trial of lightweight composite mesh for Lichtenstein inguinal hernia repair. *British Journal of Surgery* 2004; 91: 44-48
- 135 Ramanand B., Fayaz M., Patil P. Experience with the preperitoneal 'plug and patch' inguinal hernia repair *British Journal of Surgery (Letter)* 2000; 87:517
- 136 Ramshaw B., Abiad F., Voeller G., Wilson R., Mason E. Polyester (Parietex) mesh for total extraperitoneal laparoscopic inguinal hernia repair. *Surgical Endoscopy* 2003; 17: 498-501

- 137 Ravichandran D., Kalambe B.G., Pain J.A. Pilot randomized controlled study of preservation or division of ilioinguinal nerve in open mesh repair of inguinal hernia. *British Journal of Surgery* 2000; 87: 1166-1167
- 138 Read R.C. British contributions to modern herniology of the groin. *Hernia* 2005; 9: 6-11
- 139 Richards S.K., Vipond M.N., Earnshaw J.J. Review of the management of recurrent inguinal hernia. *Hernia* 2003; (Accessed April 7, 2006, at <http://springerlink.metapress.com/media03X1MB085LaRXN9E>)
- 140 Richards S.K., Earnshaw J.J. Management of primary and recurrent inguinal hernia by surgeons from the South West of England. *Annals of the Royal College of Surgery England* 2003; 85: 402-404
- 141 Ridings P., Evans D.S. The transabdominal pre-peritoneal (TAPP) inguinal hernia repair: a trip along the learning curve. *Journal of Royal College of Surgeons Edinburgh* 2000; 45: 29-32
- 142 Robbins A.W., Rutkow I.M. Mesh plug repair and groin hernia surgery. *Surgical Clinics of North America* 1998; 78(6): 1007-1023
- 143 Rutkow I.M. Demographic and socioeconomic aspects of hernia repair in the United States in 2003. *Surgical Clinics of North America* 2003; 83:1045-1051
- 144 Sandblom G., Gruber G., Kald A., Nilsson E. Audit and recurrence rates after hernia surgery. *European Journal of Surgery* 2000; 166: 154-158
- 145 Sanchez-Manuel F.J., Seco-Gil J.L. Antibiotic prophylaxis for hernia repair (Cochrane review) In: *The Cochrane Library, Issue 2 Oxford, England* 2003
- 146 Santoro E., Agresta F., Aloisi P. et al. Is minilaparoscopic inguinal hernia repair feasible? A preliminary experience. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2005; 15(3): 294-297
- 147 Sarli L., Villa F., Marchesi F. Simultaneous repair of bilateral inguinal hernias: a prospective randomized study of open, tension-free versus laparoscopic approach. *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques* 2001; 11: 262-267
- 148 Schardey H.M., Schopf S., Rudert W., Knappich P., Hernandez-Richter T. Titanisiertes Polypropylenetz: Erste klinische Beobachtungen nach Implantation in TAPP-Technik und Ergebnisse einer Umfrage an 22 Chirurgischen Kliniken. *Zentralblatt Chirurgie* 2004; 129: 363-368
- 149 Scheyer M., S. Arnold, G. Zimmermann Minimally invasive operation techniques for inguinal hernia: spectrum of indications in Austria. *Hernia* 2001; 5: 73-79
- 150 Schmedt C.G., Sauerland S., Bittner R. Comparison of endoscopic procedures versus Lichtenstein and other open mesh techniques for inguinal hernia repair: a meta-analysis of randomized controlled trials. *Surgical Endoscopy* 2005; 19: 188-199

- 151 Schmedt C.G., Leibl B.J., Bittner R. Endoscopic inguinal hernia repair in comparison with Shouldice and Lichtenstein repair: a systematic review of randomized trials. *Dig Surg* 2002; 19: 511-517
- 152 Schmedt C.G., Leibl B.J., Bittner R. Zugangsbedingte Komplikationen in der laparoskopischen Chirurgie. *Chirurg* 2002; 73: 863-879
- 153 Schmedt C.G., Däubler P., Leibl B.J., Kraft K., Bittner R. Simultaneous bilateral laparoscopic inguinal hernia repair. *Surgical Endoscopy* 2002; 16: 240-244
- 154 Schultz C., Baca I., Götzen V. Laparoscopic inguinal hernia repair. A review of 2500 cases. *Surgical Endoscopy* 2001; 15: 582-584
- 155 Schumpelick V., Klinge U., Schwab R. Stellenwert verschiedener Operationsverfahren in der Versorgung des Leistenbruches. *Viszeralchirurgie* 2004; 39: 13-19
- 156 Schumpelick V. Does every hernia demand a mesh repair? A critical review. *Hernia* 2001; 5: 5-8
- 157 Schumpelick V. In: *Hernien*; 4. Auflage Thieme Verlag 2000: 394-416
- 158 Silen W. Chronic pain and quality of life following open inguinal hernia repair. *British Journal of Surgery (Letter)* 2002; 89: 123
- 159 Singhal T., Balakrishnan S., Paix A., El-Hasani S. Early experience with laparoscopic inguinal hernia repair in a district General National Health Services Hospital. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2005; 15(3): 285-289
- 160 Smith J., Demers M.L., Pollack R., Gregory S. Prospective comparison between laparoscopic preperitoneal herniorraphy and open mesh herniorraphy. *American Surgeon* 2001; 32: 115-118
- 161 Sondenaa K., Nesvik I., Breivik K., Korner H. Long-term follow-up of 1059 consecutive primary and recurrent inguinal hernias in a teaching hospital. *European Journal of Surgery* 2001; 167: 125-129
- 162 Stephenson B.M. Complication of groin hernia repairs. *Surgical Clinics of North America* 2003; 83: 1255-1278
- 163 Stoppa R. About biomaterials and how they work in groin hernia repairs. *Hernia* 2003; 7: 57-60
- 164 Stylopoulos N., Gazelle G.S., Rattner D.W. A cost-utility analysis of treatment options for inguinal hernia in 1,513,008 adult patients. *Surgical Endoscopy* 2003; 17: 180-189
- 165 Susmallian S., Gewurtz G., Ezri T., Charuzi I. Seroma after laparoscopic repair of hernia with PTFE patch: is it really a complication?. *Hernia* 2001; 5: 139-141

- 166 Taylor S.G., O'Dwyer P.J. Chronic groin sepsis following tension-free inguinal hernioplasty. *British Journal of Surgery* 1999; 86: 562-565
- 167 Tschudi J.F., Wagner M., Klaiber C. et al. Randomized controlled trial of laparoscopic transabdominal preperitoneal hernioplasty versus Shouldice repair. *Surgical Endoscopy* 2001; 15: 1263-1266
- 168 Vale L., Ludbrook A., Grant A. Assessing the cost and consequences of laparoscopic versus open methods of groin hernia repair. *Surgical Endoscopy* 2003; 17: 844-849
- 169 Vatansev C., Belviranli M., Belviranli M. et al. The effect of different hernia repair methods on post-operative pain medication and CRP levels. *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques* 2002; 12: 243-246
- 170 Voyles C.R. Outcome analysis for groin hernia repairs. *Surgical Clinics of North America* 2003; 83: 1279-1287
- 171 Voyles C.R., Hamilton B. J., Johnson W. D., Kano N. Meta-analysis of laparoscopic inguinal hernia trials favors open hernia repair with preperitoneal mesh prosthesis. *American Journal of Surgery* 2002; 184: 6-10
- 172 Vrijland W., van den Tol M.P., Luijendijk R.W. et al. Randomized clinical trial of non-mesh versus mesh repair of primary inguinal hernia. *British Journal of Surgery* 2002; 89: 293-297
- 173 Wara P., Bay-Nielsen M., Juul P., Bendix J., Kehlet H. Prospective nationwide analysis of laparoscopic versus Lichtenstein repair of inguinal hernia. *British Journal of Surgery* 2005 (Accessed October 9, 2005, at <http://springerlink.metapress.com/mediaX02768542052>)
- 174 Wennström I., Berggren P., Akerud L., Järhult J. Equal results with laparoscopic and Shouldice repairs of primary inguinal hernia in men. Report from a prospective randomised study. *Scandinavian Journal of Surgery* 2004; 93: 34-36
- 175 Winslow E.R., Quasebarth M., Brunt L.M. Perioperative outcomes and complications of open versus laparoscopic extraperitoneal inguinal hernia repair in a mature surgical practice. *Surgical Endoscopy* 2003 (Accessed January 21, 2004, at www.springerlink.com/media/g19150459j7wnhc8te27)
- 176 Woolf C., Max M.B. Mechanism-based pain diagnosis. Issues for analgesic drug development. *Anesthesiology* 2003; 95: 241-249
- 177 Wright D., Paterson C., Scott N., Hair A., O'Dwyer P. J. Five-year follow-up of patients undergoing laparoscopic or open groin hernia repair: a randomized controlled trial. *Annals of Surgery* 2002; 235: 333-337
- 178 Yerdel M.A., Akin E.B., Dolalan S. et al. Effect of single-dose prophylactic Ampicillin and Sulbactam on wound infection after tension-free hernia repair with polypropylene mesh. *Annals of Surgery* 2001; 233:26-33

- 179 Zieren J. Inguinal hernia: obligatory indication for elective surgery? A prospective assessment of quality of life before and after plug and patch inguinal hernia repair. *Langenbecks Archives of Surgery* 2003; 387: 417-420
- 180 Zieren J., Neuss H., Philipp A. W., Müller J. M. Postoperative comfort after plug-and-patch repair of recurrent inguinal hernia. *European Journal of Surgery* 2002; 168: 18-21
- 181 Zieren J., Beyersdorff D., Beier K. M., Müller J. M. Sexual function and testicular perfusion after inguinal hernia repair with mesh. *American Journal of Surgery* 2001; 181: 204-206
- 182 Zieren J., Zieren H.U., Wenger F., Müller J.M. Leistenhernienreparation beim alten Menschen – Ergebnisse der Plug-und-Patch-Reparation unter besonderer Berücksichtigung der Lebensqualität. *Chirurg* 2000; 71: 564-567
- 183 Zieren J., Zieren H.U., Jacobi C.A., Wenger F., Müller J.M. Prospective randomized study comparing laparoscopic and tension-free inguinal hernia repair with Shouldice's operation. *American Journal of Surgery* 1998; 175: 330-333