

# Contents

<b>Zusammenfassung</b>	<b>5</b>
<b>Abstract</b>	<b>7</b>
<b>1 Introduction</b>	<b>11</b>
<b>2 Theory</b>	<b>19</b>
2.1 Optimal control of quantum systems over finite time interval	19
2.1.1 The Lagrangian formalism . . . . .	19
2.1.2 Derivation of the differential equation for the optimal control field . . . . .	22
2.1.3 An approximate analytical solution for the case of a two level system . . . . .	24
2.1.4 The analytical solution for the optimal control field with a simplified Lagrangian . . . . .	25
2.1.5 Optimal control at a given time . . . . .	27
2.1.6 Estimation of the absolute bound for the control due to relaxation and dephasing effects . . . . .	28
2.2 Calculation of the excitation spectrum of quantum systems using genetic algorithms . . . . .	30
2.2.1 Introduction to evolutionary algorithms . . . . .	30
2.2.2 Definition of the representation, fitness and genetic operators for quantum ground-state problem . . . . .	33
2.2.3 Extension of the QGA for the solution of quantum statistical problems . . . . .	38
2.3 Photon assisted tunneling between quantum dots: optimal control approach . . . . .	41
2.3.1 Equations of motion in terms of density matrix . . . . .	41
2.3.2 The control problem . . . . .	43
2.4 Explosion of noble gas clusters in strong laser fields . . . . .	46
2.4.1 Time and length scales for the cluster explosion . . . . .	46
2.4.2 Explosion scenario . . . . .	48
2.4.3 Equations of motion . . . . .	50

2.4.4	Integration of the equations of motion: the split operator method . . . . .	52
<b>3</b>	<b>Results</b>	<b>55</b>
3.1	Optimal control of a time averaged occupation of the excited level in a two-level system . . . . .	56
3.1.1	The influence of relaxation on the optimal control . . . . .	56
3.1.2	Optimal control field calculated using a simplified Lagrangian . . . . .	60
3.2	Application of the QGA to the eigenstate problem for interacting electrons in quantum dots . . . . .	63
3.2.1	The ground state problem in one and two dimensions . . . . .	63
3.2.2	Formation of a “Wigner molecule” and its “melting” . . . . .	70
3.3	Optimal field for control of the photon assisted tunneling between quantum dots . . . . .	75
3.3.1	Numerical solution using the genetic algorithm: the influence of relaxation on the optimal control . . . . .	75
3.4	Explosion of $Xe_N$ clusters . . . . .	83
3.4.1	Ionization and explosion dynamics . . . . .	83
3.4.2	Dependence of the kinetic energy distribution on the cluster size . . . . .	85
3.4.3	Isotropic inhomogeneous 3D Coulomb model . . . . .	89
<b>4</b>	<b>Conclusions and outlook</b>	<b>91</b>
<b>A</b>	<b>The adiabatic approximation</b>	<b>97</b>
<b>B</b>	<b>Derivation of an approximate solution for a two level system</b>	<b>99</b>
<b>C</b>	<b>Generalization of the QGA for higher dimensions</b>	<b>101</b>
<b>D</b>	<b>Integration of the pendulum equation</b>	<b>103</b>
	<b>Bibliography</b>	<b>105</b>
	<b>Publications</b>	<b>117</b>
	<b>Acknowledgments</b>	<b>121</b>
	<b>Curriculum vitae</b>	<b>123</b>