

## 6. References

- 1 Albers B, Bray D, Lewis J, et al. Molekularbiologie der Zelle. 3. Auflage. Weinheim: VCH Verlagsgesellschaft, 1997
- 2 Gilbert S Fertilization: Beginning a New Organism. In: Developmental biology 7th edition ed. Sunderland MA: Sinauer Associates; 2003
- 3 Weissman IL, Anderson DJ, Gage F. Stem and progenitor cells: origins, phenotypes, lineage commitments, and transdifferentiations. Annu Rev Cell Dev Biol 2001; 17:387-403
- 4 Watt FM, Hogan BL. Out of Eden: stem cells and their niches. Science 2000; 287:1427-1430
- 5 Blau HM, Brazelton TR, Weimann JM. The evolving concept of a stem cell: entity or function? Cell 2001; 105:829-841
- 6 Stem Cells: Scientific Progress and Future Research Directions. Department of Health and Human Services. June 2001. <http://stemcells.nih.gov/info/scireport>
- 7 Weissman IL. Translating stem and progenitor cell biology to the clinic: barriers and opportunities. Science 2000; 287:1442-1446
- 8 Weissman IL. Stem cells: units of development, units of regeneration, and units in evolution. Cell 2000; 100:157-168
- 9 Toma JG, Akhavan M, Fernandes KJ, et al. Isolation of multipotent adult stem cells from the dermis of mammalian skin. Nat Cell Biol 2001; 3:778-784
- 10 Moore KA, Ema H, Lemischka IR. In vitro maintenance of highly purified, transplantable hematopoietic stem cells. Blood 1997; 89:4337-4347
- 11 Glimm H, Eaves CJ. Direct evidence for multiple self-renewal divisions of human in vivo repopulating hematopoietic cells in short-term culture. Blood 1999; 94:2161-2168
- 12 Pagano SF, Impagnatiello F, Girelli M, et al. Isolation and characterization of neural stem cells from the adult human olfactory bulb. Stem Cells 2000; 18:295-300

- 13 Smith JR, Pochampally R, Perry A, et al. Isolation of a highly clonogenic and multipotential subfraction of adult stem cells from bone marrow stroma. *Stem Cells* 2004; 22:823-831
- 14 Morszeck C, Gotz W, Schierholz J, et al. Isolation of precursor cells (PCs) from human dental follicle of wisdom teeth. *Matrix Biol* 2005; 24:155-165
- 15 Fu SL, Ma ZW, Yin L, et al. [Isolation and cultivation of neural stem cells from the embryonic rat brain and spinal cord]. *Sheng Li Xue Bao* 2003; 55:278-283
- 16 Alessandri G, Pagano S, Bez A, et al. Isolation and culture of human muscle-derived stem cells able to differentiate into myogenic and neurogenic cell lineages. *Lancet* 2004; 364:1872-1883
- 17 Papini S, Cecchetti D, Campani D, et al. Isolation and clonal analysis of human epidermal keratinocyte stem cells in long-term culture. *Stem Cells* 2003; 21:481-494
- 18 Dekaney CM, Rodriguez JM, Graul MC, et al. Isolation and characterization of a putative intestinal stem cell fraction from mouse jejunum. *Gastroenterology* 2005; 129:1567-1580
- 19 Chen Z, de Paiva CS, Luo L, et al. Characterization of putative stem cell phenotype in human limbal epithelia. *Stem Cells* 2004; 22:355-366
- 20 Coles BL, Angenieux B, Inoue T, et al. Facile isolation and the characterization of human retinal stem cells. *Proc Natl Acad Sci U S A* 2004; 101:15772-15777
- 21 Faris RA, Konkin T, Halpert G. Liver stem cells: a potential source of hepatocytes for the treatment of human liver disease. *Artif Organs* 2001; 25:513-521
- 22 Abraham EJ, Kodama S, Lin JC, et al. Human pancreatic islet-derived progenitor cell engraftment in immunocompetent mice. *Am J Pathol* 2004; 164:817-830
- 23 De Felice M, Di Lauro R. Thyroid development and its disorders: genetics and molecular mechanisms. *Endocr Rev* 2004; 25:722-746
- 24 Nussey SS, Whitehead SA, 2001 *Endocrinology: An integrated approach*. BIOS Scientific Publishers Ltd.

<http://www.ncbi.nlm.nih.gov/books/bv.fcgi?call=bv.View.ShowTOC&rid=endocrin.TOC&deph=1>

- 25 Appendix E: Markers Commonly Used to Identify Stem Cells and to Characterize Differentiated Cell Types. In: Stem Cells: Scientific Progress and Future Research Directions  
Department of Health and Human Services. June 2001 <http://stemcells.nih.gov/info/scireport>
- 26 Reis-Filho JS, Preto A, Soares P, et al. p63 expression in solid cell nests of the thyroid: further evidence for a stem cell origin. *Mod Pathol* 2003; 16:43-48
- 27 Preto A, Cameselle-Teijeiro J, Moldes-Boullosa J, et al. Telomerase expression and proliferative activity suggest a stem cell role for thyroid solid cell nests. *Mod Pathol* 2004; 17:819-826
- 28 Pesce M, Scholer HR. Oct-4: control of totipotency and germline determination. *Mol Reprod Dev* 2000; 55:452-457
- 29 Kehler J, Tolkunova E, Koschorz B, et al. Oct4 is required for primordial germ cell survival. *EMBO Rep* 2004; 5:1078-1083
- 30 Pesce M, Anastassiadis K, Scholer HR. Oct-4: lessons of totipotency from embryonic stem cells. *Cells Tissues Organs* 1999; 165:144-152
- 31 Pesce M, Scholer HR. Oct-4: gatekeeper in the beginnings of mammalian development. *Stem Cells* 2001; 19:271-278
- 32 Palmieri SL, Peter W, Hess H, et al. Oct-4 transcription factor is differentially expressed in the mouse embryo during establishment of the first two extraembryonic cell lineages involved in implantation. *Dev Biol* 1994; 166:259-267
- 33 Reim G, Brand M. Spiel-ohne-grenzen/pou2 mediates regional competence to respond to Fgf8 during zebrafish early neural development. *Development* 2002; 129:917-933
- 34 Nichols J, Zevnik B, Anastassiadis K, et al. Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 1998; 95:379-391

- 35 Niwa H, Miyazaki J, Smith AG. Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet* 2000; 24:372-376
- 36 Cheng L, Thomas A, Roth LM, et al. OCT4: a novel biomarker for dysgerminoma of the ovary. *Am J Surg Pathol* 2004; 28:1341-1346
- 37 Clark AT, Bodnar MS, Fox M, et al. Spontaneous differentiation of germ cells from human embryonic stem cells in vitro. *Hum Mol Genet* 2004; 13:727-739
- 38 Gidekel S, Pizov G, Bergman Y, et al. Oct-3/4 is a dose-dependent oncogenic fate determinant. *Cancer Cell* 2003; 4:361-370
- 39 Jin T, Branch DR, Zhang X, et al. Examination of POU homeobox gene expression in human breast cancer cells. *Int J Cancer* 1999; 81:104-112
- 40 Wang P, Branch DR, Bali M, et al. The POU homeodomain protein OCT3 as a potential transcriptional activator for fibroblast growth factor-4 (FGF-4) in human breast cancer cells. *Biochem J* 2003; 375:199-205
- 41 Steingart RA, Heldenberg E, Pinhasov A, et al. A vasoactive intestinal peptide receptor analog alters the expression of homeobox genes. *Life Sci* 2002; 71:2543-2552
- 42 Looijenga LH, Stoop H, de Leeuw HP, et al. POU5F1 (OCT3/4) identifies cells with pluripotent potential in human germ cell tumors. *Cancer Res* 2003; 63:2244-2250
- 43 Tai MH, Chang CC, Kiupel M, et al. Oct4 expression in adult human stem cells: evidence in support of the stem cell theory of carcinogenesis. *Carcinogenesis* 2005; 26:495-502
- 44 Ryffel GU. Mutations in the human genes encoding the transcription factors of the hepatocyte nuclear factor (HNF)1 and HNF4 families: functional and pathological consequences. *J Mol Endocrinol* 2001; 27:11-29
- 45 Sladek F, Seidel S Hepatocyte nuclear factor 4 alpha. *Nuclear Receptors and Genetic Disease*, San Diego: Academic Press, McCabe. TBE. 2001 309-361

- 46 Zhong W, Sladek FM, Darnell JE, Jr. The expression pattern of a Drosophila homolog to the mouse transcription factor HNF-4 suggests a determinative role in gut formation. *Embo J* 1993; 12:537-544
- 47 Holewa B, Zapp D, Drewes T, et al. HNF4beta, a new gene of the HNF4 family with distinct activation and expression profiles in oogenesis and embryogenesis of *Xenopus laevis*. *Mol Cell Biol* 1997; 17:687-694
- 48 Nakhei H, Lingott A, Lemm I, et al. An alternative splice variant of the tissue specific transcription factor HNF4alpha predominates in undifferentiated murine cell types. *Nucleic Acids Res* 1998; 26:497-504
- 49 Chen WS, Manova K, Weinstein DC, et al. Disruption of the HNF-4 gene, expressed in visceral endoderm, leads to cell death in embryonic ectoderm and impaired gastrulation of mouse embryos. *Genes Dev* 1994; 8:2466-2477
- 50 Duncan SA, Manova K, Chen WS, et al. Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. *Proc Natl Acad Sci U S A* 1994; 91:7598-7602
- 51 Levinson-Dushnik M, Benvenisty N. Involvement of hepatocyte nuclear factor 3 in endoderm differentiation of embryonic stem cells. *Mol Cell Biol* 1997; 17:3817-3822
- 52 Sladek FM. Orphan receptor HNF-4 and liver-specific gene expression. *Receptor* 1993; 3:223-232
- 53 LaVoie HA. The role of GATA in mammalian reproduction. *Exp Biol Med (Maywood)* 2003; 228:1282-1290
- 54 Molkenin JD. The zinc finger-containing transcription factors GATA-4, -5, and -6. Ubiquitously expressed regulators of tissue-specific gene expression. *J Biol Chem* 2000; 275:38949-38952
- 55 Laitinen MP, Anttonen M, Ketola I, et al. Transcription factors GATA-4 and GATA-6 and a GATA family cofactor, FOG-2, are expressed in human ovary and sex cord-derived ovarian tumors. *J Clin Endocrinol Metab* 2000; 85:3476-3483

- 56 Arceci RJ, King AA, Simon MC, et al. Mouse GATA-4: a retinoic acid-inducible GATA-binding transcription factor expressed in endodermally derived tissues and heart. *Mol Cell Biol* 1993; 13:2235-2246
- 57 Kuo CT, Morrisey EE, Anandappa R, et al. GATA4 transcription factor is required for ventral morphogenesis and heart tube formation. *Genes Dev* 1997; 11:1048-1060
- 58 Molkenin JD, Lin Q, Duncan SA, et al. Requirement of the transcription factor GATA4 for heart tube formation and ventral morphogenesis. *Genes Dev* 1997; 11:1061-1072
- 59 Kiiveri S, Liu J, Westerholm-Ormio M, et al. Differential expression of GATA-4 and GATA-6 in fetal and adult mouse and human adrenal tissue. *Endocrinology* 2002; 143:3136-3143
- 60 Ketola I, Pentikainen V, Vaskivuo T, et al. Expression of transcription factor GATA-4 during human testicular development and disease. *J Clin Endocrinol Metab* 2000; 85:3925-3931
- 61 Heikinheimo M, Ermolaeva M, Bielinska M, et al. Expression and hormonal regulation of transcription factors GATA-4 and GATA-6 in the mouse ovary. *Endocrinology* 1997; 138:3505-3514
- 62 Siltanen S, Anttonen M, Heikkila P, et al. Transcription factor GATA-4 is expressed in pediatric yolk sac tumors. *Am J Pathol* 1999; 155:1823-1829
- 63 Reis-Filho JS, Schmitt FC. Taking advantage of basic research: p63 is a reliable myoepithelial and stem cell marker. *Adv Anat Pathol* 2002; 9:280-289
- 64 McKeon F. p63 and the epithelial stem cell: more than status quo? *Genes Dev* 2004; 18:465-469
- 65 Pellegrini G, Dellambra E, Golisano O, et al. p63 identifies keratinocyte stem cells. *Proc Natl Acad Sci U S A* 2001; 98:3156-3161
- 66 Yang A, Kaghad M, Wang Y, et al. p63, a p53 homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. *Mol Cell* 1998; 2:305-316

- 67 Yang A, Schweitzer R, Sun D, et al. p63 is essential for regenerative proliferation in limb, craniofacial and epithelial development. *Nature* 1999; 398:714-718
- 68 Mills AA, Zheng B, Wang XJ, et al. p63 is a p53 homologue required for limb and epidermal morphogenesis. *Nature* 1999; 398:708-713
- 69 Mizukami Y, Nonomura A, Michigishi T, et al. Solid cell nests of the thyroid. A histologic and immunohistochemical study. *Am J Clin Pathol* 1994; 101:186-191
- 70 Harach HR. Solid cell nests of the thyroid. An anatomical survey and immunohistochemical study for the presence of thyroglobulin. *Acta Anat (Basel)* 1985; 122:249-253
- 71 Preto A, Reis-Filho JS, Ricardo S, et al. P63 expression in papillary and anaplastic carcinomas of the thyroid gland: lack of an oncogenetic role in tumorigenesis and progression. *Pathol Res Pract* 2002; 198:449-454
- 72 McLeod JF, Cooke NE. The vitamin D-binding protein, alpha-fetoprotein, albumin multigene family: detection of transcripts in multiple tissues. *J Biol Chem* 1989; 264:21760-21769
- 73 Gitlin D, Perricelli A, Gitlin GM. Synthesis of -fetoprotein by liver, yolk sac, and gastrointestinal tract of the human conceptus. *Cancer Res* 1972; 32:979-982
- 74 Mares V, Kovaru F, Kovaru H. Alpha-fetoprotein in the brain of developing rats and pigs. An immunofluorescent study of cell-and-tissue differentiation. *Basic Appl Histochem* 1982; 26:53-63
- 75 Bergstrand CG, Czar B. Paper electrophoretic study of human fetal serum proteins with demonstration of a new protein fraction. *Scand J Clin Lab Invest* 1957; 9:277-286
- 76 Mizejewski GJ. Biological roles of alpha-fetoprotein during pregnancy and perinatal development. *Exp Biol Med (Maywood)* 2004; 229:439-463
- 77 Gabant P, Forrester L, Nichols J, et al. Alpha-fetoprotein, the major fetal serum protein, is not essential for embryonic development but is required for female fertility. *Proc Natl Acad Sci U S A* 2002; 99:12865-12870

- 78 Sala-Trepat JM, Dever J, Sargent TD, et al. Changes in expression of albumin and alpha-fetoprotein genes during rat liver development and neoplasia. *Biochemistry* 1979; 18:2167-2178
- 79 Sell S, Longley MA, Boulter J. alpha-Fetoprotein and albumin gene expression in brain and other tissues of fetal and adult rats. *Brain Res* 1985; 354:49-53
- 80 Nahon JL, Tratner I, Poliard A, et al. Albumin and alpha-fetoprotein gene expression in various nonhepatic rat tissues. *J Biol Chem* 1988; 263:11436-11442
- 81 Wagh PV, Lippes J. Human oviductal fluid proteins. V. Identification of human oviductin-I as alpha-fetoprotein. *Fertil Steril* 1993; 59:148-156
- 82 Yao F, Guo JM, Xu CF, et al. Detecting AFP mRNA in peripheral blood of the patients with hepatocellular carcinoma, liver cirrhosis and hepatitis. *Clin Chim Acta* 2005; 361:119-127
- 83 Omori J, Niimi A, Furuichi Y, et al. [A case of AFP-producing cholangiocarcinoma with difficulty in differential diagnosis from hepatocellular carcinoma]. *Nippon Shokakibyō Gakkai Zasshi* 2004; 101:1106-1111
- 84 Talerman A, Haije WG, Baggerman L. Serum alphafetoprotein (AFP) in patients with germ cell tumors of the gonads and extragonadal sites: correlation between endodermal sinus (yolk sac) tumor and raised serum AFP. *Cancer* 1980; 46:380-385
- 85 Dziadek M, Adamson E. Localization and synthesis of alphafoetoprotein in post-implantation mouse embryos. *J Embryol Exp Morphol* 1978; 43:289-313
- 86 Janzen RG, Mably ER, Tamaoki T, et al. Synthesis of alpha-fetoprotein by the pre-implantation and post-implantation bovine embryo. *J Reprod Fertil* 1982; 65:105-110
- 87 Smith KM, Lai PC, Robertson HA, et al. Distribution of alpha 1-fetoprotein in fetal plasma, allantoic fluid, amniotic fluid and maternal plasma of cows. *J Reprod Fertil* 1979; 57:235-238
- 88 Thomson JA, Itskovitz-Eldor J, Shapiro SS, et al. Embryonic stem cell lines derived from human blastocysts. *Science* 1998; 282:1145-1147



- 89 Schuldiner M, Yanuka O, Itskovitz-Eldor J, et al. Effects of eight growth factors on the differentiation of cells derived from human embryonic stem cells. *Proc Natl Acad Sci U S A* 2000; 97:11307-11312
- 90 Shambloott MJ, Axelman J, Littlefield JW, et al. Human embryonic germ cell derivatives express a broad range of developmentally distinct markers and proliferate extensively in vitro. *Proc Natl Acad Sci U S A* 2001; 98:113-118
- 91 Sell S, Dunsford HA. Evidence for the stem cell origin of hepatocellular carcinoma and cholangiocarcinoma. *Am J Pathol* 1989; 134:1347-1363
- 92 Lajtha LG. Stem cell concepts. *Differentiation* 1979; 14:23-34
- 93 Sherley JL. Asymmetric cell kinetics genes: the key to expansion of adult stem cells in culture. *Stem Cells* 2002; 20:561-572
- 94 Potten CS, Morris RJ. Epithelial stem cells in vivo. *J Cell Sci Suppl* 1988; 10:45-62
- 95 Liu Y, Bohn SA, Sherley JL. Inosine-5'-monophosphate dehydrogenase is a rate-determining factor for p53-dependent growth regulation. *Mol Biol Cell* 1998; 9:15-28
- 96 Sherley JL, Stadler PB, Johnson DR. Expression of the wild-type p53 antioncogene induces guanine nucleotide-dependent stem cell division kinetics. *Proc Natl Acad Sci U S A* 1995; 92:136-140
- 97 Lee HS, Crane GG, Merok JR, et al. Clonal expansion of adult rat hepatic stem cell lines by suppression of asymmetric cell kinetics (SACK). *Biotechnol Bioeng* 2003; 83:760-771
- 98 Rapoport B, Filetti S, Takai N, et al. Studies on the cyclic AMP response to thyroid stimulating immunoglobulin (TSI) and thyrotropin (TSH) in human thyroid cell monolayers. *Metabolism* 1982; 31:1159-1167
- 99 Derwahl M, Manole D, Sobke A, et al. Pathogenesis of toxic thyroid adenomas and nodules: relevance of activating mutations in the TSH-receptor and Gs-alpha gene, the possible role of iodine deficiency and secondary and TSH-independent molecular mechanisms. *Exp Clin Endocrinol Diabetes* 1998; 106 Suppl 4:S6-9

- 100 Heldin NE, Cvejic D, Smeds S, et al. Coexpression of functionally active receptors for thyrotropin and platelet-derived growth factor in human thyroid carcinoma cells. *Endocrinology* 1991; 129:2187-2193
- 101 Derwahl M, Kuemmel M, Goretzki P, et al. Expression of the human TSH receptor in a human thyroid carcinoma cell line that lacks an endogenous TSH receptor: growth inhibition by cAMP. *Biochem Biophys Res Commun* 1993; 191:1131-1138
- 102 Jones HW, Jr., McKusick VA, Harper PS, et al. George Otto Gey. (1899-1970). The HeLa cell and a reappraisal of its origin. *Obstet Gynecol* 1971; 38:945-949
- 103 Masters JR. HeLa cells 50 years on: the good, the bad and the ugly. *Nat Rev Cancer* 2002; 2:315-319
- 104 Derwahl M, Seto P, Rapoport B. Complete nucleotide sequence of the cDNA for thyroid peroxidase in FRTL5 rat thyroid cells. *Nucleic Acids Res* 1989; 17:8380
- 105 Derwahl M, Seto P, Rapoport B. An abnormal splice donor site in one allele of the thyroid peroxidase gene in FRTL5 rat thyroid cells introduces a premature stop codon: association with the absence of functional enzymatic activity. *Mol Endocrinol* 1990; 4:793-799
- 106 Sambrook J, Fritsch EF, Maniatis T *Molecular cloning - a laboratory manual*, second edition. Cold Spring Harbour, New York: Cold spring harbour laboratory press, 1989
- 107 Saiki RK, Gelfand DH, Stoffel S, et al. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 1988; 239:487-491
- 108 Manole D, Schildknecht B, Gosnell B, et al. Estrogen promotes growth of human thyroid tumor cells by different molecular mechanisms. *J Clin Endocrinol Metab* 2001; 86:1072-1077
- 109 Hagiwara K, McMenamin MG, Miura K, et al. Mutational analysis of the p63/p73L/p51/p40/CUSP/KET gene in human cancer cell lines using intronic primers. *Cancer Res* 1999; 59:4165-4169
- 110 Stryer L *Biochemie*. 4.Auflage. Heidelberg, Berlin, Oxford: Akademischer Verlag GmbH, 1996

- 111 Polak JM, and Van Noorden S Immunocytochemistry: Modern methods and applications. Bristol:Wright: 1986
- 112 Hsu S-M, Raine L, DeLellis RA Advances in Immunohistochemistry. New York:Mason: 1984
- 113 Fujikawa T, Hirose T, Fujii H, et al. Purification of adult hepatic progenitor cells using green fluorescent protein (GFP)-transgenic mice and fluorescence-activated cell sorting. *J Hepatol* 2003; 39:162-170
- 114 Taniguchi H, Suzuki A, Zheng Y, et al. Usefulness of flow-cytometric cell sorting for enrichment of hepatic stem and progenitor cells in the liver. *Transplant Proc* 2000; 32:249-251
- 115 Dontu G, Wicha MS. Survival of mammary stem cells in suspension culture: implications for stem cell biology and neoplasia. *J Mammary Gland Biol Neoplasia* 2005; 10:75-86
- 116 Baldwin HS, Artman M. Recent advances in cardiovascular development: promise for the future. *Cardiovasc Res* 1998; 40:456-468
- 117 Sasai Y, De Robertis EM. Ectodermal patterning in vertebrate embryos. *Dev Biol* 1997; 182:5-20
- 118 Hill M, 2005 UNSW Embryology Vers. 4.9 0 7334 2108 3
- 119 Larsen JW Human Embryology. 3d Edition. New York: Churchill Livingstone, 2001
- 120 Ambesi-Impiombato FS, Villone G. The FRTL-5 thyroid cell strain as a model for studies on thyroid cell growth. *Acta Endocrinol Suppl (Copenh)* 1987; 281:242-245
- 121 Schwartz RE, Reyes M, Koodie L, et al. Multipotent adult progenitor cells from bone marrow differentiate into functional hepatocyte-like cells. *J Clin Invest* 2002; 109:1291-1302
- 122 Nagy P, Bisgaard HC, Thorgeirsson SS. Expression of hepatic transcription factors during liver development and oval cell differentiation. *J Cell Biol* 1994; 126:223-233
- 123 Stull RA, Hyun WC, Pallavicini MG. Simultaneous flow cytometric analyses of enhanced green and yellow fluorescent proteins and cell surface antigens in doubly transduced immature hematopoietic cell populations. *Cytometry* 2000; 40:126-134

- 124 Gratama JW, Orfao A, Barnett D, et al. Flow cytometric enumeration of CD34+ hematopoietic stem and progenitor cells. European Working Group on Clinical Cell Analysis. *Cytometry* 1998; 34:128-142
- 125 Lin RY, Kubo A, Keller GM, et al. Committing embryonic stem cells to differentiate into thymocyte-like cells in vitro. *Endocrinology* 2003; 144:2644-2649
- 126 Vetere A, Marsich E, Di Piazza M, et al. Neurogenin3 triggers beta-cell differentiation of retinoic acid-derived endoderm cells. *Biochem J* 2003; 371:831-841
- 127 Wang L, Schultz GA. Expression of Oct-4 during differentiation of murine F9 cells. *Biochem Cell Biol* 1996; 74:579-584
- 128 Reya T, Morrison SJ, Clarke MF, et al. Stem cells, cancer, and cancer stem cells. *Nature* 2001; 414:105-111
- 129 Li CD, Zhang WY, Li HL, et al. Isolation and Identification of a Multilineage Potential Mesenchymal Cell from Human Placenta. *Placenta* 2005;
- 130 Shih DT, Lee DC, Chen SC, et al. Isolation and characterization of neurogenic mesenchymal stem cells in human scalp tissue. *Stem Cells* 2005; 23:1012-1020
- 131 Jiang Y, Jahagirdar BN, Reinhardt RL, et al. Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 2002; 418:41-49
- 132 Derwahl M, Broecker M, Kraiem Z. Clinical review 101: Thyrotropin may not be the dominant growth factor in benign and malignant thyroid tumors. *J Clin Endocrinol Metab* 1999; 84:829-834
- 133 Reya T, Clevers H. Wnt signalling in stem cells and cancer. *Nature* 2005; 434:843-850
- 134 Ponti G, Conti L, Cataudella T, et al. Comparative expression profiles of ShcB and ShcC phosphotyrosine adapter molecules in the adult brain. *Neuroscience* 2005; 133:105-115
- 135 Burstein DE, Nagi C, Wang BY, et al. Immunohistochemical detection of p53 homolog p63 in solid cell nests, papillary thyroid carcinoma, and hashimoto's thyroiditis: A stem cell hypothesis of papillary carcinoma oncogenesis. *Hum Pathol* 2004; 35:465-473

- 136 Green H, Easley K, Iuchi S. Marker succession during the development of keratinocytes from cultured human embryonic stem cells. *Proc Natl Acad Sci U S A* 2003; 100:15625-15630
- 137 Ando T, Davies TF. Self-recognition and the role of fetal microchimerism. *Best Pract Res Clin Endocrinol Metab* 2004; 18:197-211
- 138 Bianchi DW, Zickwolf GK, Weil GJ, et al. Male fetal progenitor cells persist in maternal blood for as long as 27 years postpartum. *Proc Natl Acad Sci U S A* 1996; 93:705-708
- 139 Nelson JL, Furst DE, Maloney S, et al. Microchimerism and HLA-compatible relationships of pregnancy in scleroderma. *Lancet* 1998; 351:559-562
- 140 Artlett CM, Smith JB, Jimenez SA. Identification of fetal DNA and cells in skin lesions from women with systemic sclerosis. *N Engl J Med* 1998; 338:1186-1191
- 141 Johnson KL, Nelson JL, Furst DE, et al. Fetal cell microchimerism in tissue from multiple sites in women with systemic sclerosis. *Arthritis Rheum* 2001; 44:1848-1854
- 142 Ando T, Imaizumi M, Graves PN, et al. Intrathyroidal fetal microchimerism in Graves' disease. *J Clin Endocrinol Metab* 2002; 87:3315-3320
- 143 Badenhoop K. Intrathyroidal microchimerism in Graves' disease or Hashimoto's thyroiditis: regulation of tolerance or alloimmunity by fetal-maternal immune interactions? *Eur J Endocrinol* 2004; 150:421-423
- 144 Klintschar M, Schwaiger P, Mannweiler S, et al. Evidence of fetal microchimerism in Hashimoto's thyroiditis. *J Clin Endocrinol Metab* 2001; 86:2494-2498
- 145 Ando T, Davies TF. Clinical Review 160: Postpartum autoimmune thyroid disease: the potential role of fetal microchimerism. *J Clin Endocrinol Metab* 2003; 88:2965-2971
- 146 Srivatsa B, Srivatsa S, Johnson KL, et al. Microchimerism of presumed fetal origin in thyroid specimens from women: a case-control study. *Lancet* 2001; 358:2034-2038
- 147 Renne C, Ramos Lopez E, Steimle-Grauer SA, et al. Thyroid fetal male microchimerisms in mothers with thyroid disorders: presence of Y-chromosomal immunofluorescence in thyroid-infiltrating lymphocytes is more prevalent in Hashimoto's thyroiditis and Graves' disease than in follicular adenomas. *J Clin Endocrinol Metab* 2004; 89:5810-5814

- 148 Postiglione MP, Parlato R, Rodriguez-Mallon A, et al. Role of the thyroid-stimulating hormone receptor signaling in development and differentiation of the thyroid gland. *Proc Natl Acad Sci U S A* 2002; 99:15462-15467
- 149 Siepe M, Heilmann C, von Samson P, et al. Stem cell research and cell transplantation for myocardial regeneration. *Eur J Cardiothorac Surg* 2005; 28:318-324
- 150 Pluchino S, Martino G. The therapeutic use of stem cells for myelin repair in autoimmune demyelinating disorders. *J Neurol Sci* 2005; 233:117-119
- 151 Schwab ME. Repairing the injured spinal cord. *Science* 2002; 295:1029-1031
- 152 Roche E, Santana A, Vicente-Salar N, et al. From stem cells to insulin-producing cells: towards a bioartificial endocrine pancreas. *Panminerva Med* 2005; 47:39-51
- 153 Morasso MI, Tomic-Canic M. Epidermal stem cells: the cradle of epidermal determination, differentiation and wound healing. *Biol Cell* 2005; 97:173-183
- 154 Fausto N. Liver regeneration and repair: hepatocytes, progenitor cells, and stem cells. *Hepatology* 2004; 39:1477-1487
- 155 Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 1997; 3:730-737
- 156 Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell* 1997; 91:661-672
- 157 Takano T, Amino N. Fetal cell carcinogenesis: a new hypothesis for better understanding of thyroid carcinoma. *Thyroid* 2005; 15:432-438
- 158 Hochedlinger K, Yamada Y, Beard C, et al. Ectopic expression of Oct-4 blocks progenitor-cell differentiation and causes dysplasia in epithelial tissues. *Cell* 2005; 121:465-477
- 159 Thellin O, Zorzi W, Lakaye B, et al. Housekeeping genes as internal standards: use and limits. *J Biotechnol* 1999; 75:291-295