

6. Literaturverzeichnis

Agnew HW Jr, Webb WB, Williams RL: The first night effect: An EEG study of sleep. Psychophysiology, 1966; 2, 263-266.

Anderer P, Klösch G, Gruber G, Trenker E, Pascual-Marqui RD, Zeithofer J, Barbanoj MJ, Rappelsberger P, Saletu B: Low-resolution brain electromagnetic tomography revealed simultaneously active frontal and parietal sleep spindle sources in the human cortex. Neuroscience. 2001; 103(3), 581-92.

Aserinsky E, Kleitman N: Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. Science, 1953; 118, 273-274.

Bastien C, Campbell KB: The evoked K-complex: all-or-none phenomenon ? Sleep, 1992; 15, 236-245.

Baumgart-Schmitt R, Herrmann WM, Eilers R, Bes F: On the use of neural network techniques to analyze sleep EEG data. First communication: application of evolutionary and genetic algorithms to reduce the feature space and to develop classification rules. Neuropsychobiology, 1997; 36(4), 194-210.

Baumgart-Schmitt R, Eilers R, Herrmann WM: On the use of neural network techniques to analyze sleep EEG data. Second communication: Training of evolutionary optimized neural networks on the basis of multiple subjects data and the application of context rules according to Rechtschaffen and Kales. Somnologie, 1997; 1, 171-183.

Baumgart-Schmitt R, Herrmann WM, Eilers R: On the use of neural network techniques to analyze sleep EEG data. Third communication: Robustification of the classifier by applying an algorithm obtained from 9 different networks. Neuropsychobiology, 1998; 37(1), 49-58.

Beier KM, Kubicki St: Kortikale Verteilung zweier Delta-Frequenzen im langsamen Schlaf. Z EEG-EMG, 1987; 18, 47-51.

Berger H: Über das Elektrenkephalogramm des Menschen. Arch Psychiat Nervenkrankh, 1929; 87, 527-570.

Browman CP, Cartwright RD: The first night effect on sleep and dreams. Biological Psychiatry, 1980; 15, 809-812.

Cohen J: A coefficient of agreement for nominal scales. Educational and Psychological Measurement, 1960; 20, 37-46.

Colrain IM, Webster KE, Hirst G, Campbell KB: The role of vertex sharp waves and K-complexes in the generation of N300 in auditory and respiratory-related evoked potentials during early stage 2 NREM sleep. Sleep, 2000; 23, 97-106.

Cote KA, de Ligt DR, Langley SD, Campbell KB: Scalp topography of the auditory evoked K-Complex in Stage 2 and slow wave sleep. J Sleep Res, 1999; 8, 263-272.

Danker-Hopfe H, Herrmann WM: Interrater-Reliabilität visueller Schlafstadienklassifikation nach Rechtschaffen und Kales-Regeln: Review und methodische Erwägungen. Klin Neurophysiol, 2001; 32, 89-99.

Danker-Hopfe H, Kunz D, Gruber G, Klösch G, Lorenzo JL, Himanen SL, Kemp B, Penzel T, Röschke J, Dorn H, Schlögl A, Trenker E, Dorffner G: Interrater Reliability between scorers from eight European Sleep laboratories in subjects with different sleep disorders. J Sleep Res, 2004; 13, 63-69.

Dement WC, Kleitman N: Cyclic variations in EEG during sleep and their relation to eye movements, body motility, and dreaming. Electroencephalography and Clinical Neurophysiology, 1957; 9, 673-690.

Ehlert I, Danker-Hopfe H, Höller L, von Rickenbach P, Baumgart-Schmitt R, Herrmann WM: A comparison between EEG-Recording and Scoring by QUISI Version 1.0 and Standard PSG with Visual Scoring. Somnologie, 1998; 2, 104-116.

Feinberg I, March JD, Fein G, Aminoff MJ: Log amplitude is a linear function of log frequency in NREM sleep EEG of young and elderly normal subjects. Electroencephalogr Clin Neurophysiol, 1984; 58, 158-160.

Gfullner F, Siemon G: Studies with the fully automated EEG sleep analysis system QUISI Pneumologie, 2000; 54 (12), 580-583.

Happe S, Anderer P, Gruber G, Klösch G, Saletu B, Zeitlhofer J: Scalp Topography of the Spontaneous K-Complex and of Delta-Waves in Human Sleep. Brain Topography, Fall 2002; 15, 1, 43-49.

Himanen SL: A new visual adaptive scoring system for sleep recordings. University of Tampere, 2000.

Himanen SL, Hasan J: Limitations of Rechtschaffen and Kales. Sleep Medicine Reviews, 2000; Vol.4, No.2, 149-167.

Hori T, Hayashi M, Kato K: Changes of EEG patterns and reaction time during hypnagogic state. Sleep Res, 1991; 20, 20.

Hori T, Hayashi M, Morikawa T: Topographical EEG changes and the hypnagogic experience. In: Sleep Onset: Normal and Abnormal Processes, 1994; 237-253. Eds. RD Ogilvie and JR Harsh, American Psychological Association, Washington, DC.

Hori T, Sugita Y, Koga E, Shirakawa S, Inoue K, Uchida S, Kuwahara H, Kousaka M, Kobayashi T, Tsuji Y, Terashima M, Fukuda K, Fukuda N: Proposed supplements and amendments to "A manual of standardized terminology, techniques and scoring system for

sleep stages of human subjects”, the Rechtschaffen and Kales (1968) standard. Psychiatry and Clinical Neurosciences, 2001; 55, 305-310.

Huupponen E, Himanen SL, Varri A, Hasan J, Lehtokangas M, Saarinen J: A study on gender and age differences in sleep spindles. Neuropsychobiology, 2002; 45, 99-105.

Jasper HH: The ten twenty electrode system of the international federation. Electroenceph Clin Neurophysiol, 1958; 10, 371-375.

Kim Y, Kurachi M, Horita M, Matsuura K, Kamikawa Y: Agreement of visual scoring of sleep stages among many laboratories in Japan: effect of a supplementary definition of slow wave on scoring of slow wave sleep. Jpn J Psychiatry Neurol, 1993; Mar, 47(1), 91-97.

Kubicki S, Herrmann WM, Holler L, Scheuler W: Comments on the rules by Rechtschaffen and Kales about the visual scoring of sleep EEG recordings. EEG EMG Z Elektroenzephalogr Elektromyogr Verwandte Geb, 1982; Jun, 13(2), 51-60.

Kubicki St, Holler L, Berg I, Pastelak-Price C, Dorow R: A comparison of results obtained by visual scoring and automatic analysis with the Oxford Sleep stager. Sleep, 1989; 12, 140-149.

Kubicki St, Ehlert-Spieweg I, Herrmann WM: Das Westend-Schlaf-Glossar. Z. EEG-EMG, 1997; 28, 218-253.

Kuhwahara H, Higashi H, Mizuki Y, Matsunari S, Tanaka M, Inanaga K: Automatic real-time analysis of human sleep stages by an interval histogramm method. Electroencephalogr Clin Neurophysiol, 1988; 70, 220-229.

Kunz D, Danker-Hopfe H, Gruber G, Klösch G, Lorenzo JL, Himanen SL, Kemp B, Penzel T, Röschke J, Dorffner G: Interrater-Reliabilität bei gesunden Probanden aller Altersgruppen. Somnologie, 2000; 4 (Suppl. 1), 33.

Landis JR, Koch GG: The measurement of observer agreement for categorical data. *Biometrics*, 1977; 33, 159-174.

Loomis AL, Harvey EN, Hobart GA: Cerebral sites during sleep as studied by human brain potential. *J Exp Psychol*, 1937; 21, 127-144.

Lorenzo JL, Barbanoj MJ: Variability of sleep parameters across multiple laboratory sessions in healthy young subjects: The “very first night effect”. *Psychophysiology*, 2002; 39, 409-413.

Mahowald MM, Eiken T, Cashman D, Jokinen D, Cohen D, Fitzgerald R: Clinical validation of CASS, a computer assisted sleep system. *Sleep Res*, 1989; 18, 394.

Martin WB, Johnson LC, Viglione SS, Naitoh P, Joseph RD, Moses JD: Pattern recognition of EEG-EOG as a technique for all night sleep stage scoring. *Electroencephalogr Clin Neurophysiol*, 1972; 32, 417-427.

McDannold MD, Scharf MB, Fletcher MS, Rucosky G, Hux G, Winchester T: A comparison of a computer assisted scoring system to manual methods. *Sleep Res*, 1990; 19, 371.

Monroe LJ: Inter-rater reliability of scoring EEG records. Paper read at the Association Psychophysiological Study of Sleep Meeting, Santa Monica, California, April 1967. Abstract in *Psychophysiology*, 1968; 4, 370-371.

Naitoh P, Lubin A, Johnson LC, Wyborney G: Brain wave “generating” process during waking and sleeping. Amer. EEG Soc. Ann. Meet, Washington, 1970.

Norman RG, Pal I, Stewart C, Walsleben JA Rapoport DM: Interobserver agreement among sleep scorers from different centers in a large data set. *Sleep*, 2000; 23, 901-908.

Orr WC, Hansotia P, Anderson MW, Cohen D: Computer assisted sleep staging and visual scoring: a comparison CNS “CASS” and manual methods. *Sleep Res*, 1988; 17, 344.

Pastelak-Price C: Das internationale 10-20-System zur Elektrodenplatzierung: Begründung, praktische Anleitung zu den Messschritten und Hinweisen zum Setzen der Elektroden. EEG-Labor, 1983; 5, 49-72.

Penzel T, Behler PG, von Buttlar M, Conradt R, Meier M, Möller A, Danker-Hopfe H: Reliabilität der visuellen Schlafauswertung nach Rechtschaffen und Kales von acht Aufzeichnungen durch neun Schlaflabore. Somnologie, 2003; 7, 49-58.

Rechtschaffen A, Kales A: A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. National Institutes of Health, Publication 204, US Government Printing Office, Washington DC, 1968.

Schaltenbrand N, Lengelle R, Macher JP: Neural Network Model: Application to automatic analysis of human sleep. Computers and Biomed Res, 1993; 26, 157-171.

Schaltenbrand N, Lengelle R, Toussaint M, Luthringer R, Carelli G, Jacqmin A, Lainey E, Muzet A, Macher JP: Sleep Stage scoring using neural network model: comparison between visual and automatic analysis in normal subjects and patients. Sleep, 1996; 19, 26-35.

Schenk C: Automatische Schlafstadienklassifikation mit einem einfachen, einkanaligen EEG-Gerät (Quisi). Atmung und Schlaf, 1999; Vol 3, Nr. 3, 99, 48.

Schmidt HS, Kaelbling R: The differential laboratory adaptation of sleep parameters. Biological Psychiatry, 1971; 3, 319-324.

Sharpley AL, Solomon RA, Cowen PJ: Evaluation of first night effect using ambulatory monitoring and automatic sleep stage analysis. Sleep, 1988; 11, 273-276.

Ueda K, Nittono H, Hayashi M, Hori T: Estimation of generator sources of human sleep spindles by dipole tracing method. Psychiatry and Clinical Neurosciences, 2000; 54, 270-271.
Webb WB und Dreblow LM: A modified method for scoring slow wave sleep in older subjects. Sleep, 1982; 5, 195-199.

Webb WB: Recording methods and visual scoring criteria of sleep records: Comments and recommendations. *Percept Mot Skills*, 1986; 62, 664-66.