

8. LITERATURVERZEICHNIS

Acharya R., Fry E., Stuart D., Fox G., Rowlands D., Brown F. (1989). The three-dimensional structure of foot-and-mouth disease virus at 2.9 Å resolution. *Nature* **377**: 709-716.

Albrecht M. (1991). Coxsackievirus B3 Infektion der Maus: Untersuchungen zum akuten und chronischen Verlauf der CVB3-induzierten Herzmuskelerkrankung in unterschiedlichen Mausstämmen. *Vet. Med. Diss.*, München.

Ambros V., Petterson R.F., Baltimore D. (1978). An enzymatic activity in uninfected cells that cleaves the linkage between poliovirion RNA and the 5' terminal protein. *Cell* **15**: 1439-1446.

Anderson D.R., Wilson J.E., Carthy J.M., Yang D., Kandolf R., McManus B.M. (1996). Direct interactions of coxsackievirus B3 with immune cells in the splenic compartment of mice susceptible or resistant to myocarditis. *J. Virol.* **70**: 4632-45

Appelberg R., Castro A.G., Pedrosa J., Silva R.A., Orme I.M., Minoprio P. (1994). Role of γ -interferon and TNF- α during T-cell-independent and T-cell-dependent phase of mycobacterium avium infection. *Infect. Immun.* **62**: 3962-3971.

Arase H., Arase N., Ogasawara K., Good R.A., Onoe K. (1992). An NK1.1⁺CD4⁺8⁻ single-positive thymocyte subpopulation that expresses a highly skewed T-cell antigen receptor V beta family. *Proc. Natl. Acad. Sci. USA* **89**: 6506-6510.

Arase H., Arase N., Saito T. (1995). Fas-mediated cytotoxicity by freshly isolated natural killer cells. *J. Exp. Med.* **181**: 1235-1238.

Austyn J. M., Wood K. J. (1993). Principles of cellular and molecular immunology. Oxford University Press, Oxford.

Bendelac A., Lantz O., Quimby M.E., Yewdell J.W., Bennink J.R., Brutkiewicz R.R. (1995). CD1 recognition by mouse NK1⁺ T lymphocytes. *Science* **268**: 863-865.

Bergelson J.M., Cunningham J.A., Droguett G., Kurt-Jones E.A., Krithivias A., Hong J.S., Horwitz M.S., Crowell R.L., Finberg R.W. (1997). Isolation of a common receptor for coxsackie B viruses and adenoviruses 2 and 5. *Science* **275**: 1320-1323.

Bix M., Raulet D. (1992). Functionally conformed free class I heavy chains exist on the surface of β_2 microglobulin negative cells. *J. Exp. Med.* **176**: 829-834.

Blay R., Simpson K., Leslie K., Huber S. (1989). Coxsackievirus-induced disease. CD4⁺ cells initiate both myocarditis and pancreatitis in DBA/2 mice. *Am. J. Pathol.* **135**: 899-907.

Brossay L., Jullien D., Cardell S., Sydora B.C., Burdin N., Modlin R.L., Kronenberg M. (1997). Mouse CD1 is mainly expressed on hemopoietic-derived cells. *J. Immunol.* **159**: 1216-1224.

Brown D.R., Fowell D.J., Corry D.B., Wynn T.A., Moskowitz N.H., Cheever A.W., Locksley R.M., Reiner S.L. (1996). β_2 -microglobulin-dependent NK1.1⁺ T cells are not essential for T helper cell 2 immune response. *J. Exp. Med.* **184**: 1295-1304.

Calabi F., Bradbury A. (1991). The CD1 system. *Tissue Antigens* **37**: 1-9.

Castano A.R., Tangri S., Miller J.E.W., Holcombe H.R., Jackson M.R., Huse W.D., Kronenberg M., Peterson P.A. (1995). Peptide binding and presentation by mouse CD1. *Science* **269**: 223-226.

Chow L.H., Gauntt C.J., McManus B.M. (1991). Differential effects of myocarditic variants of coxsackievirus B3 in inbred mice. A pathologic characterization of heart tissue damage. *Lab. Invest.* **64**: 55-64.

Chow L.H., Beisel K.W., McManus B.M. (1992). Enteroviral infection of mice with severe combined immunodeficiency. Evidence for direct pathogenesis of myocardial injury. *Lab. Invest.* **66**: 24-31.

Cocchi F., DeVico A.L., Garzino-Demo A., Arya S.K., Gallo R.C., Lusso P. (1995). Identification of RANTES, MIP-1 α , and MIP-1 β as major HIV-suppressive factors produced by CD8⁺ T cells. *Science* **270**: 1811-1815.

Colter J.S., Campbell J.B., Hatch R.L. (1964). The pathogenicity to mice of three variants of Mengo encephalomyelitis virus. *J. Cell. Comp. Physiol.* **65**: 229-236.

Cordell-Stewart B., Taylor M.W. (1971). Effect of double-stranded viral RNA on mammalian cells in culture. *Proc. Natl. Acad. Sci. USA* **68**: 1326-1330.

Dales S., Eggers H.J., Tamm I., Palade G.E. (1965). Electron microscopic study of the formation of poliovirus. *Virology* **26**: 379-389.

Dalton D.K., Pitts-Meek S., Keshav S., Figari I.S., Bradley A., Stewart T.A. (1993). Multiple defects of immune cell function in mice with disrupted interferon- γ genes. *Science* **259**: 1739-1742.

Deguchi H., Kitaura Y., Morita H., Kotaka M., Kawamura K. (1985). Cell-mediated immunity in coxsackie B3 virus myocarditis in mice - *in situ* characterization by monoclonal antibody of mononuclear cell infiltrates. *Heart Vessels Suppl.* **1**: 221-227.

Dethlefs S., Brahic M., Larsson-Sciard E.-L. (1997). An early, abundant cytotoxic T-lymphocyte response against Theiler's virus is critical for preventing viral persistence. *J. Virol.* **71**: 8875-8878.

Feduchi E., Carrasco L. (1991). Mechanisms of inhibition of HSV-1 replication by tumor necrosis factor and interferon gamma. *Virology* **180**: 822-825.

Fiette L., Aubert C., Brahic M., Rossi C.P. (1993). Theiler's virus infection of β_2 -microglobulin-deficient mice. *J. Virol.* **67**: 589-592.

Foulis A.K., Farquharson M.A., Cameron S.O., McGill M., Schönke H., Kandolf R. (1990). A search for the presence of the enteroviral capsid protein VP1 in pancreases of patients with type 1 (insulin-dependent) diabetes and pancreases and hearts of infants who died of coxsackieviral myocarditis. *Diabetologica* **33**: 290-298.

Franklin R.M. & Baltimore D. (1962). Patterns of macromolecular synthesis in normal and virus-infected mammalian cells. *Cold Spring Harbor Symp. Quant. Biol.* **27**: 175-194.

Franco M.A., Tin C., Rott L.S., VanCott J.L., McGhee J.R., Greenberg H.B. (1996). Evidence for CD8⁺ T-cell immunity to murine rotavirus in the absence of perforin, fas, and gamma interferon. *J. Virol.* **71**: 479-486.

Freeman H. (1970). *Picture Processing and Psychopictories*. Academic Press Inc., New York.

Fung-Leung W.-P., Kündig T.M., Zinkernagel R.M., Mak T.W. (1991). Immune response against lymphocytic choriomeningitis virus infection in mice without CD8 expression. *J. Exp. Med.* **174**: 1425-1429.

Germain R.N. (1994). MHC-dependent antigen processing and peptide presentation: providing ligands for T-lymphocyte activation. *Cell* **71**: 287-299.

Gifford R., Dalldorf G. (1951). The morbid anatomy of experimental coxsackie virus infection. *Am. J. Pathol.* **27**: 1047-1064.

Gimenez B., Amarasekera D., Argo E., Cash P. (1995). Analysis of protein synthesis by two-dimensional gel electrophoresis in T-cells persistently infected with coxsackie B virus. *Electrophoresis* **16**: 317-321.

Godeny E.K., Gauntt C.J. (1987). *In situ* immune autoradiographic identification of cells in the heart tissues of mice with coxsackievirus B3-induced myocarditis. *Am. J. Pathol.* **129**: 267-276.

Green S.J., Nancy C.A. (1993). Antimicrobial and immunopathologic effects of cytokine-induced nitric oxide synthase. *Cur. Opin. Infect. Dis.* **6**: 384-389.

Gregorio S.B., Nakao J.C., Beran G.W. (1972). Human enteroviruses in animals and arthropods in the central Philippines. *Southeast Asian J. Trop. Med. Public Health* **3**: 45-51.

Guthrie M., Lodge P.A., Huber S.A. (1984). Cardiac injury in myocarditis induced by coxsackievirus group B, type 3 in Balb/c mice is mediated by Lyt 2⁺ cytolytic lymphocytes. *Cell. Immunol.* **88**: 558-567.

Hahn S., Gehri R., Erb P. (1995). Mechanism and biological significance of CD4-mediated cytotoxicity. *Immunol. Rev.* **146**: 57-79.

Heinzel F.P., Sadick M.D., Holaday B.J., Coffman R.L., Locksley R.M. (1989). Reciprocal expression of interferon- γ or interleukin-4 during the resolution or progression of murine Leishmaniasis. *J. Exp. Med.* **169**: 59-72.

Hesse M. (1997). The role of CD4⁺ T cells in infection-associated immunopathology. Diss. rer. nat., Berlin.

Hogle J.M., Chow M., Filman D.J. (1985). Three-dimensional structure of poliovirus at 2.9 Å resolution. *Science* **229**: 1358-1365.

Hohenadl C., Klingel K., Mertsching J., Hofschneider P.H., Kandolf R. (1991). Strand-specific detection of enteroviral RNA in myocardial tissue by *in situ* hybridization. *Moll. Cell. Probes* **5**: 11-20.

Hou S., Doherty P.C., Zijlstra M., Jaenisch R., Katz J.M. (1992). Delayed clearance of Sendai virus in mice lacking class I MHC-restricted CD8⁺ T cells. *J. Immunol.* **149**: 1319-1325.

Huang S., Hendriks W., Althage A., Hemmi S., Bluethmann H., Kamijo R., Vilcek J., Zinkernagel R.M., Aguet M. (1993). Immune response in mice that lack the interferon- γ receptor. *Science* **259**: 1742-1745.

Huber S.A., Job L.P., Woodruff J.F. (1984). In vitro culture of coxsackievirus group B, type 3 immune spleen cells on infected endothelial cells and biological activity of the cultured cells *in vivo*. *Infect. Immun.* **43**: 567-573.

Huber S.A., Lodge P.A. (1984). Coxsackievirus B3 myocarditis in BALB/c mice: Evidence for autoimmunity to myocyte antigens. *Am. J. Pathol.* **116**: 21-29.

Huber S.A., Polgar J., Schultheiss P., Schwimmbeck P. (1994). Augmentation of pathogenesis of coxsackievirus B3 infections in mice by exogenous administration of interleukin-1 and interleukin-2. *J. Virol.* **68**: 195-206.

Itoh N., Yonehara S., Ishii A., Yonehara M., Mizushima S., Sameshima M., Hase A., Set Y., Nagata S. (1991). The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell* **66**: 233-243.

Janeway C. A., Travers P. (1997). Host defense against infection. In: *Immunobiology - The immune system in health and disease*. Current Biology Ltd. and Garland Publishing Inc., London.

Jenne D.E., Tschopp J. (1988). Granzymes: A family of serine proteases in granules of cytolytic T lymphocytes. *Curr. Top. Microbiol. Immunol.* **140**: 33-47.

Kägi D., Ledermann B., Bürki K., Seiler P., Odermatt B., Olsen K.J., Podack E.R., Zinkernagel R.M., Hengartner H. (1994). Cytotoxicity mediated by T cells and natural killer cells is greatly impaired in perforin-deficient mice. *Nature* **369**: 31-37.

Kägi D., Seiler P., Pavlovic J., Ledermann B., Bürki K., Zinkernagel R.M., Hengartner H. (1995). The roles of perforin- and fas-dependent cytotoxicity in protection against cytopathic and noncytopathic viruses. *Eur. J. Immunol.* **25**: 3256-3262.

Kanagawa O., Vaupel B.A., Gayama S., Koehler G., Kopf M. (1993). Resistance of mice deficient in IL-4 to retrovirus-induced immunodeficiency syndrome (MAIDS). *Science* **262**: 240-242.

Kandolf R., Canu A., Hofschneider P.H. (1985). Coxsackie B3 virus can replicate in cultured human foetal heart cells and is inhibited by interferon. *J. Mol. Cell. Cardiol.* **17**: 167-181.

Kandolf R., Hofschneider P.H. (1985). Molecular cloning of the genome of a cardiotropic coxsackie B3 virus: full-length reverse transcribed recombinant cDNA generates infectious virus in mammalian cells. *Proc. Natl. Acad. Sci. USA* **82**: 4818-4822.

Kandolf R., Ameis D., Kirschner P., Canu A., Hofschneider P.H. (1987). *In situ* detection of enteroviral genomes in myocardial cells by nucleic acid hybridization: An approach to the diagnosis of viral heart disease. *Proc. Natl. Acad. Sci. USA* **84**: 6272-6276.

Kandolf R., Hofschneider P.H. (1989). Enteroviral heart disease. *Springer Semin. Immunopathol.* **11**: 1-13.

Kandolf R., Klingel K., Zell R., Selinka H.C., Raab U., Schneider-Brachert W., Bültmann B. (1993). Molecular pathogenesis of enterovirus-induced myocarditis: virus persistence and chronic inflammation. *Intervirology* **35**: 140-151.

Karupiah G., Xie Q.-W., Buller M.L., Nathan C., Duarte C., MacMicking J.D. (1993). Inhibition of viral replication by interferon- γ -induced nitric oxide synthase. *Science* **261**: 1445-1448.

Kitamura N., Semler B.L., Rothberg P.G., Larsen G.R., Adler C.J., Dorner A.J., Emini E.A., Hanecak R., Lee J.J., van der Werf S., Anderson C.W., Wimmer E. (1981). Primary structure, gene organization and polypeptide expression of poliovirus RNA. *Nature* **291**: 547-553.

Klingel K., Hohenadl C., Canu A., Albrecht M., Seemann M., Mall G., Kandolf R. (1992). Ongoing enterovirus-induced myocarditis is associated with persistent heart muscle infection: Quantitative analysis of virus replication, tissue damage and inflammation. *Proc. Natl. Acad. Sci. USA* **89**: 314-318.

Klingel K., Kandolf R. (1993). The role of enterovirus replication in the development of acute and chronic heart muscle disease in different immunocompetent mouse strains. *Scand. J. Infect. Dis. Suppl.* **88**: 79-85.

Klingel K., Stephan S., Sauter M., Zell R., McManus B., Bültmann B., Kandolf R. (1996). Pathogenesis of murine enterovirus myocarditis: Virus dissemination and immune cell targets. *J. Virol.* **70**: 8888-8895.

Klingel K., Sauter M., Selinka H.C., Bültmann B., Kandolf R. (1998a). Comparative assessment of cytokine mRNA expression in hearts of CVB3-infected permissive and resistant mice. *Path. Res. Pract.* **194**.

Klingel K., Rieger P., Mall G., Selinka H.C., Huber M., Kandolf R. (1998b). Visualization of enteroviral replication in myocardial tissue by ultrastructural *in situ* hybridization: Identification of target cells and cytopathic effects. *Lab. Invest.* **78**: 1227-1237.

Koller B.H., Marrack P., Kappler J.W., Smithies O. (1990). Normal development of mice deficient in β_2m , MHC class I proteins, and CD8⁺ T cells. *Science* **248**: 1227-1230.

Konig R., Huang L.Y., Germain R.N. (1992). MHC class II interaction with CD4 mediated by a region analogous to the MHC class I binding site for CD8. *Nature* **356**: 796-798.

Lantz O., Bendelac A. (1994). An invariant T cell receptor alpha chain is used by a unique subset of MHC class I-specific CD4⁺ and CD4⁻CD8⁻ T cells in mice and humans. *J. Exp. Med.* **180**: 1097-1106.

Larsen G.R., Dorner A.J., Harris T.J., Wimmer E. (1980). The structure of poliovirus replicative form. *Nucleic Acids Res.* **8**: 1217-1229.

Lehmann-Grube F., Löhler J., Utermöhlen O., Gegin C. (1993). Antiviral immune responses of lymphocytic choriomeningitis virus-infected mice lacking CD8⁺ T lymphocytes because of disruption of the β_2 -microglobulin gene. *J. Virol.* **67**: 332-339.

Lehmann-Grube F., Dralle H., Utermöhlen O., Löhler J. (1994). MHC class I molecule-restricted presentation of viral antigen in β_2 -microglobulin-deficient mice. *J. Immunol.* **153**: 595-603.

Leopardi R., Ilonen J., Mattila L., Salmi A.A. (1993). Effect of measles virus infection on MHC class II expression and antigen presentation in human monocytes. *Cell. Immunol.* **147**: 388-396.

Leslie K., Blay R., Haisch C., Lodge A., Weller A., Huber S. (1989). Clinical and experimental aspects of viral myocarditis. *Clin. Microbiol. Rev.* **2**: 191-203.

Liao N.-S., Bix M., Zijlstra M., Jaenisch R., Raulet D. (1991). MHC class I deficiency: susceptibility to natural killer (NK) cells and impaired NK activity. *Science* **253**: 199-202.

Lim K.A., Benyesh-Melnick M. (1960). Typing of viruses by combinations of antiserum pools: Application to typing of enteroviruses (coxsackie and echo). *J. Immunol.* **84**: 309-317.

Littman D.R. (1987). The structure of the CD4 and CD8 genes. *Annu. Rev. Immunol.* **5**: 561-584.

Ljunggren H.-G., Karre K. (1991). In search of the "missing-self": MHC molecules and NK cell recognition. *Immunol. Today* **11**: 237-244.

Longson M., Cole F., Davies D. (1969). Isolation of a coxsackie virus group B, type 5 from the heart of a fatal case of myocarditis in an adult. *J. Clin. Pathol.* **22**: 654-658.

Lowenstein C.A., Hill S.L., Lafond-Walker A., Wu J., Allen G., Landavere M., Rose N.R., Herskowitz A. (1996). Nitric oxide inhibits viral replication in murine myocarditis. *J. Clin. Invest.* **97**: 1837-1843.

Lundgren D.L., Clapper W.E., Sanchez A. (1968). Isolation of human enteroviruses from Beagle dogs. *Proc. Soc. Exp. Biol. Med.* **128**: 463-467.

Luo M., Vriend G., Kamer G., Minor I., Arnold E., Rossmann M.G., Boege U., Scraba D.G., Duke G.M., Palmenberg A.C. (1987). The atomic structure of Mengo virus at 3.0 Å resolution. *Science* **235**: 182-191.

Mall G., Klingel K., Albrecht M., Seemann M., Rieger P., Kandolf R. (1991). Natural history of Coxsackievirus B3-induced myocarditis in ACA/Sn mice: viral persistence demonstrated by quantitative *in situ* hybridization histochemistry. *Eur. Heart J. 12 Suppl. D*: 121-123.

Maniatis T., Fritsch E.F., Sambrook J. (1982). *Molecular cloning. A laboratory manual.* Cold Spring Harbor, New York.

Matteucci D., Paglianti M., Giangregorio A.M., Capobianchi M.R., Dianzani F., Bendinelli M. (1985). Group B coxsackieviruses readily establish persistent infections in human lymphoid cell lines. *J. Virol.* **56**: 651-654.

McManus B.M., Kandolf R. (1991). Evolving concepts of cause, consequence, and control in myocarditis. *Curr. Opin. Cardiol.* **6**: 418-427.

McManus B.M., Chow L.H., Wilson J.E., Anderson D.R., Gulizia J.M., Gauntt C.J., Klingel K., Beisel K.W., Kandolf R. (1993). Direct myocardial injury by enterovirus: a central role in the evolution of murine myocarditis. *Clin. Immunol. Immunopathol.* **68**: 159-169.

Melnick J.C., Dow R.P. (1953). Poliomyelitis in Hidalgo County, Texas, 1948. Poliomyelitis and coxsackie viruses from flies. *Am. J. Hyg.* **58**: 288-309.

Melnick J.L. (1996). Enteroviruses: polioviruses, coxsackieviruses, echoviruses, and newer enteroviruses. In: *Fields Virology.* Lippincott-Raven Publishers, Philadelphia.

Mendelson C., Wimmer E., Racaniello V.R. 1989. Cellular receptor for poliovirus: molecular cloning, nucleotide sequence and expression of a new member of the immunoglobulin superfamily. *Cell* **56**: 855-865.

Morita H. (1981). Experimental coxsackie B3 virus myocarditis in golden hamsters. Light and electron microscopy findings in a long-term follow-up study. *Jpn. Circ. J.* **45**: 713-729.

Mosmann T.R., Coffman R.L. (1989). TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu. Rev. Immunol.* **7**: 145-173.

Muckelbauer J., Kremer M., Minor I., Diana G., Dutko F.J., Groarke J., Pevear D.C., Rossmann M.G. (1995). The structure of coxsackievirus B3 at 3.5 Å resolution. *Structure* **3**: 653-667.

Nathan C. (1992). Nitric oxide as a secretory product of mammalian cells. *FASEB J.* **6**: 3051-3064.

Parker D.C. (1993). T-cell dependent B-cell activation. *Annu. Rev. Immunol.* **11**: 331-360.

Peng T., Sadusky T., Li Y., Coulton G. R., Zhang H., Archard L. C. (2001). Altered expression of Bag-1 in Coxsackievirus B3 infected mouse heart. *Cardiovasc. Res.* **50**: 46-55

Podack E.R., Young J.D.E., Cohn Z.A. (1985). Isolation and biochemical and functional characterization of perforin 1 from cytolytic T-cell granules. *Proc. Natl. Acad. Sci. USA* **82**: 8629-8633.

Porcelli S.A. (1995). The CD1 family: a third lineage of antigen-presenting molecules. *Adv. Immunol.* **59**: 1-98.

Roitt I.M., Brostoff J., Male D. (1985). *Immunology*. Gower Medical Publishing, London.

Rose N.R., Neumann D.A., Herskowitz A. (1988). Genetics of susceptibility to viral myocarditis in mice. *Pathol. Immunopathol. Res.* **7**: 266-278.

Rossmann M.G., Arnold E., Erickson J.W., Frankenberger E.A., Griffith J.P., Hecht H.J., Johnson J.E., Kamer G., Luo M., Mosser A.G., Rueckert R.R., Sherry B., Vriend G. (1985). Structure of a human common cold virus and functional relationship to other picornaviruses. *Nature* **317**: 145-153.

Rueckert R.R. (1996). Picornaviridae: The viruses and their replication. In: *Fields Virology*. Lippincott-Raven Publishers, Philadelphia.

Sato S., Tsutsumi R., Burke A., Carlson G., Porro V., Seko Y., Oksumura K., Kawana R., Virmani R. (1994). Persistence of replicating coxsackievirus B3 in the athymic murine heart is associated with development of myocarditic lesions. *J. Gen. Virol.* **75**: 2911-2924.

Shiver J.W., Su L., Henkart P.A. (1992). Cytotoxicity with target DNA breakdown by rat basophilic leukemia cells expressing both cytolysin and granzyme A. *Cell* **71**: 315-322.

Schnurr D.P., Schmidt N.J. (1984). Persistent infection of mouse fibroblasts with coxsackievirus. *Arch. Virol.* **81**: 91-101.

Schnurr D.P., Schmidt N.J. (1984). Coxsackievirus B3 persistence and myocarditis in NFR nu/nu and +/nu mice. *Med. Microbiol. Immunol.* **173**: 1-7.

Scott P., Trinchieri G. (1995). The role of natural killer cells in host-parasite interactions. *Curr. Opin. Immunol.* **7**: 34-40.

Seko Y., Tsuchimochi H., Nakamura T., Okumura K., Naito S., Imataka K., Fujii J., Takaku F., Yazaki J. (1990). Expression of major histocompatibility complex class I antigen in murine ventricular myocytes infected with coxsackievirus B3. *Circul. Res.* **67**: 360-367.

Seko Y., Shinkai Y., Kawasaki A., Yagita H., Okumura K., Yazaki Y. (1992). Evidence of perforin-mediated cardiac myocyte injury in acute murine myocarditis caused by coxsackie virus B3. *J. Pathol.* **170**: 53-58.

Seko Y., Shinkai Y., Kawasaki A., Yagita H., Okumura K., Takaku F., Yazaki Y. (1991). Expression of perforin in infiltrating cells in murine hearts with acute myocarditis caused by coxsackievirus B3. *Circulation* **84**: 788-795.

Snell G. (1978). Congenic resistant strain of mice. In: *Origins of inbred mice*. Academic Press Inc., New York.

Stalder T., Hahn S., Erb P. (1994). Fas antigen is the major target molecule for CD4⁺ T cell-mediated cytotoxicity. *J. Immunol.* **152**: 1127-1133.

Stewart B.W. (1994). Mechanisms of apoptosis: integration of genetic, biochemical, and cellular indicators. *J. Natl. Cancer Inst.* **86**: 1286-1295.

Tang Y.-W., Graham B.S. (1994). Anti-IL-4 treatment at immunization modulates cytokine expression, reduces illness, and increases cytotoxic T lymphocyte activity in mice challenged with respiratory syncytial virus. *J. Clin. Invest.* **94**: 1953-1958.

Tay C.-H., Welsh R.M., Brutkiewicz R.R. (1995). NK cell response to viral infections in β_2 -microglobulin-deficient mice. *J. Immunol.* **154**: 780-789.

Tay C.-H., Welsh R.M. (1997). Distinct organ-dependent mechanisms for the control of murine cytomegalovirus infection by natural killer cells. *J. Virol.* **71**: 267-275:

Timonen T., Ortaldo J.R., Herberman R.B. (1981). Characteristics of human large granular lymphocytes and relationship to natural killer and K cells. *J. Exp. Med.* **153**: 569-582.

Tomko R.P., Xu R., Philipson L. (1997). HCAR and MCAR: The human and mouse cellular receptors for subgroup C adenoviruses and group B coxsackieviruses. *Proc. Natl. Acad. Sci. USA* **94**: 3352-3356.

Toniolo A.T., Onodera T., Jordan G., Yoon J.W., Notkins A.L. (1982). Virus-induced diabetes mellitus. Glucose abnormalities produced in mice by six members of the coxsackie B virus group. *Diabetes* **31**: 496-499.

Tripp C.S., Wolf S.F., Unanue E.R. (1993). Interleukin-12 and tumor necrosis factor-alpha are costimulators of interferon-gamma production by natural killer cells in severe combined immunodeficiency mice with listeriosis and interleukin-10 is a physiologic antagonist. *Proc. Natl. Acad. Sci. USA* **90**: 3725-3729.

Vikingsson A., Pederson K., Müller D. (1996). Altered kinetics of CD4⁺ T cell proliferation and interferon- γ production in the absence of CD8⁺ T lymphocytes in virus-infected β_2 -microglobulin-deficient mice. *Cell. Immunol.* **173**: 261-268.

Vuorinen T., Vainionpaa R., Kettinen H., Hyypia T. (1994). Coxsackievirus B3 infection in human leukocytes and lymphoid cell lines. *Blood* **84**: 823-829.

Watanabe-Fukunaga R., Brannan C.I., Itoh N., Yonehara S., Copeland N.G., Jenkins N.A., Nagata S. (1992). The cDNA structure, expression and chromosomal assignment of the mouse Fas antigen. *J. Immunol.* **148**: 1274-1279.

Wolfgram L.J., Beisel K.W., Herskowitz A., Rose N.R. (1986). Variations in the susceptibility to coxsackievirus B3-induced myocarditis among different strains of mice. *J. Immunol.* **136**: 1846-1852.

Woodruff J.F. (1980). Viral myocarditis. A review. *Am. J. Pathol.* **101**: 425-484.

Yagi S., Schnurr D., Lin J. (1992). Spectrum of monoclonal antibodies to coxsackievirus B3 includes type- and group-specific antibodies. *J. Clin. Microbiol.* **30**: 2498-2501.

Yamamura M., Uyemura K., Deans R.J., Weinberg K., Rea T.H., Bloom B.R., Modlin R.L. (1991). Defining protective responses to pathogens: cytokine profiles in leprosy lesions. *Science* **254**: 277-279.

Yang Y., Xiang Z., Ertl H.C.J., Wilson J.M. (1995). Upregulation of class I major histocompatibility complex antigens by interferon- γ is necessary for T-cell-mediated elimination of recombinant adenovirus-infected hepatocytes *in vivo*. *Proc. Natl. Acad. Sci. USA* **92**: 7257-7251.

Yoo J.W., Austin M., Onodera T., Notkins A.L. (1979). Virus-induced diabetes mellitus. Isolation of a virus from the pancreas of a child with diabetic ketoacidosis. *N. Engl. J. Med.* **300**: 1173-1179.

Yoshimoto T., Paul W.E. (1994). CD4⁺, NK1⁺ T cells promptly produce interleukin 4 in response to *in vivo* challenge with anti-CD3. *J. Exp. Med.* **179**: 1285-1295.

Yoshimoto T., Bendelac A., Watson C., Hu-Li J., Paul W.E. (1995). Role of NK1.1⁺ T cells in a TH2 response and in immunoglobulin E production. *Science* **270**: 1845-1847.

Young J., D.-E., Hengartner H., Podack E.R., Cohn Z.A. (1986). Purification and characterization of a cytolytic pore-forming protein from granules of cloned lymphocytes with natural killer activity. *Cell* **44**: 849-859.

Young L.H.Y., Joag S.V., Zheng L.-M., Lee C.-H., Lee Y.-S., Young j. D.-E. (1990). Perforin-mediated myocardial damage in acute myocarditis. *Lancet* **336**: 1019-1021.

Zaragoza C., Ocampo C., Saura M., Leppo M., Wei X.-Q., Quick R., Moncada S., Liew F.Y., Lowenstein C.J. (1998). The role of inducible nitric oxide synthase in the host response to coxsackievirus myocarditis. *Proc. Natl. Acad. Sci. USA* **95**: 2469-2474.

Zijlstra M., Bix M., Simister N.E., Loring J.M., Raulet D.H., Jaenisch R. (1990). β_2 -microglobulin deficient mice lack CD4⁻CD8⁺ cytolytic T cells. *Nature* **344**: 742-746.

Zinkernagel R.M., Doherty P.C. (1974). Restriction of in vitro T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature* **248**: 701-702.