

6 Literaturverzeichnis

- Amann K, Gross ML, Ritz E. Pathophysiology underlying accelerated atherogenesis in renal disease: Closing in on the target. *J Am Soc Nephrol*. 2004;15:1664-1666.
- Anavekar NS, McMurray JJ, Velazquez EJ, Solomon SD, Kober L, Rouleau JL, White HD, Nordlander R, Maggioni A, Dickstein K, Zelenkofske S, Leimberger JD, Califf RM, Pfeffer MA. Relation between renal dysfunction and cardiovascular outcomes after myocardial infarction. *N Engl J Med*. 2004;351:1285-1295.
- Annuk M, Lind L, Linde T, Fellstrom B. Impaired endothelium-dependent vasodilation in renal failure in humans. *Nephrol Dial Transplant*. 2001;16:302-306.
- Arnett DK, Glasser SP, McVeigh G, Prineas R, Finkelstein S, Donahue R, Cohn JN, Sinaiko A. Blood pressure and arterial compliance in young adults: the Minnesota Children's Blood Pressure Study. *Am J Hypertens*. 2001;14:200-205.
- Bachmann J, Tepel M, Raidt H, Riezler R, Graefe U, Langer K, Zidek W. Hyperhomocysteinemia and the risk for vascular disease in hemodialysis patients. *J Am Soc Nephrol*. 1995;6:121-125.
- Baigent C, Burbury K, Wheeler D. Premature cardiovascular disease in chronic renal failure. *Lancet*. 2000;356:147-152.
- Becker BN, Himmelfarb J, Henrich WL, Hakim RM. Reassessing the cardiac risk profile in chronic hemodialysis patients: a hypothesis on the role of oxidant stress and other non-traditional cardiac risk factors. *J Am Soc Nephrol*. 1997;8:475-486.
- Beltran A, McVeigh G, Morgan D, Glasser SP, Neutel JM, Weber M, Finkelstein SM, Cohn JN. Arterial compliance abnormalities in isolated systolic hypertension. *Am J Hypertens*. 2001;14:1007-1011.
- Bhagat K, Vallance P. Inflammatory cytokines impair endothelium-dependent dilatation in human veins in vivo. *Circulation*. 1997;96:3042-3047.
- Bianchi G. Hypertension in chronic renal failure and end-stage renal disease patients treated with haemodialysis or peritoneal dialysis. *Nephrol Dial Transplant*. 2000;15:105-110.
- Blake GJ, Ridker PM. Novel clinical markers of vascular wall inflammation. *Circ Res*. 2001;89:763-771.
- Blacher J, Guerin AP, Pannier B, Marchais SJ, Safar ME, London GM. Impact of aortic stiffness on survival in end-stage renal disease. *Circulation*. 1999;99:2434-2439.
- Block GA, Hulbert-Shearon TE, Levin NW, Port FK. Association of serum phosphorus and calcium x phosphate product with mortality risk in chronic hemodialysis patients: a national study. *Am J Kidney Dis*. 1998;31:607-617.
- Boaz M, Smetana S, Weinstein T, Matas Z, Gafter U, Iaina A, Knecht A, Weissgarten Y, Brunner D, Fainaru M, Green MS. Secondary prevention with antioxidants of cardiovascular disease in endstage renal disease (SPACE): randomised placebo-controlled trial. *Lancet*. 2000;356:1213-1218.
- Cameron JD, McGrath BP, Dart AM. Use of radial artery applanation tonometry and a generalized transfer function to determine aortic pressure augmentation in subjects with treated hypertension. *J Am Coll Cardiol*. 1998;32:1214-1220.
- Cavalcanti S, Ciandrini A, Severi S, Badiali F, Bini S, Gattiani A, Cagnoli L, Santoro A. Model-based study of the effects of the hemodialysis technique on the compensatory response to hypovolemia. *Kidney Int*. 2004;65:1499-1510.
- Chakraborti S, Chakraborti T. Oxidant-mediated activation of mitogen-activated protein kinases and nuclear transcription factors in the cardiovascular system: a brief overview. *Cell Signal*. 1998;10:675-683.

- Charra B, Calemard M, Laurent G. Importance of treatment time and blood pressure control in achieving long-term survival on dialysis. *Am J Nephrol.* 1996;16:35-44.
- Chen CH, Nevo E, Fetcs B, Pak PH, Yin FC, Maughan WL, Kass DA. Estimation of central aortic pressure waveform by mathematical transformation of radial tonometry pressure: validation of generalized transfer function. *Circulation.* 1997;95:1827-1836.
- Chen CH, Ting CT, Nussbacher A, Nevo E, Kass DA, Pak P, Wang SP, Chang MS, Yin FC. Validation of carotid artery tonometry as a means of estimating augmentation index of ascending aortic pressure. *Hypertension.* 1996;27:168-175.
- Cohen DL, Townsend RR. Large and small artery compliance changes during hemodialysis. *Am J Hypertens.* 2002;15:236-239.
- Cooke JP, Rossitch E Jr, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest.* 1991;88:1663-1671.
- Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, Deanfield J, Drexler H, Gerhard-Herman M, Herrington D, Vallance P, Vita J, Vogel R; International Brachial Artery Reactivity Task Force. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol.* 2002;39:257-265.
- Covic A, Goldsmith DJ, Panaghiu L, Covic M, Sedor J. Analysis of the effect of hemodialysis on peripheral and central arterial pressure waveforms. *Kidney Int.* 2000;57:2634-2643
- Daugirdas JT. Simplified equations for monitoring Kt/V, PCRn, eKt/V, and ePCRn. *Adv Ren Replace Ther.* 1995;2:295-304.
- Donauer J. Hemodialysis-induced hypotension: impact of technologic advances. *Semin Dial.* 2004;17:333-335.
- Drzewiecki GM, Melbin J, Noordergraaf A. Arterial tonometry: review and analysis. *J Biomech.* 1983;16:141-152.
- Eknoyan G, Beck GJ, Cheung AK, Daugirdas JT, Greene T, Kusek JW, Allon M, Bailey J, Delmez JA, Depner TA, Dwyer JT, Levey AS, Levin NW, Milford E, Ornt DB, Rocco MV, Schulman G, Schwab SJ, Teehan BP, Toto R; Hemodialysis (HEMO) Study Group. Effect of dialysis dose and membrane flux in maintenance hemodialysis. *N Engl J Med.* 2002;347:2010-2019.
- Finkelstein SM, Cohn JN. First- and third-order models for determining arterial compliance. *J Hypertens Suppl.* 1992;10:11-14.
- Frei U, Schober-Halstenberg HJ. Nierenersatztherapie in Deutschland. *QuaSi-Niere Jahresbericht 2003/2004*, Berlin, Deutschland.
- Gatzka CD, Cameron JD, Dart AM, Berry KL, Kingwell BA, Dewar EM, Reid CM, Jennings GL. Correction of carotid augmentation index for heart rate in elderly essential hypertensives. ANBP2 Investigators. Australian Comparative Outcome Trial of Angiotensin-Converting Enzyme Inhibitor- and Diuretic-Based Treatment of Hypertension in the Elderly. *Am J Hypertens.* 2001 ;14:573-577.
- Goldwasser P, Mittman N, Antignani A, Burrell D, Michel MA, Collier J, Avram MM. Predictors of mortality in hemodialysis patients. *J Am Soc Nephrol.* 1993;3:1613-1622.
- Goodman WG, Goldin J, Kuizon BD, Yoon C, Gales B, Sider D, Wang Y, Chung J, Emerick A, Greaser L, Elashoff RM, Salusky IB. Coronary-artery calcification in young adults with end-stage renal disease who are undergoing dialysis. *N Engl J Med.* 2000;342:1478-1483.

- Griendling KK, Sorescu D, Lassegue B, Ushio-Fukai M. Modulation of protein kinase activity and gene expression by reactive oxygen species and their role in vascular physiology and pathophysiology. *Arterioscler Thromb Vasc Biol.* 2000;20:2175-2183.
- Guerin AP, Blacher J, Pannier B, Marchais SJ, Safar ME, London GM. Impact of aortic stiffness attenuation on survival of patients in end-stage renal failure. *Circulation.* 2001;103:987-992.
- Guerin AP, London GM, Marchais SJ, Metivier F. Arterial stiffening and vascular calcifications in end-stage renal disease. *Nephrol Dial Transplant.* 2000;15:1014-1021.
- Herzog CA, Ma JZ, Collins AJ. Poor long-term survival after acute myocardial infarction among patients on long-term dialysis. *N Engl J Med.* 1998;339:799-805.
- Himmelfarb J, Hakim RM. Oxidative stress in uremia. *Curr Opin Nephrol Hypertens.* 2003;12:593-598.
- Iseki K, Kawazoe N, Fukiyama K. Serum albumin is a strong predictor of death in chronic dialysis patients. *Kidney Int.* 1993;44:115-119.
- Iseki K, Miyasato F, Tokuyama K, Nishime K, Uehara H, Shiohira Y, Sunagawa H, Yoshihara K, Yoshi S, Toma S, Kowatari T, Wake T, Oura T, Fukiyama K. Low diastolic blood pressure, hypoalbuminemia, and risk of death in a cohort of chronic hemodialysis patients. *Kidney Int.* 1997;51:1212-1217.
- Joannides R, Haefeli WE, Linder L, Richard V, Bakkali EH, Thuillez C, Luscher TF. Nitric oxide is responsible for flow-dependent dilatation of human peripheral conduit arteries in vivo. *Circulation.* 1995;91:1314-1349.
- Kalantar-Zadeh K, Block G, Humphreys MH, McAllister CJ, Kopple JD. A low, rather than a high, total plasma homocysteine is an indicator of poor outcome in hemodialysis patients. *J Am Soc Nephrol.* 2004a;15:442-453.
- Kalantar-Zadeh K, McAllister CJ, Lehn RS, Liu E, Kopple JD. A low serum iron level is a predictor of poor outcome in hemodialysis patients. *Am J Kidney Dis.* 2004b;43:671-684.
- Kalousova M, Sulkova S, Fialova L, Soukupova J, Malbohan IM, Spacek P, Braun M, Mikulikova L, Fortova M, Horejsi M, Tesar V, Zima T. Glycooxidation and inflammation in chronic haemodialysis patients. *Nephrol Dial Transplant.* 2003;18:2577-2581.
- Kamata H, Hirata H. Redox regulation of cellular signalling. *Cell Signal.* 1999;11:1-14.
- Karamanoglu M, O'Rourke MF, Avolio AP, Kelly RP. An analysis of the relationship between central aortic and peripheral upper limb pressure waves in man. *Eur Heart J.* 1993;14:160-167.
- Kelly R, Hayward C, Avolio A, O'Rourke M. Noninvasive determination of age-related changes in the human arterial pulse. *Circulation.* 1989;80:1652-1659.
- Klassen PS, Lowrie EG, Reddan DN, DeLong ER, Coladonato JA, Szczech LA, Lazarus JM, Owen WF Jr. Association between pulse pressure and mortality in patients undergoing maintenance hemodialysis. *JAMA.* 2002;287:1548-1555.
- Kosch M, Levers A, Barenbrock M, Matzkies F, Schaefer RM, Kisters K, Rahn KH, Hausberg M. Acute effects of haemodialysis on endothelial function and large artery elasticity. *Nephrol Dial Transplant.* 2001;16:1663-1668.
- Lassegue B, Sorescu D, Szocs K, Yin Q, Akers M, Zhang Y, Grant SL, Lambeth JD, Griendling KK. Novel gp91(phox) homologues in vascular smooth muscle cells: nox1 mediates angiotensin II-induced superoxide formation and redox-sensitive signaling pathways. *Circ Res.* 2001;88:888-894.
- Locatelli F, Aljama P, Barany P, Canaud B, Carrera F, Eckardt KU, Horl WH, Macdougall IC, Macleod A, Wiecek A, Cameron S; European Best Practice Guidelines

- Working Group. Revised European best practice guidelines for the management of anaemia in patients with chronic renal failure. *Nephrol Dial Transplant*. 2004;19:1-47.
- Locatelli F, Marcelli D, Conte F, D'Amico M, DelVecchio L, Limido A, Malberti F, Spotti D. Cardiovascular disease in chronic renal failure: the challenge continues. Registro Lombardo Dialisi e Trapianto. *Nephrol Dial Transplant*. 2000;15:69-80.
- London GM, Blacher J, Pannier B, Guerin AP, Marchais SJ, Safar ME. Arterial wave reflections and survival in end-stage renal failure. *Hypertension*. 2001;38:434-438.
- Mallick NP, Gokal R. Haemodialysis. *Lancet*. 1999;353:737-742.
- Marchais SJ, Guerin AP, Pannier BM, Delavaud G, London GM. Arterial compliance and blood pressure. *Drugs*. 1993;46:82-87.
- McVeigh GE, Bratteli CW, Morgan DJ, Alinder CM, Glasser SP, Finkelstein SM, Cohn JN. Age-related abnormalities in arterial compliance identified by pressure pulse contour analysis: aging and arterial compliance. *Hypertension*. 1999;33:1392-1398.
- Miura H, Wachtel RE, Liu Y, Loberiza FR Jr, Saito T, Miura M, Gutterman DD. Flow-induced dilation of human coronary arterioles: important role of Ca(2+)-activated K(+) channels. *Circulation*. 2001;103:1992-1998.
- Miyazaki H, Matsuka H, Itabe H, Usui M, Ueda S, Okuda S, Imaizumi T. Hemodialysis impairs endothelial function via oxidative stress: effects of vitamin E-coated dialyzer. *Circulation*. 2000;101:1002-1006.
- Millasseau SC, Kelly RP, Ritter JM, Chowienczyk PJ. The vascular impact of aging and vasoactive drugs: comparison of two digital volume pulse measurements. *Am J Hypertens*. 2003;16:467-472.
- Nichols WW, O'Rourke MF. Sphygmocardiography. In: Nichols WW, O'Rourke MF, eds. *McDonald's Blood Flow in Arteries*. London, England: Arnold; 1998:450-477.
- Noris M, Todeschini M, Casiraghi F, Roccatello D, Martina G, Minetti L, Imberti B, Gaspari F, Atti M, Remuzzi G. Effect of acetate, bicarbonate dialysis, and acetate-free biofiltration on nitric oxide synthesis: implications for dialysis hypotension. *Am J Kidney Dis*. 1998;32:115-124.
- Nurnberger J, Keflioglu-Scheiber A, Opazo Saez AM, Wenzel RR, Philipp T, Schafers RF. Augmentation index is associated with cardiovascular risk. *J Hypertens*. 2002;20:2407-2414.
- Olesen SP, Clapham DE, Davies PF. Haemodynamic shear stress activates a K⁺ current in endothelial cells. *Nature*. 1988;331:168-170.
- Outinen PA, Sood SK, Pfeifer SI, Pamidi S, Podor TJ, Li J, Weitz JI, Austin RC. Homocysteine-induced endoplasmic reticulum stress and growth arrest leads to specific changes in gene expression in human vascular endothelial cells. *Blood*. 1999;94:959-967.
- Owen WF Jr, Lew NL, Liu Y, Lowrie EG, Lazarus JM. The urea reduction ratio and serum albumin concentration as predictors of mortality in patients undergoing hemodialysis. *N Engl J Med*. 1993;329:1001-1006.
- Parfrey PS. Cardiac disease in dialysis patients: diagnosis, burden of disease, prognosis, risk factors and management. *Nephrol Dial Transplant*. 2000;15:58-68.
- Parthasarathy S, Santanam N, Ramachandran S, Meilhac O. Oxidants and antioxidants in atherogenesis: an appraisal. *J Lipid Res*. 1999;40:2143-2157.
- Pastan S, Bailey J. Dialysis therapy. *N Engl J Med*. 1998;338:1428-1437.
- Pohl U, Holtz J, Busse R, Bassenge E. Crucial role of the endothelium in the vasodilator response to flow in vivo. *Hypertension*. 1985;8:37-44.
- Pressman GL, Newgard PM. A transducer for the continuous external measurement of arterial blood pressure. *IEEE Trans Biomed Eng*. 1963;10:73-81.

- Raza K, Thambyrajah J, Townend JN, Exley AR, Hortas C, Filer A, Carruthers DM, Bacon PA. Suppression of inflammation in primary systemic vasculitis restores vascular endothelial function: lesson for atherosclerotic disease? *Circulation*. 2000;102:1470-1472.
- Rietzschel ER, Boeykens E, De Buyzere ML, Duprez DA, Clement DL. A comparison between systolic and diastolic pulse contour analysis in the evaluation of arterial stiffness. *Hypertension*. 2001;37:15-22.
- Safar ME. Augmentation Index, Cardiovascular Risk and Drug Treatment of Hypertension. *Symposium Report-New Strategies for Cardiovascular Disease Management: Blood-Pressure & Augmentation Index Monitoring*. 2003, Florida, USA.
- Safar ME, Blacher J, Pannier B, Guerin AP, Marchais SJ, Guyonvarc'h PM, London GM. Central pulse pressure and mortality in end-stage renal disease. *Hypertension*. 2002;39:735-738.
- Sarnak MJ, Levey AS, Schoolwerth AC, Coresh J, Culeton B, Hamm LL, McCullough PA, Kasiske BL, Kelepouris E, Klag MJ, Parfrey P, Pfeffer M, Raij L, Spinosa DJ, Wilson PW; American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation*. 2003;108:2154-2169.
- Sato T, Nishinaga M, Kawamoto A, Ozawa T, Takatsuji H. Accuracy of a continuous blood pressure monitor based on arterial tonometry. *Hypertension*. 1993;21:866-874.
- Saw J, Levin A, Gin K. Coronary artery disease in chronic kidney disease patients: assessing the evidence for diagnosis, screening and revascularization. *Can J Cardiol*. 2004;20:807-813.
- Smulyan H, Marchais SJ, Pannier B, Guerin AP, Safar ME, London GM. Influence of body height on pulsatile arterial hemodynamic data. *J Am Coll Cardiol*. 1998;31:1103-1109.
- Stenvinkel P. Inflammatory and atherosclerotic interactions in the depleted uremic patient. *Blood Purif*. 2001;19:53-61.
- Stenvinkel P, Barany P. Anaemia, rHuEPO resistance, and cardiovascular disease in end-stage renal failure; links to inflammation and oxidative stress. *Nephrol Dial Transplant*. 2002;17:32-37.
- Takazawa K, Tanaka N, Fujita M, Matsuoka O, Saiki T, Aikawa M, Tamura S, Ibukiyama C. Assessment of vasoactive agents and vascular aging by the second derivative of photoplethysmogram waveform. *Hypertension*. 1998;32:365-370.
- Tarng DC, Huang TP, Liu TY, Chen HW, Sung YJ, Wei YH. Effect of vitamin E-bonded membrane on the 8-hydroxy 2'-deoxyguanosine level in leukocyte DNA of hemodialysis patients. *Kidney Int*. 2000;58:790-799.
- Tepel M. Oxidative stress - does it play a role in the genesis of essential hypertension and hypertension of uremia? *Nephrol Dial Transplant*. 2003;18:1439-1442.
- Tepel M, Echelmeyer M, Orie NN, Zidek W. Increased intracellular reactive oxygen species in patients with end-stage renal failure: effect of hemodialysis. *Kidney Int*. 2000;58:867-872.
- Tepel M, van der Giet M, Statz M, Jankowski J, Zidek W. The antioxidant, acetylcysteine, reduces cardiovascular events in patients with end-stage renal failure: a randomized controlled trial. *Circulation*. 2003;107:992-995.

- van Guldener C, Janssen MJ, Lambert J, Steyn M, Donker AJ, Stehouwer CD. Endothelium-dependent vasodilation is impaired in peritoneal dialysis patients. *Nephrol Dial Transplant*. 1998;13:1782-1786.
- Vaziri ND, Ni Z, Oveisi F, Liang K, Pandian R. Enhanced nitric oxide inactivation and protein nitration by reactive oxygen species in renal insufficiency. *Hypertension*. 2002;39:135-141.
- Wilkinson IB, MacCallum H, Flint L, Cockcroft JR, Newby DE, Webb DJ. The influence of heart rate on augmentation index and central arterial pressure in humans. *J Physiol*. 2000;525:263-270.
- Winer N, Sowers JR, Weber MA. Gender differences in vascular compliance in young, healthy subjects assessed by pulse contour analysis. *J Clin Hypertens*. 2001;3:145-152.
- Wolin MS. Interactions of oxidants with vascular signaling systems. *Arterioscler Thromb Vasc Biol*. 2000;20:1430-1442.
- Yasmin, Brown MJ. Similarities and differences between augmentation index and pulse wave velocity in the assessment of arterial stiffness. *QJM*. 1999; 92:595-600.
- Yeun JY, Kaysen GA. C-reactive protein, oxidative stress, homocysteine, and troponin as inflammatory and metabolic predictors of atherosclerosis in ESRD. *Curr Opin Nephrol Hypertens*. 2000;9:621-630.
- Zoccali C, Bode-Boger S, Mallamaci F, Benedetto F, Tripepi G, Malatino L, Cataliotti A, Bellanuova I, Fermo I, Frolich J, Boger R. Plasma concentration of asymmetrical dimethylarginine and mortality in patients with end-stage renal disease: a prospective study. *Lancet*. 2001;358:2113-2117.