

## 7 Literaturverzeichnis

1. Abuhamad AZ, Shaffer W, Mari G, Copel JA, Hobbins JC, Evans AT.  
Single umbilical artery: does it matter which artery is missing?  
Am J Obstet Gynecol 173 (1995) 3 Pt 1: 728-32
2. Achiron R, Lipitz S, Gabbay U, Yagel S.  
Prenatal ultrasonographic diagnosis of fetal heart echogenic foci: no correlation with Down syndrome.  
Obstet Gynecol 89 (1997): 945-8
3. Alfirevic Z.  
Early amniocentesis versus transabdominal chorion villus sampling for prenatal diagnosis.  
Cochrane Database Syst Rev 2000; (2): CD000077.
4. Alfirevic Z, Sundberg K, Brigham S.  
Amniocentesis and chorionic villus sampling for prenatal diagnosis.  
Cochrane Database Syst Rev 2003; (3): CD003252
5. Ammala P, Hiilesmaa VK, Liukkonen S, Saisto T, Teramo K, von Koskull H.  
Randomized trial comparing first-trimester transcervical chorionic villus sampling and second-trimester amniocentesis.  
Prenat Diagn 13 (1993) 10: 919-27
6. Apodaca CC, Hume RF; Evans WJ, Martin LS, Evans MI, Calhoun BC.  
Parental decision-making differences between patients in two health care systems for choroid plexus cysts.  
Fetal Diagn Ther 2000 Nov-Dec; 15(6):338-41

7. Becker R, Fuhrmann W, Holzgreve W, Sperling K.

Pränatale Diagnostik und Therapie.

Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart (1995)

8. Blumberg BD, Golbus MS, Hanson KH.

The psychological sequelae of abortion performed for a genetic indication.

Am J Obstet Gynecol 122 (1975) : 799

9. Bollmann R, Chaoui R, Zienert A, Korner H.

Choroid plexus cysts in the 2d trimester- an indication for trisomie 18.

Zentralbl Gynakol 114 (1992)4:171-4

10. Brand-Saberi B, Floel H, Christ B, Schulte-Vallentin M, Schindler H.

Alterations of the fetal extracellular matrix in the nuchal oedema of Down's syndrome.

Ann Anat 1994 Dec; 176 (6): 539-47.

11. Bromley B, Liebermann E, Laboda L, Benacerraf B.

Echogenic intracardiac focus: a sonographic sign for fetal Down syndrome.

Obstet Gynecol 86 (1995): 998-1001

12. Bronshtein M, Zimmer EZ, Blazer S.

A characteristic cluster of fetal sonographic markers that are predictive of fetal Turner syndrome in early pregnancy.

Am J Obstet Gynecol 2003 Apr; 188(4): 1016-20

13. Brumback BA, Holmes LB, Ryan LM.

Adverse effects of chorionic villus sampling: a meta-analysis.

Statistics in Medicine 1999, 18 (16): 2163-75

14. Brun L, Dufour P, Savary JB, Valat AS, Boute O, Subtil D, Vaast P, Bourgeot P, Manouvrier S, de Martinville B, Puech F.

Trisomie 18: ultrasound aspects. Report of 40 cases.

Presse Med 2000 Dec 9; 29 (38): 2082-6

15. Brun JL, Saura R, Horovitz J, Maugey B, Taine L, Roux D, Vergnaud A.

First trimester diagnosis of fetal nuchal edema. Report of 29 cases.

Fetal Diagn Ther 9 (1994)4: 246-51

16. Budorick NE, Kelly TF, Dunn JA, Scioscia AL.

The single umbilical artery in a high-risk patient population: what should be offered?

J Ultrasound Med 2001 Jun; 20 (6): 619-27

17. Byrne J, Blanc WA.

Malformations and chromosome anomalies in spontaneously aborted fetuses with single umbilical artery.

Am J Obstet Gynecol 153 (1985): 340-3

18. Campbell S, Johnstone FD, Holm EM et al..

Anencephaly: early ultrasonic diagnosis and active management.

Lancet 2 (1972): 1226

19. Caughey AB, Lyell DJ, Filly RA, Washington AE, Norton ME.

The impact of the use of the isolated echogenic intracardiac focus as a screen for Down syndrome in women under the age of 35 years.

Am J Obstet Gynecol 2001 Nov; 185 (5): 1021-7

20. Coco C, Jeanty P, Jeanty C.

An isolated echogenic heart focus is not an indication for amniocentesis in 12672 unselected patients (In Process Citation).

J Ultrasound Med 2004 Apr; 23 (4): 489-96

21. Cowan RS.

Aspects of the History of Prenatal Diagnosis.

Fetal Diagn Ther (1993) Apr 8 Suppl 1: 10-7

22. Daffos F, Capella-Povolsky M, Forestier F.

Fetal blood sampling during pregnancy with use of a needle guided by ultrasound: a study of 606 consecutive cases.

Am J Obstet Gynecol 153 (1985): 665

23. Daffos F, Forestier F, Capella-Povolsky M et al..

Fetal blood sampling via the umbilical cord using a needle guided by ultrasound: a report of 66 cases.

Prenat Diagn 3 (1983): 271

24. De Vore GR.

Second trimester ultrasonography may identify 77 to 97% of fetuses with trisomy 18.

J Ultrasound Med 2000 Aug; 19(8):565-76

25. Egan JF.

The role of ultrasound in screening for fetal Down syndrome.

Curr Womens Health Rep 2002 Feb; 2(1):52-8

26. Evans MI, Hume RF, Johnson MP, Treadwell MC, Krivchenia EL, Zador IE, Sokol RJ.

Integration of genetics and ultrasonography in prenatal diagnosis: just looking is not enough.

Am J Obstet Gynecol 174 (1996) 6: 1925-31;  
discussion 1931-3

27. Falk S., Stutte HJ.

Extrazelluläre Veränderungen. In: Pathologie.

Hrsg.: Blümcke S. Walter de Gruyter-Verlag, Berlin, New York (1995)

28. Fletcher JC.

Ethical issues in clinical trials of first trimester prenatal diagnosis.

In: Brambati B, Simoni G (eds) Chorionic villus sampling.

Basel, p2 (1986)

29. Fotuny A, Borrell A, Soler A, Casals E, Costa D, Carrio A, Puerto B, Seres A, Cararach J, Delgado R.

Chorionis villus sampling by biopsie forceps. Result of 1580 procedures from a single centre.

Prenat Diagn 15 (1995) 6: 541-50

30. Fuchs KM, Peipert, JF.

First trimester Down syndrome screening: public health implications.

Semin Perinatol. 2005 Aug, 29 (4): 267-71

31. Gabrielli S, Reece EA, Piiliu G, Perolo A, Rizzo N, Bovicelli L, Hobbins JC.

The clinical significance of prenatally diagnosed choroid plexus cysts.

Am J Obstet Gynecol 160 (1989) 5 Pt 1: 1207-10

32. Gagnon S, Fraser W, Fouquette B, Bastide A, Bureau M, Fontaine JY, Huot C.  
Nature and frequency of chromosomal abnormalities in pregnancies with abnormal ultrasound findings: an analysis of 117 cases with review of the literature.  
*Prenat Diagn* 12 (1992) 1: 9-18
33. Geipel A, Germer U, Welp T, Schwinger E, Gembruch U.  
Prenatal diagnosis of single umbilical artery: determination of the absent side, associated anomalies, Doppler findings and perinatal outcome.  
*Ultrasound Obstet Gynecol* 2000 Feb; 15 (2): 114-7
34. Ghidini A, Strobelt N, Locatelli A, Mariani E, Piccoli MG, Vergani P.  
Isolated fetal choroid plexus cysts: role of ultrasonography in establishment of the risk of trisomy 18.  
*Am J Obstet Gynecol* 2000 Apr; 182(4):972-7
35. Gonen R, Dar H, Degani S.  
The karyotype of fetuses with anomalies detected by second trimester ultrasonography.  
*Eur J Obstet Gynecol Reprod Biol* 58 (1995)2:153-5
36. Grati FR, Grimi B, Frascoli G, Di Meco AM, Liuti R, Milani S, Trotta A, Dulcetti F, Grosso E, Miozzo M, Maggi F, Simoni G.  
Confirmation of mosaicism and uniparental disomy in amniocytes, after detection of mosaic chromosome abnormalities in chorionic villi.  
*Eur J Hum Genet* 2006 Mar; 14 (3): 282-8
37. Hahnemann N, Mohr J.  
Genetic diagnosis in the embryo by means of biopsy from extra embryonic membranes.  
*Bull Eur Soc Hum Genet* 2 (1968): 23

38. Hahnemann JM, Vejerslev LO.

Accuracy of cytogenetic findings on chorionic villus sampling (CVS)—diagnostic consequences of CVS mosaicism and non-mosaic discrepancy in centres contributing to EURCROMIC 1986-1992.

Prenat Diagn 1997 Sep, 17 (9): 801-20

39. Hobbins JC, Lezotte DC, Persutte WH, De Vorer GR, Benacerraf BR, Nyberg DA, Vintzileos AM, Platt LD, Carlson DE, Bahado-Singh RO, Abu hamad AZ.

An 8-center study to evaluate the utility of mid-term genetic sonograms among high-risk pregnancies.

J Ultrasound Med 2003 Jan; 22(1):33-8

40. Holzgreve W.

Pränatale Medizin.

Thieme-Verlag, Stuttgart- New York (1987)

41. Holzgreve W, Louwen F.

Schwangerenbetreuung. In: Frauenheilkunde und Geburtshilfe .

Hrsg.: Dudenhausen JW, Schneider HPG. Walter de Gruyter-Verlag, Berlin, New York (1994)

42. Huggon IC, Cook AC, Simpson JM, Smeeton NC, Sharland GK.

Isolated echogenic foci in the fetal heart as marker of chromosomal abnormality.

Ultrasound Obstet Gynecol 2001 Jan; 17 (1): 11-6

43. International Fetoscopy Group.

Special report: the status of fetoscopy and fetal tissue sampling.

Prenat Diagn 4 (1984) : 79

44. Kainer F.

Intrauterine Mangelentwicklung. In: Frauenheilkunde und Geburtshilfe.

Hrsg.: Dudenhausen JW, Schneider HPG.

Walter de Gruyter-Verlag, Berlin, New York (1994)

45. von Kaisenberg CS, Brand-Saberi B, Christ B, Vallian S, Farzaneh F, Nicolaides KH.

Collagen type VI expression in the skin of trisomy 21 fetuses.

Obstet Gynecol 1998 Mar; 91 (3): 319-23.

46. Khong TY, George K.

Chromosomal abnormalities associated with a single umbilical artery.

Prenat Diagn 12 (1992) 11: 965-8

47. Kohler H.

Fehlende Nabelschnurarterie (solitäre Nabelschnurarterie, SNA).

In: Die Plazenta des Menschen.

Hrsg.: Becker V, Schiebler ThH, Kubli F.

Georg Thieme-Verlag, Stuttgart (1981)

48. Kuliev A, Jackson L, Froster U, Brambati B, Simpson JL, Verlinski Y, Ginsberg N, Smidt-Jensen S, Zakut H.

Chorionic villus sampling safety. Report of World Health

Organization/ EURO meeting in association with the Seventh

International Conference on Early Prenatal Diagnosis of

Genetic Diseases, Tel-Aviv, Israel, May 21, 1994.

Am J Obstet Gynecol 174 (1996) 3: 807-11

49. Kuliev AM, Modell B, Jackson L, Simpson B, Brambati B, Rhoads G, Froster U, Verlinsky Y, Smidt-Jensen S, Holzgreve W, Ginsberg N, Ammala P, Dumez Y.

Risk evaluation of CVS.

Prenat Diagn 13 (1993): 197-209

50. Kupferminc MJ, Tamura RK, Sabbagh RE, Parilla BV, Cohen LS, Pergament E.

Isolated choroid plexus cyst(s): an indication for amniocentesis.

Am J Obstet Gynecol 171 (1994)4:1068-71

51. Lau KT, Leung YT, Chan LW, Sahota DS, Leung NT.

Outcome of 1355 consecutive transabdominal chorionic villus sampling in 1355 patients.

Chin Med J (Engl.) 2005 Oct 20; 118 (20): 1675-81

52. Levy DW, Mintz MC.

Technical note. The left ventricular echogenic focus: a normal finding.

A J R 150 (1988): 85-6

53. Monney C, Pescia G, Addor MC.

Turner Syndrome.

Schweiz Med Wochenschr 2000 Sep 23; 130(38): 1339-43

54. Moran CJ, Tay JB, Morrison JJ.

Ultrasound detection and perinatal outcome of fetal trisomies 21, 18 and 13 in the absence of a routine fetal anomaly scan or biochemical screening.

Ultrasound obstet Gynecol 2002 Nov; 20 (5): 482-5

55. Nanal R, Kyle P, Soothill PW

A classification of pregnancy losses after invasive prenatal diagnostic procedures: an approach to allow comparison of units with a different case mix.

Prenat Diagn 2003 Jun; 23 (6): 488-92]

56. Nicolaides KH, Snijders RJ; Gosden CM, Berry C, Campbell S.  
Ultrasonographically detectable markers of fetal chromosomal abnormalities (see comments).  
*Lancet* 340 (1992) 8821: 704-7
57. Nikkila A, Valentin L, Thelin A, Jorgensen C.  
Early amniocentesis and congenital foot deformities.  
*Fetal Diagn Ther* 2002 May-Jun; 17 (3): 129-32.
58. Nyberg DA, Mahony BS, Luthy D, Kaspur R.  
Single umbilical artery. Prenatal detection of concurrent anomalies.  
*J Ultrasound Med* 10 (1991) 5: 247-53
59. Obladen M.  
Versorgung des Neugeborenen mit gestörter Adaptation.  
In: Frauenheilkunde und Geburtshilfe.  
Hrsg.: Dudenhausen JW, Schneider HPG.  
Walter de Gruyter-Verlag, Berlin, New York (1994)
60. Papp C, Beke A, Mezei G, Toth-Pal E, Papp Z.  
Chorionic villus sampling: a 15-year experience.  
*Fetal Diagn Ther* 2002 Jul-Aug; 17 (4): 218-27
61. Pavriilla BV, Tamura RK, MacGregor SN, Geibel LJ, Sabbagh RE.  
The clinical significance of a single umbilical artery as an isolated finding on prenatal ultrasound.  
*Obstet Gynecol* 85 (1995): 570-2
62. Petrikovski BM, Challenger M, Wyse LJ.  
Natural history of echogenic foci within ventricles of the fetal heart.

Ultrasound Obstet Gynecol 5 (1995) 2: 92-4

63. Platt LD, Carlson DE, Medeaaris AL, Walla CA.

Fetal choroid plexus cysts in the second trimester of pregnancy: a cause for concern (see comments).

Am J Obstet Gynecol 164 (1991)6Pt1:1652-5, discussion 1655-6

64. Prucka S, Clemens M, Craven C, McPherson E.

Single umbilical artery: what does it mean for the fetus? A case-control analysis of pathologically ascertained cases (In Process Citation).

Genet Med 2004 Jan-Feb; 6 (1): 54-7

65. Ranzini AC, Guzman ER, Ananth CV, Day-Salvatore D, Fisher AJ, Vintzileos AM.

Sonographic identification of fetuses with Down syndrome in the third trimester: a matched control study.

Obstet Gynecol 1999 May; 93 (5Pt1):702-6

66. Rembouskos G, Cicero S, Longo D, Sacchini C, Nicolaides KH.

Single umbilical artery at 11-14 weeks` gestation: relation to chromosomal defects.

Ultrasound Obstet Gynecol 2003 Dec; 22 (6): 567-70

67. Riis P, Fuchs F.

Antenatal determination of foetal sex in prevention  
of hereditary diseases.

Lancet 2 (1966): 180-2

68. Schechter AG, Fakhry J, Shapiro LR, Gewitz MH.

In utero thickening of the chordae tendineae. A cause of intracardiac echogenic foci.

J Ultrasound Med 6 (1987) 12: 691-5

69. Scott F, Peters H, Boogert T, Robertson R, Anderson J, McLennan A, Kesby G, Edelman D.

The loss rate for invasive prenatal testing in a specialised obstetric ultrasound practice.

Aust N Z J Obstet Gynaecol 2002 Feb; 42 (1): 55-8

70. Snijders RJM, Nicolaides KH.

Ultrasound markers for fetal chromosomal defects

The Parthenon Publishing Group, New York-London (1996)

71. Sohn C, Gast AS.

Hygroma colli (cervical neck edema) as an ultrasonographic parameter of fetal chromosome anomalies.

Zentralbl Gynakol 119 (1997) 5: 232-6

72. Sohn C, Holzgreve W.

Ultraschall in Gynäkologie und Geburtshilfe.

Springer-Verlag, Berlin- Heidelberg- New York- London-

Paris- Tokyo (1995)

73. Sotiriadis A, Makrydimas G, Ioannidis JP.

Diagnostik performance of intracardiac echogenic foci for Down syndrome: a meta-analysis.

Obstet Gynecol 2003 May; 101 (5Pt1): 1009-16

74. Tanriverdi HA, Hendrik HJ, Ertan AK, Axt R, Schmidt W.

Hygroma colli cysticum: prenatal diagnosis and prognosis.

Am J Perinatol 2001 Dec; 18(8): 415-20

75. Tercanli S, Holzgreve W, Batukan C, Gerber A, Ermis H, Miny P.

Screening for aneuploidy by first trimester nuchal translucency measurement: result from a prospective trial including 1980 cases in a single center in Switzerland.

Ultraschall Med 2002 Feb; 23 (1): 22-6

76. Tietung Group, Anshan, China.

Fetal sex prediction by sex chromatin of chorionic villi cells during early pregnancy.

China Med J 1 (1975): 117

77. Yilmaztürk A, Deppermann F.

Hygroma colli- sonographic indication of fetal chromosome abnormality. A case report (see comments).

Zentralbl Gynakol 113 (1991) 15-16: 883-7

78. Zienert A, Bollmann R, Chaoui R, Bartho S.

Single umbilical artery- consequences of prenatal diagnosis.

Zentralbl Gynakol 114 (1992) 3: 131-5

79. Zoppi MA, Ibba RM, Floris M, Manca F, Axiana C, Monni G.

Changes in nuchal translucency thickness in normal and abnormal karyotype fetuses.

BJOG 2003 Jun; 110 (6): 5584-8

80. Zoppi MA, Ibba RM, Floris M, Monni G.

Fetal nuchal translucency screening in 12495 pregnancies in Sardinia.

Ultrasound Obstet Gynecol 2001 Dec; 18(6): 649-51