

5 Diskussion

- 5.1 Vom Röntgenbild zum Operationsroboter:
Navigationstechniken in der Neurochirurgie
- 5.2 Sensor-basierte elektromagnetische Navigation: Pro und Kontra
- 5.3 Einsatzgebiete und Indikationen
- 5.4 Probleme, Fehlerquellen und Genauigkeiten
- 5.5 Ausblick

6 Schlussfolgerungen

7 Technische Begriffe und Abkürzungen

8 Literaturverzeichnis

9 Anhang

- Patienteninformation - Seite 1 und 2 (zu Kapitel 4.1)
- Patienteneinverständniserklärung (zu Kapitel 4.1)
- Erklärung zum Datenschutz (zu Kapitel 4.1)
- Statistische Ergebnisse zu den Klinischen Anwendungen (zu Kapitel 4.3)
- Legende zu den Navigationsprotokollen (zu Kapitel 4.3)

10 Danksagung

8 Literaturverzeichnis

1. Abe M, Udono H, Tabuchi K, et al. (2001): Transsphenoidal surgery assisted by navigation system. *No Shinkei Geka* 29: 31-38.
2. Alberstone CD, Skirboll SL, Benzel EC, Sanders JA, Hart BL, Baldwin NG, Tessman CL, Davis JT, Lee RR (2000): Magnetic source imaging and brain surgery: presurgical and intraoperative planning in 26 patients. *J Neurosurg* 92: 79-90.
3. Alberti O, Dorward NL, Kitchen ND, Thomas DG (1997): Neuronavigation - impact on operating time. *Stereotact Funct Neurosurg* 68: 44-48.
4. Alonso-Vanegas MA, Olivier A, Quesney LF (1998): Applications of image-guided surgery to intraoperative electrophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 48: 140-156.
5. Alp MS, Dujovny M, Misra M, Charbel FT, Ausman JI (1998): Head registration techniques for image-guided surgery. *Neurol Res* 20: 31-37.
6. Bale RJ, Burtscher J, Eisner W, Obwegeser AA, Riegr M, Sweeney RA, Dessl A, Giacomuzzi SM, Twerdy K, Jaschke W (2000): Computer assisted neurosurgery by using a noninvasive vacuum-affixed dental cast that acts as a reference base: another Step towards a unified approach in the treatment of brain tumors. *J Neurosurg* 93: 208-213.
7. Bale RJ, Vogele M, Freysinger W, Gunkel AR, Martin A, Bumm K, Thumfart WF (1997): Minimally invasive head holder to improve the performance of frameless stereotactic surgery. *Laryngoscope* 107: 373-377.
8. Bale RJ, Vogele M, Martin A, Auer T, Hensler E, Eichberger P, Freysinger W, Sweeney R, Gunkel AR, Lukas PH (1997): VBH head holder to improve frameless stereotactic brachytherapy of cranial tumors. *Comput Aided Surg* 2: 286-291.
9. Barnett GH, Kormos DW, Steiner PC, Weisenberger J (1993): Use of a frameless, armless stereotactic wand for brain tumor localization with two-dimensional and three-dimensional neuroimaging. *Neurosurgery* 33: 674-678.
10. Barnett GH, McKenzie RL, Ramos L, Palmer J (1993): Nonvolumetric Stereotaxy-assisted craniotomy. Results in 50 consecutive cases. *Stereotac Funct Neurosurg* 61: 80-95.
11. Barnett GH, Miller DW, Weisenberger J (1999): Frameless stereotaxy with scalp-applied fiducial markers for brain biopsy procedures: experience in 218 cases. *J Neurosurg* 91: 569-576.
12. Barnett GH, Steiner CP, Weisenberger J (1995): Intracranial meningioma resection using frameless stereotaxy. *J Image Guid Surg* 1: 46-52.
13. Barnett GH, Steiner CP, Weisenberger J (1996): Target and trajectory guidance for interactive surgical navigation systems. *Stereotact Funct Neurosurg* 66: 91-95.

14. Benabid AL, Cinquin P, Lavalley S, Le Bas JF, Demongeot J, de Rougemont J (1987): Computer-driven robot for stereotactic surgery connected to CT scan and magnetic resonance imaging. Technological design and preliminary results. *Appl Neurophysiol* 50: 153-154.
15. Benabid AL, Hoffmann D, Le Bas JF, Lavalley S (1995): [Value of image guided neurosurgery in neuro-oncology]. *Bull Cancer* 82 Suppl 5: 573-580.
16. Berger MS (1994): Lesions in functional ("eloquent") cortex and subcortical white matter. *Clin Neurosurg* 41: 444-63.
17. Berger MS, Cohen WA, Ojemann GA (1990): Correlation of motor cortex brain mapping data with magnetic resonance imaging. *J Neurosurg* 72: 383-387.
18. Berger MS, Kincaid J, Ojemann GA et al. (1989): Brain mapping techniques to maximize resection, safety, and seizure control in children with brain tumors. *Neurosurgery* 25: 786-792.
19. Berkenstadt H, Perel A, Hadani M, Unofrievich I, Ram Z (2001): Monitored anesthesia care using remifentaniol and propofol for awake craniotomy. *J Neurosurg Anesthesiol* 13: 246-249.
20. Black PML, Alexander E III, et al. (1999): Craniotomy for tumor treatment in an intraoperative magnetic resonance imaging unit. *Neurosurgery* 45: 423-433.
21. Bradbury J (2004): Shine the magnetic resonance light on neurosurgery. *Lancet Oncol* 5: 643.
22. Braun V, Dempf S, Tomczak R, Wunderlich A, Weller R, Richter HP (2000): Functional cranial neuronavigation. Direct integration of fMRI and PET data. *J Neuroradiol* 27:157-163.
23. Brommeland T, Hennig R. (2000): A new procedure for frameless computer navigated stereotaxy. *Acta Neurochir (Wien)* 142: 443-447.
24. Brown RA (1979): A computerized tomography-computer graphics approach to stereotaxic localization. *J Neurosurg* 50: 715-720.
25. Bucholz R, Macneil W, Fewings P, Ravindra A, McDurmont L, Baumann C (2000): Automated rejection of contaminated surface measurements for improved surface registration in image guided neurosurgery. *Stud Health Technol Inform* 70: 39-45.
26. Bucholz RD (1993): The central sulcus and surgical planning. *AJNR Am J Neuroradiol* 14: 926-927.
27. Bullard DE, Nashold BS (1995): Evolution of principles of stereotactic neurosurgery. *Neurosurg Clin N Am* 6: 27-41.
28. Cappabianca P, de Divitiis E (2003): Image guided endoscopic transnasal removal of recurrent pituitary adenomas. *Neurosurgery*, 52: 483-484.
29. Castilla JM, Martin V, Fernandez-Arconada O, Delgado P, Rodriguez-Salazar A (2003): [First steps in neuronavigation]. *Neurocirugia (Astur)* 14: 398-408.

30. Caversaccio M, Zulliger D, Bachler R, et al (2000): Practical aspects for optimal registration (matching) on the lateral skull base with an optical frameless computer-aided pointer system. *Am J Otol* 21: 863-870.
31. Cedzich C, Taniguchi M, Schäfer S, Schramm J (1996): Somatosensory evoked potential phase reversal and direct motor cortex stimulation during surgery in and around the central region. *Neurosurgery* 38: 962-970.
32. Chen HJ (2001): Clinical experiences in neuronavigation. *Stereotact Funct Neurosurg* 76: 145-147.
33. Coenen VA, Dammert S, Reinges MH, Mull M, Gilsbach JM, Rohde V (2005): Image-guided microneurosurgical management of small cerebral arteriovenous malformations: the value of navigated computed tomographic angiography. *Neuroradiology* 47: 66-72.
34. Comeau RM, Sadikot AF, Fenster A, Peters TM (2000): Intraoperative ultrasound for guidance and tissue shift correction in image-guided neurosurgery. *Med Phys* 27: 787-800.
35. Couldwell WT, Apuzzo ML (1990): Initial experience related to the use of the Cosman-Roberts-Wells stereotactic instrument: Technical note. *J Neurosurg* 72: 145-148.
36. Damadian R (1971): Tumor detection by nuclear magnetic resonance. *Science* 171: 1151-1163.
37. Dandy W (1918): Ventriculography following the injection of air into the cerebral ventricles. *Annals of Surgery* 70: 397-403.
38. Danks RA, Rogers M, Aglio LS, Gugino LD, Black PM (1998): Patient tolerance of craniotomy performed with the patient under local anesthesia and monitored conscious sedation. *Neurosurgery* 42: 28-34.
39. Derosier C, Delege G, Munier T, Pharaboz C, Cosnard G (1991): [MRI, geometric distortion of the image and stereotaxy]. *J Radiol* 72, 349-353.
40. Diethelm L, Henck F, Olsson O, Strnad F, Vieten H, Zuppinger A (1977). In *Handbuch der medizinischen Radiologie*. Band 14. Berlin, Heidelberg, Springer Verlag: 246.
41. Dohrmann GJ, Rubin JM (1985): Dynamic intraoperative imaging and instrumentation of brain and spinal cord using ultrasound. *Neurol Clin* 3: 425-437.
42. Dorward NL, Alberti O, Palmer JD, Kitchen ND, Thomas DG (1999): Accuracy of true frameless stereotaxy: in vivo measurement and laboratory phantom studies. Technical note. *J Neurosurg* 90: 160-8.
43. Dorward NL, Alberti O, Velani B, Gerritsen FA, Harkness WFJ, Kitchen ND, Thomas DGT (1998): Postimaging brain distortion: magnitude, correlates, and impact on neuronavigation. *J Neurosurg* 88: 656-662.
44. Du G, Zhou L, Mao Y (2003): Neuronavigator-guided glioma surgery. *Chin Med J* 116: 1484-1487.

45. Duffau H, Capelle L, Denvil D, et al (2003): The role of dominant premotor cortex in language: a study using intraoperative functional mapping in awake patients. *Neuroimage* 20: 1903-1914.
46. Duffau H, Capelle L, Denvil D, Sichez N, Gatignol P, Taillandier L, Lopes M, Mitchell MC, Roche S, Muller JC, Bitar A, Sichez JP, van Effenterre R (2003): Usefulness of intraoperative electrical subcortical mapping during surgery for low-grade gliomas located within eloquent brain regions: functional results in a consecutive series of 103 patients. *J Neurosurg* 98: 764-778.
47. Duffau H, Capelle L, Sichez N, Denvil D, Lopes M, Sichez JP, Bitar A, Fohanno D (2002): Intraoperative mapping of the subcortical language pathways using direct stimulations. An anatomo-functional study. *Brain* 125: 199-214.
48. Duffau H, Fontaine D (2005): Successful resection of a left insular cavernous angioma using neuronavigation and intraoperative language mapping. *Acta Neurochir (Wien)* 147: 205-208.
49. Dulou R, De Soultrait F, Blondet E, Dutertre G, Pernot P, Desgeorges M (2000): [Neuronavigation in third ventricle tumors]. *Neurochirurgie* 46: 282-285.
50. Ebel H, Ebel M, Schillinger G, et al (2000): Surgery of intrinsic cerebral neoplasms in eloquent areas under local anesthesia. *Minim Invasive Neurosurg* 43: 192-196.
51. Edwards PJ, Hill DL, Little JA, Hawkes DJ (1998): A three-component deformation model for image-guided surgery. *Med Image Anal* 2: 355-367.
52. Elias WJ, Chaddock JB, Alden TD, Laws ER Jr. (1999): Frameless stereotaxy for transsphenoidal surgery. *Neurosurgery* 45: 271-275.
53. Eljamel MS (1999): Developments in image-guided neurosurgery. *Hosp Med* 60: 686.
54. Enzmann DR, Irwin KM, Marshall WH, et al. (1984): Intraoperative sonography through a burr hole: guide for brain biopsy. *AJNR Am Neuroradiol* 5: 243-246.
55. Fahlbusch R, Ganslandt O, Nimsky C (2000): Intraoperative imaging with open magnetic resonance imaging and neuronavigation. *Childs Nerv Syst* 16: 829-831.
56. Farcot JM, Haberer JP, Laugner B, et al (1981): Transnasal and transsphenoidal neuroadenolyses with alcohol. Indications. Technics. Results. Remarks about action mechanism (27 cases). *Anesth Analg* 38: 361-364.
57. FitzGerald DB, Cosgrove GR, Ronner S, Jiang H, Buchbinder BR, Belliveau JW, Rosen BR, Benson RR (1997): Location of language in the cortex: a comparison between functional MR imaging and electrocortical stimulation. *AJNR Am J Neuroradiol* 18: 1529-1539.
58. Fitzpatrick JM, West JB (2001): The distribution of target registration error in rigid-body, point-based registration. *IEEE Transactions on Medical Imaging* 20: 917-927.
59. Fitzpatrick JM, West JB, Maurer CR (1998): Predicting error in rigid-body, point-based registration. *IEEE Transactions on Medical Imaging* 17: 694-702.

60. Flegel T, Podell M, March PA, et al (2002): Use of a disposable real-time CT stereotactic navigator device for invasive dog brain biopsy through a mini-burr hole. *AJNR Am J Neuroradiol* 23: 1160-1163.
61. Foerster O (1931): The cerebral cortex of man. *Lancet* 38: 309-312.
62. Fried I, Nenov V, Ojemann SG et al. (1995): Functional MR and PET imaging of rolandic and visual cortices for neurosurgical planning. *J Neurosurg* 83: 854-861.
63. Fukaya C, Katayama Y, Yoshino A, Kobayashi K, Kasai M, Yamamoto T (2001): Intraoperative wake-up procedure with propofol and laryngeal mask for optimal excision of brain tumour in eloquent areas. *J Clin Neurosci* 8: 253-255.
64. Galloway RL, Maciunas RJ, Edwards CA (1992): Interactive image-guided neurosurgery. *IEEE Trans Biomed Eng* 39: 1226-1231.
65. Ganslandt O, Steinmeier R, Kober H, Vieth J, Kassubek J, Romstock J, Strauss C, Fahlbusch R (1997): Magnetic source imaging combined with image-guided frameless stereotaxy: a new method in surgery around the motor strip. *Neurosurgery* 41: 621-627.
66. Gildenberg PL, Kaufman HH, Murthy KS (1982): Calculation of stereotactic coordinates from the computed tomographic scan. *Neurosurgery* 10: 580-586.
67. Giorni C, Casolino SD, Branzini A, Servello D, Passerini A, Brogli G, Plachino F (1989): Computer-assisted planning of stereotactic neurosurgical procedures. *Childs Nerv Syst* 5: 299-302.
68. Golfinos JG, Fitzpatrick BC, Smith LR, Spetzler RF (1995): Clinical use of a frameless stereotactic arm: results of 325 cases. *J Neurosurg* 83: 197-205.
69. Gomez H, Barnett GH, Estes ML, Palmer J, Magdinec M (1993): Stereotactic and computer-assisted neurosurgery at the Cleveland Clinic: review of 501 consecutive cases. *Cleve Clin J Med* 60: 399-410.
70. Gonzalez-Gonzalez JL, Lopez-Arbolay O, Morales-Sabina O, et al (2005): Transnasal-transsphenoidal endoscopic surgery of the sellar region. *Neurocirugia (Astur)* 16: 27-33.
71. Gouda KI, Freidberg SR, Larsen CR, Baker RA, Silverman ML (1983): Modification of the Gouda frame to allow stereotactic biopsy of the brain using the GE 8800 computed tomographic scanner. *Neurosurgery* 13: 176-181.
72. Gralla J, Ganslandt O, Lober H, et al (2003): Image-guided removal of supratentorial cavernomas in critical brain areas: application of neuronavigation and intraoperative magnetic resonance imaging. *Minim Invasive Neurosurg* 46: 72-77.
73. Gralla J, Guzman R, Brekenfeld C, Remonda L, Kiefer C (2005): High-resolution three-dimensional T2-weighted sequence for neuronavigation: a new setup and clinical trial. *J Neurosurg* 102: 658-663.

74. Grunert P, Charalampaki K, Kassem M, Boecher-Schwarz H, Filippi R, Grunert P Jr. (2002): Frame-based and frameless stereotaxy in the localization of cavernous angiomas. *Neurosurg Rev* 26: 53-61.
75. Grunert P, Espinosa J, Busert C, et al (2002): Stereotactic biopsies guided by an optical navigation system: technique and clinical experience. *Minim Invasive Neurosurg* 45: 11-15.
76. Gumprecht H, Ebel GK, Auer DP, Lumenta CB (2002): Neuronavigation and functional MRI for surgery in patients with lesion in eloquent brain areas. *Minim Invasive Neurosurg* 45:151-153.
77. Gumprecht H, Lumenta CB (1998): The operating microscope guided by a neuronavigation system: a technical note. *Minim Invasive Neurosurg* 41: 141-143.
78. Gumprecht H, Trost HA, Lumenta CB (2000): Neuroendoscopy combined with frameless neuronavigation. *Br J Neurosurg* 14: 129-131.
79. Gumprecht H, Widenka DC, Lumenta CB (1999): BrainLab VectorVision Neuronavigation System: Technology and clinical experiences in 131 cases. *Neurosurgery* 44: 97-105.
80. Guthrie BL, Adler JR (1992): Computer-assisted preoperative planning, interactive surgery, and frameless stereotaxy. *Clin Neurosurg* 38: 112-131.
81. Haberland N, Ebmeier K, Hliscs R, Grnewald JP, Silbermann J, Steenbeck J, Nowak H, Kalff R (2000): Neuronavigation in surgery of intracranial and spinal tumors. *J Cancer Res Clin Oncol* 126: 529-541.
82. Hardenack M, Bucher N, Falk A, et al. (1998): Preoperative planning and intraoperative navigation: status quo and perspectives. *Comput Aided Surg* 3: 153-158.
83. Hardy J (2000): Frameless stereotaxy for transsphenoidal surgery. *Neurosurgery* 46: 1269-1270.
84. Har-El G (2005): Endoscopic transnasal transsphenoidal pituitary surgery-comparison with the traditional sublabial transseptal approach. *Otolaryngol Clin North Am* 38: 723-735.
85. Hassfeld S, Zoller J, Albert FK, et al (1998): Preoperative planning and intraoperative navigation in skull base surgery. *J Craniomaxillofac Surg* 26: 220-225.
86. Hastreiter P, Rezk-Salama C, et al. (2004): Strategies for brain shift evaluation. *Med Image Anal* 8(4): 447-464.
87. Hata N, Nabavi A, Wells WM III, Warsfield SK, Kikinis R, Black PM, Jolesz A (2000): Three-dimensional optical flow method for measurement of volumetric brain deformation from intraoperative MR images. *J Comput Assist Tomogr* 24: 531-538.
88. Hattingen E, Good C, Weidauer S, Herminghaus S, Raab P, Marquardt G, Raabe A, Seifert V, Zanella FE (2005): Brain surface reformatted images for fast and easy localization of perirolandic lesions. *J Neurosurg* 102: 302-310.
89. Heilbrun MP, McDonald JD (2000): The future of image-guided surgery. *Clin Neurosurg* 46: 89-101.

90. Henry RG, Berman JI, Nagarajan SS, et al. (2004): Subcortical pathways serving cortical language sites: initial experience with diffusion tensor imaging fiber tracking combined with intraoperative language mapping. *Neuroimage* 21: 616-622.
91. Hirschberg H, Samset E (1999): Intraoperative image-directed dye marking of tumor margins. *Minim Invasive Neurosurg* 42: 123-127.
92. Hoell T, Oltmanns F, Schilling A, Brock M (1997): Precise localisation of single-word retrieval in the left cerebral hemisphere. *Lancet* 349: 104.
93. Hoell T, Warschewske U, Stockhausen v HM, Brock M (1999): DC-electromagnetic neuro-navigation – clinical experience. *Zentralbl Neurochir (Suppl)* 74: (P59).
94. Hopf NJ, Grunert P, Darabi K, Busert C, Bettag M (1999): Frameless neuronavigation applied to endoscopic neurosurgery. *Minim Invasive Neurosurg* 42: 187-193.
95. Horsley V, Clarke R (1908): The structure and function of the cerebellum examined by a new method. *Brain* 31: 45-124.
96. Hounsfield GN (1973): Computerized transverse axial scanning (tomography): I. Description of the system. *Br J Radiol* 46: 1016-1022.
97. Huber P (1979). In *Zerebrale Angiographie für Klinik und Praxis*. 3.Auflage. Stuttgart: Thieme Verlag: 111.
98. Huncke K, Van de Wiele B, Fried I, Rubinstein EH (1998): The asleep-awake-asleep anesthetic technique for intraoperative language mapping. *Neurosurgery* 42: 1312-1316.
99. Jannin P, Fleig OJ, Seigneuret E, Grova C, Morandi X, Scarabin JM (2000): A data fusion environment for multimodal and multi-informational neuronavigation. *Comput Aided Surg* 5: 1-10.
100. Jodicke A, Springer T, Boker DK (2004): Real-time integration of ultrasound into neuronavigation: technical accuracy using a light-emitting-diode-based navigation system. *Acta Neurochir (Wien)* 146: 1211-1220.
101. Jolesz FA, Kikinis R, Talos IF (2001): Neuronavigation in interventional MR imaging. Frameless stereotaxy. *Neuroimaging Clin N Am* 11: 685-693.
102. Kall BA, Goerss SJ, Stiving SO, Davis DH, Kelly PJ (1996): Quantitative analysis of a noninvasive stereotactic image registration technique. *Stereotact Funct Neurosurg* 66: 69-74.
103. Kamada K, Houkin K, Takeuchi F, Ishii N, Ikeda J, Sawamura Y, Kuriki S, Kawaguchi H, Iwasaki Y (2003): Visualization of the eloquent motor system by integration of MEG, functional, and anisotropic diffusion-weighted MRI in functional neuronavigation. *Surg Neurol* 59: 352.
104. Kamada K, Todo T, Masutani Y, Aoki S, Ino K, Takano T, Kirino T, Kawahara N, Morita A (2005): Combined use of tractography-integrated functional neuronavigation and direct fiber stimulation. *J Neurosurg* 102: 664-672.

105. Kanpolat Y, Deda H, Akyar S, Caglar S, Bilgic S (1989): CT-guided trigeminal tractotomy. *Acta Neurochir (Wien)* 100: 112-114.
106. Kato A, Yoshimine T, Hayakawa T, et al (1991): Computer assisted neurosurgery: development of a frameless and armless navigation system (CNS navigator). *No Shinkei Geka* 19: 137-142.
107. Kato A, Yoshimine T, Hayakawa T, Tomita Y, Ikeda T, Mitomo M, Harada K, Mogami H (1991): A frameless, armless navigational system for computer-assisted neurosurgery. *J Neurosurg* 74: 845-849.
108. Kelly PJ (1988): Volumetric stereotactic surgical resection of intra-axial brain mass lesions. *Mayo Clin Proc* 63: 1186-1198.
109. Kelly PJ, Alker GJ, Goerss S (1982): Computer-assisted stereotactic microsurgery for the treatment of intracranial neoplasms. *Neurosurgery* 10: 324-331.
110. Kelly PJ, Kall BA, Goerss S, Earnest F IV (1986): Computer-assisted stereotaxic laser resection of intra-axial brain neoplasms. *J Neurosurg* 64: 427-439.
111. Kim IY, Jung S, Moon KS, Jung TY, Kang SS (2004): Neuronavigation-guided endoscopic surgery for pineal tumors with hydrocephalus. *Minim Invasive Neurosurg* 47: 365-368.
112. King RB, Schell GR (1987): Cortical localization and monitoring during cerebral operations. *J Neurosurg* 67: 210-219.
113. Kleinpeter G, Lothaller C (2003): Frameless neuronavigation using the ISG-system in practice: from craniotomy to delineation of lesion. *Minim Invasive Neurosurg* 46: 257-264.
114. Klisch J, Schipper J, Husstedt H, et al (2001): Transsphenoidal computer-navigation-assisted deflation of a balloon after endovascular occlusion of a direct carotid cavernous sinus fistula. *AJNR Am J Neuroradiol* 22: 537-540.
115. Knauth M, Wirtz CR, Tronnier VM, et al (1999): Intraoperative MR imaging increases the extent of tumor resection in patients with high-grade gliomas. *AJNR AM J Neuroradiol* 20: 1642-1646.
116. Kober H, Nimsky C, Vieth J, Fahlbusch R, Ganslandt O (2002): Co-registration of function and anatomy in frameless stereotaxy by contour fitting. *Stereotact Funct Neurosurg* 79: 272-283.
117. Koele W, Stammberger H, Lackner A, et al. (2002): Image guided surgery of paranasal sinuses and anterior skull base – five years experience with the InstaTrak-System. *Rhinology* 40: 1-9.
118. Koivukangas J, Louhisalmi Y, Alakuijala J, Oikarinen J (1993): Ultrasound-controlled neuronavigator-guided brain surgery. *J Neurosurg* 79: 36-42.
119. Kollias SS, Bernays R, Marugg RA, et al. (1998): Target definition and trajectory optimization for interactive MR-guided biopsies of brain tumors in an open configuration MRI system. *J Magn Reson Imaging* 8: 143-159.

120. Kombos T, Suess O, Funk T, Kern BC, Brock M (2000): Intra-operative mapping of the motor cortex during surgery in and around the motor cortex. *Acta Neurochir (Wien)* 142: 263-268.
121. Kopitnik TA, Kaufman HH, Haid RW, et al. (1987): Technique of stereotactic biopsy of two cranial target employing spherical coordinates to define a single trajectory. *Appl Neurophysiol* 50: 188-194.
122. Kosugi Y, Watanabe E, Goto J, et al. (1988): An articulated neurosurgical navigation system using MRI and CT images. *IEEE Trans Biomed Eng* 35: 147-152.
123. Kreth FW, Muacevic A, Medele R, Bise K, Meyer T, Reulen HJ (2001): The risk of haemorrhage after image guided stereotactic biopsy of intra-axial brain tumours – A prospective study. *Acta Neurochir (Wien)* 143: 539-546.
124. Krings T, Reul J, Spetzger U, Klusmann A, Roessler F, Gilsbach JM, Thron A (1998): Functional magnetic resonance mapping of sensor motor cortex for image guided neurosurgical intervention. *Acta Neurochir* 140: 215-222.
125. Kucharczyk W, Bernstein M (1997): Do the benefits of image guidance in neurosurgery justify the costs? From stereotaxy to intraoperative MR. *AJNR Am J Neuroradiol* 18: 1855-1859.
126. Kurimoto M, Hayashi N, Kamiyama H, Nagai S, Shibata T, Asahi T, Matsumura N, Hirashima Y, Endo S (2004): Impact of neuronavigation and image-guided extensive resection for adult patients with supratentorial malignant astrocytomas: a single-institution retrospective study. *Minim Invasive Neurosurg* 47: 278-283.
127. Kuroiwa T, Ohta T (2001): Operations using a frameless stereotactic system with a marker: Technical note. *Minim Invas Neurosurg* 44: 163-166.
128. Kurtsoy A, Menku A, Tucer B, et al. (2005): Neuronavigation in skull base tumors. *Minim Invasive Neurosurg* 48: 7-12.
129. Lehericy S, Duffau H, Cornu P, Capelle L, Pidoux B, Carpentier A, Auliac S, Clemenceau S, Sichez JP, Bitar A, Valery CA, Van Effenterre R, Faillot T, Srour A, Fohanno D, Philippon J, Le Bihan D, Marsault C (2000): Correspondence between functional magnetic resonance imaging somatotopy and individual brain anatomy of the central region: comparison with intraoperative stimulation in patients with brain tumors. *J Neurosurg* 92: 589-598.
130. Leksell L (1949): A stereotaxic apparatus for intracerebral surgery. *Acta Chir Scand* 99: 229-233.
131. Leksell L, Jernberg B (1980): Stereotaxis and tomography. A technical note. *Acta Neurochir (Wien)* 52: 1-7.
132. Letteboer MM, Willems PW, Viergever MA, Niessen WJ (2005): Brain shift estimation in image-guided neurosurgery using 3-D ultrasound. *IEEE Trans Biomed Eng* 52: 268-276.
133. Leuthardt EC, Fox D, Ojemann GA, Dacey RG, Grubb RL, Rich KM, Ojemann JG (2002): Frameless stereotaxy without rigid pin fixation during awake craniotomies. *Stereotact Funct Neurosurg* 79: 256-61.

134. Lewis JT, Galloway RL, Schreiner S (1998): An ultrasonic approach to localization of fiducial markers for interactive, image-guided neurosurgery-Part I: Principles. *IEEE Trans Biomed Eng* 45: 620-630.
135. Li QH, Zamorano L, Pandya A, Perez R, Gong J, Diaz F (2002): The application accuracy of the NeuroMate robot--A quantitative comparison with frameless and frame-based surgical localization systems. *Comput Aided Surg* 7: 90-98.
136. Lindseth F, Lango T, Bang J, Nagelhus Hernes TA (2002): Accuracy evaluation of a 3D ultrasound-based neuronavigation system. *Comput Aided Surg* 7: 197-222.
137. Lumenta CB, Gumprecht HK, Leonardi MA, Gerstner W, Brehm-Häuser B (1997): Three-dimensional computer-assisted stereotactic-guided microneurosurgery combined with cortical mapping of the motor area by direct electrostimulation. *Minim Invasive Neurosurg* 40: 50-54.
138. Lunsford LD, Rosenbaum AE, Perry J (1982): Stereotactic surgery using the "therapeutic" CT scanner. *Surg Neurol* 18: 116-122.
139. Lurito JT, Lowe MJ, Sartorius C, et al (2000): Comparison of fMRI and intraoperative direct cortical stimulation in localization of receptive language areas. *J Comput Assist Tomogr* 24: 99-105.
140. Maciunas RJ, Berger MS, Copeland B, Mayberg MR, Selker R, Allen GS (1996): A technique for interactive image-guided neurosurgical intervention in primary brain tumors. *Neurosurg Clin N Am* 7: 245-266.
141. Marinho AM, Barua M, Haller J, Ryken TC (2002): Enhanced anatomic visualization with ultrasound-assisted intracranial image-guidance in neurosurgery. *Technol Cancer Res Treat* 1: 181-186.
142. Maurer CR, Hill DL, Martin AJ, Liu H, McCue M, Rueckert D, Lloret D, Hall WA, Maxwell RE, Hawkes DJ, Truwit CL (1998): Investigation of intraoperative brain deformation using a 1.5-T interventional MR system: Preliminary results. *IEEE Trans Med Imaging* 17: 817-825.
143. McCutcheon IE, Kitagawa RS, Demasi PF, et al. (2004): Frameless stereotactic navigation in transsphenoidal surgery: comparison with fluoroscopy. *Stereotact Funct Neurosurg* 82: 43-48.
144. McDonald JD, Chong BW, Lewine JD, Jones G, Burr RB, McDonald PR, Koehler SB, Tsuruda J, Orrison WW, Heilbrun MP (1999): Integration of preoperative and intraoperative functional brain mapping in a frameless stereotactic environment for lesions near eloquent cortex. *J Neurosurg* 90: 591-598.
145. McNerney J, Roberts DW (2000): Frameless stereotaxy of the brain. *Mt Sinai J Med* 67: 300-310.
146. Mehdorn HM, Schrader B, Nabavi A, et al. (2000): Neuronavigation in the region of the skull base. *Laryngorhinootologie* 79: 404-411.
147. Messmer P, Baumann B, Suhm N, Jacob AL (2001): Navigationsverfahren für die bildgesteuerte Therapie – ein Überblick. *Fortschr Röntgenstr* 173: 777-784.

148. Meyer FB, Bates LM, Goerss SJ, Friedman JA, Windschitl WL, Duffy JR, Perkins WJ, O'Neill BP (2001): Awake craniotomy for aggressive resection of primary gliomas located in eloquent brain. *Mayo Clin Proc* 76: 677-87.
149. Miga MI, Paulsen KD, Lemery JM, Eisner SD, Hartov A, Kennedy FE, Roberts DW (1999): Model-updated image guidance: Initial clinical experiences with gravity-induced brain deformation. *IEEE Transactions on Medical Imaging* 18: 866-874.
150. Modayur B, Prothero J, Ojemann G, et al (1997): Visualization-based mapping of language function in the brain. *Neuroimage* 6: 245-258.
151. Moniz E (1927): L'encéphalographie artérielle, son importance dans la localisation des tumeurs cérébrales. *Revue neurologique* 2: 72-90.
152. Nabavi A, Manthei G, Blomer U, Kumpf L, Klinge H, Mehdorn HM (1995): [Neuronavigation. Computer-assisted surgery in neurosurgery]. *Radiologe* 35: 573-577.
153. Nieves MN, Haas E, Hollerhage HG, Schneider H, Pollath A, Archavlis E (2004): Combined minimal invasive techniques in deep supratentorial intracerebral haematomas. *Minim Invasive Neurosurg* 47: 294-298.
154. Nimsky C, Buchfelder M (2003): Neuronavigation in epilepsy surgery. *Arq Neuropsiquiatr* 61 Suppl 1: 109-114.
155. Nimsky C, Ganslandt O, Fahlbusch R (2004): Functional neuronavigation and intraoperative MRI. *Adv Tech Stand Neurosurg* 29: 229-263.
156. Nimsky C, Grummich P, Sorensen AG, Fahlbusch R, Ganslandt O (2005): Visualization of the pyramidal tract in glioma surgery by integrating diffusion tensor imaging in functional neuronavigation. *Zentralbl Neurochir* 66: 133-141.
157. Nimsky Ch, Rachinger J, Iro H, et al (2004): Adaptation of a hexapod-based robotic system for extended endoscope-assisted transsphenoidal skull base surgery. *Minim Invasive Neurosurg* 47: 41-6.
158. Nowinski WL (2005): The cerefy brain atlases: continuous enhancement of the electronic talairach-tournoux brain atlas. *Neuroinformatics* 3: 293-300.
159. Oertel J, Gaab MR, Runge U, Schroeder HW, Wagner W, Piek J (2004): Neuronavigation and complication rate in epilepsy surgery. *Neurosurg Rev* 27: 214-217.
160. Ohhashi G, Kamio M, Abe T, Otori N, Haruna S (2002): Endoscopic transnasal approach to the pituitary lesions using a navigation system (InstaTrak system): technical note. *Minim Invasive Neurosurg* 45: 120-123.
161. Ojemann G, Mateer C (1979): Human language cortex: localization of memory, syntax, and sequential motor-phoneme identification systems. *Science* 28: 1401-1403.
162. Ojemann G, Ojemann J, Lettich E, Berger M (1989): Cortical language localization in left, dominant hemisphere. An electrical stimulation mapping investigation in 117 patients. *J Neurosurg* 71: 316-326.

-
163. Ojemann GA (1979): Individual variability in cortical localization of language. *J Neurosurg* 50: 164-169.
 164. Olivier A, Alonso-Vanegas M, Comeau R, Peters TM (1996): Image-guided surgery of epilepsy. *Neurosurg Clin N Am* 7: 229-243.
 165. Olson JJ, Shepherd S, Bakay RA (1997): The EasyGuide Neuro image-guided surgery system. *Neurosurgery* 40: 1092-1096.
 166. Ostertag CB, Warnke PC (1999): [Neuronavigation. Computer-assisted neurosurgery]. *Nervenarzt* 70: 517-521.
 167. Pena A, Bolton MD, Whitehouse H, Pickard JD (1999): Effects of brain ventricular shape on periventricular biomechanics: A finite-element analysis. *Neurosurgery* 45: 107-118.
 168. Pouratian N, Cannestra AF, Bookheimer SY, et al. (2004): Variability of intraoperative electrocortical stimulation mapping parameters across and within individuals. *J Neurosurg* 101: 458-466.
 169. Quinones-Hinojosa A, Ojemann SG, Sanai N, et al (2003): Preoperative correlation of intraoperative cortical mapping with magnetic resonance imaging landmarks to predict localization of the Broca area. *J Neurosurg* 99: 311-318.
 170. Raabe A, Krishnan R, Zimmermann M, Seifert V (2003): Neuronavigation oder Stereotaxie? Eine Standortbestimmung zur Wahl des Verfahrens. *Zentralbl Neurochir* 64: 1-5.
 171. Reinges MH, Krings T, Nguyen HH, Kuker W, Spetzger U, Rohde V, Hutter BO, Thron A, Gilsbach JM (2000): Virtual pointer projection of the central sulcus to the outside of the skull using frameless neuronavigation - accuracy and applications. *Acta Neurochir (Wien)* 142: 1385-1390.
 172. Reinges MH, Nguyen HH, Krings T, Hutter BO, Rohde V, Gilsbach JM (2004): Course of brain shift during microsurgical resection of supratentorial cerebral lesions: limits of conventional neuronavigation. *Acta Neurochir (Wien)* 146: 369-377.
 173. Reinhardt HF, Zweifel HJ (1990): Interactive sonar-operated device for stereotactic and open surgery. *Stereotact Funct Neurosurg* 54-55: 393-397.
 174. Reithmeier T, Krammer M, Gumprecht H, Gerstner W, Lumenta CB (2003): Neuronavigation combined with electrophysiological monitoring for surgery of lesions in eloquent brain areas in 42 cases: a retrospective comparison of the neurological outcome and the quality of resection with a control group with similar lesions. *Minim Invasive Neurosurg* 46: 65-71.
 175. Rengarachy S, Wilkins R (1994). In *Principles of Neurosurgery*. London, Mosby Year Book Europe: 49.2.
 176. Rezai AR, Mogilner AY, Capell J et al. (1997): Integration of functional brain mapping in image-guided surgery. *Acta Neurochir (suppl)* 68: 85-89.

177. Roberts DW, Miga MI, Hartov A, Eisner S, Lemery JM, Kennedy FE, Paulsen KD (1999): Intraoperatively updated neuroimaging using brain modelling and sparse data. *Neurosurgery* 45: 1199-1207.
178. Roberts DW, Nakajima T, Brodwater B, Pavlidis J, Friets E, Fagan E, Hartov A, Strohbehn J (1992): Further development and clinical application of the stereotactic operating microscope. *Stereotact Funct Neurosurg* 58: 114-117.
179. Roberts DW, Strohbehn JW, Hatch JF, Murray W, Kettenberger H (1986): A frameless stereotactic integration of computerized tomographic imaging and the operating microscope. *J Neurosurg* 65: 545-549.
180. Roessler K, Ungersboeck K, Dietrich W, Aichholzer M, Hittmeir K, Matula C, Czech T, Koos WT (1997): Frameless stereotactic guided neurosurgery: Clinical experiences with an infrared based pointer device navigation system. *Acta Neurochir (Wien)* 139: 551-559.
181. Rohde V, Spangenberg P, Mayfrank L, et al (2005): Advanced neuronavigation in skull base tumors and vascular lesions. *Minim Invasive Neurosurg* 48: 13-18.
182. Rohling R, Munger P, Hollerbach JM, Peter T (1995): Comparison of relative accuracy between a mechanical and an optical position tracker for image-guided neurosurgery. *J Image Guid Surg* 1: 30-34.
183. Röntgen WC (1895): Über eine neue Art von Strahlen (Vorläufige Mitteilung). *Aus den Sitzungsberichten der Würzburger physikalisch-medizinischen Gesellschaft*. 132-141.
184. Rudnik A, Zawadzki T, Wojtacha M, et al. (2005): Endoscopic transnasal transsphenoidal treatment of pathology of the sellar region. *Minim Invasive Neurosurg* 48: 101-107.
185. Russell SM, Woo HH, Joseffer SS, Jafar JJ (2002): Role of frameless stereotaxy in the surgical treatment of cerebral arteriovenous malformations: technique and outcomes in a controlled study of 44 consecutive patients. *Neurosurgery* 51: 1108-1116.
186. Ryan MJ, Erickson RK, Levin DN, Pelizzari CA, Macdonald RL, Dohrmann GJ (1996): Frameless stereotaxy with real-time tracking of patient head movement and retrospective patient-image registration. *J Neurosurg* 85: 287-92.
187. Sabbah P, Foehrenbach H, Dutertre G, et al. (2002): Multimodal anatomic, functional, and metabolic brain imaging for tumor resection. *Clin Imaging* 26: 6-12.
188. Sarang A, Dinsmore J (2003): Anaesthesia for awake craniotomy - evolution of a technique that facilitates awake neurological testing. *Br J Anaesth* 90: 161-165.
189. Schaller C, Meyer B, Van Roost D, Schramm J (1997): Image guided microsurgery with a semifreehand neuronavigational device. *Comput Aided Surg* 2: 162-171.
190. Schiffbauer H, Berger MS, Ferrari P, Freudenstein D, Rowley HA, Roberts TP (2003): Preoperative magnetic source imaging for brain tumor surgery: a quantitative comparison with intraoperative sensory and motor mapping. *Neurosurg Focus* 15: E7.

191. Schlaier JR, Warnat J, Dorenbeck U, Proescholdt M, Schebesch KM, Brawanski A.(2004): Image fusion of MR images and real-time ultrasonography: evaluation of fusion accuracy combining two commercial instruments, a neuronavigation system and a ultrasound system. *Acta Neurochir (Wien)* 146: 271-277.
192. Schreiner S, Galloway RL, Lewis JT, Bass WA, Muratore DM (1998): An ultrasonic approach to localization of fiducial markers for interactive, image-guided neurosurgery-Part II: Implementation and automation. *IEEE Trans Biomed Eng* 45: 631-641.
193. Schul C, Wassmann H, Skopp GB, et al. (1998): Surgical management of intraosseous skull base tumors with aid of Operating Arm System. *Comput Aided Surg* 3: 12-19.
194. Schulder M, Fontana P, Lavenhar MA, Carmel PW (1999): The relationship of imaging techniques to the accuracy of frameless stereotaxy. *Stereotact Funct Neurosurg* 72: 136-141.
195. Signorelli F, Guyotat J, Schneider F, et al. (2003): Technical refinements for validating functional MRI-based neuronavigation data by electrical stimulation during cortical language mapping. *Minim Invasive Neurosurg* 46: 265-268.
196. Simos PG, Papanicolaou AC, Breier JI, Wheless JW, Constantinou JE, Gormley WB, Maggio WW (1999): Localization of language-specific cortex by using magnetic source imaging and electrical stimulation mapping. *J Neurosurg* 91: 787-796.
197. Sipos EP, Tebo SA, Zinreich SJ, Long DM, Brem H (1996): In vivo accuracy testing and clinical experience with the ISG viewing wand. *Neurosurgery* 39: 194-204.
198. Sobottka SB, Bredow J, Beuthien-Baumann B, Reiss G, Schackert G, Steinmeier R (2002): Comparison of functional brain PET images and intraoperative brain-mapping data using image-guided surgery. *Comput Aided Surg* 7: 317-325.
199. Spetzger U, Gilsbach JM, Mösger R, Schlöndorff G, Laborde G (1997): The computer-assisted-localizer (CAL): a navigation help in microneurosurgery. *Eur Surg Res* 29: 481-487.
200. Spetzger U, Laborde G, Gilsbach JM (1995): Frameless neuronavigation in modern neurosurgery. *Minim Invasive Neurosurg* 38: 163-166.
201. Spiegel E, Wycis HT, Marks M (1947): Stereotaxic apparatus for operations on the human brain. *Science* 106: 349-350.
202. Stapleton SR, Kiriakopoulos E, Mikulis D, Drake JM, Hoffman HJ, Hymphreys R, Hwang P, Otsubo H, Holowka S, Logan W, Rutka JT (1997): Combined utility of functional MRI, cortical mapping, and frameless stereotaxy in the resection of lesions in eloquent areas of brain in children. *Pediatr Neurosurg* 26: 68-82.
203. Steinmeier R, Sobottka SB, Reiss G, et al (2002): Surgery of low-grade gliomas near speech-eloquent regions: brainmapping versus preoperative functional imaging. *Onkologie* 25: 552-7.
204. Suess O, Kombos T, Ciklatekerlio Ö, Stendel R, Suess S, Brock M (2002): Impact of brain shift on intraoperative neurophysiological monitoring with cortical strip electrodes. *Acta Neurochir (Wien)* 144: 1279-1289.

205. Suess O, Kombos T, Ciklatekerlio Ö, Suess S, Da Silva C, Brock M (2003): Klinische Studie zur Anwendung der hochfrequenten monopolaren Kortexstimulation (MKS) für die intraoperative Ortung und Überwachung motorischer Hirnareale bei Eingriffen in der Nähe der Zentralregion. *Klin Neurophysiol* 34: 127-137.
206. Suess O, Kombos T, Hoell T, Baur S, Brock M (2000): A new cortical electrode for neuronavigation-guided intraoperative neurophysiological monitoring: Technical note. *Acta Neurochir (Wien)* 142: 329-332.
207. Suess O, Kombos T, Kurth R, Suess S, Mularski S, Hammersen S, Brock M (2001): Intracranial image-guided neurosurgery: Experience with a new electromagnetic navigation system. *Acta Neurochir (Wien)* 143: 927-934.
208. Suess O, Schönherr S, Schilling A, Kühn B, Mularski SO, Suess S, Brock M, Kombos T (2005): [Sensor-based detection of skull positioning for image-guided cranial navigation under free head mobility]. *Rofo* 177: 1000-1008.
209. Sure U, Alberti O, Petermeyer M, et al. (2000): Advanced image-guided skull base surgery. *Surg Neurol* 53: 563-72.
210. Sure U, Benes L, Bozinov O, Woydt M, Tirakotai W, Bertalanffy H (2005): Intraoperative landmarking of vascular anatomy by integration of duplex and Doppler ultrasonography in image-guided surgery. Technical note. *Surg Neurol* 63: 133-132.
211. Sure U, Benes L, Riegel T, Schulte DM, Bertalanffy H (2002): Image fusion for skull base neuronavigation. Technical note. *Neurol Med Chir (Tokyo)* 42: 458-462.
212. Sutton D (1993). In *A Textbook of Radiology and Imaging*. 5. Auflage. Band 2. London, Churchill Livingstone: 1391.
213. Takeuchi H, Yoshida M, Kubota T, Ishii H, Sato K, Handa Y, Itoh H (2003): Frameless stereotactic radiosurgery with mobile CT, mask immobilization and micro-multileaf collimators. *Minim Invasive Neurosurg* 46: 82-85.
214. Taniguchi M, Cedzich C, Schramm J (1993): Modification of cortical stimulation for motor evoked potentials under general anesthesia: technical description. *Neurosurgery* 32: 219-226.
215. Taylor E, Houghton W (1900): Some recent researchers on the topography of the convolutions and fissures of the brain. *Trans R Acad Med Ireland* 18: 511-522.
216. Taylor MD, Bernstein M (1999): Awake craniotomy with brain mapping as the routine surgical approach to treating patients with supratentorial intraaxial tumors: a prospective trial of 200 cases. *J Neurosurg* 90: 35-41.
217. Thomale UW, Stover JF, Unterberg AW (2005): The use of neuronavigation in transnasal transsphenoidal pituitary surgery. *Zentralbl Neurochir* 66: 126-132.
218. Tomczak RJ, Wunderlich AP, Wang Y, et al. (2000): fMRI for preoperative neurosurgical mapping of motor cortex and language in a clinical setting. *J Comput Assist Tomogr* 24: 927-934.

-
219. Tronnier VM, Wirtz CR, Knauth M, Bonsanto MM, Hassfeld S, Albert FK, Kunze S (1996): Intraoperative computer-assisted neuronavigation in functional neurosurgery. *Stereotact Funct Neurosurg* 66: 65-68.
220. Tseng CS, Chung CW, Chen HH, Wang SS, Tseng HM (1999): Development of a robotic navigation system for neurosurgery. *Stud Health Technol Inform* 62: 358-359.
221. Unsgaard G, Rygh OM, Selbekk T, Müller TB, Kolstad F, Lindseth F, Nagelhus Hernes TA (2006): Intra-operative 3D ultrasound in neurosurgery. *Acta Neurochir (Wien)* 148: 235-253.
222. Varma TR, Eldridge PR, Forster A, Fox S, Fletcher N, Steiger M, Littlechild P, Byrne P, Sinnott A, Tyler K, Flintham S (2003): Use of the NeuroMate stereotactic robot in a frameless mode for movement disorder surgery. *Stereotact Funct Neurosurg* 80: 132-135.
223. Vlieger EJ, Majoie CB, Leenstra S, Den Heeten GJ (2004): Functional magnetic resonance imaging for neurosurgical planning in neurooncology. *Eur Radiol* 14: 1143-1153.
224. Vougioukas VI, Hubbe U, Hochmuth A, Gellrich NC, van Velthoven V (2003): Perspectives and limitations of image-guided neurosurgery in pediatric patients. *Childs Nerv Syst* 19: 783-791.
225. Vougioukas VI, Hubbe U, Schipper J, et al (2003): Navigated transoral approach to the cranial base and the craniocervical junction: technical note. *Neurosurgery* 52: 247-250.
226. Vougioukas VI, Hubbe U, van Velthoven V, Freiman TM, Schramm A, Spetzger U (2004): Neuronavigation-assisted cranial reconstruction. *Neurosurgery* 55: 162-167.
227. Wagner W, Gaab MR, Schroeder HWS, Piek J, Niendorf WR (2000): Neuronavigation in der Zentralregion: Bedeutung für einzelne Operationsphasen bei verschiedenen Prozesslokalisationen. *Zentralbl Neurochir* 61: 188-193.
228. Walker DG, Ohaegbulam C, Black PM (2002): Frameless stereotaxy as an alternative to fluoroscopy for transsphenoidal surgery: use of the InstaTrak-3000 and a novel headset. *J Clin Neurosci* 9: 294-297.
229. Watanabe E, Mayanagi Y, Kosugi Y, Manaka S, Takakura K (1991): Open surgery assisted by the Navigator, a stereotactic articulated, sensitive arm. *Neurosurgery* 28: 792-800.
230. Watanabe E, Watanabe T, Manaka S, Mayanagi Y, Takakura K (1987): Three-dimensional digitizer (neuronavigator): new equipment for computed tomography-guided stereotaxic surgery. *Surgical Neurology* 27: 543-547.
231. West JB, Fitzpatrick JM, Toms SA, Maurer CR, Maciunas RJ (2001): Fiducial point placement and the accuracy of point based, rigid body registration. *Neurosurgery* 48: 810-816.
232. Wharen RE, Putzke JD, Uitti RJ (2005): Deep brain stimulation lead fixation : a comparative study of the Navigus and Medtronic burr hole fixation device. *Clin Neurol Neurosurg* 107: 393-395.
233. Wiltfang J, Rupprecht S, Ganslandt O, et al (2003): Intraoperative Image-Guided Surgery of the Lateral and Anterior Skull Base in Patients with Tumors or Trauma. *Skull Base* 13: 21-29.

-
234. Wirtz CR, Albert FK, Schwaderer M, Heuer C, Staubert A, Tronnier VM, Knauth M, Kunze S (2000): The benefit of neuronavigation for neurosurgery analyzed by its impact on glioblastoma surgery. *Neurol Res* 22: 354-360.
235. Wirtz CR, Bonsanto MM, et al. (1997): Intraoperative magnetic resonance imaging to update interactive navigation in neurosurgery: method and preliminary experience. *Comput Aided Surg* 2(3-4): 172-179.
236. Wirtz CR, Tronnier VM, Bonsanto MM, Hassfeld S, Knauth M, Kunze S (1998): Neuronavigation: Methoden und Ausblick. *Nervenarzt* 69: 1029-036.
237. Wong GK, Poon WS, NG SC, et al (2004): Fluoroscopic frameless computer-assisted navigation for transsphenoidal surgery: a clinical assessment of accuracy in spatial position and trajectory. *Minim Invasive Neurosurg* 47: 29-31.
238. Woolsey CN, Erickson TC, Gilson WE (1979): Localization in somatic, sensory and motor areas of human cerebral cortex as determined by direct recording of evoked potentials and electrical stimulation. *J Neurosurg* 51: 476-506.
239. Wurm G, Ringler H, Knogler F, Schnizer M (2003): Evaluation of neuronavigation in lesional and non-lesional epilepsy surgery. *Comput Aided Surg* 8: 204-214.
240. Zaaroor M, Bejerano Y, Weinfeld Z, Ben-Haim S (2001): Novel magnetic technology for intraoperative intracranial frameless navigation: in vivo and in vitro results. *Neurosurgery* 48: 1100-1108.
241. Zamorano L, Jiang Z, Kadi AM (1994): Computer-assisted neurosurgery system: Wayne State University hardware and software configuration. *Comput Med Imaging Graph* 18: 257-271.
242. Zamorano L, Kadi M, Dong A (1992): Computer-assisted neurosurgery. Simulation and automation. *Stereotact Funct Neurosurg* 59: 115-122.
243. Zamorano LJ, Nolte L, Kadi AM, Jiang Z (1994): Interactive intraoperative localization using an infrared-based system. *Stereotact Funct Neurosurg* 63: 84-88.
244. Zhang P, Lin W, Yao X (2004): Resection of giant pituitary adenomas through transnasal and transsphenoidal approach with nasoendoscope. *Lin Chuang Er Bi Yan Hou Ke Za Zhi* 18: 463-465.
245. Zhang X, Zhang J, Fei Z, et al (2002): Neuronavigator-guided microsurgery for resection of brain tumors. *Zhonghua Yi Xue Za Zhi* 82: 219-221.