

8. Literatur

- Abu-Jawdeh, G., Comella, N., Tomita, Y., Brown, L. F., Tognazzi, K., Sokol, S. Y. and Kocher, O. (1999). Differential expression of frpHE: a novel human stromal protein of the secreted frizzled gene family, during the endometrial cycle and malignancy. *Lab Invest* 79, 439-47.
- Alers, J. C., Rochat, J., Krijtenburg, P. J., Hop, W. C., Kranse, R., Rosenberg, C., Tanke, H. J., Schroder, F. H. and van Dekken, H. (2000). Identification of genetic markers for prostatic cancer progression. *Lab Invest* 80, 931-42.
- Alon, U., Barkai, N., Notterman, D. A., Gish, K., Ybarra, S., Mack, D. and Levine, A. J. (1999). Broad patterns of gene expression revealed by clustering analysis of tumor and normal colon tissues probed by oligonucleotide arrays. *Proc Natl Acad Sci U S A* 96, 6745-50.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* 215, 403-10.
- Arias, A. M., Brown, A. M. and Brennan, K. (1999). Wnt signalling: pathway or network? *Curr Opin Genet Dev* 9, 447-54.
- Baker, S. J., Fearon, E. R., Nigro, J. M., Hamilton, S. R., Preisinger, A. C., Jessup, J. M., vanTuinen, P., Ledbetter, D. H., Barker, D. F., Nakamura, Y. et al. (1989). Chromosome 17 deletions and p53 gene mutations in colorectal carcinomas. *Science* 244, 217-21.
- Bateman, A., Birney, E., Durbin, R., Eddy, S. R., Finn, R. D. and Sonnhammer, E. L. (1999). Pfam 3.1: 1313 multiple alignments and profile HMMs match the majority of proteins. *Nucleic Acids Res* 27, 260-2.
- Baud, E., Catilina, P., Boiteux, J. P. and Bignon, Y. J. (1998). Human bladder cancers and normal bladder mucosa present the same hot spot of heterozygous chromosome-9 deletion. *Int J Cancer* 77, 821-4.
- Baugh, L. R., Hill, A. A., Brown, E. L. and Hunter, C. P. (2001). Quantitative analysis of mRNA amplification by in vitro transcription. *Nucleic Acids Res* 29, E29.
- Bettuzzi, S., Davalli, P., Astancolle, S., Carani, C., Madeo, B., Tampieri, A., Corti, A., Saverio, B., Pierpaola, D., Serenella, A. et al. (2000). Tumor progression is accompanied by significant changes in the levels of expression of polyamine metabolism regulatory genes and clusterin (sulfated glycoprotein 2) in human prostate cancer specimens. *Cancer Res* 60, 28-34.
- Bhattacharjee, A., Richards, W. G., Staunton, J., Li, C., Monti, S., Vasa, P., Ladd, C., Beheshti, J., Bueno, R., Gillette, M. et al. (2001). Classification of human lung carcinomas by mRNA expression profiling reveals distinct adenocarcinoma subclasses. *Proc Natl Acad Sci U S A* 98, 13790-5.
- Birck, A., Ahrenkiel, V., Zeuthen, J., Hou-Jensen, K. and Guldberg, P. (2000). Mutation and allelic loss of the PTEN/MMAC1 gene in primary and metastatic melanoma biopsies. *J Invest Dermatol* 114, 277-80.
- Birnboim, H. C. and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* 7, 1513-23.

- Bonfield, J. K., Smith, K. and Staden, R. (1995). A new DNA sequence assembly program. *Nucleic Acids Res* 23, 4992-9.
- Bonkhoff, H. and Remberger, K. (1996). Differentiation pathways and histogenetic aspects of normal and abnormal prostatic growth: a stem cell model. *Prostate* 28, 98-106.
- Bos, J. L. (1989). ras oncogenes in human cancer: a review. *Cancer Res* 49, 4682-9.
- Bridges, L. C., Tani, P. H., Hanson, K. R., Roberts, C. M., Judkins, M. B. and Bowditch, R. D. (2001). The lymphocyte metalloprotease MDC-L (ADAM 28) is a ligand for the integrin α 4 β 1. *J Biol Chem* 27, 27.
- Brown, J. D. and Moon, R. T. (1998). Wnt signaling: why is everything so negative? *Curr Opin Cell Biol* 10, 182-7.
- Cairns, P., Okami, K., Halachmi, S., Halachmi, N., Esteller, M., Herman, J. G., Jen, J., Isaacs, W. B., Bova, G. S. and Sidransky, D. (1997). Frequent inactivation of PTEN/MMAC1 in primary prostate cancer. *Cancer Res* 57, 4997-5000.
- Cairns, P., Polascik, T. J., Eby, Y., Tokino, K., Califano, J., Merlo, A., Mao, L., Herath, J., Jenkins, R., Westra, W. et al. (1995). Frequency of homozygous deletion at p16/CDKN2 in primary human tumours. *Nat Genet* 11, 210-2.
- Cappellen, D., Gil Diez de Medina, S., Chopin, D., Thiery, J. P. and Radvanyi, F. (1997). Frequent loss of heterozygosity on chromosome 10q in muscle-invasive transitional cell carcinomas of the bladder. *Oncogene* 14, 3059-66.
- Cawthon, R. M., Weiss, R., Xu, G. F., Viskochil, D., Culver, M., Stevens, J., Robertson, M., Dunn, D., Gesteland, R., O'Connell, P. et al. (1990). A major segment of the neurofibromatosis type 1 gene: cDNA sequence, genomic structure, and point mutations. *Cell* 62, 193-201.
- Chen, B., He, L., Savell, V. H., Jenkins, J. J. and Parham, D. M. (2000). Inhibition of the interferon-gamma/signal transducers and activators of transcription (STAT) pathway by hypermethylation at a STAT-binding site in the p21WAF1 promoter region. *Cancer Res* 60, 3290-8.
- Choi, C., Kim, M. H., Juhng, S. W. and Oh, B. R. (2000). Loss of heterozygosity at chromosome segments 8p22 and 8p11.2-21.1 in transitional-cell carcinoma of the urinary bladder. *Int J Cancer* 86, 501-5.
- Christensen, M., Jensen, M. A., Wolf, H. and Orntoft, T. F. (1998). Pronounced microsatellite instability in transitional cell carcinomas from young patients with bladder cancer. *Int J Cancer* 79, 396-401.
- Christensen, M., Sunde, L., Bolund, L. and Orntoft, T. F. (1999). Comparison of three methods of microsatellite detection. *Scand J Clin Lab Invest* 59, 167-77.
- De Marzo, A. M., Putzi, M. J. and Nelson, W. G. (2001). New concepts in the pathology of prostatic epithelial carcinogenesis. *Urology* 57, 103-14.
- Del Sal, G., Collavin, L., Ruaro, M. E., Edomi, P., Saccone, S., Valle, G. D. and Schneider, C. (1994). Structure, function, and chromosome mapping of the growth-suppressing human homologue of the murine gas1 gene. *Proc Natl Acad Sci U S A* 91, 1848-52.

- Dennis, S., Aikawa, M., Szeto, W., d'Amore, P. A. and Papkoff, J. (1999). A secreted frizzled related protein, FrzA, selectively associates with Wnt-1 protein and regulates wnt-1 signaling. *J Cell Sci* 122, 3815-20.
- Dhanasekaran, S. M., Barrette, T. R., Ghosh, D., Shah, R., Varambally, S., Kurachi, K., Pienta, K. J., Rubin, M. A. and Chinnaiyan, A. M. (2001). Delineation of prognostic biomarkers in prostate cancer. *Nature* 412, 822-6.
- Dong, J. T., Sipe, T. W., Hyytinen, E. R., Li, C. L., Heise, C., McClintock, D. E., Grant, C. D., Chung, L. W. and Frierson, H. F., Jr. (1998). PTEN/MMAC1 is infrequently mutated in pT2 and pT3 carcinomas of the prostate. *Oncogene* 17, 1979-82.
- Dong, J. T., Suzuki, H., Pin, S. S., Bova, G. S., Schalken, J. A., Isaacs, W. B., Barrett, J. C. and Isaacs, J. T. (1996). Down-regulation of the KAI1 metastasis suppressor gene during the progression of human prostatic cancer infrequently involves gene mutation or allelic loss. *Cancer Res* 56, 4387-90.
- Dunn, J. J. and Studier, F. W. (1983). Complete nucleotide sequence of bacteriophage T7 DNA and the locations of T7 genetic elements. *J Mol Biol* 166, 477-535.
- Eastham, J. A., Grafton, W., Martin, C. M. and Williams, B. J. (2000). Suppression of primary tumor growth and the progression to metastasis with p53 adenovirus in human prostate cancer. *J Urol* 164, 814-9.
- Eisen, M. B., Spellman, P. T., Brown, P. O. and Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci U S A* 95, 14863-8.
- Emmert-Buck, M. R., Bonner, R. F., Smith, P. D., Chuquui, R. F., Zhuang, Z., Goldstein, S. R., Weiss, R. A. and Liotta, L. A. (1996). Laser capture microdissection. *Science* 274, 998-1001.
- Eto, K., Puzon-McLaughlin, W., Sheppard, D., Sehara-Fujisawa, A., Zhang, X. P. and Takada, Y. (2000). RGD-independent binding of integrin alpha9beta1 to the ADAM-12 and -15 disintegrin domains mediates cell-cell interaction. *J Biol Chem* 275, 34922-30.
- Fair, W. R., Aprikian, A. G., Cohen, D., Sogani, P. and Reuter, V. (1993). Use of neoadjuvant androgen deprivation therapy in clinically localized prostate cancer. *Clin Invest Med* 16, 516-22.
- Fambrough, D., McClure, K., Kazlauskas, A. and Lander, E. S. (1999). Diverse signaling pathways activated by growth factor receptors induce broadly overlapping, rather than independent, sets of genes. *Cell* 97, 727-41.
- Fearnhead, N. S., Britton, M. P. and Bodmer, W. F. (2001). The ABC of APC. *Hum Mol Genet* 10, 721-33.
- Feilotter, H. E., Nagai, M. A., Boag, A. H., Eng, C. and Mulligan, L. M. (1998). Analysis of PTEN and the 10q23 region in primary prostate carcinomas. *Oncogene* 16, 1743-8.
- Finch, P. W., He, X., Kelley, M. J., Uren, A., Schaudies, R. P., Popescu, N. C., Rudikoff, S., Aaronson, S. A., Varmus, H. E. and Rubin, J. S. (1997). Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action. *Proc Natl Acad Sci U S A* 94, 6770-5.

- Fink, L., Seeger, W., Ermert, L., Hanze, J., Stahl, U., Grimminger, F., Kummer, W. and Bohle, R. M. (1998). Real-time quantitative RT-PCR after laser-assisted cell picking. *Nat Med* 4, 1329-33.
- Fodor, S. P., Rava, R. P., Huang, X. C., Pease, A. C., Holmes, C. P. and Adams, C. L. (1993). Multiplexed biochemical assays with biological chips. *Nature* 364, 555-6.
- Fodor, S. P., Read, J. L., Pirrung, M. C., Stryer, L., Lu, A. T. and Solas, D. (1991). Light-directed, spatially addressable parallel chemical synthesis. *Science* 251, 767-73.
- Fritz, G., Just, I. and Kaina, B. (1999). Rho GTPases are over-expressed in human tumors. *Int J Cancer* 81, 682-7.
- Gao, A. C., Lou, W., Dong, J. T. and Isaacs, J. T. (1997). CD44 is a metastasis suppressor gene for prostatic cancer located on human chromosome 11p13. *Cancer Res* 57, 846-9.
- Gao, C., Zou, Z., Xu, L., Moul, J., Seth, P. and Srivastava, S. (2000). p53-dependent induction of heat shock protein 27 (HSP27) expression. *Int J Cancer* 88, 191-4.
- Gibas, Z. and Gibas, L. (1997). Cytogenetics of bladder cancer. *Cancer Genet Cytogenet* 95, 108-15.
- Gleason, D. F. (1992). Histologic grading of prostate cancer: a perspective. *Hum Pathol* 23, 273-9.
- Golub, T. R., Slonim, D. K., Tamayo, P., Huard, C., Gaasenbeek, M., Mesirov, J. P., Coller, H., Loh, M. L., Downing, J. R., Caligiuri, M. A. et al. (1999). Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science* 286, 531-7.
- Gormley, G. J. (1991). Role of 5 alpha-reductase inhibitors in the treatment of advanced prostatic carcinoma. *Urol Clin North Am* 18, 93-8.
- Guehenneux, F., Duret, L., Callanan, M. B., Bouhas, R., Hayette, S., Berthet, C., Samarut, C., Rimokh, R., Birot, A. M., Wang, Q. et al. (1997). Cloning of the mouse BTG3 gene and definition of a new gene family (the BTG family) involved in the negative control of the cell cycle. *Leukemia* 11, 370-5.
- Guelstein, V. I., Tchypysheva, T. A., Ermilova, V. D., Litvinova, L. V., Troyanovsky, S. M. and Bannikov, G. A. (1988). Monoclonal antibody mapping of keratins 8 and 17 and of vimentin in normal human mammary gland, benign tumors, dysplasias and breast cancer. *Int J Cancer* 42, 147-53.
- Hall, C., Sin, W. C., Teo, M., Michael, G. J., Smith, P., Dong, J. M., Lim, H. H., Manser, E., Spurr, N. K., Jones, T. A. et al. (1993). Alpha 2-chimerin, an SH2-containing GTPase-activating protein for the ras-related protein p21rac derived by alternate splicing of the human n-chimerin gene, is selectively expressed in brain regions and testes. *Mol Cell Biol* 13, 4986-98.
- Harrington, K. J., Spitzweg, C., Bateman, A. R., Morris, J. C. and Vile, R. G. (2001). Gene therapy for prostate cancer: current status and future prospects. *J Urol* 166, 1220-33.
- Heidenberg, H. B., Sesterhenn, I. A., Gaddipati, J. P., Weghorst, C. M., Buzard, G. S., Moul, J. W. and Srivastava, S. (1995). Alteration of the tumor suppressor gene p53 in a high fraction of hormone refractory prostate cancer. *J Urol* 154, 414-21.

- Henshall, S. M., Quinn, D. I., Lee, C. S., Head, D. R., Golovsky, D., Brenner, P. C., Delprado, W., Stricker, P. D., Grygiel, J. J. and Sutherland, R. L. (2001). Altered expression of androgen receptor in the malignant epithelium and adjacent stroma is associated with early relapse in prostate cancer. *Cancer Res* 61, 423-7.
- Hermanek, P. and Sobin, L.H. (1992). TMN classification of malignant tumours UICC, 5th edn. Springer, Berlin Heidelberg New York Tokyo.
- Herren, B., Garton, K. J., Coats, S., Bowen-Pope, D. F., Ross, R. and Raines, E. W. (2001). ADAM15 Overexpression in NIH3T3 Cells Enhances Cell-Cell Interactions. *Exp Cell Res* 271, 152-60.
- Hsieh, J. C., Kodjabachian, L., Rebbert, M. L., Rattner, A., Smallwood, P. M., Samos, C. H., Nusse, R., Dawid, I. B. and Nathans, J. (1999). A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature* 398, 431-6.
- Huang, C. H. and Liu, P. Z. (2001). New insights into the Rh superfamily of genes and proteins in erythroid cells and nonerythroid tissues. *Blood Cells Mol Dis* 27, 90-101.
- Hudson, D. L., Guy, A. T., Fry, P., O'Hare, M. J., Watt, F. M. and Masters, J. R. (2001). Epithelial cell differentiation pathways in the human prostate: identification of intermediate phenotypes by keratin expression. *J Histochem Cytochem* 49, 271-8.
- Huggins, C. and Hodges, C.V. (1941). Studies on prostate cancer. I. The effect of castration, of estrogen and of androgen injection on serum phosphatases in metastatic carcinoma of the prostate, *Cancer Res.*, I, 293.
- Ittmann, M. (1996). Allelic loss on chromosome 10 in prostate adenocarcinoma. *Cancer Res* 56, 2143-7.
- Janes, P. W., Daly, R. J., deFazio, A. and Sutherland, R. L. (1994). Activation of the Ras signalling pathway in human breast cancer cells overexpressing erbB-2. *Oncogene* 9, 3601-8.
- Johnston, R. F., Pickett, S. C. and Barker, D. L. (1990). Autoradiography using storage phosphor technology. *Electrophoresis* 11, 355-60.
- Kallakury, B. V., Sheehan, C. E. and Ross, J. S. (2001). Co-downregulation of cell adhesion proteins alpha- and beta-catenins, p120CTN, E-cadherin, and CD44 in prostatic adenocarcinomas. *Hum Pathol* 32, 849-55.
- Katoh, M. (2001). Frequent up-regulation of WNT2 in primary gastric cancer and colorectal cancer. *Int J Oncol* 19, 1003-7.
- Katsuma, S., Nishi, K., Tanigawara, K., Ikawa, H., Shiojima, S., Takagaki, K., Kaminishi, Y., Suzuki, Y., Hirasawa, A., Ohgi, T. et al. (2001). Molecular monitoring of bleomycin-induced pulmonary fibrosis by cDNA microarray-based gene expression profiling. *Biochem Biophys Res Commun* 288, 747-51.
- Kim, H. S., Woo, D. K., Bae, S. I., Kim, Y. I. and Kim, W. H. (2001). Allelotype of the adenoma-carcinoma sequence of the stomach. *Cancer Detect Prev* 25, 237-44.
- Kimura, A., Baumann, C. A., Chiang, S. H. and Saltiel, A. R. (2001). The sorbin homology domain: a motif for the targeting of proteins to lipid rafts. *Proc Natl Acad Sci U S A* 98, 9098-103.

- Klenow, H. and Henningsen, I. (1970). Selective elimination of the exonuclease activity of the deoxyribonucleic acid polymerase from Escherichia coli B by limited proteolysis. *Proc Natl Acad Sci U S A* *65*, 168-75.
- Knowles, M. A. (1999). Identification of novel bladder tumour suppressor genes. *Electrophoresis* *20*, 269-79.
- Knowles, M. A., Elder, P. A., Williamson, M., Cairns, J. P., Shaw, M. E. and Law, M. G. (1994). Allelotype of human bladder cancer. *Cancer Res* *54*, 531-8.
- Knudson, A. G., Jr. (1971). Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci U S A* *68*, 820-3.
- Kostic, C. and Shaw, P. H. (2000). Isolation and characterization of sixteen novel p53 response genes. *Oncogene* *19*, 3978-87.
- Kozma, R., Ahmed, S., Best, A. and Lim, L. (1996). The GTPase-activating protein n-chimaerin cooperates with Rac1 and Cdc42Hs to induce the formation of lamellipodia and filopodia. *Mol Cell Biol* *16*, 5069-80.
- Kuczyk, M. A., Bokemeyer, C., Hartmann, J., Schubach, J., Walter, C., Machtens, S., Knuchel, R., Kollmannsberger, C., Jonas, U. and Serth, J. (2001). Predictive value of altered p27Kip1 and p21WAF/Cip1 protein expression for the clinical prognosis of patients with localized prostate cancer. *Oncol Rep* *8*, 1401-7.
- Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W. et al. (2001). Initial sequencing and analysis of the human genome. *Nature* *409*, 860-921.
- Lee, L. G., Connell, C. R. and Bloch, W. (1993). Allelic discrimination by nick-translation PCR with fluorogenic probes. *Nucleic Acids Res* *21*, 3761-6.
- Lennon, G., Auffray, C., Polymeropoulos, M. and Soares, M. B. (1996). The I.M.A.G.E. Consortium: an integrated molecular analysis of genomes and their expression. *Genomics* *33*, 151-2.
- Li, J., Yen, C., Liaw, D., Podsypanina, K., Bose, S., Wang, S. I., Puc, J., Miliarexis, C., Rodgers, L., McCombie, R. et al. (1997). PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* *275*, 1943-7.
- Lin, W. H., Huang, C. J., Liu, M. W., Chang, H. M., Chen, Y. J., Tai, T. Y. and Chuang, L. M. (2001). Cloning, mapping, and characterization of the human sorbin and SH3 domain containing 1 (SORBS1) gene: a protein associated with c-Abl during insulin signaling in the hepatoma cell line Hep3B. *Genomics* *74*, 12-20.
- Liu, H. C., He, Z. and Rosenwaks, Z. (2001). Application of complementary DNA microarray (DNA chip) technology in the study of gene expression profiles during folliculogenesis. *Fertil Steril* *75*, 947-55.
- Luo, L., Salunga, R. C., Guo, H., Bittner, A., Joy, K. C., Galindo, J. E., Xiao, H., Rogers, K. E., Wan, J. S., Jackson, M. R. et al. (1999). Gene expression profiles of laser-captured adjacent neuronal subtypes. *Nat Med* *5*, 117-22.
- Luzzi, V., Holtschlag, V. and Watson, M. A. (2001). Expression profiling of ductal carcinoma in situ by laser capture microdissection and high-density oligonucleotide arrays. *Am J Pathol* *158*, 2005-10.

- Ma, J. F., Takito, J., Vijayakumar, S., Peehl, D. M., Olsson, C. A. and Al-Awqati, Q. (2001). Prostatic expression of hensin, a protein implicated in epithelial terminal differentiation. *Prostate* 49, 9-18.
- Mandai, K., Nakanishi, H., Satoh, A., Takahashi, K., Satoh, K., Nishioka, H., Mizoguchi, A. and Takai, Y. (1999). Ponsin/SH3P12: an l-afadin- and vinculin-binding protein localized at cell-cell and cell-matrix adherens junctions. *J Cell Biol* 144, 1001-17.
- Martin, K. J., Kritzman, B. M., Price, L. M., Koh, B., Kwan, C. P., Zhang, X., Mackay, A., O'Hare, M. J., Kaelin, C. M., Mutter, G. L. et al. (2000). Linking gene expression patterns to therapeutic groups in breast cancer. *Cancer Res* 60, 2232-8.
- Melkonyan, H. S., Chang, W. C., Shapiro, J. P., Mahadevappa, M., Fitzpatrick, P. A., Kiefer, M. C., Tomei, L. D. and Umansky, S. R. (1997). SARP: a family of secreted apoptosis-related proteins. *Proc Natl Acad Sci U S A* 94, 13636-41.
- Miyamoto, H., Shuin, T., Ikeda, I., Hosaka, M. and Kubota, Y. (1996). Loss of heterozygosity at the p53, RB, DCC and APC tumor suppressor gene loci in human bladder cancer. *J Urol* 155, 1444-7.
- Monaco, C., Negrini, M., Sozzi, G., Veronese, M. L., Vorechovsky, I., Godwin, A. K. and Croce, C. M. (1997). Molecular cloning and characterization of LOH11CR2A, a new gene within a refined minimal region of LOH at 11q23. *Genomics* 46, 217-22.
- Morote, J., de Torres, I., Caceres, C., Vallejo, C., Schwartz, S., Jr. and Reventos, J. (1999). Prognostic value of immunohistochemical expression of the c-erbB-2 oncoprotein in metastatic prostate cancer. *Int J Cancer* 84, 421-5.
- Mostofi, F. K., Davis, C. J., Jr. and Sesterhenn, I. A. (1992). Pathology of carcinoma of the prostate. *Cancer* 70, 235-53.
- Murphy, M., Stinnakre, M. G., Senamaud-Beaufort, C., Winston, N. J., Sweeney, C., Kubelka, M., Carrington, M., Brechot, C. and Sobczak-Thepot, J. (1997). Delayed early embryonic lethality following disruption of the murine cyclin A2 gene. *Nat Genet* 15, 83-6.
- Narla, G., Heath, K. E., Reeves, H. L., Li, D., Giono, L. E., Kimmelman, A. C., Glucksman, M. J., Narla, J., Eng, F. J., Chan, A. M. et al. (2001). KLF6, a candidate tumor suppressor gene mutated in prostate cancer. *Science* 294, 2563-6.
- Naz, R. (1997). *Prostate: Basic and Clinical Aspects*, CRC Press New York.
- Niederacher, D., Picard, F., van Roeyen, C., An, H. X., Bender, H. G. and Beckmann, M. W. (1997). Patterns of allelic loss on chromosome 17 in sporadic breast carcinomas detected by fluorescent-labeled microsatellite analysis. *Genes Chromosomes Cancer* 18, 181-92.
- Nupponen, N. and Visakorpi, T. (1999). Molecular biology of progression of prostate cancer. *Eur Urol* 35, 351-4.
- Nupponen, N. N. and Visakorpi, T. (2000). Molecular cytogenetics of prostate cancer. *Microsc Res Tech* 51, 456-63.
- Nusse, R. and Varmus, H. E. (1982). Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. *Cell* 31, 99-109.
- Obata, K. and Hoshiai, H. (2000). Common genetic changes between endometriosis and ovarian cancer. *Gynecol Obstet Invest* 50, 39-43.

- Ohyama, H., Zhang, X., Kohno, Y., Alevizos, I., Posner, M., Wong, D. T. and Todd, R. (2000). Laser capture microdissection-generated target sample for high-density oligonucleotide array hybridization. *Biotechniques* 29, 530-6.
- Orikasa, K., Fukushige, S., Hoshi, S., Orikasa, S., Kondo, K., Miyoshi, Y., Kubota, Y. and Horii, A. (1998). Infrequent genetic alterations of the PTEN gene in Japanese patients with sporadic prostate cancer. *J Hum Genet* 43, 228-30.
- Otsuka, T., Kohno, T., Mori, M., Noguchi, M., Hirohashi, S. and Yokota, J. (1996). Deletion mapping of chromosome 2 in human lung carcinoma. *Genes Chromosomes Cancer* 16, 113-9.
- Pease, A. C., Solas, D., Sullivan, E. J., Cronin, M. T., Holmes, C. P. and Fodor, S. P. (1994). Light-generated oligonucleotide arrays for rapid DNA sequence analysis. *Proc Natl Acad Sci U S A* 91, 5022-6.
- Peifer, M. and Polakis, P. (2000). Wnt signaling in oncogenesis and embryogenesis--a look outside the nucleus. *Science* 287, 1606-9.
- Pennisi, E. (1998). How a growth control path takes a wrong turn to cancer. *Science* 281, 1438-9, 1441.
- Pesche, S., Latil, A., Muzeau, F., Cussenot, O., Fournier, G., Longy, M., Eng, C. and Lidereau, R. (1998). PTEN/MMAC1/TEP1 involvement in primary prostate cancers. *Oncogene* 16, 2879-83.
- Petersen, S., Wolf, G., Bockmuhl, U., Gellert, K., Dietel, M. and Petersen, I. (1998). Allelic loss on chromosome 10q in human lung cancer: association with tumour progression and metastatic phenotype. *Br J Cancer* 77, 270-6.
- Pilarsky, C. P., Schmitt, A. O., Dahl, E. and Rosenthal A. (1999): Microarrays – chances and challenges. *Current Opinion in Molecular Therapeutics* 1: 727-736.
- Pollack, J. R., Perou, C. M., Alizadeh, A. A., Eisen, M. B., Pergamenschikov, A., Williams, C. F., Jeffrey, S. S., Botstein, D. and Brown, P. O. (1999). Genome-wide analysis of DNA copy-number changes using cDNA microarrays. *Nat Genet* 23, 41-6.
- Prochownik, E. V., Eagle Grove, L., Deubler, D., Zhu, X. L., Stephenson, R. A., Rohr, L. R., Yin, X. and Brothman, A. R. (1998). Commonly occurring loss and mutation of the MXI1 gene in prostate cancer. *Genes Chromosomes Cancer* 22, 295-304.
- Pruitt, K. and Der, C. J. (2001). Ras and Rho regulation of the cell cycle and oncogenesis. *Cancer Lett* 171, 1-10.
- Rattner, A., Hsieh, J. C., Smallwood, P. M., Gilbert, D. J., Copeland, N. G., Jenkins, N. A. and Nathans, J. (1997). A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc Natl Acad Sci U S A* 94, 2859-63.
- Remmele, W. (1997a), Pathologie Band 5, Springer Verlag Berlin Heidelberg New York Tokyo, 68-94.
- Remmele, W. (1997b), Pathologie Band 5, Springer Verlag Berlin Heidelberg New York Tokyo, 195-250.
- Ribon, V., Herrera, R., Kay, B. K. and Saltiel, A. R. (1998). A role for CAP, a novel, multifunctional Src homology 3 domain-containing protein in formation of actin stress fibers and focal adhesions. *J Biol Chem* 273, 4073-80.

- Richter, J., Jiang, F., Gorog, J. P., Sartorius, G., Egenter, C., Gasser, T. C., Moch, H., Mihatsch, M. J. and Sauter, G. (1997). Marked genetic differences between stage pTa and stage pT1 papillary bladder cancer detected by comparative genomic hybridization. *Cancer Res* 57, 2860-4.
- Rosin, M. P., Cairns, P., Epstein, J. I., Schoenberg, M. P. and Sidransky, D. (1995). Partial allelotype of carcinoma in situ of the human bladder. *Cancer Res* 55, 5213-6.
- Ross, D. T. and Perou, C. M. (2001). A comparison of gene expression signatures from breast tumors and breast tissue derived cell lines. *Dis Markers* 17, 99-109.
- Rübben, H. (1997a), Uroonkologie, Springer Verlag Berlin Heidelberg New York Tokyo, 233-399.
- Rübben, H. (1997b), Uroonkologie, Springer Verlag Berlin Heidelberg New York Tokyo, 87-178.
- Sager, R. (1997). Expression genetics in cancer: shifting the focus from DNA to RNA. *Proc Natl Acad Sci U S A* 94, 952-5.
- Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., Mullis, K. B. and Erlich, H. A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239, 487-91.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. and Arnheim, N. (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230, 1350-4.
- Sanger, F., Nicklen, S. and Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* 74, 5463-7.
- Sanseau, P. (2001). Impact of human genome sequencing for in silico target discovery. *Drug Discov Today* 6, 316-323.
- Sar, M., Lubahn, D. B., French, F. S. and Wilson, E. M. (1990). Immunohistochemical localization of the androgen receptor in rat and human tissues. *Endocrinology* 127, 3180-6.
- Sattler, H. P., Rohde, V., Bonkhoff, H., Zwergel, T. and Wullich, B. (1999). Comparative genomic hybridization reveals DNA copy number gains to frequently occur in human prostate cancer. *Prostate* 39, 79-86.
- Schena, M., Shalon, D., Davis, R. W. and Brown, P. O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270, 467-70.
- Schena, M., Shalon, D., Heller, R., Chai, A., Brown, P. O. and Davis, R. W. (1996). Parallel human genome analysis: microarray-based expression monitoring of 1000 genes. *Proc Natl Acad Sci U S A* 93, 10614-9.
- Schmitt, A. O., Specht, T., Beckmann, G., Dahl, E., Pilarsky, C. P., Hinzenmann, B. and Rosenthal, A. (1999). Exhaustive mining of EST libraries for genes differentially expressed in normal and tumour tissues. *Nucleic Acids Res* 27, 4251-60.
- Schmitz, A. A., Govek, E. E., Bottner, B. and Van Aelst, L. (2000). Rho GTPases: signaling, migration, and invasion. *Exp Cell Res* 261, 1-12.

- Seraj, M. J., Harding, M. A., Gildea, J. J., Welch, D. R. and Theodorescu, D. (2000). The relationship of BRMS1 and RhoGDI2 gene expression to metastatic potential in lineage related human bladder cancer cell lines. *Clin Exp Metastasis* 18, 519-25.
- Shigyo, M., Sugano, K., Fukayama, N., Taniguchi, T., Tobisu, K., Fujimoto, H., Doi, N., Niwakawa, M., Shimamura, K., Tsukamoto, T. et al. (1998). Allelic loss on chromosome 9 in bladder cancer tissues and urine samples detected by blunt-end single-strand DNA conformation polymorphism. *Int J Cancer* 78, 425-9.
- Simon, R., Burger, H., Brinkschmidt, C., Bocker, W., Hertle, L. and Terpe, H. J. (1998). Chromosomal aberrations associated with invasion in papillary superficial bladder cancer. *J Pathol* 185, 345-51.
- Simoneau, M., Aboulkassim, T. O., LaRue, H., Rousseau, F. and Fradet, Y. (1999). Four tumor suppressor loci on chromosome 9q in bladder cancer: evidence for two novel candidate regions at 9q22.3 and 9q31. *Oncogene* 18, 157-63.
- Singer, V. L., Jones, L. J., Yue, S. T. and Haugland, R. P. (1997). Characterization of PicoGreen reagent and development of a fluorescence-based solution assay for double-stranded DNA quantitation. *Anal Biochem* 249, 228-38.
- Sirivatanauskorn, Y., Drury, R., Crnogorac-Jurcevic, T., Sirivatanauskorn, V. and Lemoine, N. R. (1999). Laser-assisted microdissection: applications in molecular pathology. *J Pathol* 189, 150-4.
- Soloway, M. S., Hachiya, T., Ruiz, H. E., Gomez, C. C. and Civantos, F. (1993). Significance of androgen deprivation prior to radical prostatectomy, with special reference to prostate-specific antigen. *World J Urol* 11, 221-6.
- Sorlie, T., Perou, C. M., Tibshirani, R., Aas, T., Geisler, S., Johnsen, H., Hastie, T., Eisen, M. B., van de Rijn, M., Jeffrey, S. S. et al. (2001). Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 98, 10869-74.
- Southern, E. M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 98, 503-17.
- Sparks, A. B., Hoffman, N. G., McConnell, S. J., Fowlkes, D. M. and Kay, B. K. (1996). Cloning of ligand targets: systematic isolation of SH3 domain-containing proteins. *Nat Biotechnol* 14, 741-4.
- Spruck, C. H., 3rd, Ohneseit, P. F., Gonzalez-Zulueta, M., Esrig, D., Miyao, N., Tsai, Y. C., Lerner, S. P., Schmutte, C., Yang, A. S., Cote, R. et al. (1994). Two molecular pathways to transitional cell carcinoma of the bladder. *Cancer Res* 54, 784-8.
- Stamey, T. A., Warrington, J. A., Caldwell, M. C., Chen, Z., Fan, Z., Mahadevappa, M., McNeal, J. E., Nolley, R. and Zhang, Z. (2001). Molecular genetic profiling of Gleason grade 4/5 prostate cancers compared to benign prostatic hyperplasia. *J Urol* 166, 2171-7.
- Steiner, M. S., Anthony, C. T., Lu, Y. and Holt, J. T. (1998). Antisense c-myc retroviral vector suppresses established human prostate cancer. *Hum Gene Ther* 9, 747-55.
- Su, A. I., Welsh, J. B., Sapinoso, L. M., Kern, S. G., Dimitrov, P., Lapp, H., Schultz, P. G., Powell, S. M., Moskaluk, C. A., Frierson, H. F., Jr. et al. (2001). Molecular classification of human carcinomas by use of gene expression signatures. *Cancer Res* 61, 7388-93.

- Takle, L. A. and Knowles, M. A. (1996). Deletion mapping implicates two tumor suppressor genes on chromosome 8p in the development of bladder cancer. *Oncogene 12*, 1083-7.
- Tanaka, S., Akiyoshi, T., Mori, M., Wands, J. R. and Sugimachi, K. (1998). A novel frizzled gene identified in human esophageal carcinoma mediates APC/beta-catenin signals. *Proc Natl Acad Sci U S A 95*, 10164-9.
- Ugolini, F., Adelaide, J., Charafe-Jauffret, E., Nguyen, C., Jacquemier, J., Jordan, B., Birnbaum, D. and Pebusque, M. J. (1999). Differential expression assay of chromosome arm 8p genes identifies Frizzled-related (FRP1/FRZB) and Fibroblast Growth Factor Receptor 1 (FGFR1) as candidate breast cancer genes. *Oncogene 18*, 1903-10.
- Ugolini, F., Charafe-Jauffret, E., Bardou, V. J., Geneix, J., Adelaide, J., Labat-Moleur, F., Penault-Llorca, F., Longy, M., Jacquemier, J., Birnbaum, D. et al. (2001). WNT pathway and mammary carcinogenesis: Loss of expression of candidate tumor suppressor gene SFRP1 in most invasive carcinomas except of the medullary type. *Oncogene 20*, 5810-7.
- Van Gelder, R. N., von Zastrow, M. E., Yool, A., Dement, W. C., Barchas, J. D. and Eberwine, J. H. (1990). Amplified RNA synthesized from limited quantities of heterogeneous cDNA. *Proc Natl Acad Sci U S A 87*, 1663-7.
- Velculescu, V. E., Zhang, L., Vogelstein, B. and Kinzler, K. W. (1995). Serial analysis of gene expression. *Science 270*, 484-7.
- Verma, R. S., Manikal, M., Conte, R. A. and Godec, C. J. (1999). Chromosomal basis of adenocarcinoma of the prostate. *Cancer Invest 17*, 441-7.
- Vogelstein, B., Fearon, E. R., Hamilton, S. R., Kern, S. E., Preisinger, A. C., Leppert, M., Nakamura, Y., White, R., Smits, A. M. and Bos, J. L. (1988). Genetic alterations during colorectal-tumor development. *N Engl J Med 319*, 525-32.
- Vogelstein, B. and Kinzler, K. W. (1993). The multistep nature of cancer. *Trends Genet 9*, 138-41.
- Wang, L., Darling, J., Zhang, J. S., Liu, W., Qian, J., Bostwick, D., Hartmann, L., Jenkins, R., Bardenhauer, W., Schutte, J. et al. (2000). Loss of expression of the DRR 1 gene at chromosomal segment 3p21.1 in renal cell carcinoma. *Genes Chromosomes Cancer 27*, 1-10.
- Wang, M., Liu, Y. E., Greene, J., Sheng, S., Fuchs, A., Rosen, E. M. and Shi, Y. E. (1997). Inhibition of tumor growth and metastasis of human breast cancer cells transfected with tissue inhibitor of metalloproteinase 4. *Oncogene 14*, 2767-74.
- Weinstein, J., Jacobsen, F. W., Hsu-Chen, J., Wu, T. and Baum, L. G. (1994). A novel mammalian protein, p55CDC, present in dividing cells is associated with protein kinase activity and has homology to the *Saccharomyces cerevisiae* cell division cycle proteins Cdc20 and Cdc4. *Mol Cell Biol 14*, 3350-63.
- Welsh, J. B., Sapino, L. M., Su, A. I., Kern, S. G., Wang-Rodriguez, J., Moskaluk, C. A., Frierson, H. F., Jr. and Hampton, G. M. (2001). Analysis of gene expression identifies candidate markers and pharmacological targets in prostate cancer. *Cancer Res 61*, 5974-8.
- Wistuba, II, Tang, M., Maitra, A., Alvarez, H., Troncoso, P., Pimentel, F. and Gazdar, A. F. (2001). Genome-wide allelotyping analysis reveals multiple sites of allelic loss in gallbladder carcinoma. *Cancer Res 61*, 3795-800.

- Wodarz, A. and Nusse, R. (1998). Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 14, 59-88.
- Yan, P. S., Chen, C. M., Shi, H., Rahmatpanah, F., Wei, S. H., Caldwell, C. W. and Huang, T. H. (2001). Dissecting Complex Epigenetic Alterations in Breast Cancer Using CpG Island Microarrays. *Cancer Res* 61, 8375-8380.
- Yang, R., Muller, C., Huynh, V., Fung, Y. K., Yee, A. S. and Koeffler, H. P. (1999). Functions of cyclin A1 in the cell cycle and its interactions with transcription factor E2F-1 and the Rb family of proteins. *Mol Cell Biol* 19, 2400-7.
- Zhang, M., Martin, K. J., Sheng, S. and Sager, R. (1998). Expression genetics: a different approach to cancer diagnosis and prognosis. *Trends Biotechnol* 16, 66-71.
- Zhao, J., Richter, J., Wagner, U., Roth, B., Schraml, P., Zellweger, T., Ackermann, D., Schmid, U., Moch, H., Mihatsch, M. J. et al. (1999). Chromosomal imbalances in noninvasive papillary bladder neoplasms (pTa). *Cancer Res* 59, 4658-61.
- Zhou, Z., Wang, J., Han, X., Zhou, J. and Linder, S. (1998). Up-regulation of human secreted frizzled homolog in apoptosis and its down-regulation in breast tumors. *Int J Cancer* 78, 95-9.
- Zhu, X., Kanai, Y., Saito, A., Kondo, Y. and Hirohashi, S. (2000). Aberrant expression of beta-catenin and mutation of exon 3 of the beta-catenin gene in renal and urothelial carcinomas. *Pathol Int* 50, 945-52.
- Zincke, H., Bergstrahl, E. J., Blute, M. L., Myers, R. P., Barrett, D. M., Lieber, M. M., Martin, S. K. and Oesterling, J. E. (1994). Radical prostatectomy for clinically localized prostate cancer: long-term results of 1,143 patients from a single institution. *J Clin Oncol* 12, 2254-63.
- Zitzelsberger, H., Engert, D., Walch, A., Kulka, U., Aubele, M., Hofler, H., Bauchinger, M. and Werner, M. (2001). Chromosomal changes during development and progression of prostate adenocarcinomas. *Br J Cancer* 84, 202-8.
- Zohn, I. M., Campbell, S. L., Khosravi-Far, R., Rossman, K. L. and Der, C. J. (1998). Rho family proteins and Ras transformation: the RHOad less traveled gets congested. *Oncogene* 7, 1415-38.