

Bibliography

- [1] Adams J. P., Anderson A. E., Varga A. W., Dineley K. T., Cook R. G., Pfaffinger P. J., Sweatt J. D.: The A-type potassium channel Kv4.2 is a substrate for the mitogen-activated protein kinase ERK. *J Neurochem* **75** (6): 2277-2287 (2000)
- [2] Adams J. P., Sweatt J. D.: Molecular psychology: Roles for the ERK MAP Kinase cascade in memory. *Annu Rev Pharmacol Toxicol* **42**: 135-163 (2002)
- [3] Ahi J., Radulovic J., Spiess J.: The role of hippocampal signaling cascades in consolidation of fear memory. *Behav Brain Res* **149**: 17-31 (2004)
- [4] Ahn N. G. and Krebs E. G.: Evidence for an epidermal growth factor-stimulated protein kinase cascade in Swiss 3T3 cells. Activation of serine peptide kinase activity by myelin basic protein kinases *in vitro*. *J Biol Chem* **265**:11495-11501 (1990)
- [5] Ahn N. G., Seger R., Bratlien R. L., Krebs E. G.: Growth factor-stimulated phosphorylation cascades: activation of growth factor-stimulated MAP kinase. *Ciba Found Symp.* **164**: 113-126; discussion 126-31 (1992)
- [6] Alberini C. M., Ghirardi M., Metz R., Kandel E. R.: C/EBP is an immediate-early gene required for the consolidation of long-term facilitation in Aplysia. *Cell* **76** (6): 1099-1114 (1994)
- [7] Alessi D. R., Cuenda A., Cohen P., Dudley D. T., Saltiel A. R.: PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase *in vitro* and *in vivo*. *The Journal of Biological Chemistry* **270** (46): 27489-27494 (1995)
- [8] Alonso M., Viola H., Izquierdo I., Medina J. H.: Aversive experiences are associated with a rapid and transient activation of ERKs in the rat hippocampus. *Neurobiol Learn Mem* **77** (1): 119-124 (2002)
- [9] Anderson N. G., Maller J. L., Tonks N. K., Sturgill T. W.: Requirement for integration of signals from two distinct phosphorylation pathways for activation of MAP kinase. *Nature (London)* **343**: 651-653 (1990)
- [10] Anderson A. E., Adams J. P., Qian Y., Cook R. G., Pfaffinger P. J., Sweatt J. D.: Kv4.2 phosphorylation by cyclic AMP-dependent protein kinase. *J Biol Chem* **275** (8): 5337-5346 (2000)

Bibliography

- [11] Atkins C. M., Selcher J. C., Petraitis J. J., Trzaskos J. M., Sweatt J. D.: The MAPK cascade is required for mammalian associative learning. *Nature (Neuroscience)* **1** (7): 602-608 (1998)
- [12] Bailey C. H., Kaang B. K., Chen M., Martin K. C., Lim C. S., Casadio A., Kandel E.R.: Mutation in the phosphorylation sites of MAP kinase blocks learning-related internalization of apCAM in Aplysia sensory neurons. *Neuron* **18** (6): 913-924 (1997)
- [13] Bailey C. H., Chen M.: Long-term sensitization in *Aplysia* increases the number of presynaptic contacts onto the identified gill motor neuron L7. *Proc Natl Acad Sci USA* **85**: 9356-9359 (1988)
- [14] Bailey C. H., Kandel E.R.: Structural changes accompanying memory storage. *Annu Rev Physiol* **55**: 397-426 (1993)
- [15] Barnier J. P., Papin C., Eychéne A., Lecoq O., Calothy G.: The mouse B-raf gene encodes multiple protein isoforms with tissue-specific expression. *J Biol Chem* **270**: 23381-23389 (1995)
- [16] Bartsch D., Ghirardi M., Skehel P. A., Karl K. A., Herder S. P., Chen M., Bailey C. H., Kandel E. R.: *Aplysia* CREB2 represses long-term facilitation: relief of repression converts transient facilitation into long-term functional and structural change. *Cell* **83**(6): 979-92 (1995)
- [17] Bartsch D., Casadio A., Karl K. A., Serodio P., Kandel E. R.: CREB1 encodes a nuclear activator, a repressor, and a cytoplasmic modulator that form a regulatory unit critical for long-term facilitation. *Cell* **95** (2): 211-223 (1998)
- [18] Bartsch D., Ghirardi M., Casadio A., Giustetto M., Karl K. A., Zhu H., Kandel E. R.: Enhancement of memory-related long-term facilitation by ApAF, a novel transcription factor that acts downstream from both CREB1 and CREB2. *Cell* **103** (4): 595-608 (2000)
- [19] Berman D.E., Hazvi S., Rosenblum K., Seger R., Dudai Y.: Specific and differential activation of mitogen-activated protein kinase cascades by unfamiliar taste in the insular cortex of the behaving rat. *J Neurosci* **18** (23): 10037-10044 (1998)
- [20] Bernier L., Castellucci V. F., Kandel E. R., Schwarz J. H.: Facilitatory transmitter causes a selective and prolonged increase in adenosine 3:5'-monophosphate in sensory neurons mediating the gill anf siphon withdrawal reflex in Aplysia. *J Neurosci* **2** (12): 1682-1691 (1982)
- [21] Bitterman M. E., Menzel R., Fietz A., Schafer S.: Classical conditioning of proboscis extension in honeybees (*Apis mellifera*). *J Comp Psychol* **97** (2): 107-119 (1983)

Bibliography

- [22] Bliss T. V. P., Lomo T.: Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Psychol* **232**: 331-356 (1973)
- [23] Bliss T. V., Collingridge G. L.: A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* **361**: 31-39 (1993)
- [24] Blum S., Moore A. N., Adams F., Dash P. K.: A mitogen-activated protein kinase cascade in the CA1/CA2 subfield of the dorsal hippocampus is essential for long-term spatial memory. *The Journal of Neuroscience* **19** (9): 3535-3544
- [25] Bos J. L.: All in the family? New insights and questions regarding interconnectivity of Ras, Rap1 and Ral. *J EMBO* **17**: 6776-6782 (1998)
- [26] Boulton T. G., Cobb M. H.: Identification of multiple extracellular signal-regulated kinases (ERKs) with antipeptide antibodies. *Cell Regul* **2**: 357-371 (1991)
- [27] Bozon B., Kelly A., Josselyn S. A., Silva A. J., Davis S., Laroche S.: MAPK, CREB and *zif268* are all required for the consolidation of recognition memory. *Philos Trans R Soc Lond B Biol Sci* **358** (1432): 805-814 (2003)
- [28] Brambilla R., Gnesutta N., Minichiello L., White G., Roylance A. J., Herron C. E., Ramsey M., Wolfer D. P., Cestari V., Rossi-Arnaud C., Grant S. G., Chapman P. F., Lipp H. P., Sturani E., Klein R.: A role for the Ras signalling pathway in synaptic transmission and long-term memory. *Nature* **390** (6657): 281-286 (1997)
- [29] Brugg B., Matus A.: Phosphorylation determines the binding of microtubule-associated protein 2 (MAP2) to microtubules in living cells. *J Cell Biol* **114**: 735-743 (1991)
- [30] Busca R., Abbe P., Mantoux F., Aberdam E., Peyssonaux C., Eychéne A., Ortonne J. P., Ballotti R.: Ras mediates the cAMP-dependent activation of extracellular signal-regulated kinases (ERKs) in melanocytes. *J EMBO* **19**: 2900-2910 (2000)
- [31] Castellucci V. F., Kandel E. R., Schwartz J. H., Wilson F. D., Nairn A. C., Greengard P.: Intracellular injection of the catalytic subunit of cyclic AMP-dependent protein kinase simulates facilitation of transmitter release underlying behavioral sensitization in Aplysia. *Proc Natl Acad Sci U S A* **77**(12): 7492-7496 (1980)
- [32] Catling A. D., Reuter C. W., Cox M. E., Parsons S. J., Weber M. J.: Partial purification of a mitogen-activated protein kinase kinase activator from bovine brain. Identification as B-Raf or a B-Raf-associated activity. *J Biol Chem* **269**: 30014-30021 (1994)
- [33] Cobb M. H.: MAP kinase pathways. *Progress in Biophysics & Molecular Biology* **71**: 479-500 (1999)

Bibliography

- [34] Coogan A. N., O'Leary D. M., O'Connor J. J.: P42/44 MAP kinase inhibitor PD98059 attenuates multiple forms of synaptic plasticity in rat dentate gyrus *in vitro*. *J Neurophysiol* **81** (1): 103-110 (1999)
- [35] Crews C. M., Alessandrini A., Erikson R. L.: The primary structure of MEK, a protein kinase that phosphorylates the *ERK* gene product. *Science* **258**: 478-480 (1992)
- [36] Crow T. G., Xue-Bian J.-J., Siddiqi V., Neary J. T.: Serotonin activation of the ERK pathway in *Hermisenda*: contribution of calcium-dependent protein kinase C. *J Neurochem* **78**: 358-364 (2001)
- [37] Cuenda A., Rouse J., Doza Y. N., Meier R., Cohen P., Gallagher T. F., Young P. R., Lee J. C.: SB 203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1. *FEBS Lett* **364** (2): 229-233 (1995)
- [38] Dash P. K., Hochner B., Kandel E.R.: Injection of the cAMP-responsive element into the nucleus of Aplysia sensory neurons blocks long-term facilitation. *Nature* **345** (6277): 718-721 (1990)
- [39] Davis H. P., Squire L. R.: Protein synthesis and memory: a review. *Psychol Bull* **96**: 518-559 (1984)
- [40] Davis R. J.: The Mitogen-activated protein kinase signal transduction pathway. *The Journal of Biological Chemistry* **268** (20): 14553-14556 (1993a)
- [41] Davis R. L.: Mushroom bodies and *Drosophila* learning. *Neuron* **11**: 1-14 (1993b)
- [42] Davis S., Vanhoutte P., Pages C., Caboche J., Laroche S.: The MAPK/ERK cascade targets both Elk-1 and cAMP response element-binding protein to control long-term potentiation-dependent gene expression in the dentate gyrus *in vivo*. *J Neurosci* **20** (12): 4563-4572 (2000)
- [43] de Belle J. S., Heisenberg M.: Associative odor learning in *Drosophila* abolished by chemical ablation of mushroom bodies. *Science* **263**: 692-695 (1994)
- [44] De Cesare D., Sassone-Corsi P.: Transcriptional regulation by cyclic AMP-responsive factors. *Prog Nucleic Acid Res Mol Biol* **64**: 343-369 (2000)
- [45] haka A., Costa R. M., Hu H., Irvin D. K., Patel A., Kornblum H. I., Silva A. J., O'Dell T. J., Colicelli J.: The RAS effector RIN1 modulates the formation of aversive memories. *J Neurosci* **23** (3): 748-757 (2003)
- [46] Dubnau J., Tully T.: Gene discovery in *Drosophila*: new insights for learning and memory. *Annu Rev Neurosci* **21**: 407-444 (1998)

Bibliography

- [47] Dudley D. T., Pang L., Decker S. J., Bridges A. J., Saltiel A. R.: A synthetic inhibitor of the mitogen-activated protein kinase cascade. *Proc Natl Acad Sci USA* **92**: 7686-7689 (1995)
- [48] Dugan L. L., Kim J. S., Zhang Y., Bart R. D., Sun Y., Holzman D. M., Gutmann D. H.: Differential effects of cAMP in neurons and astrocytes. Role of B-Raf. *J Biol Chem* **274**: 25842-25848 (1999)
- [49] Edwards T. M., Rickard N. S., Ng K. T.: Inhibition of guanylate cyclase and protein kinase G impairs retention for the passive avoidance task in the day-old chick. *Neurobiol Learn Mem* **77** (3): 313-326 (2002)
- [50] Eisenhardt D., Friedrich A., Stollhof N., Müller U., Kress H., Menzel R.: The *AmCREB* gene is an ortholog of the mammalian CREB/CREM family of transcription factors and encodes several splice variants in the honeybee brain. *Insect Molecular Biology* **12** (4): 373-382 (2003)
- [51] Englano W., Rezzonico R., Durand-Clement M., Lallemand D., Ortonne J. P., Ballotti R.: Mitogen-activated protein kinase pathway and AP-1 are activated during cAMP-induced melanogenesis in B-16 melanoma cells. *J Biol Chem* **270** (41): 24315-24320 (1995)
- [52] English J.D., Sweatt J. D.: Activation of p42 Mitogen-activated protein kinase in hippocampal long term potentiation. *The Journal of Biological Chemistry* **271** (40): 24329-24332 (1996)
- [53] English J. D., Sweatt J. D.: A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J Biol Chem* **272** (31): 19103-19106 (1997)
- [54] Erber J., Masuhr T. H., Menzel R.: Localization of short-term memory in the brain of the bee, *Apis mellifera*. *Physiological Entomology* **5**: 343-358 (1980)
- [55] Favata M. F., Horiuchi K. Y., Manos E. J., Daulerio A. J., Stradley D. A., Feeser W. S., Van Dyk D. E., Pitts W. J., Earl R. A., Hobbs F., Copeland R. A., Magolda R. L., Scherle P. A., Trzaskos J. M.: Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J Biol Chem* **273** (29): 18623-18632 (1998)
- [56] Feany M. B., Quinn W. G.: A neuropeptide gene defined by the *Drosophila* memory mutant *amnesiac*. *Science* **268** (5212): 869-873 (1995)
- [57] Fiala A., Müller U., Menzel R.: Reversible downregulation of protein kinase A during olfactory learning using antisense technique impairs long-term memory formation in the honeybee, *Apis mellifera*. *J Neurosci* **19** (22): 10125-10134 (1999)

Bibliography

- [58] Fiore R. S., Bayer V. E., Pelech S. L., Posada J., Cooper J. A., Baraban J. M.: p42 mitogen-activated protein kinase in brain: prominent localization in neuronal cell bodies and dendrites. *Neuroscience* **55** (2): 463-472 (1993)
- [59] Fisher C. E., Michael L., Bartnett M. W., Davies J. A.: Erk MAPkinase regulates branching morphogenesis in the developing mouse kidney. *Development* **128**: 4329-4338 (2001)
- [60] Frank D. A., Greenberg M. E.: CREB: a mediator of long-term memory from mollusks to mammals. *Cell* **79**: 5-8 (1994)
- [61] Frazier W. T., Kandel E. R., Kupfermann I., Waziri R., Coggshall R. E.: Direct and common connections among identified neurons in Aplysia. *J Neurophysiol* **30**(6):1352-76 (1967)
- [62] Frey U., Krug M., Reymann K. G., Matthies H.: Anisomysine, an inhibitor of protein synthesis, blocks late phases of LTP phenomena in hippocampal CA1 neurons. *Brain Res* **452**: 57-65 (1988)
- [63] FrodinM, Peraldi P., Van Obberghen E.: Cyclic AMP activates the mitogen-activated protein kinase cascade in PC12 cells. *J Biol Chem* **269**: 6207-6214 (1994)
- [64] Froese A., Stollhoff N., Eisenhardt D.: AmCREB in memory consolidation of the honeybee *Apis Mellifera*. CD-Supplement to Göttinger Neuroforum, ISSN 0947-0875 (2005)
- [65] Gerber B., Wüstenberg D., Schutz A., Menzel R.: Temporal determinants of olfactory long-term retention in honeybee classical conditioning: nonmonotonous effects of the training trial interval. *Neurobiol Learn Mem* **69** (1): 71-78 (1998)
- [66] Giovannini M. G., Blitzer R. D., Wong T., Asoma K., Tsokas P., Morrison J. H., Iyengar R., Landau E.M.: Mitogen-activated protein kinase regulates early phosphorylation and delayed expression of Ca^{2+} /calmodulin-dependent protein kinase II in long-term potentiation. *J Neurosci* **21** (18): 7053-7062 (2001)
- [67] Giovannini M. G., Efoudebe M., Passani M. B., Baldi E., Bucherelli C., Giachi F., Corradetti R., Blandina P.: Improvement in fear memory by histamine-elicited ERK2 activation in hippocampal CA3 cells. *J Neurosci* **23** (27): 9016-9023 (2003)
- [68] Greengard P.: Possible role for cyclic nucleotides and phosphorylated membrane proteins in postsynaptic actions of neurotransmitters. *Nature* **260** 101 (1976)
- [69] Groves P. M., Thompson R. F.: Habituation: a dual-process theory. *Psychol Rev* **77** (5): 419-450 (1970)

- [70] Grünbaum L., Müller U.: Induction of a specific olfactory memory leads to a long-lasting activation of protein kinase C in the antennal lobe of the Honeybee. *J Neurosci* **18** (11): 4384-4392 (1998)
- [71] Guan Z., Kim J-H., Lomvardas S., Holick K., Xu S., Kandel E. R., Schwartz J. H.: p38 MAP kinase mediates both short-term and long-term synaptic depression in *Aplysia*. *J Neurosci* **23** (19): 7317-7325 (2003)
- [72] Guay J., Lambert H., Gingras-Breton G., Lavoie J. N., Huot J., Landry J.: Regulation of actin filament dynamics by p38 map kinase-mediated phosphorylation of heat shock protein 27. *J Cell Sci* **110** (Pt 3): 357-368 (1997)
- [73] Hammer M., Menzel R.: Learning and memory in the honeybee. *J Neurosci* **3** Pt 1: 1617-1630 (1995)
- [74] Hammer M., Menzel R.: Multiple sites of associative odor learning as revealed by local brain microinjections of octopamine in honeybees. *Learning & Memory* **5** (1): 146-156 (1998)
- [75] Hammer M.: The neural basis of associative reward learning in honeybees. *Trends Neurosci* **20**: 245-252 (1997)
- [76] Han P. L., Levin L. R., Reed R. R., Davis R. L.: Preferential expression of the *Drosophila rutabaga* gene in mushroom bodies, neural centers for learning in insects. *Neuron* **9** (4): 619-627 (1992)
- [77] Haystead T. A. J., Dent P., Wu, J., Haystead, C. M. M., Sturgill T. W.: Ordered phosphorylation of p42 MAPK by MAP kinase kinase. *FEBS Lett* **306**: 17-22 (1992)
- [78] Heisenberg M., Borst A., Wagner S., Byers D.: *Drosophila* mushroom body mutants are deficient in olfactory learning. *J Neurogenetics* **2**: 1-30 (1985)
- [79] Hendron E., Stockand J. D.: Activation of Mitogen-activated protein kinase (Mitogen-activated protein kinase/Extracellular signal-regulated kinase) cascade by aldosterone. *Molecular Biology of the Cell* **13**: 3042-3054 (2002)
- [80] Hildebrandt H., Müller U.: Octopamine mediates rapid stimulation of protein kinase A in the antennal lobe of honeybees. *J Neurobiol* **27**: 44-50 (1995a)
- [81] Hildebrandt H., Müller U.: PKA activity in the antennal lobe of honeybees is regulated by chemosensory stimulation in vivo. *Brain Res* **679**: 281-288 (1995b)
- [82] Hoshi M., Nishida E., Sakai H.: Activation of a Ca^{2+} -inhibitable protein kinase that phosphorylates microtubule-associated protein 2 in vitro by growth factors, phorbol esters, and serum in quiescent cultured human fibroblasts. *J Biol Chem* **263**: 5396-5401 (1988)

Bibliography

- [83] Huang Y. Y., Nguyen P. V., Abel T., Kandel E. R.: Long-lasting forms of synaptic potentiation in the mammalian hippocampus. *Learn Mem* **3** (2-3): 74-85 (1996)
- [84] Impey S., Mark M., Villacres E. C., Poser S., Chavkin C., Storm D. R.: Induction of CRE-mediated gene expression by stimuli that generate long-lasting LTP in area CA1 of the hippocampus. *Neuron* **16**: 973-982 (1996)
- [85] Impey S., Obrietan K., Wong S. T., Poser S., Yano S., Wayman G., Deloulme J. C., Chan G., Storm D. R.: Cross talk between ERK and PKA is required for Ca^{2+} stimulation of CREB-dependent transcription and ERK nuclear translocation. *Neuron* **21**: 869-883 (1998)
- [86] Jones M. W., French P. J., Bliss T. V., Rosenblum K.: Molecular mechanisms of long-term potentiation in the insular cortex *in vivo*. *J Neurosci* **19** (21): RC36 (1999)
- [87] Jovanovic J. N., Benfenati F., Siow Y. L., Sihra T. S., Sanghera J. S., Pelech S. L., Greengard P., Cazernik A. J.: Neurotrophins stimulate phosphorylation of synapsin I by MAP kinase and regulate synapsin I-actin interactions. *Proc Natl Acad Sci* **93**: 3679-3683 (1996)
- [88] Kaang B. K., Kandel E. R., Grant S. G.: Activation of cAMP-responsive genes by stimuli that produce long-term facilitation in Aplysia sensory neurons. *Neuron* **10**(3):427-35 (1993)
- [89] Kanamoto T., Mota M., Takeda K., Rubin L. L., Miyazono K., Ichijo H., Bazenet C. E.: Role of apoptosis signal-regulating kinase in regulation of the c-Jun N-terminal kinase pathway and apoptosis in sympathetic neurons. *Mol Cell Biol* **20** (1): 196-204 (2000)
- [90] Kandel E. R.: The molecular biology of memory storage: A dialogue between genes and synapses. *Science* **294**: 1030-1038 (2001)
- [91] Kandel E. R., Schwartz J. H.: Molecular biology of learning: modulation of transmitter release. *Science* **218**: 433-443 (1982)
- [92] Kane N. S., Robichon A., Dickinson J. A., Greenspan R. J.: Learning without performance in PKC-deficient *Drosophila*. *Neuron* **18** (2): 307-314 (1997)
- [93] Kanterewicz B. I., Urban N. N., McMahon D. B., Norman E. D., Giffen L. J., Favata M. F., Scherle P. A., Trzaskos J. M., Barrionuevo G., Klann E.: The extracellular signal-regulated kinase cascade is required for NMDA receptor-independent LTP in area CA1 but not area CA3 of the hippocampus. *J Neurosci* **20**: 3057-3066 (2000)

Bibliography

- [94] Kawasaki H., Springett G. M., Mochizuki N., Toki S., Nakaya M., Matsuda M., Housman D. E., Graybiel A. M.: A family of cAMP-binding proteins that directly activate Rap1. *Science* **282**: 2275-2279 (1998)
- [95] Kawasaki H., Springett G. M., Toki S., Canales J. J., Harlan P., Blumenstiel J. P., Chen E. J., Bany I. A., Mochizuki N., Ashbacher A., Matsuda M., Housman D. E., Graybiel A. M.: A Rap guanine nucleotide exchange factor enriched highly in the basal ganglia. *Proc Natl Acad Sci U S A* **95** (22): 13278-13283 (1998)
- [96] Kayne P. S., Sternberg P. W.: Ras pathways in *Caenorhabditis elegans*. *Curr Opin Genet Dev* **5**: 38-43 (1995)
- [97] Kelleher R. J. 3rd, Govindarajan A., Jung H. Y., Kang H., Tonegawa S.: Translational control by MAPK signaling in long-term synaptic plasticity and memory. *Cell* **116** (3): 467-479 (2004)
- [98] Klein M., Kandel E. R.: Mechanism of calcium current modulation underlying presynaptic facilitation and behavioral sensitization in Aplysia. *Proc Natl Acad Sci* **77**(11): 6912-6916 (1980)
- [99] Kornhauser J. M., Greenberg M. E.: A kinase to remember: Dual roles for MAP Kinase in long-term memory. *Neuron* **18**: 839-842 (1997)
- [100] Kummer J. L., Rao P. K., Heidenreich K. A.: Apoptosis induced by withdrawal of trophic factors is mediated by p38 mitogen-activated protein kinase. *J Biol Chem* **272** (33): 20490-20494 (1997)
- [101] Larson J., Wong D., Lynch G.: Patterned stimulation of the theta frequency is optimal for the induction of hippocampal long-term potentiation. *Brain Res* **368**: 347-350 (1986)
- [102] Lee J. C., Laydon J. T., McDonnell P. C., Gallagher T. F., Kumar S., Green D., McNulty D., Blumenthal M. J., Heys J. R., Landvatter S. W., et al.: A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* **372** (6508): 739-746 (1994)
- [103] Lee J. C., Kassis S., Kumar S., Badger A., Adams J. L.: p38 mitogen-activated protein kinase inhibitors - mechanisms and therapeutic potentials. *Pharmacol Ther* **82**: 389-397 (1999)
- [104] Lee S. H., Park J., Che Y., Han P. L., Lee J. K.: Constitutive activity and differential localization of p38alpha and p38beta MAPKs in adult mouse brain. *J Neurosci Res* **60** (5): 623-631 (2000)
- [105] Levin L. R., Han P. L., Hwang P. M., Feinstein P. G., Davis R. L., Reed R. R.: The *Drosophila* learning and memory gene *rutabaga* encodes a Ca²⁺/Calmodulin-responsive adenylyl cyclase. *Cell* **68** (3):479-489 (1992)

Bibliography

- [106] Lin L.-L., Wartmann M., Lin A. Y., Knopf J. L., Seth A., Davis R. J.: cPLA₂ is phosphorylated and activated by MAP kinase. *Cell* **72**: 269-278 (1993)
- [107] Lindauer M.: General sensory physiology. Orientation in space. *Fortschr Zool* **16**: 58-140 (1963)
- [108] Liu S., Rao Y., Daw N.: Roles of protein kinase A and protein kinase G in synaptic plasticity in the visual cortex. *Cereb Cortex* **13** (8):864-869 (2003)
- [109] Lu K.-T., Walker D. L., Davis M.: Mitogen-activated protein kinase cascade in the basolateral nucleus of amygdala is involved in extinction of fear-potentiated startle. *The Journal of Neuroscience* **21**: 1-5 (2001)
- [110] Malenka R. C., Nicoll R. A.: Long-term potentiation - a decade of progress? *J Science* **285**: 1870-1874 (1999)
- [111] Mansour S. J., Matten W. T., Hermann A. S., Candia J. M., Rong S., Fukasawa K., Vande Woude G. F., Ahn N. G.: Transformation of mammalian cells by constitutively active MAP kinase kinase. *Science* **265**: 966-970 (1994)
- [112] Marais R., Light Y., Paterson H. F., Mason C. S., Marshall C. J.: Differential regulation of Raf-1, A-Raf and B-Raf by oncogenic ras and tyrosine kinases. *J Biol Chem* **272**: 4378-4383 (1997)
- [113] Marinesco S., Carew T. J.: Serotonin release evoked by tail nerve stimulation in the CNS of *Aplysia*: characterization and relationship to heterosynaptic plasticity. *J Neurosci* **22** (6): 2299-2312 (2002)
- [114] Marshall C. J.: Specificity of receptor tyrosine kinase signalling: Transient versus sustained extracellular signal-regulated kinase activation. *Cell* **80**: 179-185 (1995)
- [115] Martin K. C., Michael D., Rose J. C., Barad M., Casadio A., Zhu H., Kandel E. R.: MAP kinase translocates into nucleus of the presynaptic cell and is required for long-term facilitation in *Aplysia*. *Neuron* **18**: 899-912 (1997)
- [116] Masuhr T., Menzel R.: Learning experiments on the use of sidespecific information in the olfactory and visual system in the honeybee *Apis mellifera*. In: *Information processing in the Visual Systems of Arthropods*, ed: R. Wehner. pp. 315-322 Berlin: Springer (1972)
- [117] Mauelshagen J.: Neural correlates of olfactory learning paradigms in an identified neuron in the honeybee brain. *J Neurophysiol* **69** (2): 609-625 (1993)
- [118] Mauelshagen J., Parker G. R., Carew T. J.: Dynamics of induction of long-term synaptic facilitation in *Aplysia*. *J Neurosci* **16** (22): 7099-7108 (1996)

- [119] McAllister A. K., Katz L. C., Lo D. C.: Neurotrophins and synaptic plasticity. *Annu Rev Neurosci* **22**: 295-318 (1999)
- [120] McBride S. M., Giuliani G., Choi C., Krause P., Correale D., Watson K., Baker G., Siwicki K. K.: Mushroom body ablation impairs short-term memory and long-term memory of courtship conditioning in *Drosophila melanogaster*. *Neuron* **24** (4): 967-977 (1999)
- [121] Menzel R.: Das Gedächtnis der Honigbiene für Spectralfarben. I. Kurzzeitiges und langzeitiges Behalten. *Z. vergl. Physiol.* **60**: 82-102 (1968)
- [122] Menzel R., Erber J., Masuhr T.: Learning and memory in the honeybee. *Experimental Analysis Of Insect Behavior*: 195-217. Berlin: Springer (1974)
- [123] Menzel R.: Learning, memory and “cognition” in honeybees. In: *Neurobiology of Comparative Cognition*: 237-292 New Your: Hillsdale (1990)
- [124] Menzel R.: Memory dynamics in the honeybee. *J Comp Physiol* **185**: 323-340 (1999)
- [125] Menzel R., Müller U.: Learning and memory in honeybees: From behavior to neural substrates. *Annu Rev Neurosci* **19**: 379-404 (1996)
- [126] Menzel R., Erber J., Masuhr T.: Learning and memory in the honeybee. In: *Ex-parimental Analysis of Insect Behaviour*, ed. L. Barton-Browne. pp. 195-217 Berlin: Springer (1974)
- [127] Menzel R., Heyne A., Kinzel C., Gerber B., Fiala A.: Pharmacological dissociation between the reinforcing, sensitizing, and response-releasing functions of reward in honeybee classical conditioning. *Behav Neurosci* **113** (4): 744-754 (1999)
- [128] Menzel R., Manz G., Menzel R., Greggers U.: Massed and spaced learning in honeybees: The role of CS, US, the intertrial interval, and the test interval. *Learning & Memory* **8**: 198-208 (2001)
- [129] Michael D., Martin K. C., Seger R., Ning M.-M., Baston R., Kandel E.: Repeated pulses of serotonin required for long-term facilitation activate mitogen-activated protein kinase in sensory neurons of *Aplysia*. *Proc Natl Acad Sci USA* **95**: 1864-1869 (1998)
- [130] Michels B, Diegelmann S, Tanimoto H, Schwenkert I, Buchner E, Gerber B.: A role for Synapsin in associative learning: The *Drosophila* larva as a study case. *Learn Mem* **12** (3): 224-231 (2005)
- [131] Mikula M., Schreiber M., Husak Z., Kucerova L., Ruth J., Wieser R., Zatloukal K., Beug H., Wagner E. F., Baccarini M.: Embryonic lethality and fetal liver apoptosis in mice lacking the c-raf-1 gene. *J EMBO* **20**: (2001)

Bibliography

- [132] Micheau J., Riedel G.: Protein kinases: which one is the memory molecule? *Cell Mol Life Sci* **55** (4): 534-548 (1999)
- [133] Miyasaka T., Miyasaka J., Saltiel A. R.: Okadaic acid stimulates the activity of microtubule associated protein kinase in PC-12 pheochromocytoma cells. *Biochem Biophys Res Comm* **168**: 1237-1243 (1990)
- [134] Monfort P., Munoz M. D., Kosenko E., Felipo V.: Long-term potentiation in hippocampus involves sequential activation of soluble guanylate cyclase, cGMP-dependent protein kinase, and cGMP-degrading phosphodiesterase. *J Neurosci* **22** (23): 10116-10122 (2002)
- [135] Morris R. G., Garrud P., Rawlins J. N., O'Keefe J.: Place navigation impaired in rats with hippocampal lesions. *Nature* **297** (5868): 681-683 (1982)
- [136] Morozov A., Muzzio I. A., Bourtchouladze R., Van-Strien N., Lapidus K., Yin D., Winder D. G., Adams J. P., Sweatt J. D., Kandel E. R.: Rap1 couples cAMP signaling to a distinct pool of p42/44MAPK regulating excitability, synaptic plasticity, learning, and memory. *Neuron* **39**: 309-325
- [137] Moser E., Moser M. B., Andersen P.: Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions. *J Neurosci* **13** (9): 3916-3925 (1993)
- [138] Müller U.: Inhibition of nitric oxide synthase impairs a distinct form of long-term memory in the honeybee, *Apis mellifera*. *Neuron* **16**: 541-549 (1996)
- [139] Müller U.: Prolonged ativation of cAMP-dependent protein kinase during conditioning induces long-term memory in honeybees. *em Neuron* **27**: 159-168 (2000)
- [140] Müller U.: Learning in honeybees: from molecules to behavior. *Zoology* **105**: 313-320 (2002)
- [141] Müller U., Hildebrandt H.: The nitric oxide/cGMP system in the antennal lobe of *Apis mellifera* is implicated in integrative processing of chemosensory stimuli. *Eur J Neurosci* **7** (11): 2240-2248 (1995)
- [142] Müller U., Carew T. J.: Serotonin induces temporally and mechanistically distinct phases of persistent PKA activity in *Aplysia* sensory neurons. *Neuron* **21**: 1423-1434 (1998)
- [143] Müller U., Hildebrandt H.: Nitric oxide/cGMP-mediated protein kinase activation in the antennal lobes plays an important role in appetitive reflex habituation in the honeybee. *J Neurosci* **22** (19): 8739-8747 (2002)
- [144] New L., Han J.: The p38 MAP kinase pathway and its biological function. *Trends Cardiovasc Med* **8** (5): 220-228 (1998)

- [145] Nighorn A., Healy M. J., Davis R.L.: The cyclic AMP phosphodiesterase encoded by the *Drosophila dunce* gene is concentrated in the mushroom body neuropil. *Neuron* **6** (3): 455-467 1991
- [146] Orban P. C., Chapman P. F., Brambilla R.: Is the Ras-MAPK signalling pathway necessary for long-term memory formation? *Trends Neurosci.* **22** (1): 38-44 (1999)
- [147] Pang L., Sawada T., Decker S. J., Saltiel A. R.: Inhibition of MAP kinase kinase blocks the differentiation of PC-12 cells induced by nerve growth factor. *The Journal of Biological Chemistry* **270** (23): 13585-13588 (1995)
- [148] Papin C., Denouel A., Calothy G., Eychéne A.: Identification of signalling proteins interacting with B-Raf in the yeast two-hybrid system. *Oncogene* **12**: 2213-2221 (1996)
- [149] Papin C., Denouel-Galy A., Laugier D., Calothy G., Eychéne A.: Modulation of kinase activity and oncogenic properties by alternative splicing reveals a novel regulatory mechanism for B-Raf. *J Biol Chem* **273**: 24939-24947 (1998)
- [150] Park J.-M., Kunieda T., Kubo T.: The activity of Mblk-1, a mushroom body-selective transcription factor from the honeybee, is modulated by the Ras/MAPK pathway. *The Journal of Biological Chemistry* **278** (20): 18689-18694 (2003)
- [151] Payne D. M., Rossomando A. J., Martino P., Erickson A. K., Her J. H., Shabanowitz J., Hunt D.F., Weber M. J., Sturgill T. W.: Identification of the regulatory phosphorylation sites in pp42/mitogen-activated protein kinase (MAP kinase). *EMBO J* **10** (4): 885-892 (1991)
- [152] Pelz C., Jander J., Rosenboom H., Hammer M., Menzel R.: I_A in Kenyon cells of the mushroom body of honeybees resembles shaker currents: kinetics, modulation by K^+ , and simulation. *J Neurophysiol* **81** (4): 1749-1759 (1999)
- [153] Peyssonnaux C., Eychene A.: The Raf/MEK/ERK pathway: new concept of activation. *Biology of the Cell* **93**: 53-62 (2001)
- [154] Powell C. M., Johnston D., Sweatt J. D.: Autonomously active protein kinase C in the maintenance phase of N-methyl-D-aspartate receptor-independent long term potentiation. *J Biol Chem* **269** (45): 27958-27963 (1994)
- [155] Pritchard C. A., Samuels M. L., Bosch E., McMahon M.: Conditionally oncogenic forms of the A-Raf and B-Raf protein kinase display different biological and biochemical properties in NIH 3T3 cells. *Mol Cell Biol* **15**: 6430-6442 (1995)
- [156] Purcell A. L., Sharma S. K., Bagnall M. W., Sutton M. A., Carew T. J.: Activation of a tyrosine kinase-MAPK cascade enhances the induction of long-term synaptic facilitation and long-term memory in *Aplysia*. *Neuron* **37**: 473-484 (2003)

Bibliography

- [157] Quinn W. G., Harris W. A., Benzer S.: Conditioned behavior in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **71** (3): 708-712 (1974)
- [158] Quinn W. G., Sziber P. P., Booker R.: The *Drosophila* memory mutant *amnesiac*. *Nature* **277** (5693): 212-214 (1979)
- [159] Raingeaud J., Gupta S., Rogers J. S., Dickens M., Han J., Ulevitch R. J., Davis R. J.: Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J Biol Chem* **270** (13): 7420-7426 (1995)
- [160] Raingeaud J., Whitmarsh A. J., Barrett T., Derijard B., Davis R. J.: MKK3- and MKK6-regulated gene expression is mediated by the p38 mitogen-activated protein kinase signal transduction pathway. *Mol Cell Biol* **16**: 1247-1255 (1996)
- [161] Ray L. B., Sturgill T. W.: Rapid stimulation by insulin of a serine/threonine kinase in 3T3-L1 adipocytes that phosphorylates microtubule-associated protein 2 in vitro. *Proc Natl Acad Sci USA* **84**: 1502-1506 (1987)
- [162] Redman A. M., Johnson J. S., Dally R., Swartz S., Wild H., Paulsen H., Caringal Y., Gunn D., Renick J., Osterhout M., Kingery-Wood J., Smith R. A., Lee W., Dumas J., Wilhelm S. M., Housley T. J., Bhargava A., Ranges G. E., Shrikhande A., Young D., Bombara M., Scott W. J.: p38 kinase inhibitors for the treatment of arthritis and osteoporosis: thieryl, furyl, and pyrrolyl ureas. *Bioorg Med Chem Lett* **11**(1): 9-12 (2001)
- [163] Reuter C. W., Catling A. D., Jelinek T., Weber M. J.: Biochemical analysis of MEK activation in NIH3T3 fibroblasts. Identification of B-Raf and other activators. *J Biol Chem* **270**: 7644-7655 (1995)
- [164] Roberson E. D., English J. D., Sweatt J. D.: A biochemist's view of long-term potentiation. *Learn Mem* **3**(1): 1-24 (1996)
- [165] Robbins D. J., Zhen E., Owaki H., Vanderbilt C., Ebert D., Geppert T. D., Cobb M. H.: Regulation and properties of extracellular signal-regulated protein kinases 1 and 2 *in vitro*. *J Biol Chem* **268**: 5097-5106 (1993)
- [166] Roberson E. D., English J. D., Sweatt J.D.: A biochemist's view of long-term potentiation. *Learn Mem* **3** (1): 1-24 (1996)
- [167] Roberson E.D., English J.D., Adams J.P., Selcher J. C., Kondratick C., Sweatt J. D.: The mitogen-activated protein kinase cascade couples PKA to CREB phosphorylation in area CA1 of the hippocampus. *J Neurosci* **19**: 4337-4348 (1999)
- [168] Rosen L. B., Ginty D. D., Weber M. J., Greenberg M. E.: Membrane depolarisation and calcium Influx stimulate MEK and MAP Kinase via activation of Ras. *Neuron* **12**: 1207-1221 (1994)

Bibliography

- [169] Ruifen B., Foy M. R., Vouimba R-M., Thompson R. F., Baudry: Cyclic changes in estradiol regulate synaptic plasticity through the MAP kinase pathway. *Proc Natl Acad Sci USA* **98** (23): 13391-13395 (2001)
- [170] Rybak J., Menzel R.: Anatomy of the mushroom bodies in the honey bee brain: the neuronal connections of the alpha-lobe. *J Comp Neurol* **334** (3):444-465 (1993)
- [171] Sananbenesi F., Fischer A., Schrick C., Spiess J., Radulovic J.: Phosphorylation of hippocampal Erk-1/2, Elk-1, and p90-Rsk-1 during contextual fear conditioning: interactions between Erk-1/2 and Elk-1. *Mol Cell Neurosci* **21**: 463-476 (2002)
- [172] Sananbenesi F., Fischer A., Schrick C., Spiess J., Radulovic J.: Phosphorylation of hippocampal Erk-1/2, Elk-1, and p90-Rsk-1 during contextual fear conditioning. *Mol Cell Neurosci* **21**: 463-476 (2003)
- [173] Sato K., Morimoto K., Suemaru S., Sato T.: Increased synapsin I immunoreactivity during long-term potentiation in rat hippocampus. *Brain Res* **872** (1-2): 219-222 (2000)
- [174] Schacher S., Castellucci V. F., Kandel E. R.: cAMP evokes long-term facilitation in Aplysia sensory neurons that requires new protein synthesis. *Science* **240** (4859): 1667-1669 (1988)
- [175] Schafe G. E., Atkins C. M., Swank M. W., Bauer E. P., Sweatt J. D., LeDoux J. E.: Activation of ERK/MAPK kinase in the amygdala is required for memory consolidation of Pavlovian fear conditioning. *J Neurosci* **20** (21): 8177-8187 (2000)
- [176] Schmitt J. M., Guire E. S., Saneyoshi T., Soderling T.R.: Calmodulin-dependent kinase kinase/calmodulin kinase I activity gates extracellular-regulated kinase-dependent long-term potentiation. *J Neurosci* **25** (5): 1281-1290 (2005)
- [177] Schulman, D., Latchman, D. S., Yellon D. M.: Urocortin protects the heart from reperfusion injury via upregulation of p 42/p44 MAPK signalling pathway. *Am J Physiol Heart Circ Physiol* **283**: H1481-H1488 (2002)
- [178] Seger R., Ahn N. G., Boulton T.G., Yancopoulos G. D., Panayotatos N., Radziejewska E., Ericsson L., Bratlien R. L., Cobb M. H., Krebs E. G.: Microtubule-associated protein 2 kinases, ERK1 and ERK2, undergo autophosphorylation on both tyrosine and threonine residues: Implication for their mechanism of activation. *Proc Natl Acad Sci USA* **88**: 6142-6146 (1991)
- [179] Seger R., Krebs E. G.: The MAPK signalling cascade. *FASEB J* **9** (9): 726-735 (1995)
- [180] Selcher J. C., Atkins C. M., Trzaskos J. M., Paylor R., Sweatt J. D.: A necessity for MAP kinase activation in mammalian spatial learning. *Learning & Memory* **6**: 478-490 (1999)

Bibliography

- [181] Selcher J. C., Nekrasova T., Paylor R., Landreth G. E., Sweatt J. D.: Mice lacking the ERK1 isoform of MAP kinase are unimpaired in emotional learning. *Learning & Memory* **8**: 11-19 (2001)
- [182] Selcher J. C., Weeber E. J., Christian J., Nekrasova T., Landreth G. E., Sweatt J. D.: A role of ERK MAP kinase in physiologic temporal integration in hippocampal area CA1. *Learning & Memory* **10**: 26-39 (2003)
- [183] Sharma S. K., Sherff C. M., Shobe J., Bagnall M. W., Sutton M. A., Carew T. J.: Differential role of mitogen-activated protein kinase in three distinct phases of memory for sensitization in *Aplysia*. *J Neurosci* **23** (9): 3899-3907 (2003a)
- [184] Sharma S. K., Bagnall M. W., Sutton M. A., Carew T. J.: Inhibition of calcineurin facilitates the induction of memory for sensitization in *Aplysia*: Requirement of mitogen-activated protein kinase. *PNAS* **100** (8): 4861-4866 (2003b)
- [185] Silva A. J., Stevens C. F., Tonegawa S., Wang Y.: Deficient hippocampal long-term potentiation in alpha-calcium-calmodulin kinase II mutant mice. *Science* **257** (5067): 201-206 (1992)
- [186] Soderling T. R., Derkach V. A.: Postsynaptic protein phosphorylation and LTP. *Trends Neurosci* **23** (2): 75-80 (2000)
- [187] Suenaga T., Morinobu S., Kawano K., Sawada T., Yamawaki S.: Influence of immobilization stress on the levels of CaMKII and phospho-CaMKII in the rat hippocampus. *Int J Neuropsychopharmacol* **7** (3) :299-309 (2004)
- [188] Sung Y.J., Walters E. T., Ambron R. T.: A neuronal isoform of protein kinase G couples mitogen-activated protein kinase nuclear import to axotomy-induced long-term hyperexcitability in *Aplysia* sensory neurons. *J Neurosci* **24** (34):7583-7595 (2004)
- [189] Sutton M. A., Carew T. J.: Parallel molecular pathways mediate expression of distinct forms of intermediate-term facilitation at tail sensory-motor synapses in *Aplysia*. *Neuron* **26**: 219-231 (2000)
- [190] Sutton M. A., Masters S. E., Bagnall M. W., Carew T. J.: Molecular mechanisms underlying an unique intermediate phase of memory in *Aplysia*. *Neuron* **31**: 143-154 (2001)
- [191] Swank M. W.: Pharmacological antagonism of tyrosine kinases and MAP kinase in brainstem blocks taste aversion learning in mice. *Physiol Behav* **69** (4-5): 499-503 (2000)
- [192] Sweatt J. D.: Toward a molecular explanation for long-term potentiation. *Learn Mem* **6** (5): 399-416 (1999)

- [193] Sweatt D. J.: The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J Neurochem* **76**: 1-10 (2001)
- [194] Takeda K.: Classical conditioned response in the honeybee. *J Insect Physiol* **6**: 168-179 (1961)
- [195] Takeuchi H., Kage E., Sawata M., Kamikouchi A., Ohashi K., Ohara M., Fujiyuki T., Kunieda T., Sekimizu K., Natori S., Kubo T.: Identification of a novel gene, Mblk-1, that encodes a putative transcription factor expressed preferentially in the large-type Kenyon cells of the honeybee brain. *Insect Mol Biol* **10** (5): 487-94 (2001)
- [196] Tempel B. L., Bonini N., Dawson D. R., Quinn W. G.: Reward learning in normal and mutant *Drosophila*. *Proc Natl Acad Sci U S A* **80** (5): 1482-1486 (1983)
- [197] Thomas G. M., Huganir R. L.: MAPK cascade signalling and synaptic plasticity. *Nature Reviews/Neuroscience* **5**: 173-183 (2004)
- [198] Thomas, U.: Die Rolle der Proteinkinase C bei der Bildung von Langzeitgedächtnis: Interaktion der PKC mit Ankerproteinen im Pilzkörper der Honigbiene *Apis Mellifera*. Inaugural Dissertation am Fachbereich Biologie, Chemie und Pharmazie an der Freien Universität Berlin (2002)
- [199] Thompson R. F., Spencer W. A.: Habituation: a model phenomenon for the study of neuronal substrates of behavior. *Psychol Rev* **73** (1): 16-43 (1966)
- [200] Thurmond R. L., Wadsworth S. A., Schafer P. H., Zivin R. A., Siekierka J. J.: Kinetics of small molecule inhibitor binding to p38 kinase. *Eur J Biochem* **268**: 5747-5754 (2001)
- [201] Treisman R.: Regulation of transcription by MAP kinase cascades. *Curr Opin Cell Biol* **8** (2): 205-215 (1996)
- [202] Tyler W. J., Alonso M., Bramham C. R., Pozzo-Miller L. D.: From acquisition to consolidation: on the role of brain-derived neurotrophic factor signaling in hippocampal-dependent learning. *Learn Mem* **9**: 224-237 (2002)
- [203] Vareschi E.: Duftunterscheidung bei der Honigbiene Einzelzell-Ableitungen und Verhaltensreaktionen. *Z Vergl Physiol* **75**: 143-173 (1971)
- [204] Vossler M. R., Yao H. R., York D., Pan M. G., Rim C. S., Stork P. J.: cAMP activates MAP kinase and Elk-1 through a B-Raf- and Rap1-dependent pathway. *Cell* **89**: 73-82 (1997)
- [205] Wang X., Flynn A., Waskiewicz A. J., Webb B. L., Vries R. G., Baines I. A., Cooper J. A., Proud C. G.: The phosphorylation of eukaryotic initiation factor elF4E in response to phorbol esters, cell stresses, and cytokines is mediated by distinct MAP kinase pathways. *J Biol Chem* **273**: 9373-9377 (1998)

Bibliography

- [206] Wang J., Chen Y.-B., Zhu X.-N., Chen R.-Z.: Activation of p44/42 mitogen-activated protein kinase pathway in long-term potentiation induced by nicotine in hippocampal CA1 region in rats. *Acta Pharmacol Sin* **22** (8): 685-690 (2001)
- [207] Wang J. Q., Tang Q., Parelkar N. K., Liu Z., Samdani S., Choe E. S., Yang L., Mao L.: Glutamate signalling to Ras-MAPK in striatal neurons. *Molecular Neurobiology* **29**: 1-13 (2004)
- [208] Ward S. G., Parry R. V., Matthews J., o'Neill L.: A p38 MAP inhibitor SB 203580 inhibits CD28-dependent T cell proliferation and IL-2 production. *Biochem Soc Trans* **25** (2): 304S
- [209] Wiklund P., Ekstrom P. A., Edstrom A.: Mitogen-activated protein kinase inhibition reveals differences in signalling pathways activated by neurotrophin-3 and other growth-stimulating conditions of adult mouse root ganglia neurons. *J Neurosci Res* **67** (1): 62-68 (2002)
- [210] Witthöft W.: Absolute Anzahl und Verteilung der Zellen in Hirn der Honigbiene. *Z Morphol Tiere* **61**: 160-184 (1967)
- [211] Wittstock S., Menzel R.: Color learning and memory in honey bees are not affected by protein synthesis inhibition. *Behav Neural Biol* **62** (3): 224-229 (1994)
- [212] Wu J., Dent P., Jelinek T., Wolfman A., Weber M. J., Sturgill T. W.: Inhibition of the EGF-activated MAPK signaling pathway by adenosine 3',5'-monophosphate. *Science* **262**: 1065-1072 (1993)
- [213] Wustenberg D., Gerber B., Menzel R.: Short communication: long- but not medium-term retention of olfactory memories in honeybees is impaired by actinomycin D and anisomycin. *Eur J Neurosci* **10** (8): 2742-2745 (1998)
- [214] Yamashita M., Shinnakasu R., Asou H., Kimura M., Hasegawa A., Hashimoto K., Hatano N., Ogata M., Nakayama T.: Ras-ERK MAPK cascade regulates GATA3 stability and Th2 differentiation through ubiquitin-proteasome pathway. *J Biol Chem.* (2005) in press
- [215] Yan M., Dai T., Deak J. C., Kyriakis J. M., Zon L. I., Woodgett J. R., Templeton D. J.: Activation of stress-activated protein kinase by MEKK1 phosphorylation of its activator SEK1. *Nature* **372**: 798-800 (1994)
- [216] Zhang J. J., Okutani F., Inoue S., Kaba H.: Activation of the mitogen-activated protein kinase/extracellular signal-regulated kinase signaling pathway leading to cyclic AMP response element-binding protein phosphorylation is required for the long-term facilitation process of aversive olfactory learning in young rats. *Neuroscience* **121** (1): 9-16 (2003)

Bibliography

- [217] Zhen X., Du W., Romano A. G., Friedman E., Harvey J. A.: The p38 mitogen-activated protein kinase is involved in associative learning in rabbits. *The Journal of Neuroscience* **21** (15): 5513-5519 (2001)