

5. Literaturverzeichnis

- Aggelli, A.**, Bell, M., Boden, N., Keen, J.N., Knowles, P.F., McLeish, T.C., Pitkeathly, M. & Radford, S.E. Responsive gels formed by the spontaneous self-assembly of peptides into polymeric beta-sheet tapes. *Nature* 386, 259-262 (1997).
- Altmann, K.-H.**, Flörsheimer, A. & Mutter, M. Sequence-dependence of secondary structure formation. *Int. J. Peptide Protein Res.*, 27:314-319 (1986).
- Bada, J.L.** *In vivo* racemization of amino acids in mammalian proteins. In: *Methods in Enzymology*, vol 106 (Wald, F. & Moldave, K., eds.) Academic Press, Orlando, Florida, 98-115 (1984).
- Balcerski, J.S.**, Pysh, E.S., Bonora, G.M. & Toniolo, C. *J. Am. Chem. Soc.* 98, 3740-3473 (1976).
- Barrow, C.J.** & Zagorski, M.G. Solution structures of beta peptide and its constituent fragments: relation to amyloid deposition. *Science* 253, 179-182 (1991).
- Barrow, C.J.**, Yasuda, A., Kenny, P.T. & Zagorski, M.G. Solution conformations and aggregational properties of synthetic amyloid β -peptides of Alzheimer's disease. *J. Mol. Biol.* 225, 1075-1093 (1992).
- Baumeister, R.**, & Eimer, S. Amyloid aggregates, presenilins, and Alzheimer's disease. *Angew. Chem. Int. Edit.* 37, 2978-2982 (1998).
- Behlke, J.**, & Ristau, O. Molecular mass determination by sedimentation velocity experiments and direct fitting of the concentration profiles. *Biophys. J.* 72, 428-434 (1997).
- Berger, E.P.**, Rasmussen, K.C., Weinreb, P.H. & Lansbury, P.T., Jr. Rapid, large-scale synthesis and solubility studies of the amyloid β protein of Alzheimer's disease. In: *Peptides. Chemistry, Structure and Biology.*(Hodges R.S. & Smith, J.A, eds.) Escom, Leiden; 502-504 (1994).
- Biere, A.L.**, Ostaszewski, B., Stimson, E.R., Hyman, B.T., Maggio, J.E. & Selkoe, D.J. Amyloid beta-peptide is transported on lipoproteins and albumin in human plasma. *J. Biol. Chem.* 271, 32916-32922 (1996).
- Blaber, M.**, Zhang, X.J., Lindstrom, J.D., Pepiot, S.D., Baase, W.A. & Matthews, B.W. Determination of alpha-helix propensity within the context of a folded protein. Sites 44 and 131 in bacteriophage T4 lysozyme. *J. Mol. Biol.* 235, 600-624 (1994).
- Blanco, F.J.**, Rivas, G. & Serrano, L. A short linear peptide that folds into a native stable beta-hairpin in aqueous solution. *Nat. Struct. Biol.* 1, 584-590 (1994).
- Blondelle, S.E.** & Houghten, R.A. Design of model amphipathic peptides having potent antimicrobial activities. *Biochemistry* 31, 12688-12694 (1992).
- Blondelle, S.E.**, Forood, B., Houghten, R.A. & Perez-Paya, E. Polyalanine based peptides as models for self-associated beta-sheet complexes. *Biochemistry* 36, 8393-8400 (1997).

- Booth, D.R.**, Sunde, M., Bellotti, V., Robinson, C.V., Hutchinson, W.L., Fraser, P.E., Hawkins, P.N., Dobson, C.M., Radford, S.E., Blake, C.C. & Pepys, M.B. (1997) Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. *Nature*, 385, 787-793 (1997).
- Brack, A.** & Orgel, L.E. β -Structures of alternating polypeptides and their possible prebiotic significance. *Nature* 256, 383-387 (1975).
- Braunschweiler, L.** & Ernst, R.R. Coherence transfer by isotopic mixing: application to proton correlation spectroscopy. *J. Magn. Reson.* 53, 521-528 (1983).
- Brunden, K.R.**, Richter-Cook, N.J., Chaturvedi, N. & Frederickson, R.C. pH-dependent binding of synthetic beta-amyloid peptides to glycosaminoglycans. *J. Neurochem.* 61, 2147-2154 (1993).
- Brünger, A.T.** *X-PLOR Manual Version 3.1. A system for X-ray crystallography and NMR*, Yale University Press, New Haven, CT, USA (1993).
- Bush, A.I.**, Pettingell, W.H., Multhaup, G., d-Paradis, M., Vonsattel, J.P., Gusella, J.F., Beyreuther, K., Masters, C.L. & Tanzi, R.E. Rapid induction of Alzheimer A beta amyloid formation by zinc. *Science* 265, 1464-1467 (1994).
- Buxbaum, J.D.**, Liu, K.N., Luo, Y., Slack, J.L., Stocking, K.L., Peschon, J.J., Johnson, R.S., Castner, B.J., Cerretti, D.P. & Black, R.A. Evidence that tumor necrosis factor alpha converting enzyme is involved in regulated alpha-secretase cleavage of the Alzheimer amyloid protein precursor. *J. Biol. Chem.* 273, 27765-27767 (1998).
- Castano, S.**, Desbat, B., & Dufourcq, J. Ideally amphiphatic β -sheeted peptides at interfaces: structure, orientation, affinities for lipids and hemolytic activity of (KL)_mK peptides. *Biochim. Biophys Acta* 1463, 65-80 (2000).
- Carter, D.B.** & Chou, K.C. A model for structure-dependent binding of Congo red to Alzheimer beta-amyloid fibrils. *Neurobiol. of Aging* 19, 37-40 (1998).
- Cerpa, R.**, Cohen, F.E. & Kuntz, I.D. Conformational switching in designed peptides: the helix/sheet transition. *Fold. Des.* 1, 91-101 (1996).
- Chakrabarty, A.**, Kortemme, T. & Baldwin, R.L. Helix propensities of the amino acids measured in alanine-based peptides without helix-stabilizing side-chain interactions. *Protein Sci.* 3, 843-852 (1994).
- Chang, C.D.** & Meienhofer, J. *Int. J. Peptide Protein Res.* 11, 246 (1978).
- Chen, Y.**, Yang, J.T. & Martinez, H.M. Determination of the secondary structures of proteins by circular dichroism and optical rotatory dispersion. *Biochemistry* 11, 4120-4131 (1972).
- Chiti, F.**, Webster, P., Taddei, N., Clark, A., Stefani, M., Ramponi, G. & Dobson, C.M. Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proc. Natl. Acad. Sci. U.S.A.* 96, 3590-3594 (1999).

- Choo, D.W.**, Schneider, J.P., Graciani, N.R. & Kelly, J.W. Nucleated antiparallel beta-sheet that folds and undergoes self-assembly: a template promoted folding strategy toward controlled molecular architectures. *Macromolecules* 29, 355-366 (1996).
- Choo, L.P.**, Wetzel, D.L., Halliday, W.C., Jackson, M., LeVine, S.M. & Mantsch, H.H. In situ characterization of beta-amyloid in Alzheimer's diseased tissue by synchrotron Fourier transform infrared Microspectroscopy. *Biophys. J.* 71, 1672-1679 (1996).
- Chou, P.Y.** & Fasman, G.D. Conformational parameters for amino acids in helical, beta-sheet, and random coil regions calculated from proteins. *Biochemistry* 13, 211-222 (1974).
- Chou, P.Y.** & Fasman, G.D. Prediction of the secondary structure of proteins from their amino acid sequence. *Adv. Enzymol. Relat. Areas Mol. Biol.* 47, 45-148 (1978).
- Cohn, E.J.** & Edstall, J.T. In: *Proteins, Amino Acids and Peptides*, Academic Press, New York (1943).
- Coles, M.**, Bicknell, W., Watson, A.A., Fairlie, D.P., & Craik, D.J. Solution structure of amyloid beta-peptide (1-40) in a water-micelle environment. Is the membrane-spanning domain where we think it is? *Biochemistry* 37, 11064-11077 (1998).
- Cooper, J. H.** Selective amyloid staining as a function of amyloid composition and structure. *Lab. Invest.* 31, 232-238 (1974).
- Creighton, T.E.** (ed.) *Proteins: Structures and Molecular Principles*. Freeman, New York, 2nd Ed. (1993).
- Dalal, S.**, Blasubramanian, S. & Regan, L. Protein alchemy: changing β -sheet into α -helix. *Nat. Struct. Biol.* 4, 548-552 (1997).
- Damaschun, G.**, Damaschun, H., Gast, K. & Zirwer, D. Proteins can adopt totally different folded conformations. *J. Mol. Biol.* 291, 715-725 (1999).
- Davis, D.G.** & Bax, A. Assignments of complex ^1H NMR spectra via two-dimensional homonuclear Hartmann-Hahn spectroscopy. *J. Am. Chem. Soc.* 107, 2820-2821 (1985).
- De Alba, E.**, Santoro, J., Rico, M. & Jimenez, M.A. De novo design of a monomeric three-stranded antiparallel beta-sheet. *Protein Sci.* 8, 854-865 (1999).
- DeGrado, W.F.** & Lear, J.D. Induction of peptide conformation at apolar/water interfaces. *J. Am. Chem. Soc.* 107, 7684-7689 (1985).
- Delaglio, F.**, Grzesiek, S., Vuister, G., Zhu, G., Pfeifer, J. & Bax, A. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J. Biomolecular. NMR* 6, 277-293 (1995).
- De La Torre, G.J.** & Bloomfield, V.A. Hydrodynamic properties of complex, rigid, biological macromolecules: theory and applications. *Q. Rev. Biophys.* 14, 81-139 (1981).

- De Strooper, B.**, Annaert, W., Cupers, P., Saftig, P., Craessaerts, K., Mumm, J.S., Schroeter, E.H., Schrijvers, V., Wolfe, M.S., Ray, W.J., Goate, A. & Kopan, R. A presenilin-1-dependent gamma-secretase-like protease mediates release of Notch intracellular domain. *Nature* 398, 518-522 (1999).
- Derrick, J.P.**, and Wigley, D.B. Crystal structure of a streptococcal protein G domain bound to an Fab fragment. *Nature* 359, 752-754 (1992).
- Dill, K.A.** Dominant forces in protein folding. *Biochemistry* 29, 7133-7155 (1990).
- Dobson, C.M.** Protein misfolding, evolution and disease. *Trends Biochem. Sci.* 24, 329-332(1999).
- Dong, A.**, Prestrelski, S.J., Allison, S.D. & Carpenter, J.F. Infrared spectroscopic studies of lyophilization- and temperature-induced protein aggregation. *J. Pharm. Sci.* 84, 415-424 (1995).
- Du, Y.S.**, Bales, K.R., Dodel, R.C., Liu, X.D., Glinn, M.A., Horn, J.W., Little, S.P. & Paul, S.M. α_2 -Macroglobulin attenuates beta-amyloid peptide 1-40 fibril formation and associated neurotoxicity of cultured fetal rat cortical neurons. *J. Neurochem.* 70, 1182-1188 (1998).
- El Agnaf, O.M.A.**, Guthrie, D.J., Walsh, D.M. & Irvine, G.B. The influence of the central region containing residues 19-25 on the aggregation properties and secondary structure of Alzheimer's beta-amyloid peptide. *Eur. J. Biochem.* 256, 560-569 (1998).
- Esler, W.P.**, Stimson, E.R., Ghilardi, J.R., Lu, Y.A., Felix, A.M., Vinters, H.V., Mantyh, P.W., Lee, J.P. & Maggio, J.E. Point substitution in the central hydrophobic cluster of a human beta-amyloid congener disrupts peptide folding and abolishes plaque competence. *Biochemistry* 35, 13914-13921 (1996).
- Fabian, H.**, Choo, L.-P., Szendrei, G.I., Jackson, M., Halliday, W.C., Otvos, L. & Mantsch, H.H. Infrared spectroscopic characterization of Alzheimer plaques. *Appl. Spectrosc.* 47, 1513-1518 (1993a).
- Fabian, H.**, Schultz, C., Naumann, D., Landt, O., Hahn, U. & Saenger, W. Secondary structure and temperature-induced unfolding and refolding of ribonuclease T1 in aqueous solution. A Fourier transform infrared spectroscopic study. *J. Mol. Biol.* 232, 967-981 (1993b).
- Fauchere, J.L.**, Charton, M., Kier, L.B., Verloop, A. & Pliska, V. Amino acid side chain parameters for correlation studies in biology and pharmacology. *Int. J. Pept. Protein Res.* 32, 269-278 (1988).
- Fink, A.L.** Protein aggregation: folding aggregates, inclusion bodies and amyloid. *Fold. Des.* 3, R9-R23 (1998).
- Forood, B.**, Perez-Paya, E., Houghten, R.A. & Blondelle, S.E. Formation of an extremely stable polyalanine beta-sheet macromolecule. *Biochem. Biophys. Res. Commun.* 211, 7-13 (1995).
- Fraser, P.E.**, McLachlan, D.R., Surewicz, W.K., Mizzen, C.A., Snow, A.D., Nguyen, J.T. & Kirschner, D.A. Conformation and fibrillogenesis of Alzheimer A β peptides with selected substitution of charged residues. *J. Mol. Biol.* 244, 64-73 (1994).

- Fraser, R.D.B.** & MacRae, T.P. Conformation in fibrous proteins. Academic Press, New York (1973).
- Friedreich, N.** & Kekule, A. Zur Amyloidfrage. *Arch. Pathol. Physiol. Klin. Med.* 16, 50-65 (1859).
- Fukuda, H.**, Shimizu, T., Nakajima, M., Moro, H. & Shirasawa, T. Synthesis, aggregation, and neurotoxicity of Alzheimer's A β 1-42 amyloid peptide and its isoaspartyl isomers. *Bioorg. Med. Chem. Lett.* 9, 953-956 (1999).
- Geddes, A.J.**, Parker, K.D., Atkins, E.D.T., Beighton, E. 'Cross- β ' conformation in proteins. *J. Mol. Biol.* 32, 343-358 (1968).
- Geula, C.**, Wu, C.K., Saroff, D., Lorenzo, A., Yuan, M.L. & Yanker, B.A. Aging renders the brain vulnerable to amyloid beta-protein neurotoxicity. *Nature Med.* 4, 827-831 (1998).
- Ghiso, J.**, Calero, M., Matsubara, E., Governale, S., Chuba, J., Beavis, R., Wisniewski, T. & Frangione, B. Alzheimer's soluble amyloid beta is a normal component of human urine. *FEBS Lett.* 408, 105-108 (1997).
- Gill, S.C.** & von Hippel, P.H. Calculation of protein extinction coefficients from amino acid sequence data. *Anal. Biochem.* 182, 319-326 (1989).
- Giulian, D.**, Haverkamp, L.J., Yu, J., Karshin, W., Tom, D., Li, J., Kazanskaia, A., Kirkpatrick, J. & Roher, A.E. The HHQK domain of beta-amyloid provides a structural basis for the immunopathology of Alzheimer's disease. *J. Biol. Chem.* 273, 29719-29726 (1998).
- Glenner, G.G.**, Eanes, E.D. & Page, D.L. The relation of the properties of Congo red-stained amyloid fibrils to the β -conformation. *J. Histochem. Cytochem.* 20, 821-826 (1972).
- Glenner, G.G.** Amyloid deposits and amyloidosis. The β -fibrilloses (first of two parts). *N. Engl. J. Med.* 302, 1283-1292 (1980).
- Glenner, G.G.** & Wong, C.W. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem. Biophys. Res. Commun.* 120, 885-890 (1984).
- Guijarro, J.I.**, Sunde, M., Jones, J.A., Campbell, I.D. & Dobson, C.M. Amyloid fibril formation by an SH3 domain. *Proc. Natl. Acad. Sci. U.S.A.* 95, 4224-4228 (1998).
- Hamada, D.**, Segawa, S. & Goto, Y. Non-native alpha-helical intermediate in the refolding of beta-lactoglobulin, a predominantly beta-sheet protein. *Nat. Struct. Biol.* 3, 868-873 (1996).
- Haris, P.I.** & Chapman, D. The conformational analysis of peptides using Fourier transform IR spectroscopy. *Biopolymers* 37, 251-263 (1995).
- Harper, J.D.**, Wong, S.S., Lieber, C.M. & Lansbury, T.T., Jr. Observation of metastable A β amyloid protofibrils by atomic force microscopy. *Chem. Biol.* 4, 119-125 (1997).
- Hecht, M.H.** De novo design of beta-sheet proteins. *Proc. Natl. Acad. Sci. U.S.A.* 91, 8729-8730 (1994).

- Hendriks, L.**, Van Duijn, C.M., Cras, P., Cruts, M., van Hul, W., van Harskamp, F., Warren, A., McInnis, M.G., Antonarakis, S.E., Martin, J.J., Hofman, A. & van Broeckhoven Presenile dementia & cerebral haemorrhage linked to a mutation at codon 692 of the beta-amyloid precursor protein gene. *Nat. Genet. 1*, 218-221 (1992).
- Hilbich, C.**, Kisters Woike, B., Reed, J., Masters, C.L. & Beyreuther, K. Substitutions of hydrophobic amino acids reduce the amyloidogenicity of Alzheimer's disease beta A4 peptides. *J. Mol. Biol.* 228, 460-473 (1992).
- Holzhauser, M.** (Hrsg.) Methoden in der Proteinanalytik. Springer-Verlag, Heidelberg (1996).
- Hussain, I.**, Powell, D., Howlett, D.R., Tew, D.G., Week, T.D., Chapman, C., Gloger, I.S., Murphy, K.E., Southan, C.D., Ryan, D.M., Smith, T.S., Simmons, D.L., Walsh, F.S., Dingwall, C. & Christie, G. Identification of a novel aspartic protease (Asp 2) as beta-secretase. *Mol. Cell. Neurosci.* 14, 419-427 (1999).
- Ionescu-Zanetti, C.**, Khurana, R., Gillespie, J.R., Petrick, J.S., Trabachino, L.C., Minert, L.J., Carter, S.A. & Fink, A.L. Monitoring the assembly of Ig light-chain amyloid fibrils by atomic force microscopy. *Proc. Natl. Acad. Sci. U.S.A.* 96, 13175-13179 (1999).
- Jarrett, J.T.**, Berger, E.P. & Lansbury, P.T., Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry*, 32, 4693-4697 (1993).
- Jeener, J.**, Meier, B.H., Bachmann, P. & Ernst, R.R. Investigation of exchange processes by two dimensional NMR spectroscopy. *J. Chem. Phys.* 71, 4546-4553 (1979).
- Jimenez, J.L.**, Guijarro, J.I., Orlova, E., Zurdo, J., Dobson, C.M., Sunde, M. & Saibil, H.R. Cryo-electron microscopy structure of an SH3 amyloid fibril and model of the molecular packing. *EMBO J.* 18, 815-821 (1999).
- Johnson, B.A.** & Blevins, R.A. A computer program for the visualization and analysis of NMR data. *J. Biomol. NMR* 4, 603-614 (1994).
- Kaiser, E.T.** *Anal. Biochem.* 34, 595 (1970)
- Kaiser, E.T.** & Kezdy, F.J. Amphiphilic secondary structure: design of peptide hormones. *Science* 223, 249-255 (1984).
- Kaiser, E.T.** & Kezdy, F.J. Secondary structures of proteins and peptides in amphiphilic environments. *Proc. Natl. Acad. Sci. U.S.A.* 80, 1137-1143 (1983).
- Kang, J.**, Lemaire, H.G., Unterbeck, A, Salbaum, J.H., Masters, C.L., Grzeschik, K.H., Mulhaupt, G., Beyreuther, K. and Muller-Hill, B. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 325, 733-736 (1987).
- Kelly, J.W.** Alternative conformations of amyloidogenic proteins govern their behavior. *Curr. Opin. Struct. Biol.* 6, 11-17 (1996).

- Kelly, J.W.** The alternative conformations of amyloidogenic proteins and their multi-step assembly. *Curr. Opin. Struct. Biol.* 8, 101-108 (1998).
- Kelly, M.M.**, Pysh, E.S., Bonora, G.M. & Toniolo, C. *J. Am. Chem. Soc.* 99, 3264-3266 (1977).
- Kim, C.A.** & Berg, J.M. Thermodynamic beta-sheet propensities measured using a zinc-finger host peptide. *Nature* 362, 267-270 (1993).
- Kirschner, D.A.**, Abraham, C. & Selkoe, D.J. X-ray diffraction from intraneuronal paired helical filaments and extraneuronal amyloid fibers in Alzheimer disease indicates cross-beta conformation. *Proc. Natl. Acad. Sci. U.S.A.* 83, 503-507 (1986).
- Klunk, W.E.**, Pettegrew, J.W. & Abraham, D.J. Quantitative evaluation of congo red binding to amyloid-like proteins with a beta-pleated sheet conformation. *J. Histochem. Cytochem.* 37, 1273-1281 (1989).
- Konno, T.**, Murata, K. & Nagayama, K. Amyloid-like aggregates of a plant protein: a case of a sweet-tasting protein, monellin. *FEBS Lett.* 454, 122-126 (1999).
- Koradi, R.**, Billeter, M. & Wüthrich K. MOLMOL: a program for display and analysis of macromolecular structures. *J. Mol. Graph* 14, 51-55 (1996).
- Kortemme, T.**, Ramirez-Alvarado, M. & Serrano, L. Design of a 20-amino acid, three-stranded beta-sheet protein. *Science*, 281, 253-256 (1998).
- Kraus, M.**, Janek, K., Bienert, M. & Krause, E. Characterization of intermolecular β -sheet peptides by mass spectrometry and hydrogen isotope exchange. *Rapid Commun. Mass Spectrom.* 14, 1-11 (2000).
- Krause, E.**, Beyermann, M., Dathe, M., Rothmund, S. & Bienert, M. Location of an amphipathic alpha-helix in peptides using reversed-phase HPLC retention behavior of D-amino acid analogs. *Anal. Chem.* 67, 252-258 (1995).
- Krause, E.**, Beyermann, M., Fabian, H., Dathe, M., Rothmund, S. & Bienert, M. Conformation of a water-soluble beta-sheet model peptide. A circular dichroism and Fourier-transform infrared spectroscopic study of double D-amino acid replacements. *Int. J. Pept. Protein Res.* 48, 559-568 (1996).
- Krause, E.**, Rothmund, S., Beyermann, M. & Bienert, M. Monitoring of α -helical secondary structures in peptides by reversed-phase HPLC of replacement sets. *Anal. Chim. Acta* 352, 365-374 (1997).
- Krause, E.**, Bienert, M., Schmieder, P. & Wenschuh, H. The helix-destabilizing propensity scale of D-amino acids: the influence of side chain steric effects. *J. Am. Chem. Soc.* in press (2000).
- Krimm, S.** & Bandekar, J. Vibrational spectroscopy and conformation of peptides, polypeptides, and proteins. *Adv. Protein Chem.* 38, 181-364 (1986).

- Kuwajima, K.**, Yamaya, H. & Sugai, S. The burst-phase intermediate in the refolding of β -Lactoglobulin studied by stopped-flow circular dichroism and absorption spectroscopy. *J. Mol. Biol.* 264, 806-822 (1996).
- Kwok, S.C.**, Tripet, B., Man, J.H., Chana, M.S., Lavigne, P., Mant, C.T. & Hodges, R.S. Structural cassette mutagenesis in a de novo designed protein: proof of a novel concept for examining protein folding and stability. *Biopolymers* 47, 101-123 (1998).
- Lammich, S.**, Kojro, E., Postina, R., Gilbert, S., Pfeiffer, R., Jasionowski, M., Haass, C. & Fahrenholz, F. Constitutive and regulated alpha-secretase cleavage of Alzheimer's amyloid precursor protein by a disintegrin metalloprotease. *Proc. Natl. Acad. Sci. U.S.A.* 96, 3922-3927 (1999).
- Lansbury, P.T., Jr.** In pursuit of the molecular structure of amyloid plaque: new technology provides unexpected and critical information. *Biochemistry* 31, 6865-6870 (1992).
- Lansbury, P.T., Jr.** A reductionist view of Alzheimer's disease. *Acc.Chem. Res.* 29, 317-321 (1996).
- Lansbury, P.T., Jr.** Evolution of amyloid: what normal protein folding may tell us about fibrillogenesis and disease. *Proc. Natl. Acad. Sci. USA* 96, 3342-3344 (1999).
- Lazo, N.D.** & Downing, D.T. Beta-helical fibrils from a model peptide. *Biochem. Biophys. Res. Commun.* 235, 675-679 (1997).
- Lee, J.P.**, Stimson, E.R., Ghilardi, J.R., Mantyh, P.W., Lu, Y.A., Felix, A.M., Llanos, W., Behbin, A., Cummings, M., Van Crielinge, M., Timms, W. & Maggio, J.E. 1H NMR of A beta amyloid peptide congeners in water solution. Conformational changes correlate with plaque competence. *Biochemistry* 34, 5191-5200 (1995).
- Lehrman, S.R.**, Tuls, J.L. & Lund, M. Peptide alpha-helicity in aqueous trifluoroethanol: correlations with predicted alpha-helicity and the secondary structure of the corresponding regions of bovine growth hormone. *Biochemistry* 29, 5590-5596 (1990).
- LeVine, H., III** Thioflavine T interaction with synthetic Alzheimer's disease beta-amyloid peptides: detection of amyloid aggregation in solution. *Protein Sci.* 2, 404-410 (1993).
- Levy, E.**, Carman, M.D., Fernandez-Madrid, I.J., Power, M.D., Lieberburg, I., van Duinen, S.G., Bots, G.T., Luyendijk, W. & Frangione, B. Mutation of the Alzheimer's disease amyloid gene in hereditary cerebral hemorrhage, Dutch type. *Science* 248, 1124-1126 (1990).
- Lim, A.**, Saderholm, M.J., Makhov, A.M., Kroll, M., Yan, Y.B., Perera, L., Griffith, J.D. & Erickson, B.W. Engineering of betabellin-15D: A 64 residue beta sheet protein that forms long narrow multimeric fibrils. *Protein Sci.* 7, 1545-1554 (1998).
- Liu, Z.P.**, Rizo, J. & Gierasch, L.M. Equilibrium folding studies of cellular retinoic acid binding protein, a predominantly beta-sheet protein. *Biochemistry* 33, 134-142 (1994).

- Lomakin, A.**, Chung, D.S., Benedek, G.B., Kirschner, D.A. & Teplow, D.B. On the nucleation and growth of amyloid β -fibrils: detection of nuclei and quantitation of rate constants. *Proc. Nat. Acad. Sci. USA* 93, 1125-1129 (1996).
- Ma, J.**, Yee, A., Brewer, H.B., Jr., Das, S. & Potter, H. Amyloid-associated proteins alpha 1-antichymotrypsin and apolipoprotein E promote assembly of Alzheimer beta-protein into filaments. *Nature* 372, 92-94 (1994).
- Matsubara, E.**, Frangione, B. & Ghiso, J. Characterization of apolipoprotein J-Alzheimer's A beta interaction. *J. Biol. Chem.* 270, 7563-7567 (1995).
- Mayo, K.H.**, Ilyina, E. & Park, H. A recipe for designing water-soluble, β -sheet-forming peptide. *Protein Science* 5, 1301-1315 (1996).
- Merrifield, R.B.** The synthesis of a tetrapeptide. *J. Am. Chem. Soc.* 85, 2149-? (1963).
- Merutka, G.** & Stellwagen, E. Analysis of peptides for helical prediction. *Biochemistry* 28, 352-357 (1989).
- Merutka, G.**, Lipton, W., Shalongo, W., Park, S.H. & Stellwagen, E. Effect of central-residue replacements on the helical stability of a monomeric peptide. *Biochemistry* 29, 7511-7515 (1990).
- Mihara, H.** & Takahashi, Y. Engineering peptides and proteins that undergo alpha-to-beta transitions. *Curr. Opin. Struct. Biol.* 7, 501-508 (1997).
- Mihara, H.**, Takahashi, Y. & Ueno, A. Design of peptides undergoing self-catalytic alpha-to-beta transition and amyloidogenesis. *Biopolymers*, 47, 83-92 (1998).
- Milton, S.C.**, Milton, de R.L.C., Kates, S.A., Glabe, C. Improved synthesis and purification of Alzheimer's A β 1-42 and analogs. *Lett. Pept. Sci.* 6, 151-156 (1999).
- Minor, D.L., Jr.**, and Kim, P.S. Measurement of the beta-sheet-forming propensities of amino acids. *Nature* 367, 660-663 (1994a).
- Minor, D.L., Jr.**, and Kim, P.S. Context is a major determinant of beta-sheet propensity. *Nature* 371, 264-267 (1994b).
- Minor, D.L., Jr.**, and Kim, P.S. Context-dependent secondary structure formation of a designed protein sequence. *Nature* 380, 730-734 (1996).
- Mutter, M.** Die Konstruktion von neuen Proteinen und Enzymen-eine Zukunftsperspektive? *Angew. Chem.* 97, 639-654 (1985).
- Naiki, H.**, Higuchi, K., Hosokawa, M. & Takeda, T. Fluorometric determination of amyloid fibrils in vitro using the fluorescent dye, thioflavin T1. *Anal. Biochem.* 177, 244-249 (1989).
- Narayanan, U.**, Keiderling, T.A., Bonora, G.M. & Toniolo, C. *J. Am. Chem. Soc.* 108, 2431-2437 (1986).
- Nelson, J.W.**, & Kallenbach, N.R. Persistence of the alpha-helix stop signal in the S-peptide in trifluoroethanol solutions. *Biochemistry* 28, 5256-5261 (1989).

- Nelson, J.W.** & Kallenbach, N.R. Stabilization of the ribonuclease S-peptide alpha-helix by trifluoroethanol. *Proteins: Struct. Funct. Genet.* 1, 211-217 (1986).
- Nesloney, C.L.** & Kelly, J.W. Progress towards understanding beta-sheet structure. *Bioorg. Med. Chem.* 4, 739-766 (1996).
- O'Neil, K.T.** & DeGrado, W.F. A thermodynamic scale for the helix-forming tendencies of the commonly occurring amino acids. *Science* 250, 646-651 (1990).
- Oehlke, J.,** Krause, E., Wiesner, B., Beyermann, M. & Bienert, M. Extensive cellular uptake into endothelial cells of an amphipathic beta-sheet forming peptide. *FEBS Lett.* 415, 196-199 (1997).
- Osterman, D.G.,** and Kaiser, E.T. Design and characterization of peptides with amphiphilic beta-strand structures. *J. Cell. Biochem.* 29, 57-72 (1985).
- Pauling, L.** & Corey, R.B. Configurations of polypeptide chains with favored orientations around single bonds: two new pleated sheets. *Proc. Natl. Acad. Sci. U.S.A.* 37, 729-740 (1951).
- Perutz, M.F.,** Johnson, T., Suzuki, M. & Finch, J.T. Glutamine repeats as polar zippers: their possible role in inherited neurodegenerative diseases. *Proc. Natl. Acad. Sci. U.S.A.* 91, 5355-5358 (1994).
- Perutz, M.F.** Glutamine repeats and inherited neurodegenerative diseases: molecular aspects. *Curr. Opin. Struct. Biol.* 6, 848-858 (1996).
- Perutz, M.F.** Amyloid fibrils. Mutations make enzyme polymerize. *Nature*, 385, 773-775 (1997).
- Piantini, U.,** Sorensen, O.W. & Ernst, R.R. Multiple quantum filters elucidating NMR coupling networks. *J. Am. Chem. Soc.* 104, 6800-6801 (1982).
- Provencher, S.W.** CONTIN: A general purpose constrained regularization program for inverting noisy linear algebraic and integral equations. *Comput. Phys. Commun.* 27, 229-242 (1982).
- Prusiner, S.B.** Transgenetics of prion disease. *Curr. Top. Microbiol. Immunol.* 206, 275-304 (1996).
- Puchtler, H.,** Sweat, F. & Levine, M. On the binding of congo red by amyloid. *J. Histochem.* 10, 355-364 (1962).
- Puglisi, J.D.,** Chen, L., Blanchard, S., and Frankel, A.D. Solution structure of a bovine immunodeficiency virus Tat-TAR peptide-RNA complex. *Science* 270, 1200-1203 (1995).
- Quinn, T.P.,** Tweedy, N.B., Williams, R.W., Richardson, J.S. & Richardson, D.C. Betadoublet: de novo design, synthesis and characterization of a betasandwich protein. *Proc. Natl. Acad. Sci. U.S.A.* 91, 8747-8757 (1994).
- Rajan, R.** & Balaram, P. A model for the interaction of trifluoroethanol with peptides and proteins. *Int. J. Peptide Protein Res.* 48, 328-336 (1996).
- Richardson, J.S.** β -Sheet topology and the relatedness of proteins. *Nature* 268, 495-500, (1977).
- Rochet, J.-C.** & Lansbury, P.T., Jr. Amyloid fibrillogenesis: Themes and variations. *Curr. Opin. Struct. Biol.* 10, 60-68 (2000).

- Roher, A.E.**, Lowenson, J.D., Clarke, S., Wolkow, C., Wang, R., Cotter, R.J., Reardon, I.M., Zurcher Neely, H.A., Heinrichson, R.L., Ball, M.J. & *et al.* Structural alterations in the peptide backbone of beta-amyloid core protein may account for its deposition and stability in Alzheimer's disease. *J. Biol. Chem.* 268, 3072-3083 (1993).
- Rohl, C.A.**, Chakrabarty, A. & Baldwin, R.L. Helix propagation and N-cap propensities of the amino acids measured in alanine-based peptides in 40 volume percent trifluoroethanol. *Protein Sci.* 5, 2623-2637 (1996).
- Rothmund, S.**, Beyermann, M., Krause, E., Krause, G., Bienert, M., Hodges, R.S., Sykes, B.D. & Sonnichsen, F.D. Structure effects of double D-amino acid replacements: a nuclear magnetic resonance and circular dichroism study using amphipathic model helices. *Biochemistry*, 34, 12954-12962 (1995).
- Salomon, A.R.**, Marcinowski, K.J., Friedland, R.P. & Zagorski, M.G. Nicotine inhibits amyloid formation by the beta-peptide. *Biochemistry* 35, 13568-13578 (1996).
- Scherzinger, E.**, Lurz, R., Turmaine, M., Mangiarini, L., Hollenbach, B., Hasenbank, R., Bates, G.P., Davies, S.W., Lehrach, H. & Wanker, E.E. Huntingtin-encoded polyglutamine expansions form amyloid-like protein aggregates in vitro and in vivo. *Cell* 90, 549-558 (1997).
- Schwarzman, A.L.**, Gregori, L., Vitek, M.P., Lyubski, S., Strittmatter, W.J., Enghilde, J.J., Bhasin, R., Silverman, J., Weisgraber, K.H., Coyle, P.K., Zargorski, M.G., Talafous, J., Eisenberg, M., Saunders, A.M., Roses, A.D. & Goldgaber, D. Transthyretin sequesters amyloid β protein and prevents amyloid formation. *Proc. Natl. Acad. Sci. USA* 91, 8368-8372 (1994).
- Segawa, S.I.**, Fukuno, T., Fujiwara, K. & Noda, Y. Local Structures in unfolded lysozym and correlation with secondary structure in the native conformation. *Biopolymers* 31, 497-509 (1991).
- Selkoe, D.J.** Normal and abnormal biology of the β -amyloid precursor protein. *Annu. Rev. Neurosci.* 17, 489-517 (1994).
- Selkoe, D.J.** The cell biology of beta-amyloid precursor protein and presenilin in Alzheimer's disease. *Trends Cell Biol.* 8, 447-453 (1998).
- Selkoe, D.J.** Translating cell biology into therapeutic advances in Alzheimer's disease. *Nature* 399, A23-31 (1999).
- Serrano, L.**, Sancho, J., Hirshberg, M. & Fersht, A.R. Alpha-helix stability in proteins. I. Empirical correlations concerning substitution of side-chains at the N and C-caps and the replacement of alanine by glycine or serine at solvent-exposed surfaces. *J. Mol. Biol.* 227, 544-559 (1992).
- Seubert, P.**, Vigo Pelfrey, C., Esch, F., Lee, M., Dovey, H., Davis, D., Sinha, S., Schlossmacher, M., Whaley, J., Swindlehurst, C., McCormack, R., Wolfert, R., Selkoe, D., Lieberburg, I. & Schenk, D. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* 359, 325-327 (1992).

- Shao, H.**, Jao, S., Ma, K. & Zagorski, M.G. Solution structures of micelle-bound amyloid beta-(1-40) and beta-(1-42) peptides of Alzheimer's disease. *J. Mol. Biol.* 285, 755-773 (1999).
- Sinha, S.**, Anderson, J.P., Barbour, R., Basi, G.S., Caccavello, R., Davis, D., Doan, M., Dovey, H.F., Frigon, N., Hong, J., Jacobson- Croak, K., Jewett, N., Keim, P., Knops, J., Lieberburg, I., Power, M., Tan, H., Tatsuno, G., Tung, J., Schenk, D., Seubert, P., Suomensaaari, S.M., Wang, S.W., Walker, D., Zhao, J., McConlogue, L. & John, V. Purification and cloning of amyloid precursor protein beta-secretase from human brain. *Nature* 402, 537-540 (1999).
- Shapira, R.**, Austin, G.E. & Mirra, S.S. Neuritic plaque amyloid in Alzheimer's disease is highly racemized. *J. Neurochem.* 50, 69-74 (1988).
- Shoji, M.**, Golde, T.E., Ghiso, J., Cheung, T.T., Estus, S., Shaffer, L.M., Cai, X.D., McKay, D.M., Tintner, R., Frangione, B. & Younkin, S.G. Production of the Alzheimer amyloid beta protein by normal proteolytic processing. *Science*, 258, 126-129 (1992).
- Sipe, J.D.** Amyloidosis. *Ann. Rev. Biochem.* 61, 947-75 (1992).
- Sipe, J.D.** Amyloidosis. *Crit. Rev. Clin. Lab. Sci.* 31, 325-35 (1994).
- Smith, C.K.** & Regan, L. Construction and design of beta-sheets. *Acc. Chem. Res.* 30, 153-161 (1997).
- Smith, C.K.**, Withka, J.M. & Regan, L. A thermodynamic scale for the beta-sheet forming tendencies of the amino acids. *Biochemistry*, 33, 5510-5517 (1994).
- Somers, W.S.** & Phillips, S.E. Crystal structure of the met repressor-operator complex at 2.8 Å resolution reveals DNA recognition by beta-strands. *Nature* 359, 387-393 (1992).
- Sönnichsen, F.D.**, Van Eyk, J.E., Hodges, R.S. & Sykes, B.D. Effect of trifluoroethanol on protein secondary structure: an NMR and CD study using a synthetic actin peptide. *Biochemistry* 31, 8790-8798 (1992).
- Soto, C.**, Castano, E.M., Frangione, B. & Inestrosa, N.C. The alpha-helical to beta-strand transition in the amino-terminal fragment of the amyloid beta-peptide modulates amyloid formation. *J. Biol. Chem.* 270, 3063-3067 (1995).
- Soto, C.**, Golabek, A., Wisniewski, T. & Castano, E.M. Alzheimer's beta-amyloid peptide is conformationally modified by apolipoprotein E in vitro. *Neuroreport* 7, 721-725 (1996).
- Soto, C.**, Sigurdsson, E.M., Morelli, L., Kumar, R.A., Castano, E.M. & Frangione, B. Beta-sheet breaker peptides inhibit fibrillogenesis in a rat brain model of amyloidosis: Implications for Alzheimer's therapy. *Nature Med.* 4, 822-826 (1998).
- St George-Hyslop, P.H.** & Westaway, D.A. Antibody clears senile plaques. *Nature* 400, 116-117 (1999).

- Sticht, H.**, Bayer, P., Willbold, D., Dames, S., Hilbich, C., Beyreuther, K., Frank, R.W. & Rosch, P. Structure of amyloid A4-(1-40)-peptide of Alzheimer's disease. *Eur. J. Biochem.* 233, 293-298 (1995).
- Streyer, L.** (Hrsg.) Biochemie. Spektrum Akad. Verlag, Heidelberg, 4.Aufl. (1996)
- Sunde, M.** & Blake, C.C. From the globular to the fibrous state: protein structure and structural conversion in amyloid formation. *Q. Rev. Biophys.* 31, 1-39 (1998).
- Sunde, M.**, Serpell, L.C., Bartlam, M., Fraser, P.E., Pepys, M.B. & Blake, C.C. Common core structure of amyloid fibrils by synchrotron X-ray diffraction. *J. Mol. Biol.* 273, 729-739 (1997).
- Symmons, M.F.**, Buchanan, S.G., Clarke, D.T., Jones, G. & Gay, N.J. X-ray diffraction and far-UV CD studies of filaments formed by a leucine-rich repeat peptide: structural similarity to the amyloid fibrils of prions and Alzheimer's disease beta-protein. *FEBS Lett.* 412, 397-403 (1997).
- Teplow, D.B.** Structural and kinetic features of amyloid β -protein fibrillogenesis. *Amyloid: Int. J. Exp. Clin. Invest.* 5, 112-142 (1998).
- Terzi, E.**, Hölzemann, G. & Seelig, J. Reversible random coil-beta-sheet transition of the Alzheimer β -amyloid fragment (25-35). *Biochemistry*, 33, 1345-1350 (1994).
- Tjernberg, L.O.**, Naslund, J., Lindqvist, F., Johansson, J., Karlstrom, A.R., Thyberg, J., Terenius, L. & Nordstedt, C. Arrest of β -amyloid fibril formation by a pentapeptide ligand. *J. Biol. Chem.* 271, 8545-8548 (1996).
- Tomiyama, T.**, Asano, S., Furiya, Y., Shirasawa, T., Endo, N. & Mori, H. Racemization of Asp23 residue affects the aggregation properties of Alzheimer amyloid beta protein analogues. *J. Biol. Chem.* 269, 10205-10208 (1994).
- Turnell, W.G.** & Finch, J.T. Binding of the dye congo red to the amyloid protein pig insulin reveals a novel homology amongst amyloid-forming peptide sequences. *J. Mol. Biol.* 227, 1205-1223 (1992).
- Vassar, R.**, Bennett, B.D., Babu Khan, S., Kahn, S., Mendiaz, E.A., Denis, P., Teplow, D.B., Ross, S., Amarante, P., Loeloff, R., Luo, Y., Fisher, S., Fuller, L., Edenson, S., Lile, J., Jarosinski, M.A., Biere, A.L., Curran, E., Burgess, T., Louis, J.C., Collins, F., Treanor, J., Rogers, G. & Citron, M. Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* 286, 735-741 (1999).
- Virchow, R.** Bau und Zusammensetzung der Corpora Amylacea des Menschen. *Verh. Phys. Med. Ges. Würzburg* 2, 51-54 (1851).
- Voet, D.**, Voet, J.G. & Pratt, C. (eds.) Fundamentals of Biochemistry. John Wiley & Sons, Inc. New York (1999).
- Walsh, D.M.**, Lomakin, A., Benedek, G.B., Condron, M.M. & Teplow, D.B. Amyloid β -protein fibrillogenesis: detection of a protofibrillar intermediate. *J. Biol. Chem.* 272, 22364-22374 (1997).

- Waterhous, D.V.** & Johnson, W.C., Jr. Importance of environment in determining secondary structure in proteins. *Biochemistry* 33, 2121-2128 (1994).
- Weaver, L.**, Stagsted, J., Behnke, O., Matthews, B.W. & Olsson, L. Beta-sheet models for the ordered filamentous structure formed by a peptide that enhances the action of insulin. *J. Struct. Biol.* 117, 165-172 (1996).
- Wenschuh, H.**, Halada, P., Lamer, S., Jungblut, P. & Krause, E. The ease of peptide detection by matrix-assisted laser desorption/ionization mass spectrometry: the effect of secondary structure on signal intensity. *Rapid Commun. Mass. Spectrom.* 12, 115-119 (1998).
- West, M.W.**, Wang, W., Patterson, J., Mancias, J.D., Beasley, J.R. & Hecht, M.H. De novo amyloid proteins from designed combinatorial libraries. *Proc. Natl. Acad. Sci. U.S.A.* 96, 11211-11216 (1999).
- Westermarck, P.**, Araki, S., Benson, M.D., Cohen, A.S., Frangione, B., Masters, C.L., Saraiva, M.J., Sipe, J.D., Husby, G., Kyle, R.A. & Selkoe, D. Nomenclature of amyloid fibril proteins. Report from the meeting of the International Nomenclature Committee on Amyloidosis, August 8-9, 1998 Part 1. *Amyloid: Int. J. Clin. Invest.* 6, 63-66 (1999).
- Wilson, D.M.** & Binder, L.I. Polymerization of microtubule-associated protein tau under near-physiological conditions. *J. Biol. Chem.* 270, 2406-24341 (1995).
- Wetzel, R.** For protein misassembly, it's the "I" decade. *Cell* 86, 699-702 (1996).
- Wishard, D.S.**, Bigan, C.G., Holm, A., Hodges, R.S. & Syke, B.D. 1H, 13C and 15N random coil NMR chemical shifts of the common amino acids. Investigation of nearest-neighbour effects. *J. Biomol. NMR* 5, 67-81 (1995).
- Wolfe, M.S.**, Xia, W.M., Ostaszewski, B.L., Diehl, T.S., Kimberly, W.T. & Selkoe, D.J. Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and gamma-secretase activity. *Nature* 398, 513-517 (1999).
- Wood, S.J.**, Wetzel, R., Martin, J.D. & Hurler, M.R. Prolines and amyloidogenicity in fragments of the Alzheimer's peptide β /A4. *Biochemistry* 34, 724-730 (1995).
- Wouters, M.A.**, and Curmi, P.M. An analysis of side chain interactions and pair correlations within antiparallel beta-sheets: the differences between backbone hydrogen-bonded and non-hydrogen-bonded residue pairs. *Proteins* 22, 119-131 (1995).
- Wüthrich, K.** In: *NMR of proteins and nucleic acids* (John Wiley & Sons, New York; 1986).
- Xiong, H.**, Buckwalter, B.L., Shieh, H.M. & Hecht, M.H. Periodicity of polar and nonpolar amino acids is the major determinant of secondary structure in self-assembling oligomeric peptides. *Proc. Natl. Acad. Sci. U.S.A.* 92, 6349-6353 (1995).

- Yamada, N.**, Ariga, K., Naito, M., Matsubara, K. & Koyama, E. Regulation of beta-sheet structures within amyloid-like beta-sheet assemblage from tripeptide derivatives. *J. Am. Chem. Soc.* *120*, 12192-12199 (1998).
- Yan, R.Q.**, Bienkowski, M.J., Shuck, M.E., Miao, H.Y., Tory, M.C., Pauley, A.M., Brashler, J.R., Stratman, N.C., Mathews, W.R., Buhl, A.E., Carter, D.B., Tomasselli, A.G., Parodi, L.A., Heinrikson, R.L. & Gurney, M.E. Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity. *Nature* *402*, 533-537 (1999).
- Yan, S.M.** & Clarke, N.D. A hybrid sequence approach to praelus challenge. *Proteins* *30*, 136-143 (1998).
- Yan, Y.**, & Erickson, B.W. Engineering of betabellin 14D: disulfide-induced folding of a beta-sheet protein. *Protein Sci.* *3*, 1069-1073 (1994).
- Yankner, B.A.** Mechanisms of neuronal degeneration in Alzheimer's disease. *Neuron* *16*, 921-932 (1996).
- Zagorski, M.G.** & Barrow, C.J. NMR studies of amyloid beta-peptides: proton assignments, secondary structure, and mechanism of an alpha-helix→beta-sheet conversion for a homologous, 28-residue, N-terminal fragment. *Biochemistry* *31*, 5621-5631 (1992).
- Zhang, S.**, Holmes, T., Lockshin, C. & Rich, A. Spontaneous assembly of a self-complementary oligopeptide to form a stable macroscopic membrane. *Proc. Natl. Acad. Sci. U.S.A.* *90*, 3334-3338 (1993).
- Zhong, L.** & Johnson, W.C., Jr. Environment affects amino acid preference for secondary structure. *Proc. Natl. Acad. Sci. U.S.A.* *89*, 4462-4465 (1992).
- Zhou, N.E.**, Mant, C.T. & Hodges, R.S. (1990) Effect of preferred binding domains on peptide retention behavior in reversed-phase chromatography: amphipathic alpha-helices. *Peptide Res.* *3*, 8-20.