

## 7 Literatur

**Abou-Samra, A. B., Juppner, H., Kong, X. F., Schipani, E., Iida-Klein, A., Karga, H., Urena, P., Gardella, T. F., Potts, J. T., Jr., Kronenberg, H. M. et al.** (1994). Structure, function, and expression of the receptor for parathyroid hormone and parathyroid hormone-related peptide. *Adv Nephrol Necker Hosp* **23**, 247-64.

**Ahn, J., Ludecke, H. J., Lindow, S., Horton, W. A., Lee, B., Wagner, M. J., Horsthemke, B. and Wells, D. E.** (1995). Cloning of the putative tumour suppressor gene for hereditary multiple exostoses (EXT1). *Nat Genet* **11**, 137-43.

**Alvarez, J., Sohn, P., Zeng, X., Doetschman, T., Robbins, D. J. and Serra, R.** (2002). TGFbeta2 mediates the effects of hedgehog on hypertrophic differentiation and PTHrP expression. *Development* **129**, 1913-24.

**Amizuka, N., Warshawsky, H., Henderson, J. E., Goltzman, D. and Karaplis, A. C.** (1994). Parathyroid hormone-related peptide-depleted mice show abnormal epiphyseal cartilage development and altered endochondral bone formation. *J Cell Biol* **126**, 1611-1623.

**Ausubel, Fm, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K.** (1996). Current protocols in Molecular Biology.

**Aza-Blanc, P. and Kornberg, T. B.** (1999). Ci: a complex transducer of the hedgehog signal. *Trends Genet* **15**, 458-62.

**Baeg, G. H., Lin, X., Khare, N., Baumgartner, S. and Perrimon, N.** (2001). Heparan sulfate proteoglycans are critical for the organization of the extracellular distribution of Wingless. *Development* **128**, 87-94.

**Bai, C. B., Auerbach, W., Lee, J. S., Stephen, D. and Joyner, A. L.** (2002). Gli2, but not Gli1, is required for initial Shh signaling and ectopic activation of the Shh pathway. *Development* **129**, 4753-61.

**Bai, C. B. and Joyner, A. L.** (2001). Gli1 can rescue the in vivo function of Gli2. *Development* **128**, 5161-72.

**Bai, C. B., Stephen, D. and Joyner, A. L.** (2004). All mouse ventral spinal cord patterning by hedgehog is Gli dependent and involves an activator function of Gli3. *Dev Cell* **6**, 103-15.

**Becker, S., Wang, Z. J., Massey, H., Arauz, A., Labosky, P., Hammerschmidt, M., St-Jacques, B., Bumcrot, D., McMahon, A. and Grabel, L.** (1997). A role for Indian hedgehog in extraembryonic endoderm differentiation in F9 cells and the early mouse embryo. *Dev Biol* **187**, 298-310.

**Bellaiche, Y., The, I. and Perrimon, N.** (1998). Tout-velu is a Drosophila homologue of the putative tumour suppressor EXT-1 and is needed for Hh diffusion. *Nature* **394**, 85-8.

**Bitgood, M. J. and McMahon, A. P.** (1995). Hedgehog and Bmp genes are coexpressed at many diverse sites of cell-cell interaction in the mouse embryo. *Dev Biol* **172**, 126-138.

- Briscoe, J., Chen, Y., Jessell, T. M. and Struhl, G.** (2001). A hedgehog-insensitive form of patched provides evidence for direct long-range morphogen activity of sonic hedgehog in the neural tube. *Mol Cell* **7**, 1279-91.
- Bumcrot, D. A., Takada, R. and McMahon, A. P.** (1995). Proteolytic processing yields two secreted forms of sonic hedgehog. *Mol Cell Biol* **15**, 2294-303.
- Burke, R., Nellen, D., Bellotto, M., Hafen, E., Senti, K. A., Dickson, B. J. and Basler, K.** (1999). Dispatched, a novel sterol-sensing domain protein dedicated to the release of cholesterol-modified hedgehog from signaling cells. *Cell* **99**, 803-15.
- Cardin, A. D. and Weintraub, H. J.** (1989). Molecular modeling of protein-glycosaminoglycan interactions. *Arteriosclerosis* **9**, 21-32.
- Celeste, A. J., Rosen, V., Buecker, J. L., Kriz, R., Wang, E. A. and Wozney, J. M.** (1986). Isolation of the human gene for bone gla protein utilizing mouse and rat cDNA clones. *Embo J* **5**, 1885-90.
- Chamoun, Z., Mann, R. K., Nellen, D., von Kessler, D. P., Bellotto, M., Beachy, P. A. and Basler, K.** (2001). Skinny hedgehog, an acyltransferase required for palmitoylation and activity of the hedgehog signal. *Science* **293**, 2080-4.
- Chiang, C., Litingtung, Y., Lee, E., Young, K. E., Corden, J. L., Westphal, H. and Beachy, P. A.** (1996). Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**, 407-13.
- Cohen, M. M., Jr.** (2003). Craniofacial anomalies: Clinical and molecular perspectives. *Ann Acad Med Singapore* **32**, 244-51.
- Cook, A., Raskind, W., Blanton, S. H., Pauli, R. M., Gregg, R. G., Francomano, C. A., Puffenberger, E., Conrad, E. U., Schmale, G., Schellenberg, G. et al.** (1993). Genetic heterogeneity in families with hereditary multiple exostoses. *Am J Hum Genet* **53**, 71-9.
- Cormack, D.** (1987). Ham's histology. *Philadelphia: Pennsylvania*.
- Dai, P., Akimaru, H., Tanaka, Y., Maekawa, T., Nakafuku, M. and Ishii, S.** (1999). Sonic Hedgehog-induced activation of the Gli1 promoter is mediated by GLI3. *J Biol Chem* **274**, 8143-52.
- David, G., Bai, X. M., Van der Schueren, B., Cassiman, J. J. and Van den Berghe, H.** (1992). Developmental changes in heparan sulfate expression: in situ detection with mAbs. *J Cell Biol* **119**, 961-75.
- Ding, Q., Motoyama, J., Gasca, S., Mo, R., Sasaki, H., Rossant, J. and Hui, C. C.** (1998). Diminished Sonic hedgehog signaling and lack of floor plate differentiation in Gli2 mutant mice. *Development* **125**, 2533-43.
- Dominguez, M., Brunner, M., Hafen, E. and Basler, K.** (1996). Sending and receiving the hedgehog signal: control by the Drosophila Gli protein Cubitus interruptus. *Science* **272**, 1621-5.
- Entchev, E. V. and Gonzalez-Gaitan, M. A.** (2002). Morphogen gradient formation and vesicular trafficking. *Traffic* **3**, 98-109.
- Erlebacher, A., Filvaroff, E. H., Gitelman, S. E. and Derynck, R.** (1995). Toward a molecular understanding of skeletal development. *Cell* **80**, 371-8.

- Esko, J. D. and Lindahl, U.** (2001). Molecular diversity of heparan sulfate. *J Clin Invest* **108**, 169-73.
- Esko, J. D. and Selleck, S. B.** (2002). Order out of chaos: assembly of ligand binding sites in heparan sulfate. *Annu Rev Biochem* **71**, 435-71.
- Fujise, M., Takeo, S., Kamimura, K., Matsuo, T., Aigaki, T., Izumi, S. and Nakato, H.** (2003). Dally regulates Dpp morphogen gradient formation in the Drosophila wing. *Development* **130**, 1515-22.
- Gallet, A., Rodriguez, R., Ruel, L. and Therond, P. P.** (2003). Cholesterol modification of hedgehog is required for trafficking and movement, revealing an asymmetric cellular response to hedgehog. *Dev Cell* **4**, 191-204.
- Goodrich, L. V., Johnson, R. L., Milenkovic, L., McMahon, J. A. and Scott, M. P.** (1996). Conservation of the hedgehog/patched signaling pathway from flies to mice: induction of a mouse patched gene by Hedgehog. *Genes Dev* **10**, 301-12.
- Greco, V., Hannus, M. and Eaton, S.** (2001). Argosomes: a potential vehicle for the spread of morphogens through epithelia. *Cell* **106**, 633-45.
- Gritli-Linde, A., Lewis, P., McMahon, A. P. and Linde, A.** (2001). The whereabouts of a morphogen: direct evidence for short- and graded long-range activity of hedgehog signaling peptides. *Dev Biol* **236**, 364-86.
- Hall, C. R., Cole, W. G., Haynes, R. and Hecht, J. T.** (2002). Reevaluation of a genetic model for the development of exostosis in hereditary multiple exostosis. *Am J Med Genet* **112**, 1-5.
- Han, C., Belenkaya, T. Y., Khodoun, M., Tauchi, M. and Lin, X.** (2004). Distinct and collaborative roles of Drosophila EXT family proteins in morphogen signalling and gradient formation. *Development* **131**, 1563-75.
- Hecht, J. T., Hogue, D., Strong, L. C., Hansen, M. F., Blanton, S. H. and Wagner, M.** (1995). Hereditary multiple exostosis and chondrosarcoma: linkage to chromosome II and loss of heterozygosity for EXT-linked markers on chromosomes II and 8. *Am J Hum Genet* **56**, 1125-31.
- Hopyan, S., Gokgoz, N., Poon, R., Gensure, R. C., Yu, C., Cole, W. G., Bell, R. S., Juppner, H., Andrulis, I. L., Wunder, J. S. et al.** (2002). A mutant PTH/PTHrP type I receptor in enchondromatosis. *Nat Genet* **30**, 306-10.
- Horton, W. A.** (1993). In vitro chondrogenesis in human chondrodysplasias. *Am J Med Genet* **45**, 179-82.
- Hui, C. C. and Joyner, A. L.** (1993). A mouse model of greig cephalopolysyndactyly syndrome: the extra-toesJ mutation contains an intragenic deletion of the Gli3 gene. *Nat Genet* **3**, 241-6.
- Hui, C. C., Slusarski, D., Platt, K. A., Holmgren, R. and Joyner, A. L.** (1994). Expression of three mouse homologs of the Drosophila segment polarity gene cubitus interruptus, Gli, Gli-2, and Gli-3, in ectoderm- and mesoderm-derived tissues suggests multiple roles during postimplantation development. *Dev Biol* **162**, 402-13.

- Inatani, M., Irie, F., Plump, A. S., Tessier-Lavigne, M. and Yamaguchi, Y.** (2003). Mammalian brain morphogenesis and midline axon guidance require heparan sulfate. *Science* **302**, 1044-6.
- Incardona, J. P., Gaffield, W., Kapur, R. P. and Roelink, H.** (1998). The teratogenic Veratrum alkaloid cyclopamine inhibits sonic hedgehog signal transduction. *Development* **125**, 3553-62.
- Ingham, P. W. and McMahon, A. P.** (2001). Hedgehog signaling in animal development: paradigms and principles. *Genes Dev* **15**, 3059-87.
- Jacenko, O., LuValle, P., Solum, K. and Olsen, B. R.** (1993). A dominant negative mutation in the alpha 1 (X) collagen gene produces spondylometaphyseal defects in mice. *Prog Clin Biol Res* **383B**, 427-36.
- Jessell, T. M.** (2000). Neuronal specification in the spinal cord: inductive signals and transcriptional codes. *Nat Rev Genet* **1**, 20-9.
- Johnson, D. R.** (1967). Extra-toes: anew mutant gene causing multiple abnormalities in the mouse. *J Embryol Exp Morphol* **17**, 543-81.
- Kang, S., Graham, J. M., Jr., Olney, A. H. and Biesecker, L. G.** (1997). GLI3 frameshift mutations cause autosomal dominant Pallister-Hall syndrome. *Nat Genet* **15**, 266-8.
- Karp, S. J., Schipani, E., St-Jacques, B., Hunzelman, J., Kronenberg, H. and McMahon, A. P.** (2000). Indian hedgehog coordinates endochondral bone growth and morphogenesis via parathyroid hormone related-protein-dependent and -independent pathways. *Development* **127**, 543-8.
- Kitagawa, H., Shimakawa, H. and Sugahara, K.** (1999). The tumor suppressor EXT-like gene EXTL2 encodes an alpha1, 4-N-acetylhexosaminyltransferase that transfers N-acetylgalactosamine and N-acetylglucosamine to the common glycosaminoglycan-protein linkage region. The key enzyme for the chain initiation of heparan sulfate. *J Biol Chem* **274**, 13933-7.
- Kobayashi, S., Morimoto, K., Shimizu, T., Takahashi, M., Kurosawa, H. and Shirasawa, T.** (2000). Association of EXT1 and EXT2, hereditary multiple exostoses gene products, in Golgi apparatus. *Biochem Biophys Res Commun* **268**, 860-7.
- Kobayashi, T., Chung, U. I., Schipani, E., Starbuck, M., Karsenty, G., Katagiri, T., Goad, D. L., Lanske, B. and Kronenberg, H. M.** (2002). PTHrP and Indian hedgehog control differentiation of growth plate chondrocytes at multiple steps. *Development* **129**, 2977-86.
- Kohno, K., Martin, G. R. and Yamada, Y.** (1984). Isolation and characterization of a cDNA clone for the amino-terminal portion of the pro-alpha 1(II) chain of cartilage collagen. *J Biol Chem* **259**, 13668-73.
- Koziel, L., Kunath, M., Kelly, O. G. and Vortkamp, A.** (2004). Ext1-dependent heparan sulfate regulates the range of Ihh signaling during endochondral ossification. *Dev Cell* **6**, 801-13.
- Lander, A. D. and Selleck, S. B.** (2000). The elusive functions of proteoglycans: in vivo veritas. *J Cell Biol* **148**, 227-32.

- Lanske, B., Karaplis, A. C., Lee, K., Luz, A., Vortkamp, A., Pirro, A., Karperien, M., Defize, L. H. K., Ho, C., Mulligan, R. C. et al.** (1996). PTH/PTHrP receptor in early development and Indian hedgehog-regulated bone growth [see comments]. *Science* **273**, 663-6.
- Lawrence, P. A.** (2001). Morphogens: how big is the big picture? *Nat Cell Biol* **3**, E151-4.
- Le Merrer, M., Legeai-Mallet, L., Jeannin, P. M., Horsthemke, B., Schinzel, A., Plauchu, H., Toutain, A., Achard, F., Munnich, A. and Maroteaux, P.** (1994). A gene for hereditary multiple exostoses maps to chromosome 19p. *Hum Mol Genet* **3**, 717-22.
- Lee, J., Platt, K. A., Censullo, P. and Ruiz i Altaba, A.** (1997). Gli1 is a target of Sonic hedgehog that induces ventral neural tube development. *Development* **124**, 2537-52.
- Lee, J. J., Ekker, S. C., von Kessler, D. P., Porter, J. A., Sun, B. I. and Beachy, P. A.** (1994). Autoproteolysis in hedgehog protein biogenesis. *Science* **266**, 1528-37.
- Lee, J. J., von Kessler, D. P., Parks, S. and Beachy, P. A.** (1992). Secretion and localized transcription suggest a role in positional signaling for products of the segmentation gene hedgehog. *Cell* **71**, 33-50.
- Lee, K., Deeds, J. D. and Segre, G. V.** (1995). Expression of parathyroid hormone-related peptide and its receptor messenger ribonucleic acids during fetal development of rats. *Endocrinology* **136**, 453-463.
- Leighton, P. A., Mitchell, K. J., Goodrich, L. V., Lu, X., Pinson, K., Scherz, P., Skarnes, W. C. and Tessier-Lavigne, M.** (2001). Defining brain wiring patterns and mechanisms through gene trapping in mice. *Nature* **410**, 174-9.
- Lin, X., Wei, G., Shi, Z., Dryer, L., Esko, J. D., Wells, D. E. and Matzuk, M. M.** (2000). Disruption of gastrulation and heparan sulfate biosynthesis in EXT1-deficient mice. *Dev Biol* **224**, 299-311.
- Lind, T., Tufaro, F., McCormick, C., Lindahl, U. and Lidholt, K.** (1998). The putative tumor suppressors EXT1 and EXT2 are glycosyltransferases required for the biosynthesis of heparan sulfate. *J Biol Chem* **273**, 26265-8.
- Lindahl, U., Kusche-Gullberg, M. and Kjellen, L.** (1998). Regulated diversity of heparan sulfate. *J Biol Chem* **273**, 24979-82.
- Litingtung, Y., Dahn, R. D., Li, Y., Fallon, J. F. and Chiang, C.** (2002). Shh and Gli3 are dispensable for limb skeleton formation but regulate digit number and identity. *Nature* **418**, 979-83.
- Long, F., Chung, U. I., Ohba, S., McMahon, J., Kronenberg, H. M. and McMahon, A. P.** (2004). Ihh signaling is directly required for the osteoblast lineage in the endochondral skeleton. *Development* **131**, 1309-18. Epub 2004 Feb 18.
- Long, F., Zhang, X. M., Karp, S., Yang, Y. and McMahon, A. P.** (2001). Genetic manipulation of hedgehog signaling in the endochondral skeleton reveals a direct role in the regulation of chondrocyte proliferation. *Development* **128**, 5099-5108.
- Mann, R. K. and Beachy, P. A.** (2004). Novel lipid modifications of secreted protein signals. *Annu Rev Biochem* **73**, 891-923.

- Marigo, V., Johnson, R. L., Vortkamp, A. and Tabin, C. J.** (1996). Sonic hedgehog differentially regulates expression of GLI and GLI3 during limb development. *Dev Biol* **180**, 273-83.
- Matise, M. P., Epstein, D. J., Park, H. L., Platt, K. A. and Joyner, A. L.** (1998). Gli2 is required for induction of floor plate and adjacent cells, but not most ventral neurons in the mouse central nervous system. *Development* **125**, 2759-70.
- McCormick, C., Duncan, G., Goutsos, K. T. and Tufaro, F.** (2000). The putative tumor suppressors EXT1 and EXT2 form a stable complex that accumulates in the Golgi apparatus and catalyzes the synthesis of heparan sulfate. *Proc Natl Acad Sci U S A* **97**, 668-73.
- McCormick, C., Duncan, G. and Tufaro, F.** (1999). New perspectives on the molecular basis of hereditary bone tumours. *Mol Med Today* **5**, 481-6.
- McCormick, C., Leduc, Y., Martindale, D., Mattison, K., Esford, L. E., Dyer, A. P. and Tufaro, F.** (1998). The putative tumour suppressor EXT1 alters the expression of cell-surface heparan sulfate. *Nat Genet* **19**, 158-61.
- McMahon, A. P.** (2000). More surprises in the Hedgehog signaling pathway. *Cell* **100**, 185-8.
- Methot, N. and Basler, K.** (1999). Hedgehog controls limb development by regulating the activities of distinct transcriptional activator and repressor forms of Cubitus interruptus. *Cell* **96**, 819-31.
- Methot, N. and Basler, K.** (2000). Suppressor of fused opposes hedgehog signal transduction by impeding nuclear accumulation of the activator form of Cubitus interruptus. *Development* **127**, 4001-10.
- Minina, E., Kreschel, C., Naski, M. C., Ornitz, D. M. and Vortkamp, A.** (2002). Interaction of FGF, Ihh/Pthlh, and BMP signaling integrates chondrocyte proliferation and hypertrophic differentiation. *Dev Cell* **3**, 439-49.
- Minina, E., Wenzel, H. M., Kreschel, C., Karp, S., Gaffield, W., McMahon, A. P. and Vortkamp, A.** (2001). BMP and Ihh/PTHrP signaling interact to coordinate chondrocyte proliferation and differentiation. *Development* **128**, 4523-34.
- Mitchell, K. J., Pinson, K. I., Kelly, O. G., Brennan, J., Zupicich, J., Scherz, P., Leighton, P. A., Goodrich, L. V., Lu, X., Avery, B. J. et al.** (2001). Functional analysis of secreted and transmembrane proteins critical to mouse development. *Nat Genet* **28**, 241-9.
- Mo, R., Freer, A. M., Zinyk, D. L., Crackower, M. A., Michaud, J., Heng, H. H., Chik, K. W., Shi, X. M., Tsui, L. C., Cheng, S. H. et al.** (1997). Specific and redundant functions of Gli2 and Gli3 zinc finger genes in skeletal patterning and development. *Development* **124**, 113-23.
- Mullis, K., Faloona, F., Scharf, S., Saiki, R., Horn, G. and Erlich, H.** (1986). Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Cold Spring Harb Symp Quant Biol* **51 Pt 1**, 263-73.
- Naski, M. C., Colvin, J. S., Coffin, J. D. and Ornitz, D. M.** (1998). Repression of hedgehog signaling and BMP4 expression in growth plate cartilage by fibroblast growth factor receptor 3. *Development* **125**, 4977-88.

- Nusslein-Volhard, C. and Wieschaus, E.** (1980). Mutations affecting segment number and polarity in *Drosophila*. *Nature* **287**, 795-801.
- Nybakken, K. and Perrimon, N.** (2002). Heparan sulfate proteoglycan modulation of developmental signaling in *Drosophila*. *Biochim Biophys Acta* **1573**, 280-91.
- Ohlmeyer, J. T. and Kalderon, D.** (1998). Hedgehog stimulates maturation of Cubitus interruptus into a labile transcriptional activator. *Nature* **396**, 749-53.
- Ortega, N., Behonick, D. J. and Werb, Z.** (2004). Matrix remodeling during endochondral ossification. *Trends Cell Biol* **14**, 86-93.
- Park, H. L., Bai, C., Platt, K. A., Matisse, M. P., Beeghly, A., Hui, C. C., Nakashima, M. and Joyner, A. L.** (2000). Mouse *Gli1* mutants are viable but have defects in SHH signaling in combination with a *Gli2* mutation. *Development* **127**, 1593-605.
- Pathi, S., Pagan-Westphal, S., Baker, D. P., Garber, E. A., Rayhorn, P., Bumcrot, D., Tabin, C. J., Blake Pepinsky, R. and Williams, K. P.** (2001). Comparative biological responses to human Sonic, Indian, and Desert hedgehog. *Mech Dev* **106**, 107-17.
- Pepinsky, R. B., Zeng, C., Wen, D., Rayhorn, P., Baker, D. P., Williams, K. P., Bixler, S. A., Ambrose, C. M., Garber, E. A., Miatkowski, K. et al.** (1998). Identification of a palmitic acid - modified form of human Sonic hedgehog. *J Biol Chem* **273**, 14037-45.
- Porter, J. A., Young, K. E. and Beachy, P. A.** (1996). Cholesterol modification of hedgehog signaling proteins in animal development. *Science* **274**, 255-9.
- Pye, D. A., Vives, R. R., Hyde, P. and Gallagher, J. T.** (2000). Regulation of FGF-1 mitogenic activity by heparan sulfate oligosaccharides is dependent on specific structural features: differential requirements for the modulation of FGF-1 and FGF-2. *Glycobiology* **10**, 1183-92.
- Raskind, W. H., Conrad, E. U., Chansky, H. and Matsushita, M.** (1995). Loss of heterozygosity in chondrosarcomas for markers linked to hereditary multiple exostoses loci on chromosomes 8 and 11. *Am J Hum Genet* **56**, 1132-9.
- Rubin, J. B., Choi, Y. and Segal, R. A.** (2002). Cerebellar proteoglycans regulate sonic hedgehog responses during development. *Development* **129**, 2223-32.
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. and Arnheim, N.** (1985). Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**, 1350-4.
- Sambrook, J., Fritsch EF., Maniatis T.** (1989). Molecular cloning.
- Sasaki, H., Nishizaki, Y., Hui, C., Nakafuku, M. and Kondoh, H.** (1999). Regulation of *Gli2* and *Gli3* activities by an amino-terminal repression domain: implication of *Gli2* and *Gli3* as primary mediators of Shh signaling. *Development* **126**, 3915-24.
- Schipani, E., Lanske, B., Hunzelman, J., Luz, A., Kovacs, C. S., Lee, K., Pirro, A., Kronenberg, H. M. and Juppner, H.** (1997). Targeted expression of constitutively active receptors for parathyroid hormone and parathyroid hormone-related peptide delays endochondral bone formation and rescues mice that lack parathyroid hormone-related peptide. *Proc Natl Acad Sci U S A* **94**, 13689-94.

- Schlessinger, J., Plotnikov, A. N., Ibrahimi, O. A., Eliseenkova, A. V., Yeh, B. K., Yayon, A., Linhardt, R. J. and Mohammadi, M. (2000). Crystal structure of a ternary FGF-FGFR-heparin complex reveals a dual role for heparin in FGFR binding and dimerization. *Mol Cell* **6**, 743-50.
- Schweitzer, R., Vogan, K. J. and Tabin, C. J. (2000). Similar expression and regulation of Gli2 and Gli3 in the chick limb bud. *Mech Dev* **98**, 171-4.
- Seibel, M. J. and Woitge, H. W. (1999). Basic principles and clinical applications of biochemical markers of bone metabolism: biochemical and technical aspects. *J Clin Densitom* **2**, 299-321.
- Senay, C., Lind, T., Muguruma, K., Tone, Y., Kitagawa, H., Sugahara, K., Lidholt, K., Lindahl, U. and Kusche-Gullberg, M. (2000). The EXT1/EXT2 tumor suppressors: catalytic activities and role in heparan sulfate biosynthesis. *EMBO Rep* **1**, 282-6.
- St-Jacques, B., Hammerschmidt, M. and McMahon, A. P. (1999). Indian hedgehog signaling regulates proliferation and differentiation of chondrocytes and is essential for bone formation [published erratum appears in *Genes Dev* 1999 Oct 1;13(19):2617]. *Genes Dev* **13**, 2072-86.
- Stickens, D., Brown, D. and Evans, G. A. (2000). EXT genes are differentially expressed in bone and cartilage during mouse embryogenesis. *Dev Dyn* **218**, 452-64.
- Stickens, D., Clines, G., Burbee, D., Ramos, P., Thomas, S., Hogue, D., Hecht, J. T., Lovett, M. and Evans, G. A. (1996). The EXT2 multiple exostoses gene defines a family of putative tumour suppressor genes. *Nat Genet* **14**, 25-32.
- Stricker, S., Fundele, R., Vortkamp, A. and Mundlos, S. (2002). Role of Runx genes in chondrocyte differentiation. *Dev Biol* **245**, 95-108.
- Tabata, T., Eaton, S. and Kornberg, T. B. (1992). The Drosophila hedgehog gene is expressed specifically in posterior compartment cells and is a target of engrailed regulation. *Genes Dev* **6**, 2635-45.
- Tabata, T. and Takei, Y. (2004). Morphogens, their identification and regulation. *Development* **131**, 703-12.
- Takei, Y., Ozawa, Y., Sato, M., Watanabe, A. and Tabata, T. (2004). Three Drosophila EXT genes shape morphogen gradients through synthesis of heparan sulfate proteoglycans. *Development* **131**, 73-82.
- The, I., Bellaiche, Y. and Perrimon, N. (1999). Hedgehog movement is regulated through tout velu-dependent synthesis of a heparan sulfate proteoglycan. *Mol Cell* **4**, 633-9.
- Van Hul, W., Wuyts, W., Hendrickx, J., Speleman, F., Wauters, J., De Boulle, K., Van Roy, N., Bossuyt, P. and Willems, P. J. (1998). Identification of a third EXT-like gene (EXTL3) belonging to the EXT gene family. *Genomics* **47**, 230-7.
- Vortkamp, A., Gessler, M. and Grzeschik, K. H. (1991). GLI3 zinc-finger gene interrupted by translocations in Greig syndrome families. *Nature* **352**, 539-40.
- Vortkamp, A., Lee, K., Lanske, B., Segre, G. V., Kronenberg, H. M. and Tabin, C. J. (1996). Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein [see comments]. *Science* **273**, 613-22.



- Wang, B., Fallon, J. F. and Beachy, P. A.** (2000). Hedgehog-regulated processing of Gli3 produces an anterior/posterior repressor gradient in the developing vertebrate limb. *Cell* **100**, 423-34.
- Weir, E. C., Philbrick, W. M., Amling, M., Neff, L. A., Baron, R. and Broadus, A. E.** (1996). Targeted overexpression of parathyroid hormone-related peptide in chondrocytes causes chondrodysplasia and delayed endochondral bone formation. *Proc Natl Acad Sci U S A* **93**, 10240-5.
- Wijgerde, M., McMahon, J. A., Rule, M. and McMahon, A. P.** (2002). A direct requirement for Hedgehog signaling for normal specification of all ventral progenitor domains in the presumptive mammalian spinal cord. *Genes Dev* **16**, 2849-64.
- Wise, C. A., Clines, G. A., Massa, H., Trask, B. J. and Lovett, M.** (1997). Identification and localization of the gene for EXTL, a third member of the multiple exostoses gene family. *Genome Res* **7**, 10-6.
- Wu, Y. Q., Heutink, P., de Vries, B. B., Sandkuijl, L. A., van den Ouweland, A. M., Niermeijer, M. F., Galjaard, H., Reyniers, E., Willems, P. J. and Halley, D. J.** (1994). Assignment of a second locus for multiple exostoses to the pericentromeric region of chromosome 11. *Hum Mol Genet* **3**, 167-71.
- Wuyts, W., Van Hul, W., Hendrickx, J., Speleman, F., Wauters, J., De Boule, K., Van Roy, N., Van Agtmael, T., Bossuyt, P. and Willems, P. J.** (1997). Identification and characterization of a novel member of the EXT gene family, EXTL2. *Eur J Hum Genet* **5**, 382-9.
- Yamada, S., Busse, M., Ueno, M., Kelly, O. G., Skarnes, W. C., Sugahara, K. and Kusche-Gullberg, M.** (2004). Embryonic fibroblasts with a gene trap mutation in *Ext1* produce short heparan sulfate chains. *J Biol Chem* **279**, 32134-41.
- Yamagiwa, H., Tokunaga, K., Hayami, T., Hatano, H., Uchida, M., Endo, N. and Takahashi, H. E.** (1999). Expression of metalloproteinase-13 (Collagenase-3) is induced during fracture healing in mice. *Bone* **25**, 197-203.
- Yang, Y., Drossopoulou, G., Chuang, P. T., Duprez, D., Marti, E., Bumcrot, D., Vargesson, N., Clarke, J., Niswander, L., McMahon, A. et al.** (1997). Relationship between dose, distance and time in Sonic Hedgehog-mediated regulation of anteroposterior polarity in the chick limb. *Development* **124**, 4393-404.
- Zak, B. M., Crawford, B. E. and Esko, J. D.** (2002). Hereditary multiple exostoses and heparan sulfate polymerization. *Biochim Biophys Acta* **1573**, 346-55.
- Zhang, L. and Esko, J. D.** (1994). Amino acid determinants that drive heparan sulfate assembly in a proteoglycan. *J Biol Chem* **269**, 19295-9.
- Zou, H., Wieser, R., Massague, J. and Niswander, L.** (1997). Distinct roles of type I bone morphogenetic protein receptors in the formation and differentiation of cartilage. *Genes Dev* **11**, 2191-203.