

Literaturverzeichnis

- AHO, S. (2004):
Many faces of periplakin: domain-specific antibodies detect the protein throughout the epidermis, explaining the multiple protein-protein interactions.
Cell Tissue Res 316, 87-97
- AKIYAMA, M., I. MATSUO, H. SHIMIZU (2002):
Formation of cornified cell envelope in human hair follicle development.
Br J Dermatol 146, 968-976
- ALONSO, L., E. FUCHS (2003):
Stem cells of the skin epithelium.
Proc Natl Acad Sci USA 100, 11830-11835
- ANDRIANI, F., A. MARGULIS, N. LIN, S. GRIFFY, J.A. GARLICK (2003):
Analysis of microenvironmental factors contributing to basement membrane assembly and normalized epidermal phenotype.
J Invest Dermatol 120, 923-931
- AURELIAN, L., C. C. SMITH, R. WINCHURCH, M. KULKA, T. GYOTOKU, L. ZACCARO, F.J. CHERST, J. W. BURNETT (2001):
A novel gene expressed in human keratinocytes with long-term in vitro growth potential is required for cell growth.
J Invest Dermatol 116, 286-295
- BADEN, H.P. (1984):
The keratinocyte has become the subject of intensive investigation.
J Invest Dermatol 82, 305-307
- BADEN, H.P., J. KUBILUS (1983):
Fibrous proteins of bovine hoof.
J Invest Dermatol 81, 220-224
- BADEN, H. P., J.K. KUBILUS, S.B. PHILLIPS (1987):
Characterisation of Monoclonal Antibodies generated to the cornified envelope of human cultured keratinocytes.
J Invest Dermatol 89, 454-459
- BANKS-SCHLEGEL, S., H. GREEN (1981):
Involucrin synthesis and tissue assembly by keratinocytes in natural and cultured human epithelia.
J Cell Biol 90, 732-737
- BICKENBACH, J.R., E. CHISM (1998):
Selection and extended growth of murine epidermal stem cells in culture.
Exp Cell Res 244, 184-195
- BOHNERT, A., J. HORNING, I.C. MACKENZIE, E. FUSENIG (1986):
Epithelial-mesenchymal interactions control basement membrane production and differentiation in cultured and transplanted mouse keratinocytes.
Cell Tissue Res 244, 413-429

- BOWDEN, P.E., H.-J. STARK, D. BREITKREUTZ, N.E. FUSENIG (1987):
Expression and modification of keratins during terminal differentiation of mammalian epidermis.
In: Moscona, A.A. und A. Monroy: Current topics in developmental biology: The molecular and developmental biology of keratins, Vol. 22, Chap. 3
New York Academic Press, 35-68
- BOYCE, S.T., R.G. HAM (1983):
Calcium-regulated differentiation of normal human epidermal keratinocytes in chemically defined clonal culture and serum-free serial medium.
J Invest Dermatol 81, 33s-40s
- BRAGULLA, H., CH. MÜLLING (1992):
Light- and ultramicroscopic studies of hoof horn keratinization with special reference to aseptic inflammation of the bovine and equine digit.
Proc 7th International Symposium on Disorders of the Ruminant Digit.
21-25.06.1992, Rebild/Dk, 8
- BRUHNKE, J. (1931):
Vergleichende Untersuchungen der Hornwandstruktur des Zehenendes bei Huf- und Klautieren.
Dtsch Tierärztl Wochenschr 39, 4-10
- BUDRAS, K.-D., H. BRAGULLA (1992):
Besonderheiten des membrane coating material (MCM, Kittsubstanz zwischen Keratinozyten) im harten Horn des Pferdehufes.
Verh Anat Ges 85 (Anat Anz Suppl. 170), 435-436
- BUDRAS, K.-D., R.L. HULLINGER, W.O. SACK (1989):
Light and electron microscopy of keratinization in the laminar epidermis of the equine hoof with reference to laminitis.
Am J Vet Res 57 (4), 1150-1160
- BUDRAS, K.-D., CH. MÜLLING, A. HOROWITZ (1996):
Rate of keratinization of the wall segment of the hoof and its relation to width and structure of the zona alba (white line) with respect to claw disease in cattle.
Am J Vet Res 57 (4), 444-455
- BUDRAS, K.-D., H. GEYER J. MAIRL, CH. MÜLLING (1998):
Anatomy and tructure of hoof horn (Workshop report).
Proc 10th International Symposium on Lameness in Ruminants.
7.-10.09.1998, Luzern/CH, 176-187
- BULIĆ-JAKUŠ, F., T. STRAHINIĆ-BELOVARI, S. MARIĆ, D. JEŽEK, G. JURIĆ-LEBIĆ, M. VLAHOVIĆ, D. ŠERMAN (2001):
Chemically defined protein-free in vitro culture of mammalian embryo does not restrict its developmental potential for differentiation of skin appendanges.
Cell Tissues Organs 169, 134-143
- CALHOUM, M.L., A.W. STINSON (1981):
Integument: Digital organs and horn.
In: Dellmann, H.-D., E.M. Brown (Hrsg.): Textbook of Veterinary Histology.
Philadelphia: Verlag Lea & Febiger. 2. Auflage 1981, 404-411

- CHEN, T.A., R.E. HALLIWELL, P.B. HILL (2002):
Failure of extracts from *Malassezia pachydermisi* to stimulate canine keratinocyte proliferation in vitro.
Vet Dermatol 13, 323-329
- COOPER, D., T.-T. Sun (1986):
Monoclonal antibody analysis of bovine epithelial keratins.
J Biol Chem 261 (10), 4646-4654
- DAHM A.M., A. DE BRUIN, A. LINAT, C. VON TSCHARNER, M. WYDER, M.M. SUTER (2002):
Cultivation and characterisation of primary and subcultured equine keratinocytes.
Equine Vet J 34, 114-120
- DIRKS, C. (1985):
Makroskopische, licht- und elektronenmikroskopische Untersuchungen über den Rückenteil der Rinderklaue.
Berlin: FUB, Fb Vet Med, Diss.
- DROZDOFF, V., W.J., PLEDGER (1993):
Commitment to differentiation and expression of early differentiation markers in murine keratinocytes in vitro are regulated independently of extracellular calcium concentrations.
J Cell Biol 123, 909-919
- DUNNWALD, M., A. TOMANEK-CHALKLEY, D. ALEXANDRUNAS, J. FISHBAUGH, J.R. BICKENBACH (2001):
Isolating a pure population of epidermal stem cells for use in tissue engineering.
Exp Dermatol 10, 45-54
- EKFALCK, A., B. JONES, N. OBEL (1991):
Possible importance for laminitis research of recent studies on substances influencing the differentiation of cultured keratinocytes.
J Vet Med A Physiol Pathol Clin Med 38, 721-727
- EKFALCK, A., B. FUNKQUIST, B. JONES, N. OBEL (1988):
Presence of receptors for epidermal growth factor (EGF) in the matrix of the bovine hoof – a possible new approach to the laminitis problem.
J Vet Med A Physiol Pathol Clin Med 35, 321-330
- EKFALCK, A., H. RODRIGUEZ-MARTINEZ, N. OBEL (1990):
Cultivation of tissue from the matrix of the stratum medium of the equine and bovine hoof walls.
Am J Vet Res 51, 1852-1856
- EL-GHALBZOURI, A., E. LAMME, M. PONEC (2002a):
Crucial role of fibroblasts in regulating epidermal morphogenesis.
Cell Tissue Res 310, 189-199
- EL-GHALBZOURI, A., S. GIBBS, E. LAMME, C.A. VAN BLITTERSWIJK, M. PONEC (2002b):
Effect of fibroblasts on epidermal regeneration.
Br J Dermatol 147, 230-243

- ELIAS, P.M., S.K. AHN, B.E. BROWN, D. CRUMRINE, K.R. FEINGOLD (2002a):
Origin of epidermal calcium gradient: regulation by barrier status and role of active vs passive mechanisms.
J Invest Dermatol 119, 1269-1274
- ELIAS, P.M., S.K. AHN, M. DENDA, B.E. BROWN, D. CRUMRINE, L.K. KIMTAI, L. KÖMÜVES, S.H. LEE, K.R. FEINGOLD (2002b):
Modulation in epidermal calcium regulate the expression of differentiation-specific markers.
J Invest Dermatol 119, 1128-1136
- EUN, H.C., C. NAM (2003):
Alternative methods for evaluating skin irritation using three-dimensional cultures.
Exogenous Dermatology 2, 1-5
- FLECKMANN, P., B.A. DALE, K.A. HOLBROOK (1985):
Profilaggrin, a high-molecular-weight precursor of filaggrin in human epidermis and cultured keratinocytes.
J Invest Dermatol 85 (6), 507-512
- FORMANEK, M., A. TEMMEL, B. KNERER, M. WILLHEIM, W. MILLESI, J. KORNFELH (1998):
Magnetic cell separation for purification of human oral keratinocytes: an effective method for functional studies without prior cell subcultivation.
Eur Arch Otorhinolaryngol 255, 211-215
- FRESHNEY, R.I. (2000):
Culture of animal cell: a manual of basic technique.
New York, Chichester, Weinheim, Brisbane, Singapore, Toronto: Wiley-Liss. 4. Auflage 2000
- FUCHS, C. (1993):
Zur mikroskopischen Anatomie der Zehenendorgane von Pferd, Wiederkäuern, Schwein, Hund und Katze - eine vergleichende Literaturstudie
München: Univ, Vet Med Fak, Diss.
- FUCHS, E., H. GREEN (1978):
The expression of keratin genes in epidermis and cultured epidermal cells.
Cell 15, 887-897
- FUCHS, E., H. GREEN (1981):
Regulation of terminal differentiation of cultures human keratinocytes by Vitamin A.
Cell 25, 617-625
- FUCHS, E., J. DOWLING, J. SERGE, S.H. LO, Q.U. YU (1997)
Integrators of epidermal growth and differentiation: distinct functions of beta 1 and beta 4 integrins.
Curr Opin Genet Dev 7, 672-682
- FÜRST, A. (1992):
Makroskopische und mikroskopische Anatomie der Rinderklaue.
Zürich: Univ, Vet Med Fak, Diss

- FURUKAWA, F., K. FUJII, Y. HORIGUCHI, N. MATSUYOSHI, M. FUJITA, K. TODA, S. IMAMURA, H. WAKITA, S. SHIRAHAMA, M. TAKIGAWA (1997):
Roles of E- and P- Cadherin in human skin.
Microsc Res Tech 38, 343-352
- GAN, S.-Q., P.M. STEINERT (1993):
Filaggrins.
In: Kreis, T., R. Vale (Hrsg.): Guidebook to the Cytoskeletal and Motor Proteins.
Oxford, New York: Oxford University Press. 1 Auflage 1993, 152-153
- GEGENBAUR, C. (1885):
Zur Morphologie des Nagels.
Morphologisches Jahrbuch 10, 465-479
- GIBBS, S., E. BOELSMA, J. KEMPENAAR, M. PONEC (1998):
Temperature-sensitive regulation of epidermal morphogenesis and the expression of cornified envelope precursors by EGF and TGF α .
Cell Tissue Res 292, 107-114
- GIROUD, A., L. P. LEBLOND (1951):
The keratinization of epidermis and its derivatives especially the hair as shown by x-ray diffraction and histochemical studies.
Ann N Y Acad Sci 35, 613-625
- GREEN, H. (1977):
Terminal differentiation of cultured human keratinocytes.
Cell 11, 405-416
- GREEN, K.J., C.G. GAUDRY (2000):
Are desmosomes more than tethers for intermediate filaments?
Nat Rev Mol Cell Biol 1, 208-216
- HAASS, N.K., K.S.M. SMALLEY, M. HERLYN (2004):
The role of altered cell-cell communication in melanoma progression.
J Mol Histol 35, 309-318
- HAGER, B., J.R. BICKENBACH, P. FLECKMAN (1999):
Long-term culture of murine epidermal keratinocytes.
J Invest Dermatol 112, 971-976
- HÄKKINEN, L., L. KOIVISTO, H. LARJAVA (2002):
An improved method for culture of epidermal keratinocytes from newborn mouse skin.
Methods Cell Sci 23, 189-196
- HASHIMOTO, K. (1971):
Ultrastructure of the human toenail.III. Cell migration, keratinisation and formation of the intercellular cement.
Arch Dermatol Forsch 240, 1-22
- HAYWARD, A.F. (1979):
Membrane-coating granules.
Int Rev Cytol 59, 97-127

- HENDRY, K.A., A.J. MACCALLUM, C.H. KNIGHT, C.J. WILDE (2001):
Synthesis and distribution of cytokeratins in healthy and ulcerated claw epidermis.
J Dairy Res 68, 525-537
- HENDRY, K.A.K., A.J. MACCALLUM, C.H. KNIGHT, C.J. WILDE (1999):
Effect of endocrine and paracrine factors on protein synthesis and cell proliferation in bovine hoof tissue culture
J Dairy Res 66, 23-33
- HENDRY, K.A.K., M.J. LANCELOTT, C.H. KNIGHT, S.A. KEMPSON, C.J. WILDE (1995):
Protein synthesis in tissue cultured from bovine hoof.
Cell Tissue Res 281, 93-99
- HENNINGS, H., D. MICHAEL, C. CHENG, P. STEINERT, K.A. HOLBROOK, S.H. YUPSA (1980):
Calcium regulation of growth and differentiation of mouse epidermal cells in culture.
Cell 19, 245-254
- HERZHOFF, K., S. SOLLBERG, C. HUERKAMP, T. KRIEG, B. ECKES (1999):
Fibroblast expression of collagen integrin receptors alpha1beta1 and alpha2beta1 is not changed in systemic scleroderma.
Br J Dermatol 141, 218-223
- HINES, M.D., H.C. JIN, M.J. WHEELOCK, P.J. JENSEN (1999):
Inhibition of cadherin function differentially affects markers of terminal differentiation in cultured human keratinocytes.
J Cell Sci 112, 4569-4579
- HINTERHUBER, G., Y. MARQUARDT, E. DIEM, K. RAPPERSBERGER, K. WOLFF, D. FOEDINGER (2002):
Organotypic keratinocyte coculture using normal human serum: an immunomorphological study at light and electron microscopic levels.
Exp Dermatol 11, 413-420
- HOBBS, R.M., V. SILVA-VAGRAS, R. GROVES, F.M. WATT (2004):
Expression of activated MEK1 in differentiating epidermal cells is sufficient to generate hyperproliferative and inflammatory skin lesions.
J Invest Dermatol 123, 503-515
- HOCHSTETTER, T. (1998):
Die Hornqualität der Rinderklaue unter Einfluss einer Biotinsupplementierung.
Berlin: FUB, FB Vet Med, Diss
- HODIVALA, K.J., F.M. WATT (1994):
Evidence that cadherins play a role in the downregulation of integrin expression that occurs during keratinocyte terminal differentiation.
J Cell Biol 124, 589-600
- HOHMANN, H. (1902):
Untersuchungen über die Klauenlederhaut des Rindes.
Monatsh Prakt Tierheilkd 13, 49-96
- HOLBROOK, K.A., H. HENNINGS (1983):
Phenotypic expression of epidermal cells in vitro: a review.
J Invest Dermatol 81, 11s-24s

- HUANG, Y.C., T.W. WANG, J.S. SUN, F.H. LIN (2003):
Cultured keratinocytes and dermal fibroblasts on a double-layer scaffold with bi-medium culture system.
Biomed Sci Instrum 39, 500-505
- INOKUCHI, S., K. SHIMAMURA, H. TOHYA, M. KIDOKORO, M. TAMAKA, Y. UEYAMA, Y. SAWADA (1995):
Effects of fibroblasts of different origin on long term maintenance of xenotransplanted human epidermal keratinocytes in immunodeficient mice.
Cell Tissue Res 281, 223-229
- ISHIDAI-YAMAMOTO, A., H. TANAKA, H. NAKANE, H. TAKAHASHI, Y. HASHIMOTO, H. IIZUKI (1999):
Programmed cell death in normal epidermis and loricrin keratoderma. Multiple functions of Profilaggrin in keratinization.
J Invest Dermatol, Symposium Proceedings Vol. 4, 145-149
- ISLAM, TC., T. SKARIN, S. SUMITRAN, R. TOFTGRAD (2000):
Retinoids induce apoptosis in cultured keratinocytes.
Br J Dermatol 143, 709-719
- ISSEROFF, R.R., N.E. FUSENIG, D.B. RIFKIN (1983):
Plasminogen activator in differentiating mouse keratinocytes.
J Invest Dermatol 80, 217-222
- JONES, P.H., F.M. WATT (1993):
Separation of human epidermal stem cells from transit amplifying cells on the basis of differences in integrin function and expression.
Cell 73, 713-724
- KREMER, M., E. LANG, A.C. BERGER (2000):
Evaluation of dermal-epidermal skin equivalents ('composite skin') of human keratinocytes in a collagen-glycosaminoglycan matrix (Integra™ Artificial Skin).
Br J Plast Surg 53, 459-465
- KIM, B.M., S. SUZUKI, Y. NISHIMURA, S.C.UM, K. MOROTA, T. MARUGUCHI, Y. IKADA (1999):
Cellular artificial skin substitute produced by short period simultaneous culture of fibroblasts and keratinocytes.
Br J Plast Surg 52, 573-578
- KITAHARA, T., H. OGAWA (1997):
Cellular features of differentiation in the nail.
Microsc Res Tech 38, 436-442
- KITAHARA, T., H. OGAWA (1994):
Variation of differentiation in nail and bovine hoof cells.
J Invest Dermatol 102, 725-729
- KORTE, B. (1987):
Ein Beitrag zur Entwicklung der Kaue des Schafes mit besonderer Berücksichtigung der Hornbildung.
Berlin: FUB, FB Vet Med, Diss.

- KUBILUS, J. (1983):
Modulation of differentiation by Retinoids.
J Invest Dermatol 81, 55s-58s
- KUBILUS, J., J. KVEDAR, H.P. BADEN (1987):
Identification of new components of the cornified envelope of human and bovine epidermis.
J Invest Dermatol 89, 44-50
- KUBILUS, J., M. J. MACDONALD, H.P. BADEN (1979):
Epidermal proteins of cultured human and bovine keratinocytes.
Biochim Biophys Acta 578, 484-492
- KUBILUS, J., I. SCOTT, C.R. CLIVE, J. YENDLE, J. KVEDAR, H.P. BADEN (1985):
The occurrence of profilaggrin and its processing in cultured keratinocytes.
J Invest Dermatol 85, 513-517
- KÜNZEL, E. (1990):
Haut (Integumentum commune).
In: Mosimann, W., T. Kohler: Zytologie, Histologie und mikroskopische Anatomie der Haussäugetiere, Kap 16
Berlin, Parey Verlag, 259-287
- KVEDAR, J.C., J. KUBILUS, H.P. BADEN (1986):
Cytokeratins of the bovine hoof: Classification and studies on expression.
Biochim Biophys Acta 884, 462-473
- KYHSE-ANDERSEN, J. (1984):
Electroblotting of multiple gels: a simple apparatus without buffer tank for rapid transfer of proteins from polyacrylamide to nitrocellulose.
J Biochem Biophys Meth 10, 203-209
- LANDMANN, L. (1988):
The epidermal permeability barrier.
Anat Embryol (Berl) 178 (1), 1-13
- LAVKER, R.M., T.-T. SUN (1983):
Rapid modulation of keratinocyte differentiation by the external environment.
J Invest Dermatol 80, 228-237
- LEE, L.D., J. KUBILUS, H.P. BADEN (1979):
Intraspecies heterogeneity of epidermal keratins isolated from bovine hoof and snout.
Biochem J 177, 187-196
- LEVY, L., S. BROAD, D. DIEKMANN, R.D. EVANS, F.M. WATT (2000):
 β 1 integrins regulate keratinocyte adhesion and differentiation by distinct mechanisms.
Mol Biol Cell 11, 453-466
- LI, L., T. TENNENBAUM, S.H. YUPSA (1996):
Suspension-induced murine keratinocyte differentiation is mediated by Calcium.
J Invest Dermatol 106, 254-260

- LIEBICH, H.-G. (1999):
Funktionelle Histologie der Haussäugetiere: Lehrbuch und Farbatlas für Studium und Praxis.
Stuttgart, New York: Schattauer. 3. Auflage 1999
- MAAS-SZABOWSKI, N., A. SHIMOTOYODOME, N.E. FUSENIG (1999):
Keratinocyte growth regulation in fibroblast cocultures via a double paracrine mechanism.
J Cell Sci 112, 1843-1853
- MAAS-SZABOWSKI, N., A. SZABOWSKI, H.-J. STARK, S. ANDRECHT, A. KOLBUS, M. SCHROPP-KISTNER, P. ANGEL, N.E. FUSENIG (2001):
Organotypic cocultures with genetically modified mouse fibroblasts as a tool to dissect molecular mechanisms regulating keratinocyte growth and differentiation.
J Invest Dermatol 116, 816-820
- MACK, J.W., A.C. STEVEN, P.M. STEINERT (1993):
The mechanism of interaction of filaggrin with intermediate filaments. The ionic zipper hypothesis.
J Mol Biol 232, 50-66
- MADISON, K.C., D.C. S WARTZENDRUBER, P.W. WERTZ, D.T. DOWNING (1988):
Lamellar granule extrusion and stratum corneum intercellular lamellae in murine keratinocyte cultures.
J Invest Dermatol 90, 110-116
- MANAVES, V., W. QUIN, A.L. BAUER, S. ROSSIE, M. KOBAYASHI, S.G. RANE (2004):
Calcium and vitamin D increase mRNA levels for the growth control hIK1 channel in human epidermal keratinocytes but functional channels are not observed.
BMC Dermatol 4, doi: 10.186/1471-5945-4-7
<http://www.biomedcentral.com/1471-5945/4/7>
- MARCHESE, C., A. FELICI, V. VISCO, G. LUCANIA, M. IGARASHI, M. PICARDO, L. FRATI, M.R. TORRISI (2001):
Fibroblast growth factor 10 induces proliferation and differentiation of human primary cultured keratinocytes.
J Invest Dermatol 116, 623-628
- MATOLTSY, A.G. (1976):
Keratinization.
J Invest Dermatol 67, 20-25
- MALTOLTSY, A.G., P.F. PRAKKAL (1965):
Membrane-coating granules of keratinizing epithelia.
J Cell Biol 24, 297-307
- MEHUL, B., D. BERNARD, R. SCHMIDT (2001):
Calmodulin-like skin protein: a new marker of keratinocyte differentiation.
J Invest Dermatol 116, 905-909
- MERNE, M., S. SYRJÄNEN (2003):
The mesenchymal substrate influences the epithelial phenotype in a three-dimensional cell culture.
Arch Dermatol Res 295, 190-198

- MILSTONE, L.M. (1983):
Population dynamics in cultures of stratified squamous epithelium.
J Invest Dermatol 81, 69s-74s
- MILSTONE, L.M., J. McGUIRE (1981):
Different polypeptides from the intermediate filaments in bovine hoof and oesophageal epithelium and in aortic endothelium.
J Cell Biol 88, 312-316
- MILTENYI, S., W. MÜLLER, W. WEICHEL, A. RADBRUCH (1990):
High gradient magnetic cell separation with MACS.
Cytometry 11, 231-238
- MOLL, R., W.W. FRANKE, D.L. SCHILLER, B. GEIGER, R. KREPLER (1982):
The catalog of human cytokeratins: Patterns of expression in normal epithelia, tumors and cultured cells.
Cell 11, 11-24
- MÜLLING, CH. (1993):
Struktur, Verhornung und Hornqualität in Ballen, Sohle und weißer Linie der Rinderklaue und ihre Bedeutung für Klauenerkrankungen.
Berlin: FUB, FB Vet Med, Diss
- MÜLLING, CH. K. W. (2000):
Three-dimensional appearance of bovine keratinocytes in different stages of differentiation revealed by cell maceration and scanning electron microscopic investigation.
Folia Morphol (Warsz) 59, 239-246
- MÜLLING, CH., K.-D. BUDRAS (1998):
Der Interzellularkitt (Membrane Coating Material, MCM) in der Epidermis der Rinderklaue.
Wien Tierärztl Monatsschr 85, 216-223
- MÜLLING, CH., K.-D. BUDRAS (2002):
Klaue (Ungula)
In: Budras, K.-D., A. Wünsche: Atlas der Anatomie des Rindes, Kap 2
Hannover, Schlütersche, 1 Auflage, 2002, 26
- MÜLLING, CH., U. NEBEL, K.-D. BUDRAS (2004):
Regulation of cellular proliferation and differentiation in claw epidermis by epidermal-dermal interactions.
In: Abstract book of the 25th Congress of the European Association of Veterinary Anatomists, 28-31.07.2004 Oslo/Norway, 133
- MÜLLING, CH., H. BRAGULLA, K.-D. BUDRAS, S. REESE (1994):
Strukturelle Faktoren mit Einfluss auf die Hornqualität und Prädilektionsstellen für Erkrankungen an der Fussungsfläche der Rinderklaue.
Schweiz Archv Tierheilkd 136, 49-57

- MÜLLING, CH.K.W., H.H. BRAGULLA, S. REESE, K.-D. BUDRAS, W. STEINBERG (1999):
How structures in bovine hoof epidermis are influenced by nutritional factors.
Anat Histol Embryol 28, 103-108
Proc 13th International Symposium on Lameness in Ruminants.
11-15.02.2004, Maribor/SLO, 69-71
- NEBEL, U., D. HOFFMANN, CH.K.W. MÜLLING, K.-D. BUDRAS (2004a):
Co-culture of epidermal and dermal bovine hoof cells in a perfusion chamber -
Development and first experimental experiences.
Proc 13th International Symposium on Lameness in Ruminants.
11-15.02.2004, Maribor/SLO, 69-71
- NEBEL, U., CH. MÜLLING, M. NORDHOFF, K.-D., BUDRAS (2004b):
In vitro infection of bovine epidermal cells and bovine skin explants with
Treponemes.
Proc 13th International Symposium on Lameness in Ruminants.
11-15.02.2004, Maribor/SLO, 141-143
- NIRUNSUKSIRI, W., S.-H. ZHANG, FLECKMAN (1998):
Reduced stability and bi-allelic, coequal expression of profilaggrin mRNA in
Keratinocytes cultured from subjects with ichthyosis vulgaris.
J Invest Dermatol 110, 854-861
- O'KEEFE, E.J., M.L. CHUI, R.E. PAYNE (1988):
Stimulation of growth of keratinocytes by basic fibroblast growth factor.
J Invest Dermatol 90, 767-769
- OKU, H., C. KUMAMOTO, T. MIYAGI, T. HIYANE, J. NAGATA, I. CHINEN (1994):
Serum-free culture of rat keratinocytes.
In Vitro Cell Dev Biol Amin 30A, 496-503
- OWENS, D.W., V.G. BRUNTON, E.K. PARKINSON, M.C. FRAME (2000):
E-cadherin at the cell periphery is a determinant of keratinocyte differentiation in
vitro.
Biochem Biophys Res Commun 369, 369-376
- PARK, H.-S., H.J. KANG, C.H. KIM, C.H. KIM, E.S. HAN, K. HAN, T.H. KIM, Y.J. GIN,
Y.S. SON (2004):
Application of physical force is essential to enrich for epidermal stem cells in primary
human keratinocyte isolation.
Tissue Eng 10, 343-351
- PASONEN-SEPPÄNEN, S., S. KARVINEN, K. TÖRRÖNEN, J.M.T. HYTTINEN, T.
JOKELA, M.J. LAMMI, M.I. TAMMI, R. TAMMI (2003):
EGF upregulates, whereas TGF- β downregulates, the hyaluronan synthase Has2
and Has3 in organotypic keratinocyte cultures: correlations with epidermal
proliferation and differentiation.
J Invest Dermatol 120, 1038-1044
- PONEC, M. (2002):
Skin constructs for replacement of skin tissue for in vitro testing.
Adv Drug Deliv Rev 54 Suppl 1, 19-30

- PONEC, M., S. GIBBS, A. WEERHEIM, J. KEMPENAAR, A. MULDER, A.M. MOMMAAS (1997):
Epidermal growth factor and temperature regulate keratinocyte differentiation.
Arch Dermatol Res 289, 317-326
- POTTEN, C.S., C. BOOTH (2002):
Keratinocyte stem cells: a commentary
J Invest Dermatol 119, 888-899
- POUMAY, Y., M.R. PITTELKOW (1995):
Cell density and culture factors regulate keratinocyte commitment to differentiation and expression of suprabasal K1/K10 keratins.
J Invest Dermatol 104, 271-276
- POUMAY, Y., M. LECLERCQ-SMEKENS (1998):
In vitro models of epidermal differentiation.
Folia Med (Plodiv) 40, 5-12
- POUMAY, Y., I.H. ROLAND, M. LECLERCQ-SMEKENS, R. LELOUP (1994):
Basal detachment of the epidermis using dispase: tissue spatial organization and fate of integrin alpha-6 beta-4 and hemidesmosomes.
J Invest Dermatol 102, 11-117
- PRESLAND, R.B., M. TOMIC-CANIC, S.P. LEWIS, B.A. DALE (2001):
Regulation of human profilaggrin promoter activity in cultured cells by retinoic acid and glucocorticoids.
J Dermatol Sci 27, 192-205
- POUMAY, Y., I.H. ROLAND, M. LECLERCQ-SMEKENS, R. LELOUP (1994):
Basal detachment of epidermis using dispase: tissue spatial organization and fate of integrin $\alpha 6\beta 4$ and hemidesmosomes.
J Invest Dermatol 102, 111-117
- RHEINWALD, J.G., H. GREEN (1975a):
Formation of a keratinizing epithelium in culture by a cloned cell line derived from a teratoma.
Cell 6, 317-330
- RHEINWALD, J.G., H. GREEN (1975b):
Serial cultivation of strains of human epidermal keratinocytes formation of keratinizing colonies from single cell.
Cell 6, 331-344
- RHEINWALD, J.G., H. GREEN (1977):
Epidermal growth factor and the multiplication of cultured human epidermal keratinocytes.
Nature 265, 421-424
- RICE, R.H., H. GREEN (1979):
Presence in human epidermal cells of a soluble protein precursor of the cross-linking by calcium ions.
Cell 18, 681-694

- ROBINSON, H.W., C.G. HODGEN (1940):
The biuret reaction in the determination of serum proteins. I.A study of the conditions necessary for the production of a stable color which bears a quantitative relationship to the protein concentration.
J Biol Chem 135, 707
- ROMIES, B. (1989):
Mikroskopische Technik.
München: Verlag Urban und Schwarzenberg. 17 Auflage 1989
- SCHELFHOUT, V.R., E.D. COENE, B. DELAEY, A.A. WAEYTENS, L. DE RYCKE, M. DELEU, C. R. DE POTTER (2002):
The role of heregulin-alpha as a motility factor and amphiregulin as a growth factor in wound healing.
J Pathol 198, 523-533
- SCHILLER, D.L., W.W. FRANKE, B. GEIGER (1982):
A subfamily of relatively large and basic cytokeatin polypeptides as defined by mapping is represented by one or several polypeptides in epithelial cells.
EMBO J 1 (6), 761-769
- SHIRAKATA, Y.T., H. KOMURASAKI, Y. TOYODA, K. HANAKAWA, S. YAMASAKI, K. TOKUMARU, K. SAYAMA, K. HASHIMOTO (2000):
Epiregulin, a novel member of the epidermal growth factor family, is an autocrine growth factor in normal humankeratinocytes.
J Biol Chem 275 (8), 5748-5753
- SMOLA, H., G. THIEKÖTTER, N.E. FUSENIG (1993):
Mutual induction of growth factor gene expression by epidermal-dermal cell interaction.
J Cell Biol 122, 417-429
- STARK, H.-J., A. SZABOWSKI, N.E. FUSENIG, N. MASS-SZABOWSKI (2004):
Organotypic cocultures as skin equivalents: a complex and sophisticated *in vitro* system.
Biological Procedures Online 6, 55-60
Doi:10.1251/bpo72
- STARK, H.-J., M. BAUR, D. BREITKREUTZ, N. MIRANCEA, N.E. FUSENIG (1999):
Organotypic keratinocyte cocultures in defined medium with regular epidermal morphogenesis and differentiation.
J Invest Dermatol 112, 681-691
- STEINERT, P.M., W.W. IDLER (1975):
The polypeptide composition of bovine epidermal α -keratin.
Biochem J 151, 603-614
- STREYER, L. (1996):
Biochemie.
Heidelberg, Berlin, New York: Spektrum Akademischer Verlag. 4. Auflage
- SULTAN, K.R., H.P. HAAGSMAN (2001):
Species-specific primary cell culture: a research tool in veterinary science.
Veterinary Sciences Tomorrow 1
www.vetscite.org/issue1/reviews/sultan_0800.pdf

- SUN, T.-T., R. EICHNER, W.G. NELSON, S.C.G. TSENG, R.A. WEISS, M. JARVINEN, J. WOODCOCK-MITCHELL (1983):
Keratin classes: Molecular markers of different types of epithelial differentiation.
J Invest Dermatol 83, 109s-115s
- SZULGIT, G., R. RUDOLPH, A. WANDEL, M. TENENHAUS, R. PANOS, H. GARDNER (2002):
Alterations in fibroblast $\alpha 1\beta 1$ Integrin collagen receptor expression in keloids and hypertrophic scars.
J Invest Dermatol 118, 409-415
- THACHER, S.M., R.H. RICE (1985):
Keratinocyte-specific transglutaminase of cultured human epidermal cells: relation to cross-linked envelope formation and terminal differentiation.
Cell 40, 685-695
- TINKLE, C.L., T. LECHLER, H.A. PASOLLI, E. FUCHS (2004):
Conditional targeting of E-cadherin in skin: Insights into hyperproliferative and degenerative responses.
Proc Natl Acad Sci USA 101, 552-557
- TOMLINSON, D.J., CH. MÜLLING, T.M. FAKLER (2004):
Formation of keratins in the bovine claw: Roles of hormones, minerals, and vitamins in the functional claw integrity.
J Dairy Sci 87, 797-809
- TSAO, M.C., B.J. WALTHALL, R.G. HAM (1982):
Clonal growth of normal human epidermal keratinocytes in a defined medium.
J Cell Physiol 110, 219-229
- TSENG, S.C.G., M.J. JARVINEN, W.G. NELSON, J.-W. HUANG, J. WOODCOCK-MITCHELL, T.-T. SUN (1982):
Correlation of specific keratins with different types of epithelial differentiation: Monoclonal antibody studies.
Cell 30, 361-372
- TUTCHI, L., A.A. CHASSOT, R. REZZONICO, K. YEOW, A. LOUBAT, B. FERRUA, G. LENEGRATE, J.P. ORTONNE, G. PONZIO (2002):
Dynamic characterization of the molecular events during *in vitro* epidermal wound healing.
J Invest Dermatol 119, 56-63
- UCHI, H., H. TERAOKA, T. KOGA, M. FURUE (2000):
Cytokines and chemokines in the epidermis.
J Dermatol Sci 24 Suppl1, 29-38
- VENABLE, J.H., R. COGGESHALL (1965):
A simplified lead citrate stain for use in electron microscopy.
J Cell Biol 25, 407-408
- WAKITA, H., M. TAKIGAWA (1999):
Activation of epidermal growth factor receptor promotes late terminal differentiation of cell-matrix interaction-disrupted keratinocytes.
J Biol Chem 274, 37285-38291

- WANNER, R., B. WOLFF, F. GLOWACKI, G. KOLDE, B. WITTIG (1999):
The loss of desmosomes after retinoic acid treatment results in apparent inhibition of HaCaT keratinocyte differentiation.
Arch Dermatol Res 291, 346-353
- WARZECHA, CH. (1993):
The ruminant hoof: Morphological and histochemical findings in cattle, sheep and goat.
Zürich: Univ, Vet Med Fak, Diss
- WATSON, A., C. BAKER, J. BAILEY, T. FRAY, P. MARKWELL (2004):
A high yield method for growing primary canine keratinocytes.
Vet J 168, 81-86
- WATT, F.M. (1988):
Proliferation and terminal differentiation of human epidermal keratinocytes in culture
Biochem Soc Trans 16, 666-668
- WATT, F.M. (1983):
Involucrin and other markers of keratinocyte terminal differentiation.
J Invest Dermatol 81, 100s-103s
- WATT, F.M., H. GREN (1981):
Involucrin synthesis is correlated with cell size in human epidermal cultures.
J Cell Biol 90, 738-742
- WATT, F.M., H. GREEN (1982):
Stratification and terminal differentiation of cultured epidermal cells.
Nature 295, 434-436
- WESTERFELD, I. (2003):
Struktur und Funktion des bovinen Klauenbeinträgers.
Berlin: FUB, FB Vet Med, Diss
- WERNER, S., H. SMOLA (2001):
Paracrine regulation of keratinocyte proliferation and differentiation.
Trends Cell Biol 11, 143-146
- WESTERMEIER, R. (1990):
Elektrophoresepraktikum.
Weinheim: Verlag VHC. 1. Auflage 1990, 167-196
- WILKENS, H. (1963):
Zur makroskopischen und mikroskopischen Morphologie der Rinderklaue mit einem Vergleich der Architektur von Klauen- und Huftröhrchen.
Hannover: tierärztl. Hochsch., Habil.-Schr.
- WUNN, D., J. WARDROP, K. MEYERS, J. KRAMER, C. RAGLE (1999):
Culture and characterization of equine terminal arch endothelial cells and hoof keratinocytes.
Am J Vet Res 60, 128-132

YAMASAKI, K., N. TORIU, Y. HANAKAWA, Y. SHIRAKATA, K. SAYAMA, A. TAKAYANAGI, M. OHTSOBO, S. GAMOU, N. SHIMIZU, M. FUJII, K. MIYAZONO, K. HASHIMOTO (2003):

Keratinocyte growth inhibition by high-dose epidermal growth factor is mediated by transforming growth factor β autoinduction: A negative feedback mechanism for keratinocyte growth.

J Invest Dermatol 120, 1030-1037

YUPSA, S.H., A.E. KILKENNY, P.M. STEINERT, D.R. ROOP (1989):

Expression of murine epidermal differentiation markers is tightly regulated by restricted extracellular calcium concentrations in vitro.

J Cell Biol 109, 1207-1217

ZHANG, D., S. KARUNARATNE, M. KESSLER, D. MAHONY, J.A. ROTNAGEL (2002):

Characterization of mouse filaggrin: evidence for nuclear engulfment and translocation of profilaggrin B-domain during epidermal differentiation.

J Invest Dermatol 119, 905-912

ZHOU, S., N. MATSUYOSHI, S.-B. LIANG, T. TAKEKEUCHI, Y. OHTSUKI, Y. MIYACHI (2002):

Expression of T-cadherin in basal keratinocytes of skin.

J Invest Dermatol 118, 1080-1084

ZHU, A.J., I. HAASE, F.M. WATT (1999):

Signaling via β_1 integrins and mitogen-activated protein kinase determines human epidermal stem cell fate in vitro.

Proc Natl Acad Sci USA 96, 6728-6733

ZIETZSCHMANN, O. (1918):

Das Zehenendorgan der rezenten Säugetiere: Krallen, Nägel, Hufe.

Schweiz Arch Tierheilkd 60 (6), 241-272