

9 Literatur

- Adachi, S., Kuwata, T., Miyake, M., and Iwata, M. (2001). Induction of CCR7 expression in thymocytes requires both ERK signal and Ca²⁺-signal. *Biochem Biophys Res Commun* 288, 1188-1193.
- Albelda, S. M., and Buck, C. A. (1990). Integrins and other cell adhesion molecules. *Faseb J* 4, 2868-2880.
- Alessi, D. R., Cuenda, A., Cohen, P., Dudley, D. T., and Saltiel, A. R. (1995). PD 098059 is a specific inhibitor of the activation of mitogen- activated protein kinase kinase in vitro and in vivo. *J Biol Chem* 270, 27489-27494.
- Alt, C., Laschinger, M., and Engelhardt, B. (2002). Functional expression of the lymphoid chemokines CCL19 (ELC) and CCL 21 (SLC) at the blood-brain barrier suggests their involvement in G- protein-dependent lymphocyte recruitment into the central nervous system during experimental autoimmune encephalomyelitis. *Eur J Immunol* 32, 2133-2144.
- Ansel, K. M., and Cyster, J. G. (2001). Chemokines in lymphopoiesis and lymphoid organ development. *Curr Opin Immunol* 13, 172-179.
- Ansel, K. M., Ngo, V. N., Hyman, P. L., Luther, S. A., Förster, R., Sedgwick, J. D., Browning, J. L., Lipp, M., and Cyster, J. G. (2000). A chemokine-driven positive feedback loop organizes lymphoid follicles. *Nature* 406, 309-314.
- Arenberg, D. A., Zlotnick, A., Strom, S. R., Burdick, M. D., and Strieter, R. M. (2001). The murine CC chemokine, 6C-kine, inhibits tumor growth and angiogenesis in a human lung cancer SCID mouse model. *Cancer Immunol Immunother* 49, 587-592.
- Arvanitakis, L., Geras-Raaka, E., Varma, A., Gershengorn, M. C., and Cesarman, E. (1997). Human herpesvirus KSHV encodes a constitutively active G-protein- coupled receptor linked to cell proliferation. *Nature* 385, 347-350.
- Asada, M., Furukawa, K., Segawa, K., Endo, T., and Kobata, A. (1997). Increased expression of highly branched N-glycans at cell surface is correlated with the malignant phenotypes of mouse tumor cells. *Cancer Res* 57, 1073-1080.
- Asensio, V. C., Kincaid, C., and Campbell, I. L. (1999). Chemokines and the inflammatory response to viral infection in the central nervous system with a focus on lymphocytic choriomeningitis virus. *J Neurovirol* 5, 65-75.
- Azenshtein, E., Luboshits, G., Shina, S., Neumark, E., Shahbazian, D., Weil, M., Wigler, N., Keydar, I., and Ben-Baruch, A. (2002). The CC chemokine RANTES in breast carcinoma progression, regulation of expression and potential mechanisms of promalignant activity. *Cancer Res* 62, 1093-1102.
- Azuma, H., Takahara, S., Ichimaru, N., Wang, J. D., Itoh, Y., Otsuki, Y., Morimoto, J., Fukui, R., Hoshiga, M., Ishihara, T., et al. (2002). Marked prevention of tumor growth and metastasis by a novel immunosuppressive agent, FTY720, in mouse breast cancer models. *Cancer Res* 62, 1410-1419.
- Bacon, K. B., Flores Romo, L., Life, P. F., Taub, D. D., Premack, B. A., Arkinstall, S. J., Wells, T. N., Schall, T. J., and Power, C. A. (1995). IL-8-induced signal transduction in T lymphocytes involves receptor-mediated activation of phospholipases C and D. *J Immunol* 154, 3654-3666.
- Baekkevold, E. S., Yamanaka, T., Palframan, R. T., Carlsen, H. S., Reinholt, F. P., von Andrian, U. H., Brandtzaeg, P., and Haraldsen, G. (2001). The CCR7 ligand elc (CCL19) is transcytosed in high endothelial venules and mediates T cell recruitment. *J Exp Med* 193, 1105-1112.
- Baggioolini, M., and Dahinden, C. A. (1994). CC chemokines in allergic inflammation. *Immunol Today* 15, 127-133.
- Baggioolini, M., Dewald, B., and Moser, B. (1994). Interleukin-8 and related chemotactic cytokines-CXC and CC chemokines. *Adv Immunol* 55, 97-179.
- Baggioolini, M., Dewald, B., and Moser, B. (1997). Human chemokines: an update. *Annu Rev Immunol* 15, 675-705.

- Baird, A. M., Gerstein, R. M., and Berg, L. J. (1999). The role of cytokine receptor signaling in lymphocyte development. *Curr Opin Immunol* 11, 157-166.
- Bais, C., Santomasso, B., Coso, O., Arvanitakis, L., Raaka, E. G., Gutkind, J. S., Asch, A. S., Cesarman, E., Gershengorn, M. C., Mesri, E. A., and Gerhengorn, M. C. (1998). G-protein-coupled receptor of Kaposi's sarcoma-associated herpesvirus is a viral oncogene and angiogenesis activator. *Nature* 391, 86-89.
- Banas, B., Wornle, M., Berger, T., Nelson, P. J., Cohen, C. D., Kretzler, M., Pfirsinger, J., Mack, M., Lipp, M., Grone, H. J., and Schlondorff, D. (2002). Roles of SLC/CCL21 and CCR7 in human kidney for mesangial proliferation, migration, apoptosis, and tissue homöostasis. *J Immunol* 168, 4301-4307.
- Bancroft GJ, Kelly JP. (1994). Macrophage activation and innate resistance to infection in SCID mice. *Immunobiology* 191, 424-431.
- Bar-Eli, M. (1999). Role of interleukin-8 in tumor growth and metastasis of human melanoma. *Pathobiology* 67, 12-18.
- Bardi, G., Lipp, M., Baggolini, M., and Loetscher, P. (2001). The T cell chemokine receptor CCR7 is internalized on stimulation with ELC, but not with SLC. *Eur J Immunol* 31, 3291-3297.
- Bautz, F., Denzlinger, C., Kanz, L., and Mohle, R. (2001). Chemotaxis and transendothelial migration of CD34(+) hematopoietic progenitor cells induced by the inflammatory mediator leukotriene D4 are mediated by the 7-transmembrane receptor CysLT1. *Blood* 97, 3433-3440.
- Bell D, Chomarat P, Broyles D, Netto G, Harb GM, Lebecque S, Valladeau J, Davoust J, Palucka KA, Banchereau J. (1999). In breast carcinoma tissue, immature dendritic cells reside within the tumor, whereas mature dendritic cells are located in peritumoral areas. *J Exp Med* 190, 1417-1426.
- Belperio JA, Keane MP, Arenberg DA, Addison CL, Ehlert JE, Burdick MD, Strieter RM.(2000). CXC chemokines in angiogenesis. *J Leukoc Biol* 68, 1-8.
- Ben Baruch, A., Michiel, D. F., and Oppenheim, J. J. (1995). Signals and receptors involved in recruitment of inflammatory cells. *J. Biol. Chem.* 270, 11703-11706.
- Bennett, T. A., Maestas, D. C., and Prossnitz, E. R. (2000). Arrestin binding to the G protein-coupled N-formyl peptide receptor is regulated by the conserved "DRY" sequence. *J Biol Chem* 275, 24590-24594.
- Berstein, G., Blank, J. L., Jhon, D. Y., Exton, J. H., Rhee, S. G., and Ross, E. M. (1992). Phospholipase C-beta 1 is a GTPase-activating protein for Gq/11, its physiologic regulator. *Cell* 70, 411-418.
- Biber, K., Rappert, A., Kettenmann, H., Brouwer, N., Copray, S. C., and Boddeke, H. W. (2002). Neuronal SLC (CCL21) expression: implications for the neuron-microglial signaling system. *Ernst Schering Res Found Workshop* 39, 45-60.
- Biber, K., Sauter, A., Brouwer, N., Copray, S. C., and Boddeke, H. W. (2001). Ischemia-induced neuronal expression of the microglia attracting chemokine Secondary Lymphoid-tissue Chemokine (SLC). *Glia* 34, 121-133.
- Birkenbach, M., Josefson, K., Yalamanchili, R., Lenoir, G., and Kieff, E. (1993). Epstein-Barr virus-induced genes: first lymphocyte-specific G protein-coupled peptide receptors. *J Virol* 67, 2209-2220.
- Bleul, C. C., Fuhlbrigge, R. C., Casasnovas, J. M., Aiuti, A., and Springer, T. A. (1996). A highly efficacious lymphocyte chemoattractant, stromal cell-derived factor 1 (SDF-1) *J. Exp. Med.* 184, 1101-1109.
- Bohm, S. K., Khitin, L. M., Smeekens, S. P., Grady, E. F., Payan, D. G., and Bunnett, N. W. (1997). Identification of potential tyrosine-containing endocytic motifs in the carboxyl-tail and seventh transmembrane domain of the neurokinin 1 receptor. *J Biol Chem* 272, 2363-2372.
- Boyer, B., Bourgeois, Y., and Poupon, M. F. (2002). Src kinase contributes to the metastatic spread of carcinoma cells. *Oncogene* 21, 2347-2356.
- Boyer, B., Valles, A. M., and Edme, N. (2000). Induction and regulation of epithelial-mesenchymal transitions. *Biochem Pharmacol* 60, 1091-1099.
- Braun SE, Chen K, Foster RG, Kim CH, Hromas R, Kaplan MH, Broxmeyer HE, Cornetta K. (2000). The CC chemokine CK beta-11/MIP-3 beta/ELC/Exodus 3 mediates tumor rejection

- of murine breast cancer cells through NK cells. *J Immunol* 164, 4025-4031.
- Breitfeld D., Ohl, L., Kremmer, E., Ellwart, J., Sallusto, F., Lipp, M., and Förster, R. (2000). Follicular B helper T cells express CXC chemokine receptor 5, localize to B cell follicles, and support immunoglobulin production. *J Exp Med* 192, 1545-1552.
 - Brinkmann, V., Chen, S., Feng, L., Pinschewer, D., Nikolova, Z., and Hof, R. (2001a). FTY720 alters lymphocyte homing and protects allografts without inducing general immunosuppression. *Transplant Proc* 33, 530-531.
 - Brinkmann, V., Davis, M. D., Heise, C. E., Albert, R., Cottens, S., Hof, R., Bruns, C., Prieschl, E., Baumruker, T., Hiestand, P., et al. (2002). The immune modulator FTY720 targets sphingosine 1-phosphate receptors. *J Biol Chem* 277, 21453-21457.
 - Brinkmann, V., and Lynch, K. (2002). FTY720: targeting G-protein-coupled receptors for sphingosine 1-phosphate in transplantation and autoimmunity. *Curr Opin Immunol* 14, 569.
 - Brinkmann, V., Pinschewer, D. D., Feng, L., and Chen, S. (2001b). FTY720: altered lymphocyte traffic results in allograft protection. *Transplantation* 72, 764-769.
 - Brinkmann, V., Wilt, C., Kristofic, C., Nikolova, Z., Hof, R. P., Chen, S., Albert, R., and Cottens, S. (2001c). FTY720: dissection of membrane receptor-operated, stereospecific effects on cell migration from receptor-independent antiproliferative and apoptotic effects. *Transplant Proc* 33, 3078-3080.
 - Buras, J. A., Stahl, G. L., Svoboda, K. K., and Reenstra, W. R. (2000). Hyperbaric oxygen downregulates ICAM-1 expression induced by hypoxia and hypoglycemia: the role of NOS. *Am J Physiol Cell Physiol* 278, 292-302.
 - Burgstahler, R., Kempkes, B., Steube, K., and Lipp, M. (1995). Expression of the chemokine receptor BLR2/EBI1 is specifically transactivated by Epstein-Barr virus nuclear antigen 2. *Biochem Biophys Res Commun* 215, 737-743.
 - Butcher, E. C. (1991). Leukocyte-endothelial cell recognition: three (or more) steps to specificity and diversity. *Cell* 67, 1033-1036.
 - Butcher, E. C., and Picker, L. J. (1996). Lymphocyte homing and homöostasis. *Science* 272, 60-66.
 - Campbell, J. J., Bowman, E. P., Murphy, K., Youngman, K. R., Siani, M. A., Thompson, D. A., Wu, L., Zlotnik, A., and Butcher, E. C. (1998). 6-C-kine (SLC), a lymphocyte adhesion-triggering chemokine expressed by high endothelium, is an agonist for the MIP-3beta receptor CCR7. *J Cell Biol* 141, 1053-1059.
 - Campbell, J. J., and Butcher, E. C. (2000). Chemokines in tissue-specific and microenvironment-specific lymphocyte homing. *Curr Opin Immunol* 12, 336-341.
 - Campbell, J. J., Foxman, E. F., and Butcher, E. C. (1997). Chemoattractant receptor cross talk as a regulatory mechanism in leukocyte adhesion and migration. *Eur. J. Immunol.* 27, 2571-2578.
 - Campbell, J. J., Haraldsen, G., Pan, J., Rottman, J., Qin, S., Ponath, P., Andrew, D. P., Warnke, R., Ruffing, N., Kassam, N., et al. (1999). The chemokine receptor CCR4 in vascular recognition by cutaneous but not intestinal memory T cells. *Nature* 400, 776-780.
 - Cao, Y., Chen, C., Weatherbee, J. A., Tsang, M., and Folkman, J. (1995). gro-beta, a -C-X-C- chemokine, is an angiogenesis inhibitor that suppresses the growth of Lewis lung carcinoma in mice. *J Exp Med* 182, 2069-2077.
 - Capila I, Linhardt RJ. (2002). Heparin-protein interactions. *Angew Chem Int Ed* 41, 391-412.
 - Carey, I., Williams, C. L., Ways, D. K., and Noti, J. D. (1999). Overexpression of protein kinase C-alpha in MCF-7 breast cancer cells results in differential regulation and expression of alphavbeta3 and alphavbeta5. *Int J Oncol* 15, 127-136
 - Chabot, D. J., Chen, H., Dimitrov, D. S., and Broder, C. C. (2000). N-linked glycosylation of CXCR4 masks coreceptor function for CCR5-dependent human immunodeficiency virus type 1 isolates. *J Virol* 74, 4404-4413.
 - Chen, S., Bacon, K. B., Garcia, G., Liao, R., Pan, Z. K., Sullivan, S. K., Nakano, H., Matsuzawa, A., Brinkmann, V., and Feng, L. (2001). FTY720, a novel transplantation drug, modulates lymphocyte migratory responses to chemokines. *Transplant Proc* 33, 3057-3063.
 - Chen, S. C., Vassileva, G., Kinsley, D., Holzmann, S., Manfra, D., Wiekowski, M. T., Romani, N., and Lira, S. A. (2002). Ectopic expression of the murine chemokines CCL21a

- and CCL21b induces the formation of lymph node-like structures in pancreas, but not skin, of transgenic mice. *J Immunol* 168, 1001-1008.
- Chiba, K., Yanagawa, Y., Masubuchi, Y., Kataoka, H., Kawaguchi, T., Ohtsuki, M., and Hoshino, Y. (1998). FTY720, a novel immunosuppressant, induces sequestration of circulating mature lymphocytes by acceleration of lymphocyte homing in rats. I. FTY720 selectively decreases the number of circulating mature lymphocytes by acceleration of lymphocyte homing. *J Immunol* 160, 5037-5044.
 - Christopherson, K. W., 2nd, Campbell, J. J., and Hromas, R. A. (2001). Transgenic overexpression of the CC chemokine CCL21 disrupts T-cell migration. *Blood* 98, 3562-3568.
 - Cinamon G, Shinder V, Alon R. (2001). Shear forces promote lymphocyte migration across vascular endothelium bearing apical chemokines. *Nat Immunol*. 2, 515-22.
 - Claing, A., Laporte, S. A., Caron, M. G., and Lefkowitz, R. J. (2002). Endocytosis of G protein-coupled receptors: roles of G protein-coupled receptor kinases and beta-arrestin proteins. *Prog Neurobiol* 66, 61-79.
 - Clapham, D. E., and Neer, E. J. (1993). New roles for G-protein beta gamma-dimers in transmembrane signalling. *Nature* 365, 403-406.
 - Clark, E. A., Golub, T. R., Lander, E. S., and Hynes, R. O. (2000). Genomic analysis of metastasis reveals an essential role for RhoC. *Nature* 406, 532-535.
 - Cocchi, F., DeVico, A. L., Garzino-Demo, A., Arya, S. K., Gallo, R. C., and Lusso, P. (1995). Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV- suppressive factors produced by CD8+ T cells. *Science* 270, 1811-1815.
 - Collins, K. A., and White, W. L. (1995). Intercellular adhesion molecule 1 (ICAM-1) and bcl-2 are differentially expressed in early evolving malignant melanoma. *Am J Dermatopathol* 17, 429-438.
 - Constantin, G., Majeed, M., Giagulli, C., Piccio, L., Kim, J. Y., Butcher, E. C., and Laudanna, C. (2000). Chemokines trigger immediate beta2 integrin affinity and mobility changes: differential regulation and roles in lymphocyte arrest under flow. *Immunity* 13, 759-769.
 - Cyster, J. G., and Goodnow, C. C. (1995). Pertussis toxin inhibits migration of B and T lymphocytes into splenic white pulp cords. *J Exp Med* 182, 581-586.
 - D'Apuzzo, M., Rolink, A., Loetscher, M., Hoxie, J. A., Clark Lewis, I., Melchers, F., Bagliolini, M., and Moser, B. (1997). The chemokine SDF-1, stromal cell-derived factor 1, attracts early stage B cell precursors via the chemokine receptor CXCR4. *Eur. J. Immunol.* 27, 1788-1793.
 - Daaka, Y., Luttrell, L. M., Ahn, S., Della Rocca, G. J., Ferguson, S. S., Caron, M. G., and Lefkowitz, R. J. (1998). Essential role for G protein-coupled receptor endocytosis in the activation of mitogen-activated protein kinase. *J. Biol. Chem.* 273, 685-688.
 - Dalal, B. I., Keown, P. A., and Greenberg, A. H. (1993). Immunocytochemical localization of secreted transforming growth factor- beta 1 to the advancing edges of primary tumors and to lymph node metastases of human mammary carcinoma. *Am J Pathol* 143, 381-389.
 - De Larco, J. E., Wuertz, B. R., Rosner, K. A., Erickson, S. A., Gamache, D. E., Manivel, J. C., and Furcht, L. T. (2001). A potential role for interleukin-8 in the metastatic phenotype of breast carcinoma cells. *Am J Pathol* 158, 639-646.
 - Debes, G. F., Höpken, U. E., and Hamann, A. (2002). In vivo differentiated cytokine-producing CD4(+) T cells express functional CCR7. *J Immunol* 168, 5441-5447.
 - Deng, H., Liu, R., Ellmeier, W., Choe, S., Unutmaz, D., Burkhardt, M., Di Marzio, P., Marmon, S., Sutton, R. E., Hill, C. M., et al. (1996). Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381, 661-666.
 - Dhawan, P., and Richmond, A. (2002). Role of CXCL1 in tumorigenesis of melanoma. *J Leukoc Biol* 72, 9-18.
 - Dixon, W. T., Demetrick, D. J., Ohyama, K., Sikora, L. K., and Jerry, L. M. (1990). Biosynthesis, glycosylation and intracellular processing of the neuroglandular antigen, a human melanoma-associated antigen. *Cancer Res* 50, 4557-4565.
 - Dobner, T., Wolf, I., Emrich, T., and Lipp, M. (1992). Differentiation-specific expression of a novel G protein-coupled receptor from Burkitt's lymphoma. *Eur J Immunol* 22, 2795-2799.
 - Dohlman, H. G., and Thorner, J. (1997). RGS proteins and signaling by heterotrimeric G

- proteins. *J. Biol. Chem.* 272, 3871-3874.
- Dragic, T., Litwin, V., Allaway, G. P., Martin, S. R., Huang, Y., Nagashima, K. A., Cayanan, C., Madden, P. J., Koup, R. A., Moore, J. P., and Paxton, W. A. (1996). HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC- CKR-5. *Nature* 381, 667-673.
 - Ebrahimzadeh, P. R., Hogfors, C., and Braide, M. (2000). Neutrophil chemotaxis in moving gradients of fMLP. *J Leukoc Biol* 67, 651-661.
 - Eckly-Michel, A. E., Le Bec, A., and Lugnier, C. (1997). Chelerythrine, a protein kinase C inhibitor, interacts with cyclic nucleotide phosphodiesterases. *Eur J Pharmacol* 324, 85-88.
 - Emrich, T., Förster, R., and Lipp, M. (1994). Transmembrane topology of the lymphocyte-specific G-protein-coupled receptor BLR1: analysis by flow cytometry and immunocytochemistry. *Cell Mol Biol* 40, 413-419.
 - Engers R, Gabbert HE. (2000). Mechanisms of tumor metastasis: cell biological aspects and clinical implications. *J Cancer Res Clin Oncol* 126, 682-92.
 - Feil, C., and Augustin, H. G. (1998). Endothelial cells differentially express functional CXC-chemokine receptor-4 (CXCR-4/fusin) under the control of autocrine activity and exogenous cytokines. *Biochem Biophys Res Commun* 247, 38-45.
 - Finke D, Acha-Orbea H, Mattis A, Lipp M, Krahenbuhl J. (2002). CD4+CD3- cells induce Peyer's patch development: role of alpha4beta1 integrin activation by CXCR5. *Immunity* 17, 363-73
 - Förster, R., Emrich, T., Kremmer, E., and Lipp, M. (1994). Expression of the G-protein-coupled receptor BLR1 defines mature, recirculating B cells and a subset of T-helper memory cells. *Blood* 84, 830-840.
 - Förster, R., Kremmer E, Schubel A, Breitfeld D, Kleinschmidt A, Nerl C, Bernhardt G, Lipp M. (1998). Intracellular and surface expression of the HIV-1 coreceptor CXCR4/fusin on various leukocyte subsets: rapid internalization and recycling upon activation. *J Immunol.* 160, 1522-1531.
 - Förster, R., Mattis, A. E., Kremmer, E., Wolf, E., Brem, G., and Lipp, M. (1996). A putative chemokine receptor, BLR1, directs B cell migration to defined lymphoid organs and specific anatomic compartments of the spleen. *Cell* 87, 1037-1047.
 - Förster, R., Schubel, A., Breitfeld, D., Kremmer, E., Renner-Muller, I., Wolf, E., and Lipp, M. (1999). CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* 99, 23-33.
 - Folkman, J. (1992). The role of angiogenesis in tumor growth. *Semin Cancer Biol* 3, 65-71.
 - Foord, S. M., and Marshall, F. H. (1999). RAMPs: accessory proteins for seven transmembrane domain receptors. *Trends Pharmacol Sci* 20, 184-187.
 - Foxman, E. F., Campbell, J. J., and Butcher, E. C. (1997). Multistep navigation and the combinatorial control of leukocyte chemotaxis. *J Cell Biol* 139, 1349-1360.
 - Fitzsimmons D, Hagman J. (1996). Regulation of gene expression at early stages of B-cell and T-cell differentiation. *Curr Opin Immunol* 8, 166-174.
 - Frixen, U. H., Behrens, J., Sachs, M., Eberle, G., Voss, B., Warda, A., Lochner, D., and Birchmeier, W. (1991). E-cadherin-mediated cell-cell adhesion prevents invasiveness of human carcinoma cells. *J Cell Biol* 113, 173-185.
 - Fujisawa N, Sakao Y, Hayashi S, Hadden WA 3rd, Harmon CL, Miller EJ. (2000). alpha-Chemokine growth factors for adenocarcinomas; a synthetic peptide inhibitor for alpha-chemokines inhibits the growth of adenocarcinoma cell lines. *J Cancer Res Clin Oncol* 126 (1), 19-26
 - Ganju, R. K., Brubaker, S. A., Meyer, J., Dutt, P., Yang, Y., Qin, S., Newman, W., and Groopman, J. E. (1998). The alpha-chemokine, stromal cell-derived factor-1alpha, binds to the transmembrane G-protein-coupled CXCR-4 receptor and activates multiple signal transduction pathways. *J Biol Chem* 273, 23169-23175.
 - Ganser, A., Carlo-Stella, C., Bartram, C. R., Boehm, T., Heil, G., Henglein, B., Muller, H., Raghavachar, A., von Briesen, H., Griesinger, F., and et al. (1988). Establishment of two Epstein-Barr virus negative Burkitt cell lines from a patient with AIDS and B-cell lymphoma. *Blood* 72, 1255-1260.
 - Geran, R. I., Greenberg, N. H., Macdonald, M. M., and Abbott, B. J. (1977). Modified

- protocol for the testing of new synthetics in the L1210 lymphoid leukemia murine model in the DR&D program, DCT, NCI. *Natl Cancer Inst Monogr* 45, 151-153.
- Gerard, N. P., and Gerard, C. (1991). The chemotactic receptor for human C5a anaphylatoxin. *Nature* 349, 614-617.
 - Gerhardt CC, Romero IA, Cancello R, Camoin L, Strosberg AD. (2001). Chemokines control fat accumulation and leptin secretion by cultured human adipocytes. *Mol Cell Endocrinol* 175, 81-92.
 - Gershenwald, J. E., and Fidler, I. J. (2002). Cancer. Targeting lymphatic metastasis. *Science* 296, 1811-1812.
 - Girard, J. P., and Springer, T. A. (1995a). Cloning from purified high endothelial venules cells of hevin, a close relative of the antiadhesive extracellular matrix protein SPARC. *Immunity* 2, 113-123.
 - Girard, J. P., and Springer, T. A. (1995b). High endothelial venules (HEVs): specialized endothelium for lymphocyte migration. *Immunol. Today* 16, 449-457.
 - Godessart, N., and Kunkel, S. L. (2001). Chemokines in autoimmune disease. *Curr Opin Immunol* 13, 670-675.
 - Gordge, P. C., Hulme, M. J., Clegg, R. A., and Miller, W. R. (1996). Elevation of protein kinase A and protein kinase C activities in malignant as compared with normal human breast tissue. *Eur J Cancer* 32A, 2120-2126.
 - Gosling, J., Dairaghi, D. J., Wang, Y., Hanley, M., Talbot, D., Miao, Z., and Schall, T. J. (2000). Cutting edge: identification of a novel chemokine receptor that binds dendritic cell- and T cell-active chemokines including ELC, SLC, and TECK. *J Immunol* 164, 2851-2856.
 - Gräler, M. H., Bernhardt, G., and Lipp, M. (1998). EDG6, a novel G-protein-coupled receptor related to receptors for bioactive lysophospholipids, is specifically expressed in lymphoid tissue. *Genomics* 53, 164-169.
 - Green, A. R., Green, V. L., White, M. C., and Speirs, V. (1997). Expression of cytokine messenger RNA in normal and neoplastic human breast tissue: identification of interleukin-8 as a potential regulatory factor in breast tumours. *Int J Cancer* 72, 937-941.
 - Gunn, M. D., Kyuwa, S., Tam, C., Kakiuchi, T., Matsuzawa, A., Williams, L. T., and Nakano, H. (1999). Mice lacking expression of secondary lymphoid organ chemokine have defects in lymphocyte homing and dendritic cell localization. *J Exp Med* 189, 451-460.
 - Gunn, M. D., Tangemann, K., Tam, C., Cyster, J. G., Rosen, S. D., and Williams, L. T. (1998). A chemokine expressed in lymphoid high endothelial venules promotes the adhesion and chemotaxis of naive T lymphocytes. *Proc Natl Acad Sci USA* 95, 258-263.
 - Haribabu, B., Richardson, R. M., Fisher, I., Sozzani, S., Peiper, S. C., Horuk, R., Ali, H., and Snyderman, R. (1997). Regulation of human chemokine receptors CXCR4. Role of phosphorylation in desensitization and internalization. *J Biol Chem* 272, 28726-28731.
 - Hartl, P., and Lipp, M. (1987). Generation of a variant t(2;8) translocation of Burkitt's lymphoma by site-specific recombination via the kappa light-chain joining signals. *Mol Cell Biol* 7, 2037-2045.
 - Hasegawa, H., Nomura, T., Kohno, M., Tateishi, N., Suzuki, Y., Maeda, N., Fujisawa, R., Yoshie, O., and Fujita, S. (2000). Increased chemokine receptor CCR7/EBI1 expression enhances the infiltration of lymphoid organs by adult T-cell leukemia cells. *Blood* 95, 30-38.
 - Hasegawa, H., Utsunomiya, Y., Yasukawa, M., Yanagisawa, K., and Fujita, S. (1994). Induction of G protein-coupled peptide receptor EBI 1 by human herpesvirus 6 and 7 infection in CD4+ T cells. *J Virol* 68, 5326-5329.
 - Hay, E. D. (1995). An overview of epithelial-mesenchymal transformation. *Acta Anat* 154, 8-20.
 - Hendrix, M. J., Seftor, E. A., Chu, Y. W., Trevor, K. T., and Seftor, R. E. (1996). Role of intermediate filaments in migration, invasion and metastasis. *Cancer Metastasis Rev* 15, 507-525.
 - Henning, G., Ohl, L., Junt, T., Reiterer, P., Brinkmann, V., Nakano, H., Hohenberger, W., Lipp, M., and Förster, R. (2001). CC chemokine receptor 7-dependent and -independent pathways for lymphocyte homing: modulation by FTY720. *J Exp Med* 194, 1875-1881.
 - Hepler, J. R., Berman, D. M., Gilman, A. G., and Kozasa, T. (1997). RGS4 and GAIP are

- GTPase-activating proteins for Gq alpha and block activation of phospholipase C beta by gamma-thio-GTP-Gq alpha. Proc. Natl. Acad. Sci. USA 94, 428-432.
- Hepler, J. R., and Gilman, A. G. (1992). G proteins. Trends Biochem Sci 17, 383-387.
 - Herbert, J. M., Augereau, J. M., Gleye, J., and Maffrand, J. P. (1990). Chelerythrine is a potent and specific inhibitor of protein kinase C. Biochem Biophys Res Commun 172, 993-999.
 - Herbert, J. M., Savi, P., Laplace, M. C., Dumas, A., and Dol, F. (1993). Chelerythrine, a selective protein kinase C inhibitor, counteracts pyrogen-induced expression of tissue factor without effect on thrombomodulin down-regulation in endothelial cells. Thromb Res 71, 487-493.
 - Hirao, M., Onai, N., Hiroishi, K., Watkins, S. C., Matsushima, K., Robbins, P. D., Lotze, M. T., and Tahara, H. (2000). CC chemokine receptor-7 on dendritic cells is induced after interaction with apoptotic tumor cells: critical role in migration from the tumor site to draining lymph nodes. Cancer Res 60, 2209-2217.
 - Hirose, J., Kawashima, H., Swope Willis, M., Springer, T. A., Hasegawa, H., Yoshie, O., and Miyasaka, M. (2002). Chondroitin sulfate B exerts its inhibitory effect on secondary lymphoid tissue chemokine (SLC) by binding to the C-terminus of SLC. Biochim Biophys Acta 1571, 219-224.
 - Hirose, J., Kawashima, H., Yoshie, O., Tashiro, K., and Miyasaka, M. (2001). Versican interacts with chemokines and modulates cellular responses. J Biol Chem 276, 5228-5234.
 - Hjelmström, P. (2001). Lymphoid neogenesis: de novo formation of lymphoid tissue in chronic inflammation through expression of homing chemokines. J Leukoc Biol 69, 331-339.
 - Honda, K., Nakano, H., Yoshida, H., Nishikawa, S., Rennert, P., Ikuta, K., Tamechika, M., Yamaguchi, K., Fukumoto, T., Chiba, T., and Nishikawa, S. I. (2001). Molecular basis for hematopoietic/mesenchymal interaction during initiation of Peyer's patch organogenesis. J Exp Med 193, 621-630.
 - Höpken, U. E., Foss, H. D., Meyer, D., Hinz, M., Leder, K., Stein, H., and Lipp, M. (2002). Up-regulation of the chemokine receptor CCR7 in classical but not in lymphocyte-predominant Hodgkin disease correlates with distinct dissemination of neoplastic cells in lymphoid organs. Blood 99, 1109-1116.
 - Huang, S., Xie, K., Bucana, C. D., Ullrich, S. E., and Bar-Eli, M. (1996). Interleukin 10 suppresses tumor growth and metastasis of human melanoma cells: potential inhibition of angiogenesis. Clin Cancer Res 2, 1969-1979.
 - Iyengar, R. (1993). Molecular and functional diversity of mammalian Gs-stimulated adenylyl cyclases. Faseb J 7, 768-775.
 - Janatpour MJ, Hudak S, Sathe M, Sedgwick JD, McEvoy LM. (2001). Tumor necrosis factor-dependent segmental control of MIG expression by high endothelial venules in inflamed lymph nodes regulates monocyte recruitment. J Exp Med 194, 1375-84.
 - Jarzabek, K., Laudanski, P., Dzieciol, J., Dabrowska, M., and Wolczynski, S. (2002). Protein kinase C involvement in proliferation and survival of breast cancer cells. Folia Histochem Cytobiol 40, 193-194.
 - Jenh, C. H., Cox, M. A., Kaminski, H., Zhang, M., Byrnes, H., Fine, J., Lundell, D., Chou, C. C., Narula, S. K., and Zavodny, P. J. (1999). Cutting edge: species specificity of the CC chemokine 6Ckine signaling through the CXC chemokine receptor CXCR3: human 6Ckine is not a ligand for the human or mouse CXCR3 receptors. J Immunol 162, 3765-3769.
 - Johnson, J. P. (1991). The role of ICAM-1 in tumor development. Chem Immunol 50, 143-163.
 - Kahan, B. D. (1998). FTY720: a new immunosuppressive agent with novel mechanism(s) of action. Transplant Proc 30, 2210-2213.
 - Kaiser, E., Förster, R., Wolf, I., Ebensperger, C., Kuehl, W. M., and Lipp, M. (1993). The G protein-coupled receptor BLR1 is involved in murine B cell differentiation and is also expressed in neuronal tissues. Eur J Immunol 23, 2532-2539.
 - Karkkainen, M. J., Jussila, L., Ferrell, R. E., Finegold, D. N., and Alitalo, K. (2001). Molecular regulation of lymphangiogenesis and targets for tissue oedema. Trends Mol Med 7, 18-22.
 - Karkkainen MJ, Alitalo K. Lymphatic endothelial regulation, lymphoedema, and lymph node

- metastasis. (2002). *Semin Cell Dev Biol* 13, 9-18.
- Karpanen, T., Egeblad, M., Karkkainen, M. J., Kubo, H., Yla-Herttula, S., Jaattela, M., and Alitalo, K. (2001). Vascular endothelial growth factor C promotes tumor lymphangiogenesis and intralymphatic tumor growth. *Cancer Res* 61, 1786-1790.
 - Karpus, W. J., Lukacs, N. W., Kennedy, K. J., Smith, W. S., Hurst, S. D., and Barrett, T. A. (1997). Differential CC chemokine-induced enhancement of T helper cell cytokine production. *J Immunol* 158, 4129-4136.
 - Keane, M. P., Arenberg, D. A., Lynch, J. P., 3rd, Whyte, R. I., Iannettoni, M. D., Burdick, M. D., Wilke, C. A., Morris, S. B., Glass, M. C., DiGiovine, B., et al. (1997). The CXC chemokines, IL-8 and IP-10, regulate angiogenic activity in idiopathic pulmonary fibrosis. *J Immunol* 159, 1437-1443.
 - Kim CH, Broxmeyer HE. (1999). Chemokines: signal lamps for trafficking of T and B cells for development and effector function. *J Leukoc Biol* 65, 6-15
 - Kim, C. H., Pelus, L. M., White, J. R., Applebaum, E., Johanson, K., and Broxmeyer, H. E. (1998). CK beta-11/macrophage inflammatory protein-3 beta/EBI1-ligand chemokine is an efficacious chemoattractant for T and B cells. *J Immunol* 160, 2418-2424.
 - Kinzler, K. W., and Vogelstein, B. (1996). Lessons from hereditary colorectal cancer. *Cell* 87, 159-170.
 - Kirschmann, D. A., Seftor, E. A., Nieva, D. R., Mariano, E. A., and Hendrix, M. J. (1999). Differentially expressed genes associated with the metastatic phenotype in breast cancer. *Breast Cancer Res Treat* 55, 127-136.
 - Kligman, A. M., and Kligman, L. H. (1994). Carcinogens show comedogenic activity: a potential animal screen for tumorigenic substances. *Cancer Lett* 87, 171-178.
 - Knudson, A. G. (1993). Antioncogenes and human cancer. *Proc Natl Acad Sci USA* 90, 10914-10921.
 - Koch, A. E., Halloran, M. M., Haskell, C. J., Shah, M. R., and Polverini, P. J. (1995). Angiogenesis mediated by soluble forms of E-selectin and vascular cell adhesion molecule-1. *Nature* 376, 517-519.
 - Koch, W. J., Hawes, B. E., Allen, L. F., and Lefkowitz, R. J. (1994). Direct evidence that Gi-coupled receptor stimulation of mitogen-activated protein kinase is mediated by G beta gamma activation of p21ras. *Proc Natl Acad Sci USA* 91, 12706-12710.
 - Kolch, W., Martiny-Baron, G., Kieser, A., and Marme, D. (1995). Regulation of the expression of the VEGF/VPS and its receptors: role in tumor angiogenesis. *Breast Cancer Res Treat* 36, 139-155.
 - König, J. A., and Edwardson, J. M. (1997). Endocytosis and recycling of G protein-coupled receptors. *Trends Pharmacol. Sci.* 18, 276-287.
 - Krasel, C., Dammeier, S., Winstel, R., Brockmann, J., Mischak, H., and Lohse, M. J. (2001). Phosphorylation of GRK2 by protein kinase C abolishes its inhibition by calmodulin. *J Biol Chem* 276, 1911-1915.
 - Kriehuber, E., Breiteneder-Geleff, S., Groeger, M., Soleiman, A., Schoppmann, S. F., Stingl, G., Kerjaschki, D., and Maurer, D. (2001). Isolation and characterization of dermal lymphatic and blood endothelial cells reveal stable and functionally specialized cell lineages. *J Exp Med* 194, 797-808.
 - Kucharzik, T., Lugering, N., Pauels, H. G., Domschke, W., and Stoll, R. (1998). IL-4, IL-10 and IL-13 down-regulate monocyte-chemoattracting protein-1 (MCP-1) production in activated intestinal epithelial cells. *Clin Exp Immunol* 111, 152-157.
 - Kurschat P, Mauch C. (2000). Mechanisms of metastasis. *Clin Exp Dermatol* 25,482-489.
 - Kurschat P, Zigrino P, Nischt R, Breitkopf K, Steurer P, Klein CE, Krieg T, Mauch C. (1999). Tissue inhibitor of matrix metalloproteinase-2 regulates matrix metalloproteinase-2 activation by modulation of membrane-type 1 matrix metalloproteinase activity in high and low invasive melanoma cell lines. *J Biol Chem* 274, 1056-1062.
 - Lambertucci, J. R., Rayes, A. A., Serufo, J. C., and Nobre, V. (2001). Pyogenic abscesses and parasitic diseases. *Rev Inst Med Trop Sao Paulo* 43, 67-74.
 - Lange, R., Peng, X., Wimmer, E., Lipp, M., and Bernhardt, G. (2001). The poliovirus receptor CD155 mediates cell-to-matrix contacts by specifically binding to vitronectin. *Virology* 285,

- 218-227.
- LaRosa, G. J., Thomas, K. M., Kaufmann, M. E., Mark, R., White, M., Taylor, L., Gray, G., Witt, D., and Navarro, J. (1992). Amino terminus of the interleukin-8 receptor is a major determinant of receptor subtype specificity. *J Biol Chem* 267, 25402-25406.
 - Le, Y., Wetzel, M. A., Shen, W., Gong, W., Rogers, T. J., Henderson, E. E., and Wang, J. M. (2001). Desensitization of chemokine receptor CCR5 in dendritic cells at the early stage of differentiation by activation of formyl peptide receptors. *Clin Immunol* 99, 365-372.
 - Leek, R. D., Hunt, N. C., Landers, R. J., Lewis, C. E., Royds, J. A., and Harris, A. L. (2000). Macrophage infiltration is associated with VEGF and EGFR expression in breast cancer. *J Pathol* 190, 430-436.
 - Legler, D. F., Loetscher, M., Roos, R. S., Clark-Lewis, I., Baggolini, M., and Moser, B. (1998). B cell-attracting chemokine 1, a human CXC chemokine expressed in lymphoid tissues, selectively attracts B lymphocytes via BLR1/CXCR5. *J Exp Med* 187, 655-660.
 - Lemmon, M. A., Ferguson, K. M., and Schlessinger, J. (1996). PH domains: diverse sequences with a common fold recruit signaling molecules to the cell surface. *Cell* 85, 621-624.
 - Lev, S., Moreno, H., Martinez, R., Canoll, P., Peles, E., Musacchio, J. M., Plowman, G. D., Rudy, B., and Schlessinger, J. (1995). Protein tyrosine kinase PYK2 involved in Ca^{2+} -induced regulation of ion channel and MAP kinase functions. *Nature* 376, 737-745.
 - Li, Q., Vaingankar, S. M., Green, H. M., and Martins-Green, M. (1999). Activation of the 9E3/cCAF chemokine by phorbol esters occurs via multiple signal transduction pathways that converge to MEK1/ERK2 and activate the Elk1 transcription factor. *J Biol Chem* 274, 15454-15465.
 - Limatola, C., Giovannelli, A., Maggi, L., Ragozzino, D., Castellani, L., Ciotti, M. T., Vacca, F., Mercanti, D., Santoni, A., and Eusebi, F. (2000). SDF-1alpha-mediated modulation of synaptic transmission in rat cerebellum. *Eur J Neurosci* 12, 2497-2504.
 - Lin, T. J., Issekutz, T. B., and Marshall, J. S. (2001). SDF-1 induces IL-8 production and transendothelial migration of human cord blood-derived mast cells. *Int Arch Allergy Immunol* 124, 142-145.
 - Lissitzky, J. C., Pourreau-Schneider, N., Kopp, F., Chiarotti, Y., Charpin, C., Jacquemier, J., and Martin, P. M. (1985). Invasion, metastases of solid tumors. Interaction of tumor cells with tissue and vascular basement membranes. *Bull Cancer* 72, 367-376.
 - Loetscher, P., Pellegrino, A., Gong, J. H., Mattioli, I., Loetscher, M., Bardi, G., Baggolini, M., and Clark-Lewis, I. (2001). The ligands of CXC chemokine receptor 3, I-TAC, Mig, and IP10, are natural antagonists for CCR3. *J Biol Chem* 276, 2986-2991.
 - Lowes, V. L., Ip, N. Y., and Wong, Y. H. (2002). Integration of signals from receptor tyrosine kinases and g protein-coupled receptors. *Neurosignals* 11, 5-19.
 - Luster AD. (2002). The role of chemokines in linking innate and adaptive immunity. *Curr Opin Immunol* 14, 129-35
 - Luther, S. A., Bidgol, A., Hargreaves, D. C., Schmidt, A., Xu, Y., Paniyadi, J., Matloubian, M., and Cyster, J. G. (2002). Differing activities of homöostatic chemokines CCL19, CCL21, and CXCL12 in lymphocyte and dendritic cell recruitment and lymphoid neogenesis. *J Immunol* 169, 424-433.
 - Luther, S. A., Lopez, T., Bai, W., Hanahan, D., and Cyster, J. G. (2000a). BLC expression in pancreatic islets causes B cell recruitment and lymphotoxin-dependent lymphoid neogenesis. *Immunity* 12, 471-481.
 - Luther, S. A., Tang, H. L., Hyman, P. L., Farr, A. G., and Cyster, J. G. (2000b). Coexpression of the chemokines ELC and SLC by T zone stromal cells and deletion of the ELC gene in the plt=plt mouse. *Proc Natl Acad Sci USA* 97, 12694-12699.
 - Luttrell, L. M., Ferguson, S. S., Daaka, Y., Miller, W. E., Maudsley, S., Della Rocca, G. J., Lin, F., Kawakatsu, H., Owada, K., Luttrell, D. K., et al. (1999). Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. *Science* 283, 655-661.
 - Luttrell, L. M., Hawes, B. E., van Biesen, T., Luttrell, D. K., Lansing, T. J., and Lefkowitz, R. J. (1996). Role of c-Src tyrosine kinase in G protein-coupled receptor- and Gbetagamma subunit-mediated activation of mitogen-activated protein kinases. *J Biol Chem* 271, 19443-

- 19450.
- Machesky, L. M., and Hall, A. (1997). Role of actin polymerization and adhesion to extracellular matrix in Rac- and Rho-induced cytoskeletal reorganization. *J. Cell Biol.* 138, 913-926.
 - Mandala, S., Hajdu, R., Bergstrom, J., Quackenbush, E., Xie, J., Milligan, J., Thornton, R., Shei, G. J., Card, D., Keohane, C., et al. (2002). Alteration of lymphocyte trafficking by sphingosine-1-phosphate receptor agonists. *Science* 296, 346-349.
 - Mandriota, S. J., Jussila, L., Jeltsch, M., Compagni, A., Baetens, D., Prevo, R., Banerji, S., Huarte, J., Montesano, R., Jackson, D. G., et al. (2001). Vascular endothelial growth factor-C-mediated lymphangiogenesis promotes tumour metastasis. *Embo J* 20, 672-682.
 - Mandriota, S. J., Seghezzi, G., Vassalli, J. D., Ferrara, N., Wasi, S., Mazzieri, R., Mignatti, P., and Pepper, M. S. (1995). Vascular endothelial growth factor increases urokinase receptor expression in vascular endothelial cells. *J Biol Chem* 270, 9709-9716.
 - Mashino, K., Sadanaga, N., Yamaguchi, H., Tanaka, F., Ohta, M., Shibuta, K., Inoue, H., and Mori, M. (2002). Expression of Chemokine Receptor CCR7 Is Associated with Lymph Node Metastasis of Gastric Carcinoma. *Cancer Res* 62, 2937-2941.
 - Mazzucchelli, L., Blaser, A., Kappeler, A., Scharli, P., Laissue, J. A., Baggolini, M., and Uggioni, M. (1999). BCA-1 is highly expressed in Helicobacter pylori-induced mucosa-associated lymphoid tissue and gastric lymphoma. *J Clin Invest* 104, 49-54.
 - Mebius RE, Streeter PR, Michie S, Butcher EC, Weissman IL. (1996). A developmental switch in lymphocyte homing receptor and endothelial vascular addressin expression regulates lymphocyte homing and permits CD4+ CD3- cells to colonize lymph nodes. *Proc Natl Acad Sci* 93, 11019-11024.
 - Meiners, S., Brinkmann, V., Naundorf, H., and Birchmeier, W. (1998). Role of morphogenetic factors in metastasis of mammary carcinoma cells. *Oncogene* 16, 9-20.
 - Mellado, M., Rodriguez-Frade, J. M., Aragay, A., del Real, G., Martin, A. M., Vila-Coro, A. J., Serrano, A., Mayor, F., Jr., and Martinez, A. C. (1998). The chemokine monocyte chemotactic protein 1 triggers Janus kinase 2 activation and tyrosine phosphorylation of the CCR2B receptor. *J Immunol* 161, 805-813.
 - Merrill AH Jr, Schmelz EM, Dillehay DL, Spiegel S, Shayman JA, Schroeder JJ, Riley RT, Voss KA, Wang E. (1997). Sphingolipids--the enigmatic lipid class: biochemistry, physiology, and pathophysiology. *Toxicol Appl Pharmacol* 142, 208-225.
 - Middleton J, Neil S, Wintle J, Clark-Lewis I, Moore H, Lam C, Auer M, Hub E, Rot A. (1997). Transcytosis and surface presentation of IL-8 by venular endothelial cells. *Cell* 91, 385-95.
 - Mitsusada, M., Suzuki, S., Kobayashi, E., Enosawa, S., Kakefuda, T., and Miyata, M. (1997). Prevention of graft rejection and graft-versus-host reaction by a novel immunosuppressant, FTY720, in rat small bowel transplantation. *Transpl Int* 10, 343-349.
 - Mogi, A., Hatai, M., Soga, H., Takenoshita, S., Nagamachi, Y., Fujimoto, J., Yamamoto, T., Yokota, J., and Yaoi, Y. (1995). Possible role of protein kinase C in the regulation of intracellular stability of focal adhesion kinase in mouse 3T3 cells. *FEBS Lett* 373, 135-140.
 - Mokros, T., Rehm, A., Droese, J., Oppermann, M., Lipp, M., and Hopken, U. E. (2002). Surface expression and endocytosis of the human cytomegalovirus-encoded chemokine receptor US28 is regulated by agonist-independent phosphorylation. *J Biol Chem* 227, 45122-8.
 - Molino M, Woolkalis MJ, Prevost N, Pratico D, Barnathan ES, Taraboletti G, Haggarty BS, Hesselgesser J, Horuk, R, Hoxie JA, Brass LF. (2000) CXCR4 on human endothelial cells can serve as both a mediator of biological responses and as a receptor for HIV-2. *Biochim Biophys Acta* 1500: 227-40.
 - Montruccchio, G., Lupia, E., Battaglia, E., Del Sorbo, L., Boccellino, M., Biancone, L., Emanuelli, G., and Camussi, G. (2000). Platelet-activating factor enhances vascular endothelial growth factor- induced endothelial cell motility and neoangiogenesis in a murine matrigel model. *Arterioscler Thromb Vasc Biol* 20, 80-88.
 - Moore, B. B., Keane, M. P., Addison, C. L., Arenberg, D. A., and Strieter, R. M. (1998). CXC chemokine modulation of angiogenesis: the importance of balance between angiogenic and

- angiostatic members of the family. *J. Investig. Med.* 46, 113-120.
- Moore, M. A. (2001). The role of chemoattraction in cancer metastases. *Bioessays* 23, 674-676.
 - Mori, S., Nakano, H., Aritomi, K., Wang, C. R., Gunn, M. D., and Kakiuchi, T. (2001). Mice lacking expression of the chemokines CCL21-ser and CCL19 (plt mice) demonstrate delayed but enhanced T cell immune responses. *J Exp Med* 193, 207-218.
 - Müller, A., Homey, B., Soto, H., Ge, N., Catron, D., Buchanan, M. E., McClanahan, T., Murphy, E., Yuan, W., Wagner, S. N., et al. (2001). Involvement of chemokine receptors in breast cancer metastasis. *Nature* 410, 50-56.
 - Müller, G., and Lipp, M. (2001). Signal transduction by the chemokine receptor CXCR5: structural requirements for G protein activation analyzed by chimeric CXCR1/CXCR5 molecules. *Biol Chem* 382, 1387-1397.
 - Gerd Müller; Philipp Reiterer; Uta Höpken, Sven Golfier und Martin Lipp (2003) Role of Homeostatic Chemokine and Sphingosine-1-Phosphate Receptors in the Organization of Lymphoid Tissue. *Annals of the New York Academy of Sciences* 987, 107-116.
 - Muraki, K., and Imaizumi, Y. (2001). A novel function of sphingosine-1-phosphate to activate a non-selective cation channel in human endothelial cells. *J Physiol* 537, 431-441.
 - Murakami T, Maki W, Cardones AR, Fang H, Tun Kyi A, Nestle FO, Hwang ST. (2002). Expression of CXC chemokine receptor-4 enhances the pulmonary metastatic potential of murine B16 melanoma cells. *Cancer Res* 62: 7328-34.
 - Murphy, P. M. (1994). The molecular biology of leukocyte chemoattractant receptors. *Annu Rev Immunol* 12, 593-633.
 - Muta, H., Boise, L. H., Fang, L., and Podack, E. R. (2000). CD30 signals integrate expression of cytotoxic effector molecules, lymphocyte trafficking signals, and signals for proliferation and apoptosis. *J Immunol* 165, 5105-5111.
 - Nagasawa, T., Hirota, S., Tachibana, K., Takakura, N., Nishikawa, S., Kitamura, Y., Yoshida, N., Kikutani, H., and Kishimoto, T. (1996). Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* 382, 635-638.
 - Nagira, M., Imai, T., Hieshima, K., Kusuda, J., Ridanpaa, M., Takagi, S., Nishimura, M., Kakizaki, M., Nomiyama, H., and Yoshie, O. (1997). Molecular cloning of a novel human CC chemokine secondary lymphoid-tissue chemokine that is a potent chemoattractant for lymphocytes and mapped to chromosome 9p13. *J Biol Chem* 272, 19518-19524.
 - Nakano, H., and Gunn, M. D. (2001). Gene duplications at the chemokine locus on mouse chromosome 4: multiple strain-specific haplotypes and the deletion of secondary lymphoid-organ chemokine and EBI-1 ligand chemokine genes in the plt mutation. *J Immunol* 166, 361-369.
 - Natali, P. G., Hamby, C. V., Felding-Habermann, B., Liang, B., Nicotra, M. R., Di Filippo, F., Giannarelli, D., Temponi, M., and Ferrone, S. (1997). Clinical significance of alpha(v)beta3 integrin and intercellular adhesion molecule-1 expression in cutaneous malignant melanoma lesions. *Cancer Res* 57, 1554-1560.
 - Neote, K., DiGregorio, D., Mak, J. Y., Horuk, R., and Schall, T. J. (1993). Molecular cloning, functional expression, and signaling characteristics of a C-C chemokine receptor. *Cell* 72, 415-425.
 - Neptune, E. R., Iiri, T., and Bourne, H. R. (1999). Galphai is not required for chemotaxis mediated by Gi-coupled receptors. *J Biol Chem* 274, 2824-2828.
 - Newman, P. J., and Albelda, S. M. (1992). Cellular and molecular aspects of PECAM-1. *Nouv Rev Fr Hematol* 34, S9-13.
 - Ngo, V. N., Korner, H., Gunn, M. D., Schmidt, K. N., Riminton, D. S., Cooper, M. D., Browning, J. L., Sedgwick, J. D., and Cyster, J. G. (1999). Lymphotoxin alpha/beta and tumor necrosis factor are required for stromal cell expression of homing chemokines in B and T cell areas of the spleen. *J Exp Med* 189, 403-412.
 - Ngo, V. N., Tang, H. L., and Cyster, J. G. (1998). Epstein-Barr virus-induced molecule 1 ligand chemokine is expressed by dendritic cells in lymphoid tissues and strongly attracts naive T cells and activated B cells. *J Exp Med* 188, 181-191.
 - Nobes, C. D., and Hall, A. (1995). Rho, rac, and cdc42 GTPases regulate the assembly of

- multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* 81, 53-62.
- Nomura, T., and Hasegawa, H. (2000). Chemokines and anti-cancer immunotherapy: anti-tumor effect of EBI1- ligand chemokine (ELC) and secondary lymphoid tissue chemokine (SLC). *Anticancer Res* 20, 4073-4080.
 - Nomura T, Hasegawa H, Kohno M, Sasaki M, Fujita S. (2001). Enhancement of anti-tumor immunity by tumor cells transfected with the secondary lymphoid tissue chemokine EBI-1-ligand chemokine and stromal cell-derived factor-1alpha chemokine genes. *Int J Cancer*. 91, 597-606.
 - Oberlin, E., Amara, A., Bachelerie, F., Bessia, C., Virelizier, J. L., Arenzana Seisdedos, F., Schwartz, O., Heard, J. M., Clark Lewis, I., Legler, D. F., et al. (1996). The CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-adapted HIV-1. *Nature* 382, 833-835.
 - Ogata, M., Zhang, Y., Wang, Y., Itakura, M., Zhang, Y. Y., Harada, A., Hashimoto, S., and Matsushima, K. (1999). Chemotactic response toward chemokines and its regulation by transforming growth factor-beta1 of murine bone marrow hematopoietic progenitor cell-derived different subset of dendritic cells. *Blood* 93, 3225-3232.
 - Ogawa, Y., Hirakawa, K., Nakata, B., Fujihara, T., Sawada, T., Kato, Y., Yoshikawa, K., and Sowa, M. (1998). Expression of intercellular adhesion molecule-1 in invasive breast cancer reflects low growth potential, negative lymph node involvement, and good prognosis. *Clin Cancer Res* 4, 31-36.
 - Okada, T., Ngo, V. N., Ekland, E. H., Förster, R., Lipp, M., Littman, D. R., and Cyster, J. G. (2002). Chemokine Requirements for B Cell Entry to Lymph Nodes and Peyer's Patches. *J Exp Med* 196, 65-75.
 - Olson, T. S., and Ley, K. (2002). Chemokines and chemokine receptors in leukocyte trafficking. *Am J Physiol Regul Integr Comp Physiol* 283, 7-28.
 - Oppenheim, J. J., Zachariae, C. O., Mukaida, N., and Matsushima, K. (1991). Properties of the novel proinflammatory supergene "intercrine" cytokine family. *Annu. Rev. Immunol.* 9, 617-648.
 - Oppermann, M., Mack, M., Proudfoot, A. E., and Olbrich, H. (1999). Differential effects of CC chemokines on CC chemokine receptor 5 (CCR5) phosphorylation and identification of phosphorylation sites on the CCR5 carboxyl terminus. *J Biol Chem* 274, 8875-8885.
 - Orsini, M. J., Parent, J. L., Mundell, S. J., Benovic, J. L., and Marchese, A. (1999). Trafficking of the HIV coreceptor CXCR4. Role of arrestins and identification of residues in the c-terminal tail that mediate receptor internalization. *J Biol Chem* 274, 31076-31086.
 - Ohshima K, Tutiya T, Yamaguchi T, Suzuki K, Suzumiya J, Kawasaki C, Haraoka S, Kikuchi M. (2002). Infiltration of Th1 and Th2 lymphocytes around Hodgkin and Reed-Sternberg (H&RS) cells in Hodgkin disease: Relation with expression of CXC and CC chemokines on H&RS cells. *Int J Cancer* 98, 567-72
 - Pachynski, R. K., Wu, S. W., Gunn, M. D., and Erle, D. J. (1998). Secondary lymphoid-tissue chemokine (SLC) stimulates integrin alpha 4 beta 7-mediated adhesion of lymphocytes to mucosal addressin cell adhesion molecule-1 (MAdCAM-1) under flow. *J Immunol* 161, 952-956.
 - Pages C, Daviaud D, An S, Krief S, Lafontan M, Valet P, Saulnier-Blache JS. (2001). Endothelial differentiation gene-2 receptor is involved in lysophosphatidic acid-dependent control of 3T3F442A preadipocyte proliferation and spreading. *J Biol Chem* 276, 11599-605.
 - Paik, J. H., Chae, S., Lee, M. J., Thangada, S., and Hla, T. (2001). Sphingosine 1-phosphate-induced endothelial cell migration requires the expression of EDG-1 and EDG-3 receptors and Rho-dependent activation of alpha vbeta3- and beta1-containing integrins. *J Biol Chem* 276, 11830-11837.
 - Palczewski, K. (1997). GTP-binding-protein-coupled receptor kinases-two mechanistic models. *Eur J Biochem* 248, 261-269.
 - Patel, D. D., Koopmann, W., Imai, T., Whichard, L. P., Yoshie, O., and Krangel, M. S. (2001). Chemokines have diverse abilities to form solid phase gradients. *Clin Immunol* 99, 43-52.

- Payne, A. S., and Cornelius, L. A. (2002). The role of chemokines in melanoma tumor growth and metastasis. *J Invest Dermatol* 118, 915-922.
- Phillips, R., and Ager, A. (2002). Activation of pertussis toxin-sensitive CXCL12 (SDF-1) receptors mediates transendothelial migration of T lymphocytes across lymph node high endothelial cells. *Eur J Immunol* 32, 837-847.
- Piali, L., Weber, C., LaRosa, G., Mackay, C. R., Springer, T. A., Clark-Lewis, I., and Moser, B. (1998). The chemokine receptor CXCR3 mediates rapid and shear-resistant adhesion-induction of effector T lymphocytes by the chemokines IP10 and Mig. *Eur J Immunol* 28, 961-972.
- Picker, L. J., and Butcher, E. C. (1992). Physiological and molecular mechanisms of lymphocyte homing. *Annu Rev Immunol* 10, 561-591.
- Pinschewer, D. D., Ochsenbein, A. F., Odermatt, B., Brinkmann, V., Hengartner, H., and Zinkernagel, R. M. (2000). FTY720 immunosuppression impairs effector T cell peripheral homing without affecting induction, expansion, and memory. *J Immunol* 164, 5761-5770.
- Pitcher, J. A., Freedman, N. J., and Lefkowitz, R. J. (1998). G protein-coupled receptor kinases. *Annu Rev Biochem* 67, 653-692.
- Ploix, C., Lo, D., and Carson, M. J. (2001). A ligand for the chemokine receptor CCR7 can influence the homoeostatic proliferation of CD4 T cells and progression of autoimmunity. *J Immunol* 167, 6724-6730.
- Probst, W. C., Snyder, L. A., Schuster, D. I., Brosius, J., and Sealfon, S. C. (1992). Sequence alignment of the G-protein coupled receptor superfamily. *DNA Cell Biol.* 11, 1-20.
- Pronin, A. N., and Benovic, J. L. (1997). Regulation of the G protein-coupled receptor kinase GRK5 by protein kinase C. *J Biol Chem* 272, 3806-3812.
- Rabinovich, G. A., Rubinstein, N., and Toscano, M. A. (2002). Role of galectins in inflammatory and immunomodulatory processes. *Biochim Biophys Acta* 1572, 274-284.
- Rainard, P. (2002). Bovine milk fat globules do not inhibit C5a chemotactic activity. *Vet Res* 33, 413-419.
- Rane, S. G., and Reddy, E. P. (2000). Janus kinases: components of multiple signaling pathways. *Oncogene* 19, 5662-5679.
- Rane, S. G., and Reddy, E. P. (2002). JAKs, STATs and Src kinases in hematopoiesis. *Oncogene* 21, 3334-3358.
- Rappert, A., Biber, K., Nolte, C., Lipp, M., Schubel, A., Lu, B., Gerard, N. P., Gerard, C., Boddeke, H. W., and Kettenmann, H. (2002). Secondary lymphoid tissue chemokine (CCL21) activates CXCR3 to trigger a Cl⁻ current and chemotaxis in murine microglia. *J Immunol* 168, 3221-3226.
- Regidor, P. A., Callies, R., Regidor, M., and Schindler, A. E. (1998). Expression of the cell adhesion molecules ICAM-1 and VCAM-1 in the cytosol of breast cancer tissue, benign breast tissue and corresponding sera. *Eur J Gynaecol Oncol* 19, 377-383.
- Reif, K., and Cyster, J. G. (2000). RGS molecule expression in murine B lymphocytes and ability to down-regulate chemotaxis to lymphoid chemokines. *J Immunol* 164, 4720-4729.
- Reif, K., Ekland, E. H., Ohl, L., Nakano, H., Lipp, M., Förster, R., and Cyster, J. G. (2002). Balanced responsiveness to chemoattractants from adjacent zones determines B-cell position. *Nature* 416, 94-99.
- Reuning U, Magdolen V, Wilhelm O, Fischer K, Lutz V, Graeff H, Schmitt M. (1998). Multifunctional potential of the plasminogen activation system in tumor invasion and metastasis (review). *Int J Oncol* 13, 893-906.
- Richardson, R. M., DuBose, R. A., Ali, H., Tomhave, E. D., Haribabu, B., and Snyderman, R. (1995). Regulation of human interleukin-8 receptor A: identification of a phosphorylation site involved in modulating receptor functions. *Biochemistry* 34, 14193-14201.
- Richardson, R. M., Ali, H., Pridgen, B. C., Haribabu, B., and Snyderman, R. (1998). Multiple signaling pathways of human interleukin-8 receptor A. Independent regulation by phosphorylation. *In J. Biol. Chem.* 273, 10690-10695.
- Ridley, A. J., Paterson, H. F., Johnston, C. L., Diekmann, D., and Hall, A. (1992). The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell* 70, 401-410.

- Robinson, S., Cohen, M., Prayson, R., Ransohoff, R. M., Tabrizi, N., and Miller, R. H. (2001). Constitutive expression of growth-related oncogene and its receptor in oligodendrogiomas. *Neurosurgery* 48, 864-873.
- Robledo, M. M., Bartolome, R. A., Longo, N., Rodriguez-Frade, J. M., Mellado, M., Longo, I., van Muijen, G. N., Sanchez-Mateos, P., and Teixido, J. (2001). Expression of functional chemokine receptors CXCR3 and CXCR4 on human melanoma cells. *J Biol Chem* 276, 45098-45105.
- Röbuck, K. A. (1999). Oxidant stress regulation of IL-8 and ICAM-1 gene expression: differential activation and binding of the transcription factors AP-1 and NF-kappaB (Review). *Int J Mol Med* 4, 223-230.
- Ronnov-Jessen, L., and Petersen, O. W. (1993). Induction of alpha-smooth muscle actin by transforming growth factor- beta 1 in quiescent human breast gland fibroblasts. Implications for myofibroblast generation in breast neoplasia. *Lab Invest* 68, 696-707.
- Rosenkilde, M. M., Waldhoer, M., Luttichau, H. R., and Schwartz, T. W. (2001). Virally encoded 7TM receptors. *Oncogene* 20, 1582-1593.
- Ross, D. T., Scherf, U., Eisen, M. B., Perou, C. M., Rees, C., Spellman, P., Iyer, V., Jeffrey, S. S., Van de Rijn, M., Waltham, M., et al. (2000). Systematic variation in gene expression patterns in human cancer cell lines. *Nat Genet* 24, 227-235.
- Rossi, D., and Zlotnik, A. (2000). The biology of chemokines and their receptors. *Annu Rev Immunol* 18, 217-242.
- Salcedo R, Ponce ML, Young HA, Wasserman K, Ward JM, Kleinman HK, Oppenheim JJ, Murphy WJ. (2000). Human endothelial cells express CCR2 and respond to MCP-1: direct role of MCP-1 in angiogenesis and tumor progression. *Blood* 96, 34-40.
- Sadhu, C., Masinovsky, B., and Staunton, D. E. (1998). Differential regulation of chemoattractant-stimulated beta 2, beta 3, and beta 7 integrin activity. *J Immunol* 160, 5622-5628.
- Saeki, H., Wu, M. T., Olasz, E., and Hwang, S. T. (2000). A migratory population of skin-derived dendritic cells expresses CXCR5, responds to B lymphocyte chemoattractant in vitro, and co-localizes to B cell zones in lymph nodes in vivo. *Eur J Immunol* 30, 2808-2814.
- Sallusto, F., Lenig, D., Förster, R., Lipp, M., and Lanzavecchia, A. (1999a). Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 401, 708-712.
- Sallusto, F., Palermo, B., Lenig, D., Miettinen, M., Matikainen, S., Julkunen, I., Förster, R., Burgstahler, R., Lipp, M., and Lanzavecchia, A. (1999b). Distinct patterns and kinetics of chemokine production regulate dendritic cell function. *Eur J Immunol* 29, 1617-1625.
- Sallusto F, Kremmer E, Palermo B, Hoy A, Ponath P, Qin S, Förster R, Lipp M, Lanzavecchia A. (1999c). Switch in chemokine receptor expression upon TCR stimulation reveals novel homing potential for recently activated T cells. *Eur J Immunol* 29, 2037-45.
- Sallusto F, Schaeerli P, Loetscher P, Schaniel C, Lenig D, Mackay CR, Qin S, Lanzavecchia A. (1998). Rapid and coordinated switch in chemokine receptor expression during dendritic cell maturation. *Eur J Immunol* 28, 2760-2769.
- Salmi, M., and Jalkanen, S. (1997). How do lymphocytes know where to go: current concepts and enigmas of lymphocyte homing. *Adv. Immunol.* 64, 139-218.
- Salvucci, O., Yao, L., Villalba, S., Sajewicz, A., Pittaluga, S., and Tosato, G. (2002). Regulation of endothelial cell branching morphogenesis by endogenous chemokine stromal-derived factor-1. *Blood* 99, 2703-2711.
- Scandella, E., Men, Y., Gillessen, S., Förster, R., and Groettrup, M. (2002). Prostaglandin E2 is a key factor for CCR7 surface expression and migration of monocyte-derived dendritic cells. *Blood* 100, 1354-1361.
- Schadendorf D, Worm M, Czarnetzki BM. (1993). Determination of granulocyte /macrophage-colony-stimulating factor secretion by human melanoma cells and its effects on human melanoma cell proliferation. *J Cancer Res Clin Oncol* 119, 501-3
- Schall, T. J., and Bacon, K. B. (1994). Chemokines, leukocyte trafficking, and inflammation. *Curr. Opin. Immunol.* 6, 865-873.
- Scherf, U., Ross, D. T., Waltham, M., Smith, L. H., Lee, J. K., Tanabe, L., Kohn, K. W.,

- Reinhold, W. C., Myers, T. G., Andrews, D. T., et al. (2000). A gene expression database for the molecular pharmacology of cancer. *Nat Genet* 24, 236-244.
- Schneider GP, Salcedo R, Dong HF, Kleinman HK, Oppenheim JJ, Howard OM. (2002). Suradista NSC 651016 inhibits the angiogenic activity of CXCL12-stromal cell-derived factor 1alpha. *Clin Cancer Res* 8, 3955-60
 - Schor, S. L. (1994). Cytokine control of cell motility: modulation and mediation by the extracellular matrix. *Prog Growth Factor Res* 5, 223-248.
 - Schweickart, V. L., Raport, C. J., Godiska, R., Byers, M. G., Eddy, R. L., Jr., Shows, T. B., and Gray, P. W. (1994). Cloning of human and mouse EBI1, a lymphoid-specific G-protein-coupled receptor encoded on human chromosome 17q12-q21.2. *Genomics* 23, 643-650.
 - Sell, S., and Pierce, G. B. (1994). Maturation arrest of stem cell differentiation is a common pathway for the cellular origin of teratocarcinomas and epithelial cancers. *Lab Invest* 70, 6-22.
 - Sharma S, Stolina M, Luo J, Strieter RM, Burdick M, Zhu LX, Batra RK, Dubinett SM. (2000). Secondary lymphoid tissue chemokine mediates T cell-dependent antitumor responses in vivo. *J Immunol* 164, 4558-4563.
 - Shimonaka, M., and Yamaguchi, Y. (1994). Purification and biological characterization of epitaxin, a fibroblast- derived motility factor for epithelial cells. *J Biol Chem* 269, 14284-14289.
 - Shinomiya, T., Li, X. K., Amemiya, H., and Suzuki, S. (1997). An immunosuppressive agent, FTY720, increases intracellular concentration of calcium ion and induces apoptosis in HL-60. *Immunology* 91, 594-600.
 - Signoret, N., Rosenkilde, M. M., Klasse, P. J., Schwartz, T. W., Malim, M. H., Hoxie, J. A., and Marsh, M. (1998). Differential regulation of CXCR4 and CCR5 endocytosis. *J Cell Sci* 111, 2819-2830.
 - Sim, S. J., Ro, J. Y., Ordonez, N. G., Park, Y. W., Kee, K. H., and Ayala, A. G. (1999). Metastatic renal cell carcinoma to the bladder: a clinicopathologic and immunohistochemical study. *Mod Pathol* 12, 351-355.
 - Skobe, M., Hawighorst, T., Jackson, D. G., Prevo, R., Janes, L., Velasco, P., Riccardi, L., Alitalo, K., Claffey, K., and Detmar, M. (2001). Induction of tumor lymphangiogenesis by VEGF-C promotes breast cancer metastasis. *Nat Med* 7, 192-198.
 - Sliva, D., English, D., Lyons, D., and Lloyd, F. P., Jr. (2002). Protein kinase C induces motility of breast cancers by upregulating secretion of urokinase-type plasminogen activator through activation of AP-1 and NF-kappaB. *Biochem Biophys Res Commun* 290, 552-557.
 - Smith, J. R., Braziel, R. M., Paoletti, S., Lipp, M., Uggioni, M., and Rosenbaum, J. T. (2002). Expression of B cell-attracting chemokine-1 (CXCL13) by malignant lymphocytes and vascular endothelium in primary central nervous system lymphoma. *Blood* 101, 815-21.
 - Solecki, D., Bernhardt, G., Lipp, M., and Wimmer, E. (2000). Identification of a nuclear respiratory factor-1 binding site within the core promoter of the human polio virus receptor/CD155 gene. *J Biol Chem* 275, 12453-12462.
 - Solecki, D. J., Gromeier, M., Mueller, S., Bernhardt, G., and Wimmer, E. (2002). Expression of the human poliovirus receptor/CD155 gene is activated by sonic hedgehog. *J Biol Chem* 277, 25697-25702.
 - Soto, H., Wang, W., Strieter, R. M., Copeland, N. G., Gilbert, D. J., Jenkins, N. A., Hedrick, J., and Zlotnik, A. (1998). The CC chemokine 6Ckine binds the CXC chemokine receptor CXCR3. *Proc Natl Acad Sci USA* 95, 8205-8210.
 - Sozzani, S., Allavena, P., D'Amico, G., Luini, W., Bianchi, G., Kataura, M., Imai, T., Yoshie, O., Bonecchi, R., and Mantovani, A. (1998). Differential regulation of chemokine receptors during dendritic cell maturation: a model for their trafficking properties. *J Immunol* 161, 1083-1086.
 - Springer, T. A. (1994). Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* 76, 301-314.
 - Stein, J. V., Rot, A., Luo, Y., Narasimhaswamy, M., Nakano, H., Gunn, M. D., Matsuzawa, A., Quackenbush, E. J., Dorf, M. E., and von Andrian, U. H. (2000). The CC chemokine thymus-derived chemotactic agent 4 (TCA-4, secondary lymphoid tissue chemokine, 6Ckine, exodus-2) triggers lymphocyte function-associated antigen 1-mediated arrest of rolling T

- lymphocytes in peripheral lymph node high endothelial venules. *J Exp Med* 191, 61-76.
- Stein, J. V., Soriano, S. F., M'Rini, C., Nombela-Arrieta, C., Gonzalez De Buitrago, G., Rodriguez-Frade, J. M., Mellado, M., Girard, J. P., and Martinez, A. C. (2002). CCR7-mediated physiological lymphocyte homing involves activation of a tyrosine kinase pathway. *Blood* 101, 38-44.
 - Strader, C. D., Fong, T. M., Tota, M. R., Underwood, D., and Dixon, R. A. (1994). Structure and function of G protein-coupled receptors. *Annu Rev Biochem* 63, 101-132.
 - Streblow DN, Soderberg-Naucler C, Vieira J, Smith P, Wakabayashi E, Ruchti F, Mattison K, Altschuler Y, Nelson JA. (1999). The human cytomegalovirus chemokine receptor US28 mediates vascular smooth muscle cell migration. *Cell* 99, 511-20.
 - Strieter, R. M. (2001). Chemokines: not just leukocyte chemoattractants in the promotion of cancer. *Nat Immunol* 2, 285-286.
 - Strieter, R. M., Polverini, P. J., Arenberg, D. A., and Kunkel, S. L. (1995a). The role of CXC chemokines as regulators of angiogenesis. *Shock* 4, 155-160.
 - Strieter, R. M., Polverini, P. J., Arenberg, D. A., Walz, A., Opdenakker, G., Van Damme, J., and Kunkel, S. L. (1995b). Role of C-X-C chemokines as regulators of angiogenesis in lung cancer. *J Leukoc Biol* 57, 752-762.
 - Strieter, R. M., Polverini, P. J., Kunkel, S. L., Arenberg, D. A., Burdick, M. D., Kasper, J., Dzuiba, J., Van Damme, J., Walz, A., Marriott, D., and et al. (1995c). The functional role of the ELR motif in CXC chemokine-mediated angiogenesis. *J Biol Chem* 270, 27348-27357.
 - Sullivan, S. K., McGrath, D. A., Grigoriadis, D., and Bacon, K. B. (1999). Pharmacological and signaling analysis of human chemokine receptor CCR- 7 stably expressed in HEK-293 cells: high-affinity binding of recombinant ligands MIP-3beta and SLC stimulates multiple signaling cascades. *Biochem Biophys Res Commun* 263, 685-690.
 - Suzuki, S., Enosawa, S., Kakefuda, T., Shinomiya, T., Amari, M., Naoe, S., Hoshino, Y., and Chiba, K. (1996). A novel immunosuppressant, FTY720, with a unique mechanism of action, induces long-term graft acceptance in rat and dog allotransplantation. *Transplantation* 61, 200-205.
 - Suzuki, K., Kazui, T., Kawabe, A., Li, X. K., Amemiya, H., and Suzuki, S. (1998). Beneficial immunosuppressive effect of FTY720 combined with intrathymic injection of splenic cells on rat pancreaticoduodenal allograft. *Transplant Proc* 30, 1067-1068.
 - Szekanecz Z, and Koch AE.(2001). Chemokines and angiogenesis. *Curr Opin Rheumatol* 13, 202-8
 - Takuwa Y (2002). Subtype-specific differential regulation of Rho family G proteins and cell migration by the Edg family sphingosine-1-phosphate receptors. *Biochim Biophys Acta* 1582, 112-20.
 - Tanaka, Y., Adams, D. H., and Shaw, S. (1993). Regulation of leukocyte recruitment by proadhesive cytokines immobilized on endothelial proteoglycan. *Curr. Top. Microbiol. Immunol.* 184, 99-106.
 - Tanaka, Y., Mine, S., Figgdr, C. G., Wake, A., Hirano, H., Tsukada, J., Aso, M., Fujii, K., Saito, K., van Kooyk, Y., and Eto, S. (1998). Constitutive chemokine production results in activation of leukocyte function-associated antigen-1 on adult T-cell leukemia cells. *Blood* 91, 3909-3919.
 - Tangemann, K., Gunn, M. D., Giblin, P., and Rosen, S. D. (1998). A high endothelial cell-derived chemokine induces rapid, efficient, and subset-selective arrest of rolling T lymphocytes on a reconstituted endothelial substrate. *J. Immunol.* 161, 6330-6337.
 - Teruya-Feldstein, J., Tosato, G., and Jaffe, E. S. (2000). The role of chemokines in Hodgkin's disease. *Leuk Lymphoma* 38, 363-371.
 - Thanaraj, T. A., and Clark, F. (2001). Human GC-AG alternative intron isoforms with weak donor sites show enhanced consensus at acceptor exon positions. *Nucleic Acids Res* 29, 2581-2593.
 - Till, K. J., Lin, K., Zuzel, M., and Cawley, J. C. (2002). The chemokine receptor CCR7 and alpha4 integrin are important for migration of chronic lymphocytic leukemia cells into lymph nodes. *Blood* 99, 2977-2984.
 - Tilton, B., Ho, L., Oberlin, E., Loetscher, P., Baleux, F., Clark-Lewis, I., and Thelen, M.

- (2000). Signal transduction by CXC chemokine receptor 4. Stromal cell-derived factor 1 stimulates prolonged protein kinase B and extracellular signal-regulated kinase 2 activation in T lymphocytes. *J Exp Med* 192, 313-324.
- Townson, J. R., and Nibbs, R. J. (2002). Characterization of mouse CCX-CKR, a receptor for the lymphocyte-attracting chemokines TECK/mCCL25, SLC/mCCL21 and MIP-3beta/mCCL19: comparison to human CCX-CKR. *Eur J Immunol* 32, 1230-1241.
 - Trentin, L., Agostini, C., Facco, M., Piazza, F., Perin, A., Siviero, M., Gurrieri, C., Galvan, S., Adami, F., Zambello, R., and Semenzato, G. (1999). The chemokine receptor CXCR3 is expressed on malignant B cells and mediates chemotaxis. *J Clin Invest* 104, 115-121.
 - Ueno, T., Hara, K., Willis, M. S., Malin, M. A., Höpken, U. E., Gray, D. H., Matsushima, K., Lipp, M., Springer, T. A., Boyd, R. L., et al. (2002). Role for CCR7 ligands in the emigration of newly generated T lymphocytes from the neonatal thymus. *Immunity* 16, 205-218.
 - Vassileva, G., Soto, H., Zlotnik, A., Nakano, H., Kakiuchi, T., Hedrick, J. A., and Lira, S. A. (1999). The reduced expression of 6Ckine in the plt mouse results from the deletion of one of two 6Ckine genes. *J Exp Med* 190, 1183-1188.
 - Vicari AP, Ait-Yahia S, Chemin K, Mueller A, Zlotnik A, Caux C. (2000). Antitumor effects of the mouse chemokine 6Ckine/SLC through angiostatic and immunological mechanisms. *J Immunol* 165, 1992-2000.
 - Vieira, J., and Messing, J. (1982). The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* 19, 259-268.
 - Vila-Coro, A. J., Rodriguez-Frade, J. M., Martin De Ana, A., Moreno-Ortiz, M. C., Martinez, A. C., and Mellado, M. (1999). The chemokine SDF-1alpha triggers CXCR4 receptor dimerization and activates the JAK/STAT pathway. *Faseb J* 13, 1699-1710.
 - Volin, M. V., Joseph, L., Shockley, M. S., and Davies, P. F. (1998). Chemokine receptor CXCR4 expression in endothelium. *Biochem Biophys Res Commun* 242, 46-53.
 - Wang, J., Asensio, V. C., and Campbell, I. L. (2002). Cytokines and chemokines as mediators of protection and injury in the central nervous system assessed in transgenic mice. *Curr Top Microbiol Immunol* 265, 23-48.
 - Wang, J. F., Park, I. W., and Groopman, J. E. (2000). Stromal cell-derived factor-1alpha stimulates tyrosine phosphorylation of multiple focal adhesion proteins and induces migration of hematopoietic progenitor cells: roles of phosphoinositide-3 kinase and protein kinase C. *Blood* 95, 2505-2513.
 - Wang, J. M., Deng, X., Gong, W., and Su, S. (1998a). Chemokines and their role in tumor growth and metastasis. *J Immunol Methods* 220, 1-17.
 - Wang, M. E., Tejpal, N., Qu, X., Yu, J., Okamoto, M., Stepkowski, S. M., and Kahan, B. D. (1998b). Immunosuppressive effects of FTY720 alone or in combination with cyclosporine and/or sirolimus. *Transplantation* 65, 899-905.
 - Webb, L. M., Ehrengruber, M. U., Clark-Lewis, I., Baggiolini, M., and Rot, A. (1993). Binding to heparan sulfate or heparin enhances neutrophil responses to interleukin 8. *Proc Natl Acad Sci USA* 90, 7158-7162.
 - Wiley, H. E., Gonzalez, E. B., Maki, W., Wu, M. T., and Hwang, S. T. (2001). Expression of CC chemokine receptor-7 and regional lymph node metastasis of B16 murine melanoma. *J Natl Cancer Inst* 93, 1638-1643.
 - Wilkie, T. M., Gilbert, D. J., Olsen, A. S., Chen, X. N., Amatruda, T. T., Korenberg, J. R., Trask, B. J., de Jong, P., Reed, R. R., Simon, M. I., and et al. (1992). Evolution of the mammalian G protein alpha subunit multigene family. *Nat Genet* 1, 85-91.
 - Willimann, K., Legler, D. F., Loetscher, M., Roos, R. S., Delgado, M. B., Clark-Lewis, I., Baggiolini, M., and Moser, B. (1998). The chemokine SLC is expressed in T cell areas of lymph nodes and mucosal lymphoid tissues and attracts activated T cells via CCR7. *Eur J Immunol* 28, 2025-2034.
 - Yanagawa, Y., Masubuchi, Y., and Chiba, K. (1998). FTY720, a novel immunosuppressant, induces sequestration of circulating mature lymphocytes by acceleration of lymphocyte homing in rats, III. Increase in frequency of CD62L-positive T cells in Peyer's patches by FTY720-induced lymphocyte homing. *Immunology* 95, 591-594.
 - Ye, R. D., (2001). Regulation of nuclear factor kB activation by G-protein-coupled receptors.

- J Leukoc Biol 70, 839-848
- Yoshida, R., Imai, T., Hieshima, K., Kusuda, J., Baba, M., Kitaura, M., Nishimura, M., Kakizaki, M., Nomiyama, H., and Yoshie, O. (1997). Molecular cloning of a novel human CC chemokine EBI1-ligand chemokine that is a specific functional ligand for EBI1, CCR7. *J Biol Chem* 272, 13803-13809.
 - Yoshida, R., Nagira, M., Imai, T., Baba, M., Takagi, S., Tabira, Y., Akagi, J., Nomiyama, H., and Yoshie, O. (1998a). EBI1-ligand chemokine (ELC) attracts a broad spectrum of lymphocytes: activated T cells strongly up-regulate CCR7 and efficiently migrate toward ELC. *Int Immunol* 10, 901-910.
 - Yoshida, R., Nagira, M., Kitaura, M., Imagawa, N., Imai, T., and Yoshie, O. (1998b). Secondary lymphoid-tissue chemokine is a functional ligand for the CC chemokine receptor CCR7. *J Biol Chem* 273, 7118-7122.
 - Yoshie, O., Imai, T., and Nomiyama, H. (1997). Novel lymphocyte-specific CC chemokines and their receptors. *J Leukoc Biol* 62, 634-644.
 - Youngs, S. J., Ali, S. A., Taub, D. D., and Rees, R. C. (1997). Chemokines induce migrational responses in human breast carcinoma cell lines. *Int J Cancer* 71, 257-266.
 - Yu, G. H., Lawton, T. J., Pasha, T. L., and Reynolds, C. (2000). Intercellular adhesion molecule expression in ductal carcinoma of the breast: correlation of immunohistochemical staining with cytologic smear pattern. *Diagn Cytopathol* 23, 73-76.
 - Zhang, G. J., and Adachi, I. (1999). Serum levels of soluble intercellular adhesion molecule-1 and E-selectin in metastatic breast carcinoma: correlations with clinicopathological features and prognosis. *Int J Oncol* 14, 71-77.