

9 LITERATURVERZEICHNIS

1. Saini, S., et al., Ferrite particles: a superparamagnetic MR contrast agent for enhanced detection of liver carcinoma. *Radiology*, 1987. 162(1 Pt 1): p. 217-22.
2. Saini, S., et al., Ferrite particles: a superparamagnetic MR contrast agent for the reticuloendothelial system. *Radiology*, 1987. 162(1 Pt 1): p. 211-6.
3. Chambon, C., et al., Superparamagnetic iron oxides as positive MR contrast agents: in vitro and in vivo evidence. *Magn Reson Imaging*, 1993. 11(4): p. 509-19.
4. Anzai, Y., et al., MR angiography with an ultrasmall superparamagnetic iron oxide blood pool agent. *J Magn Reson Imaging*, 1997. 7(1): p. 209-14.
5. Reimer, P., et al., Contrast-enhanced 3D-MRA of the upper abdomen with a bolus-injectable SPIO (SH U 555 A). *J Magn Reson Imaging*, 1999. 10(1): p. 65-71.
6. Schmitz, S.A., T. Albrecht, and K.J. Wolf, MR angiography with superparamagnetic iron oxide: feasibility study. *Radiology*, 1999. 213(2): p. 603-7.
7. Reimer, P., et al., SPIO-enhanced 2D-TOF MR angiography of the portal venous system: results of an intraindividual comparison. *J Magn Reson Imaging*, 1997. 7(6): p. 945-9.
8. Schmitz, S.A., et al., SPIO-unterstützte MR-Angiographie des portalvenösen Systems. *RöFo*, 2000. 172(1): p. 51-4.
9. Allkemper, T., et al., Contrast-enhanced blood-pool MR angiography with optimized iron oxides: effect of size and dose on vascular contrast enhancement in rabbits. *Radiology*, 2002. 223(2): p. 432-8.

10. Bremer, C., et al., RES-specific imaging of the liver and spleen with iron oxide particles designed for blood pool MR-angiography. *J Magn Reson Imaging*, 1999. 10(3): p. 461-7.
11. Schmitz, S.A., et al., SPIO-unterstützte MR-Angiographie zur Detektion venöser Thromben im Tiermodell. *RöFo*, 1999. 170(3): p. 316-321.
12. Brückmann, H. and W. Reith, Intrazerebrale Blutungen und zerebrovaskuläre Erkrankungen, in *Magnetresonanztomographie*, R. M. and S. W., Editors. 2002, Springer: Berlin-Heidelberg. p. 233-272.
13. Blackmore, C.C., et al., Magnetic resonance imaging of blood and clots in vitro. *Invest Radiol*, 1990. 25(12): p. 1316-24.
14. Bradley, W.G., Jr., MR appearance of hemorrhage in the brain. *Radiology*, 1993. 189(1): p. 15-26.
15. Hoppel, B.E., et al., Measurement of regional blood oxygenation and cerebral hemodynamics. *Magn Reson Med*, 1993. 30(6): p. 715-23.
16. Detre, J.A. and J. Wang, Technical aspects and utility of fMRI using BOLD and ASL. *Clin Neurophysiol*, 2002. 113(5): p. 621-34.
17. Bush, C.H., The magnetic resonance imaging of musculoskeletal hemorrhage. *Skeletal Radiol*, 2000. 29(1): p. 1-9.
18. Bradley, W.G., Jr. and P.G. Schmidt, Effect of methemoglobin formation on the MR appearance of subarachnoid hemorrhage. *Radiology*, 1985. 156(1): p. 99-103.
19. Gomori, J.M., et al., Intracranial hematomas: imaging by high-field MR. *Radiology*, 1985. 157(1): p. 87-93.
20. Gomori, J.M. and R.I. Grossman, Mechanisms responsible for the MR appearance and evolution of intracranial hemorrhage. *Radiographics*, 1988. 8(3): p. 427-40.

21. Francis, C.W. and S. Totterman, Magnetic resonance imaging of deep vein thrombi correlates with response to thrombolytic therapy. *Thromb Haemost*, 1995. 73(3): p. 386-91.
22. Isensee, C., J. Reul, and A. Thron, Magnetic resonance imaging of thrombosed dural sinuses. *Stroke*, 1994. 25(1): p. 29-34.
23. Singer, J.R., Blood flow rates by nuclear magnetic resonance. *Science*, 1959. 130: p. 1652-1653.
24. Listerud, J. and S.W. Atals, Fundamentals of Flow and Hemodynamics, in *Magnetic Resonance Imaging of the Brain and Spine*, S.W. Atals, Editor. 1996, Lippincott-Raven. p. 65-87.
25. Parker, D.L., C. Yuan, and D.D. Blatter, MR angiography by multiple thin slab 3D acquisition. *Magn Reson Med*, 1991. 17(2): p. 434-51.
26. Liu, K., D.H. Lee, and B.K. Rutt, Systematic assessment and evaluation of sliding interleaved kY (SLINKY) acquisition for 3D MRA. *J Magn Reson Imaging*, 1998. 8(4): p. 912-23.
27. Atkinson, D., et al., Improved MR angiography: magnetization transfer suppression with variable flip angle excitation and increased resolution. *Radiology*, 1994. 190(3): p. 890-4.
28. Dagirmanjian, A., et al., High resolution, magnetization transfer saturation, variable flip angle, time-of-flight MRA in the detection of intracranial vascular stenoses. *J Comput Assist Tomogr*, 1995. 19(5): p. 700-6.
29. Edelman, R.R., et al., Improved time-of-flight MR angiography of the brain with magnetization transfer contrast. *Radiology*, 1992. 184(2): p. 395-9.
30. Pike, G.B., et al., Magnetization transfer time-of-flight magnetic resonance angiography. *Magn Reson Med*, 1992. 25(2): p. 372-9.

31. Laub, G.A. and W.A. Kaiser, MR angiography with gradient motion refocusing. *J Comput Assist Tomogr*, 1988. 12(3): p. 377-82.
32. Niendorf, H.P., T. Balzer, and P. Reimer, Kontrastmittel, in *Magnetresonanztomographie*, M. Reiser and W. Semmler, Editors. 2002, Springer: Berlin, Heidelberg, New York. p. 111-132.
33. Weinmann, H.-J., Eigenschaften von Gd-DTPA-Dimeglumin, in *Magnevist - Eine Monographie*, N. Hosten, et al., Editors. 2001, Blackwell Wissenschaftsverlag: Berlin, Wien. p. 5-12.
34. Petersein, J., S. Saini, and R. Weissleder, Liver. II: Iron oxide-based reticuloendothelial contrast agents for MR imaging. *Clinical review*. 1996.
35. Turetschek, K., et al., Tumor microvascular characterization using ultrasmall superparamagnetic iron oxide particles (USPIO) in an experimental breast cancer model. *J Magn Reson Imaging*, 2001. 13(6): p. 882-8.
36. Kresse, M., et al., Targeting of ultrasmall superparamagnetic iron oxide (USPIO) particles to tumor cells in vivo by using transferrin receptor pathways. *Magn Reson Med*, 1998. 40(2): p. 236-42.
37. Saeed, M., et al., Demarcation of myocardial ischemia: magnetic susceptibility effect of contrast medium in MR imaging. *Radiology*, 1989. 173(3): p. 763-7.
38. Schorner, W., M. Laniado, and R. Felix, [1st clinical use of gadolinium-DTPA in the nuclear magnetic resonance tomography visualization of a parapelvic kidney cyst]. *ROFO Fortschr Geb Rontgenstr Nuklearmed*, 1984. 141(2): p. 227-8.
39. Bousquet, J.C., et al., Gd-DOTA: characterization of a new paramagnetic complex. *Radiology*, 1988. 166(3): p. 693-8.
40. Sze, G., et al., Multicenter study of gadodiamide injection as a contrast agent in MR imaging of the brain and spine. *Radiology*, 1991. 181(3): p. 693-9.

41. Runge, V.M., et al., Clinical safety and efficacy of gadoteridol: a study in 411 patients with suspected intracranial and spinal disease. *Radiology*, 1991. 181(3): p. 701-9.
42. Caudana, R., et al., Focal malignant hepatic lesions: MR imaging enhanced with gadolinium benzyloxypropionictetra-acetate (BOPTA)--preliminary results of phase II clinical application. *Radiology*, 1996. 199(2): p. 513-20.
43. Hartmann, M., et al., [Does the administration of a high dose of a paramagnetic contrast medium (Gadovist) improve the diagnostic value of magnetic resonance tomography in glioblastomas?]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*, 1996. 164(2): p. 119-25.
44. Rubin, D.L., et al., A multicenter, randomized, double-blind study to evaluate the safety, tolerability, and efficacy of OptiMARK (gadoversetamide injection) compared with Magnevist (gadopentetate dimeglumine) in patients with liver pathology: results of a Phase III clinical trial. *J Magn Reson Imaging*, 1999. 9(2): p. 240-50.
45. Coche, E.E., F.D. Hammer, and P.P. Goffette, Demonstration of pulmonary embolism with gadolinium-enhanced spiral CT. *Eur Radiol*, 2001. 11(11): p. 2306-9.
46. Hamm, B., et al., Phase I clinical evaluation of Gd-EOB-DTPA as a hepatobiliary MR contrast agent: safety, pharmacokinetics, and MR imaging. *Radiology*, 1995. 195(3): p. 785-92.
47. Schuhmann-Giampieri, G., et al., Preclinical evaluation of Gd-EOB-DTPA as a contrast agent in MR imaging of the hepatobiliary system. *Radiology*, 1992. 183(1): p. 59-64.
48. Weinmann, H.J., et al., A new lipophilic gadolinium chelate as a tissue-specific contrast medium for MRI. *Magn Reson Med*, 1991. 22(2): p. 233-7; discussion 242.

49. Holman, E.R., et al., Assessment of acute myocardial infarction in man with magnetic resonance imaging and the use of a new paramagnetic contrast agent gadolinium-BOPTA. *Magn Reson Imaging*, 1996. 14(1): p. 21-9.
50. Hamm, B., et al., Focal liver lesions: MR imaging with Mn-DPDP--initial clinical results in 40 patients. *Radiology*, 1992. 182(1): p. 167-74.
51. Vassallo, P., et al., AMI-227-enhanced MR lymphography: usefulness for differentiating reactive from tumor-bearing lymph nodes. *Radiology*, 1994. 193(2): p. 501-6.
52. Weissleder, R., et al., Ultras-small superparamagnetic iron oxide: an intravenous contrast agent for assessing lymph nodes with MR imaging. *Radiology*, 1990. 175(2): p. 494-8.
53. Schmiedl, U., et al., Albumin labeled with Gd-DTPA as an intravascular, blood pool-enhancing agent for MR imaging: biodistribution and imaging studies. *Radiology*, 1987. 162(1 Pt 1): p. 205-10.
54. Bluemke, D.A., et al., Carotid MR angiography: phase II study of safety and efficacy for MS-325. *Radiology*, 2001. 219(1): p. 114-22.
55. Gaillard, S., et al., Safety and pharmacokinetics of p792, a new blood-pool agent: results of clinical testing in nonpatient volunteers. *Invest Radiol*, 2002. 37(4): p. 161-6.
56. Mahfouz, A.E., Ms-325 Epix. *Curr Opin Investig Drugs*, 2000. 1(4): p. 476-80.
57. Ruehm, S.G., et al., MR angiography with a new rapid-clearance blood pool agent: Initial experience in rabbits. *Magn Reson Med*, 2002. 48(5): p. 844-51.
58. Sharafuddin, M.J., et al., Comparison of MS-325- and gadodiamide-enhanced MR venography of ilio-caval veins. *J Vasc Interv Radiol*, 2002. 13(10): p. 1021-7.

59. Larsson, E.M., et al., MR venography using an intravascular contrast agent: results from a multicenter phase 2 study of dosage. *AJR Am J Roentgenol*, 2003. 180(1): p. 227-32.
60. Klein, C., et al., Improvement of image quality of non-invasive coronary artery imaging with magnetic resonance by the use of the intravascular contrast agent Clariscan trade mark (NC100150 injection) in patients with coronary artery disease. *J Magn Reson Imaging*, 2003. 17(6): p. 656-62.
61. Clarke, S.E., et al., Comparison of two blood pool contrast agents for 0.5-T MR angiography: experimental study in rabbits. *Radiology*, 2000. 214(3): p. 787-94.
62. Wacker, F.K., et al., Use of a blood-pool contrast agent for MR-guided vascular procedures: feasibility of ultrasmall superparamagnetic iron oxide particles. *Acad Radiol*, 2002. 9(11): p. 1251-4.
63. Prince, M.R., Gadolinium-enhanced MR aortography. *Radiology*, 1994. 191(1): p. 155-64.
64. Geftter, W.B., et al., Pulmonary thromboembolism: recent developments in diagnosis with CT and MR imaging. *Radiology*, 1995. 197(3): p. 561-74.
65. Prince, M.R., et al., Breath-hold gadolinium-enhanced MR angiography of the abdominal aorta and its major branches. *Radiology*, 1995. 197(3): p. 785-92.
66. Bongartz, G.M., Flussphänomene und MR-Angiographietechniken, in *Magnetresonanztomographie*, M. Reiser and W. Semmler, Editors. 2002, Springer: Berlin, Heidelberg, New York. p. 510-527.
67. Bosmans, H. and G. Marchal, Spatial versus temporal resolution in contrast-enhanced magnetic resonance angiography, in *Magnetic Resonance Angiography*, I.P. Arlart, G.M. Bongartz, and G. Marchal, Editors. 2001, Springer: Berlin, Heidelberg, New York. p. 105-113.

68. Prince, M., T. Grist, and J.F. Debatin, 3D Contrast MR Angiography. 3rd ed. 2003, Berlin, Heidelberg, New York: Springer.
69. Prince, M.R., et al., Contrast-enhanced abdominal MR angiography: optimization of imaging delay time by automating the detection of contrast material arrival in the aorta. *Radiology*, 1997. 203(1): p. 109-14.
70. Meaney, J.F., et al., Pulmonary magnetic resonance angiography. *J Magn Reson Imaging*, 1999. 10(3): p. 326-38.
71. Isoda, H., et al., Clinical evaluation of pulmonary 3D time-of-flight MRA with breath holding using contrast media. *J Comput Assist Tomogr*, 1995. 19(6): p. 911-9.
72. Meaney, J.F., et al., Diagnosis of pulmonary embolism with magnetic resonance angiography. *N Engl J Med*, 1997. 336(20): p. 1422-7.
73. Gupta, A., et al., Acute pulmonary embolism: diagnosis with MR angiography. *Radiology*, 1999. 210(2): p. 353-9.
74. Kreitner, K.F., et al., [Contrast media enhanced three dimensional MR angiography of the pulmonary arteries in patients with chronic recurrent pulmonary embolism--comparison with selective intra-arterial DSA]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*, 2000. 172(2): p. 122-8.
75. Kruger, S., et al., Diagnosis of pulmonary arterial hypertension and pulmonary embolism with magnetic resonance angiography. *Chest*, 2001. 120(5): p. 1556-61.
76. Goyen, M., et al., Dynamic 3D MR angiography of the pulmonary arteries in under four seconds. *J Magn Reson Imaging*, 2001. 13(3): p. 372-7.
77. Oudkerk, M., et al., Comparison of contrast-enhanced magnetic resonance angiography and conventional pulmonary angiography for the diagnosis of pulmonary embolism: a prospective study. *Lancet*, 2002. 359(9318): p. 1643-7.

78. Grist, T.M., et al., Pulmonary angiography with MR imaging: preliminary clinical experience. *Radiology*, 1993. 189(2): p. 523-30.
79. Schiebler, M.L., et al., Suspected pulmonary embolism: prospective evaluation with pulmonary MR angiography. *Radiology*, 1993. 189(1): p. 125-31.
80. Sostman, H.D., et al., Prospective comparison of helical CT and MR imaging in clinically suspected acute pulmonary embolism. *J Magn Reson Imaging*, 1996. 6(2): p. 275-81.
81. Loubeyre, P., et al., Dynamic contrast-enhanced MR angiography of pulmonary embolism: comparison with pulmonary angiography. *AJR Am J Roentgenol*, 1994. 162(5): p. 1035-9.
82. Bergin, C.J., et al., Chronic thromboembolism: diagnosis with helical CT and MR imaging with angiographic and surgical correlation. *Radiology*, 1997. 204(3): p. 695-702.
83. Lebowitz, J.A., et al., Gadolinium-enhanced body MR venography with subtraction technique. *AJR Am J Roentgenol*, 1997. 169(3): p. 755-8.
84. Fraser, D.G., et al., Deep venous thrombosis: diagnosis by using venous enhanced subtracted peak arterial MR venography versus conventional venography. *Radiology*, 2003. 226(3): p. 812-20.
85. Ruehm, S.G., K. Zimny, and J.F. Debatin, Direct contrast-enhanced 3D MR venography. *Eur Radiol*, 2001. 11(1): p. 102-12.
86. Spritzer, C.E., et al., Deep venous thrombosis: experience with gradient-echo MR imaging in 66 patients. *Radiology*, 1990. 177(1): p. 235-41.
87. Carpenter, J.P., et al., Magnetic resonance venography for the detection of deep venous thrombosis: comparison with contrast venography and duplex Doppler ultrasonography. *J Vasc Surg*, 1993. 18(5): p. 734-41.

88. Evans, A.J., et al., 1992 ARRS Executive Council Award. Detection of deep venous thrombosis: prospective comparison of MR imaging with contrast venography. *AJR Am J Roentgenol*, 1993. 161(1): p. 131-9.
89. Dupas, B., et al., Angiomagnetic resonance imaging of iliofemorocaval venous thrombosis. *Lancet*, 1995. 346(8966): p. 17-9.
90. Laissy, J.P., et al., Assessment of deep venous thrombosis in the lower limbs and pelvis: MR venography versus duplex Doppler sonography. 1996.
91. Catalano, C., et al., Role of MR venography in the evaluation of deep venous thrombosis. *Acta Radiol*, 1997. 38(5): p. 907-12.
92. Kroencke, T.J., et al., Three-dimensional gadolinium-enhanced magnetic resonance venography in suspected thrombo-occlusive disease of the central chest veins. *Chest*, 2001. 120(5): p. 1570-6.
93. Rodenwaldt, J., et al., [The MR imaging of the vena cava system: a comparison of different technics]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*, 1999. 171(2): p. 100-5.
94. Stehbens, W.E., Thrombosis and Vascular Trauma, in *Vascular Pathology*, W.E. Stehbens and J.T. Lie, Editors. 1995, Chapman & Hall: London. p. 63-87.
95. Rieger, H. and W. Theiss, Tiefe Bein- und Beckenvenenthrombose, in *Klinische Angiologie*, H. Rieger and W. Schoop, Editors. 1998, Springer: Heidelberg. p. 1037-1073.
96. Prandoni, P. and J.W. ten Cate, Epidemiology, risk factors, and natural history of venous thromboembolism, in *Pulmonary Embolism*, M. Oudkerk, P. Prandoni, and J.W. ten Cate, Editors. 1999, Blackwell Wissenschaft Berlin: Berlin. p. 2-32.
97. Biland, L., E. Zemp, and L.K. Widmer, [Epidemiology of venous thromboembolism]. *Internist (Berl)*, 1987. 28(5): p. 285-90.

98. Hansson, P.O., et al., Deep vein thrombosis and pulmonary embolism in the general population. 'The Study of Men Born in 1913'. Arch Intern Med, 1997. 157(15): p. 1665-70.
99. Silverstein, M.D., et al., Trends in the incidence of deep vein thrombosis and pulmonary embolism: a 25-year population-based study. Arch Intern Med, 1998. 158(6): p. 585-93.
101. Dorfman, G.S., et al., Occult pulmonary embolism: a common occurrence in deep venous thrombosis. AJR, 1987. 148(2): p. 263-6.
102. Huisman, M.V., et al., Unexpected high prevalence of silent pulmonary embolism in patients with deep venous thrombosis. Chest, 1989. 95(3): p. 498-502.
103. Nielsen, H.K., et al., Silent pulmonary embolism in patients with deep venous thrombosis. Incidence and fate in a randomized, controlled trial of anti-coagulation versus no anticoagulation. J Intern Med, 1994. 235(5): p. 457-61.
104. Leu, H.J., Pathologische Morphologie und Histopathologie der Venenwand, in Klinische Angiologie, H. Rieger and W. Schoop, Editors. 1998, Springer: Berlin, Heidelberg. p. 863-874.
105. Brunner, U., [The femoral-iliac thrombosis as an evolutive venous obstruction: problems from the surgical viewpoint]. Vasa, 1974. 3(1): p. 22-7.
106. Petrides, P.E., Blut, in Biochemie und Pathobiochemie, G. Löffler and P.E. Petrides, Editors. 1998, Springer: Berlin. p. 920-931.
107. Harmening, D.M. and L.D. Lemery, Introduction of Hemostasis: An Overview of Hemostatic Mechanism, Platelet Structure and Function, and Extrinsic and Intrinsic System, in Clinical Hematology and Fundamentals of Hemostasis, D.M. Harmening, Editor. 1997, Davis Company: Philadelphia. p. 481-508.
108. Andree, C., Zur Entstehung der Virchow'schen Triade. Ann. Univ. Sarav. Med., 1990/91. 26: p. 45.

109. Breddin, H.K. and M. Basic-Micic, [Pathophysiology and pathogenesis of venous thrombosis]. *Chirurg*, 1992. 63(4): p. 260-3.
110. Cooper, B.A. and J.R. Leclerc, Overview of Hemostasis and Blood Coagulation, in *Venous Thromboembolic Disorders*, J.R. Leclerc, Editor. 1991, Lea & Febiger: Philadelphia, London. p. 1-16.
111. Hamer, J.D. and P.C. Malone, Experimental deep venous thrombogenesis by a non-invasive method. *Ann R Coll Surg Engl*, 1984. 66(6): p. 416-9.
112. Aeland, R., Thrombus formation in microvascular surgery: An experimental study of the effects of surgical trauma. *Surg.*, 1973. 73(5): p. 766-771.
113. Thomas, D.P., et al., The relationship between vessel wall injury and venous thrombosis: an experimental study. *Br J Haematol*, 1985. 59(3): p. 449-57.
114. Aronson, D.L. and D.P. Thomas, Experimental studies on venous thrombosis: effect of coagulants, procoagulants and vessel contusion. *Thromb Haemost*, 1985. 54(4): p. 866-70.
115. Davies, M.J., Stability and instability: two faces of coronary atherosclerosis. The Paul Dudley White Lecture 1995. *Circulation*, 1996. 94(8): p. 2013-20.
116. Wessler, S., Thrombosis in the presence of vascular stasis. *Am J Med*, 1962. 33: p. 648-666.
117. Breddin, H.K. and V. Hach-Wunderle, [Physiologic studies of blood coagulation in deep venous thrombosis]. *Internist (Berl)*, 1987. 28(5): p. 308-16.
118. Prentice, C.R., Pathogenesis of thrombosis. *Haemostasis*, 1990. 20 Suppl 1: p. 50-9.
119. Hadfield, G., Thrombosis. *Ann. Roy. Coll. Surg. Engl.*, 1950. 6: p. 219-234.
120. Stehbens, W.E., Atherosclerosis and degenerative disease of blood vessels, in *Vascular Pathology*, W.E. Stehbens and J.T. Lie, Editors. 1995, Chapman and Hall: London. p. 175-268.

121. Leu, H.J., Histologische Altersbestimmung von arteriellen und venösen Thromben und Emboli. *Vasa*, 1973. 2(3): p. 265-74.
122. Frenzel, H. and W. Hort, Vergleichende experimentelle Untersuchungen über die Thrombenorganisation in Arterien und Venen (Organization of experimental thrombi in arteries and veins. A comparative study (author's transl)). *Basic Res Cardiol*, 1975. 70(5): p. 480-94.
123. Leu, A.J. and H.J. Leu, Spezielle Probleme bei der histologischen Altersbestimmung von Thromben und Emboli. *Pathologe*, 1989. 10(2): p. 87-92.
124. Klos, I., et al., [The lysability of thrombi obtained during operation. The degree of organisation and lysability by streptokinase in vitro of arterial and venous thrombi of different ages]. *Munch Med Wochenschr*, 1972. 114(44): p. 1927-32.
125. Dalen, J.E. and J.S. Alpert, Natural history of pulmonary embolism. *Prog Cardiovasc Dis*, 1975. 17(4): p. 257-70.
126. Abrams, H.L., Venography of the Lower Extremities, in *Abrams Angiography*, H.L. Abrams and P.H. Cook, Editors. 1983, Little, Brown: Boston. p. 1877-1921.
127. Gottlob R, M.R., Experimental thrombosis similiary clinicals conditions produced on the hind-legs of minor laboratoy animals. *angiology*, 1954. 5(1): p. 307-311.
128. Bacher, P., et al., The thrombolytic potency of LMW-heparin compared to urokinase in a rabbit jugular vein clot lysis model. *Thromb Res*, 1992. 66(2-3): p. 151-8.
129. Clozel, J.P., et al., Time course of thrombolysis induced by intravenous bolus or infusion of tissue plasminogen activator in a rabbit jugular vein thrombosis model. *Circulation*, 1989. 79(1): p. 125-33.

130. Collen, D., J.M. Stassen, and M. Verstraete, Thrombolysis with human extrinsic (tissue-type) plasminogen activator in rabbits with experimental jugular vein thrombosis. Effect of molecular form and dose of activator, age of the thrombus, and route of administration. *J Clin Invest*, 1983. 71(2): p. 368-76.
131. Doutremepuich, C., et al., Experimental venous thrombosis in rats treated with heparin and a low molecular weight heparin fraction. *Haemostasis*, 1983. 13(2): p. 109-12.
132. Doutremepuich, C., et al., Comparative study on the effects of a low molecular weight heparin (CY 216) and standard heparin in low dosage on experimental venous thrombosis. *Haemostasis*, 1987. 17(4): p. 201-5.
133. Henke, P.K., et al., Viral IL-10 gene transfer decreases inflammation and cell adhesion molecule expression in a rat model of venous thrombosis. *J Immunol*, 2000. 164(4): p. 2131-41.
134. Herbert, J.M., A. Bernat, and J.P. Maffrand, Importance of platelets in experimental venous thrombosis in the rat. *Blood*, 1992. 80(9): p. 2281-6.
135. Reyers, I., et al., Failure of aspirin at different doses to modify experimental thrombosis in rats. *Thromb Res*, 1980. 18(5): p. 669-74.
136. Wakefield, T.W., et al., Deep venous thrombosis in the baboon: an experimental model. *J Vasc Surg*, 1991. 14(5): p. 588-98.
137. Chi, L., et al., A novel model of venous thrombosis in the vena cava of rabbits. *J Pharmacol Toxicol Methods*, 1998. 39(4): p. 193-202.
138. Dale, G.L., et al., Non-isotopic method for quantitation of platelets and erythrocytes in experimental thrombi. *Thromb Haemost*, 1996. 75(4): p. 668-73.
139. Dupe, R.J., J. Green, and R.A. Smith, Acylated derivatives of streptokinase-plasminogen activator complex as thrombolytic agents in a dog model of aged venous thrombosis. *Thromb Haemost*, 1985. 53(1): p. 56-9.

140. Electricwala, A., J.J. Emeis, and T. Atkinson, Evaluation of epithelial tissue plasminogen activator as a thrombolytic agent in a rabbit model of venous thrombosis. *J Pharmacol Exp Ther*, 1986. 238(1): p. 254-8.
141. Lavelle, S.M. and M.M. Iomhair, The reliability of thrombotic tendency measured by intravascular wires in the rat. *Thromb Res*, 1997. 87(4): p. 353-7.
142. Matsuo, O., et al., Thrombolytic effect of single-chain pro-urokinase in a rabbit jugular vein thrombosis model. *Thromb Res*, 1986. 42(2): p. 187-94.
143. Eriksson, B.I., et al., Antithrombotic effect of two low molecular weight thrombin inhibitors and a low-molecular weight heparin in a caval vein thrombosis model in the rat. *Thromb Haemost*, 1997. 78(5): p. 1404-7.
144. Spokas, E.G. and T.C. Wun, Venous thrombosis produced in the vena cava of rabbits by vascular damage and stasis. *J Pharmacol Toxicol Methods*, 1992. 27(4): p. 225-32.
145. Underwood, M.J., et al., Quantifying the effect of locally delivered anticoagulant drugs: modification of an in vivo model of venous thrombosis. *J Vasc Res*, 1993. 30(6): p. 323-6.
146. Marsh Lyle, E., et al., Assessment of thrombin inhibitor efficacy in a novel rabbit model of simultaneous arterial and venous thrombosis. *Thromb Haemost*, 1998. 79(3): p. 656-62.
147. Holst, J., et al., Local application of recombinant active-site inhibited human clotting factor VIIa reduces thrombus weight and improves patency in a rabbit venous thrombosis model. *Eur J Vasc Endovasc Surg*, 1998. 15(6): p. 515-20.
148. McClanahan, T.B., et al., Antithrombotic effects of BCH 2763, a new direct thrombin inhibitor, in a canine model of venous thrombosis. *J Thromb Thrombolysis*, 1999. 7(3): p. 301-6.

149. Rebello, S.S., et al., CVS-1123, a direct thrombin inhibitor, prevents occlusive arterial and venous thrombosis in a canine model of vascular injury. *J Cardiovasc Pharmacol*, 1997. 29(2): p. 240-9.
150. Sigel, B., et al., Intimal hyperplasia producing thrombus organization in an experimental venous thrombosis model. *J Vasc Surg*, 1994. 19(2): p. 350-60.
151. Sudo, Y. and B.R. Lucchesi, Antithrombotic effect of GYKI-14766 in a canine model of arterial and venous rethrombosis: a comparison with heparin. *J Cardiovasc Pharmacol*, 1996. 27(4): p. 545-55.
152. Vorwerk, D., et al., [Laser thermal venous obliteration: experimental results in the rat model]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*, 1994. 160(1): p. 84-8.
153. Imbault, P., et al., Antithrombotic effects of aspirin and LMWH in a laser-induced model of arterials and venous thrombosis. *Thromb Res*, 1996. 82(6): p. 469-78.
154. Morishita, K. and M. Iwamoto, Synergistic antithrombotic effects of argatroban and ticlopidine in the rat venous thrombosis model. *Thromb Res*, 1998. 92(6): p. 261-6.
155. Millet, J., J. Theveniaux, and M. Pascal, A new experimental model of venous thrombosis in rats involving partial stasis and slight endothelium alterations. *Thromb Res*, 1987. 45(2): p. 123-33.
156. el Kouri, D., et al., Experimental venous thrombosis: what animal model? *Int Angiol*, 1992. 11(4): p. 304-8.
157. Hosaka, J., et al., Induced thrombosis in the pig inferior vena cava: a model of deep venous thrombosis. *J Vasc Interv Radiol*, 1996. 7(3): p. 395-400.
158. Tacke, J., et al., [Experimental model of early chronic pelvic vein thrombosis with occlusion spirals and thrombin]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*, 1998. 169(6): p. 645-8.

159. Landefeld, C.S., E. McGuire, and A.M. Cohen, Clinical findings associated with acute proximal deep vein thrombosis: a basis for quantifying clinical judgment. *Am J Med*, 1990. 88(4): p. 382-8.
160. Sanson, B.J., et al., Comparison of a clinical probability estimate and two clinical models in patients with suspected pulmonary embolism. ANTELOPE-Study Group. *Thromb Haemost*, 2000. 83(2): p. 199-203.
161. Wells, P.S., et al., Application of a diagnostic clinical model for the management of hospitalized patients with suspected deep-vein thrombosis. *Thromb Haemost*, 1999. 81(4): p. 493-7.
162. Wells, P.S., et al., A novel and rapid whole-blood assay for D-dimer in patients with clinically suspected deep vein thrombosis. *Circulation*, 1995. 91(8): p. 2184-7.
163. Ginsberg, J.S., et al., Sensitivity and specificity of a rapid whole-blood assay for D-dimer in the diagnosis of pulmonary embolism. *Ann Intern Med*, 1998. 129(12): p. 1006-11.
164. Fobbe, F. and K. Koppenhagen, Diagnostik der symptomatischen und asymptomatischen venösen Thrombose. *Hämostaseologie*, 1995(15): p. 148-55.
165. Kahn, S.R., et al., A randomized management study of impedance plethysmography vs. contrast venography in patients with a first episode of clinically suspected deep vein thrombosis. *Thromb Res*, 2001. 102(1): p. 15-24.
166. Wells, P.S., et al., A simple clinical model for the diagnosis of deep-vein thrombosis combined with impedance plethysmography: potential for an improvement in the diagnostic process. *J Intern Med*, 1998. 243(1): p. 15-23.
167. Jäger, K., R. Eichlisberger, and B. Frauchiger, Stellenwert der bildgebenden Sonographie für die Diagnostik der Venenthrombose. *Hämostaseologie*, 1993. 13: p. 116-124.

168. Douglas, M.G. and D.S. Sumner, Duplex scanning for deep vein thrombosis: has it replaced both phlebography and noninvasive testing? *Semin Vasc Surg*, 1996. 9(1): p. 3-12.
169. Baxter, G.M., The role of ultrasound in deep venous thrombosis. *Clin Radiol*, 1997. 52(1): p. 1-3.
170. Hach, W. and V. Hach-Wunderle, *Phlebographie der Bein- und Beckenvenen*. 4. Auflage ed. 1996, Konstanz: Schnetztor-Verlag.
171. Clauß, W., et al., Determination of Risk Factors Regarding the Administration of Contrast Media, in *Contrast Media in Practice*, W. Clauß and P. Dawson, Editors. 1994, Springer: Berlin, Heidelberg, New York. p. 71-87.
172. Dawson, P., et al., Influence of Contrast Media on Organs and Vessels, in *Contrast Media in Practice*, P. Dawson and W. Clauß, Editors. 1994, Springer: Berlin, Heidelberg, New York. p. 53-70.
173. Bettmann, M.A. and S. Paulin, Leg phlebography: the incidence, nature and modification of undesirable side effects. *Radiology*, 1977. 122(1): p. 101-4.
174. Hagen, B., Phlebography, in *Contrast Media in Practise*, P. Dawson and W. Clauß, Editors. 1994, Springer: Berlin, Heidelberg, New York. p. 133-138.
175. Bettmann, M.A., et al., Contrast venography of the leg: diagnostic efficacy, tolerance, and complication rates with ionic and nonionic contrast media. *Radiology*, 1987. 165(1): p. 113-6.
176. Lensing, A.W., et al., Lower extremity venography with iohexol: results and complications. *Radiology*, 1990. 177(2): p. 503-5.
177. Shehadi, W.H. and G. Toniolo, Adverse reactions to contrast media: a report from the Committee on Safety of Contrast Media of the International Society of Radiology. *Radiology*, 1980. 137(2): p. 299-302.

178. Shehadi, W.H., Contrast media adverse reactions: occurrence, recurrence, and distribution patterns. *Radiology*, 1982. 143(1): p. 11-7.
179. Stein, P.D., et al., Complications and validity of pulmonary angiography in acute pulmonary embolism. *Circulation*, 1992. 85(2): p. 462-8.
180. Hudson, E.R., et al., Pulmonary angiography performed with iopamidol: complications in 1,434 patients. *Radiology*, 1996. 198(1): p. 61-5.
181. Hull, R.D., et al., Pulmonary angiography, ventilation lung scanning, and venography for clinically suspected pulmonary embolism with abnormal perfusion lung scan. *Ann Intern Med*, 1983. 98(6): p. 891-9.
182. PIOPED, Value of the ventilation/perfusion scan in acute pulmonary embolism. Results of the prospective investigation of pulmonary embolism diagnosis (PIOPED). The PIOPED Investigators. *Jama*, 1990. 263(20): p. 2753-9.
183. Rathbun, S.W., G.E. Raskob, and T.L. Whitsett, Sensitivity and specificity of helical computed tomography in the diagnosis of pulmonary embolism: a systematic review. *Ann Intern Med*, 2000. 132(3): p. 227-32.
184. Lee, A.Y. and J. Hirsh, Diagnosis and treatment of venous thromboembolism. *Annu Rev Med*, 2002. 53: p. 15-33.
185. Schmitz, S.A., et al., USPIO-enhanced direct MR imaging of thrombus: preclinical evaluation in rabbits. *Radiology*, 2001. 221(1): p. 237-43.
186. Agresti, A., Modelling patterns of agreement and disagreement. *Stat Methods Med Res*, 1992. 1(2): p. 201-18.
187. Valavanis, A., *Interventional Neuroradiology*. Medical Radiology, ed. A. Valavanis. 1993, Berlin, Heidelberg, New York: Springer.

188. Freeny, P.C., et al., Long-term radiographic-pathologic follow-up of patients treated with visceral transcatheter occlusion using isobutyl 2-cyanoacrylate (Bucrylate). *Radiology*, 1979. 132(1): p. 51-60.
189. Kunnen, M., [New techniques for embolisation of the internal spermatic vein: intravenous tissue adhesive (author's transl)]. *ROFO Fortschr Geb Rontgenstr Nuklearmed*, 1980. 133(6): p. 625-9.
190. Erdman, W.A., et al., Venous thrombosis: clinical and experimental MR imaging. *Radiology*, 1986. 161(1): p. 233-8.
191. Lanzer, P., et al., Sequential 2D inflow venography: initial clinical observations. *Magn Reson Med*, 1991. 19(2): p. 470-6.
192. Evans, A.J., et al., Detection of deep venous thrombosis: prospective comparison of MR imaging and sonography. *J Magn Reson Imaging*, 1996. 6(1): p. 44-51.
193. Muhler, A., The future of contrast-enhanced magnetic resonance angiography. Are blood pool agents needed? *Invest Radiol*, 1998. 33(9): p. 709-14.
194. Nolte-Ernsting, C., et al., Abdominal MR angiography performed using blood pool contrast agents: comparison of a new superparamagnetic iron oxide nanoparticle and a linear gadolinium polymer. *AJR Am J Roentgenol*, 1998. 171(1): p. 107-13.
195. Weishaupt, D., et al., Equilibrium-phase MR angiography of the aortoiliac and renal arteries using a blood pool contrast agent. *AJR Am J Roentgenol*, 2000. 175(1): p. 189-95.
196. Grist, T.M., et al., Steady-state and dynamic MR angiography with MS-325: initial experience in humans. *Radiology*, 1998. 207(2): p. 539-44.
197. Stuber, M., et al., Contrast agent-enhanced, free-breathing, three-dimensional coronary magnetic resonance angiography. *J Magn Reson Imaging*, 1999. 10(5): p. 790-9.

198. Wielopolski, P.A., et al., Breath-hold coronary MR angiography with volume-targeted imaging. *Radiology*, 1998. 209(1): p. 209-19.
199. Dirksen, M.S., et al., Improved MR Coronary Angiography with Use of a New Rapid Clearance Blood Pool Contrast Agent in Pigs. *Radiology*, 2003. 3: p. 3.
200. Goyen, M., et al., Whole-body three-dimensional MR angiography with a rolling table platform: initial clinical experience. *Radiology*, 2002. 224(1): p. 270-7.
201. Omary, R.A., et al., MR-guided angioplasty of renal artery stenosis in a pig model: a feasibility study. *J Vasc Interv Radiol*, 2000. 11(3): p. 373-81.
202. Wacker, F.K., et al., MR image-guided endovascular procedures with the ultrasmall superparamagnetic iron oxide SH U 555 C as an intravascular contrast agent: study in pigs. *Radiology*, 2003. 226(2): p. 459-64.
203. Frank, H., et al., Efficacy of thrombolytic therapy in pulmonary embolism determined by MION-enhanced MRA: an experimental study in rabbits. *Invest Radiol*, 1998. 33(12): p. 853-7.