

## 7. Literaturverzeichnis

Abedi H, Zachary I (1997) Vascular endothelial growth factor stimulates tyrosine phosphorylation and recruitment to new focal adhesions of focal adhesion kinase and paxillin in endothelial cells. *J Biol Chem*; 272:15442-15451

Ahmed A, Dunk C, Kniss D, Wilkes M (1997) Role of VEGF receptor-1 (FLT-1) in mediating calcium dependent nitric oxide release and limiting DNA synthesis in human trophoblast cells. *Lab Invest*; 76:779-791

Asahara Y, Takshashi T, Masuda H et al. (1999) VEGF contributes to postnatal neovascularization by mobilizing bone marrow-derived endothelial progenitor cells. *EMBO J*; 18:3964-3972

Ashworth JR, Warren A, Johnson I, Baker P (1998) Plasma from pre-eclamptic women induces a functional change in myometrial resistance arteries. *Br J Obstet Gynecol*; 105: 459-461

Baker PN, Krasnow L, Roberts J, Yeo K (1995) Elevated serum levels of VEGF in patients with PE. *Obstet Gynecol*; 86:815-821

Baker PN, Davidge ST, Roberts JM (1996) Plasma from women with preeclampsia increases endothelial cell nitric oxide production. *Hypertension*; 26:56-61

Baker PN, Davidge ST, Pitt BR (1995) Plasma from preeclamptic women increases vascular smooth muscle cell nitric oxide production. *J Soc Gynecol Invest*; 2:178

Barleon B, Siemeister G, Martiny-Baron G, Weindel K, Herzog C, Marmé D (1997) Vascular endothelial growth factor up-regulates its receptor fms-like tyrosine kinase 1 (FLT-1) and a soluble variant of FLT-1 in human vascular endothelial cells. *Cancer Research*; 57:5421-5425

Barleon B, Sozzani S, Zhou D, Weich H, Mantovani A, Marmé D (1996) Migration of human monocytes in response to VEGF is mediated via the VEGF receptor FLT-1. *Blood*; 87:3336-3343

Brockelsby J, Hayman R, Ahmed A, Warren A, Johnson I, Baker P (1999) VEGF mimics preeclamptic plasma in inhibiting uterine blood vessel relaxation in pregnancy: Implication in the pathogenesis of preeclampsia. *Lab Invest*; 79:1101-1111

Carmeliet C et al. (1996) Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. *Nature*; 380:435-439

Charnock-Jones S, Sharkey AM, Boocock CA, Ahmed A, Plevin R, Ferrara N, Smith SK (1994) VEGF receptor localization and activation in human trophoblast and choriocarcinoma cells. *Biology of Reproduction*; 51:524-530

Clark BA, Halvorson L, Sachs B, Epstein FH (1992) Plasma endothelin levels in preeclampsia: Elevation and correlation with uric acid levels and renal impairment. *Am J Obstet Gynecol*; 166:962-968

- Clark DE, Smith SK, He Y et al. (1998) A vascular endothelial growth factor antagonist is produced by the human placenta and released into the maternal circulation. *Biology of Reproduction*; 59:1540-1548
- Clark P, Boswell F, Greer I (1998) The neutrophil and preeclampsia. *Seminars in Reproductive Endocrinology*; Vol 16
- Clauss M, Weich H, Breier G, Knies U, Roeckl W, Waltenberger J, Risau W (1996) Vascular endothelial growth factor receptor FLT-1 mediates biological activities- implications for a functional role of placenta growth factor in monocyte activation and chemotaxis. *J Biol Chem* 271:17629-17634
- Clauss M, Gerlach M, Gerlach H (1990) Vascular permeability factor: A tumor derived polypeptide that induces endothelial cell and monocyte procoagulant activity and promotes monocyte migration. *J Exp Med*; 172:1535-1545
- Cockell AP, Learmont JG, Smáráson AK et al. (1997) Human placental syncytiotrophoblast microvillous membranes impair maternal vascular endothelial function. *Br J Obstet Gynecol*; 104:235-240
- Das SK, Chakraborty I, Wang J (1997) Expression of VEGF and VEGF-receptor m-RNA in the periimplantation rabbit uterus. *Biol Reprod*; 56(6):1390-9
- Davidge ST, Baker PN, Roberts JM (1995) NOS expression is increased in endothelial cells exposed to plasma from women with preeclampsia. *Am J Physiol*; 269:H1106-1112
- Dekker G, Sibai B (1999) The immunology of preeclampsia. *Seminars in Reproductive Endocrinology*; Vol 23
- De Wolf F, Robertson WB, Brosens I (1975) The ultrastructure of acute atherosclerosis in hypertensive pregnancy. *Am L Obstet Gynecol*; 123:164-174
- Dietl J (2001) *Molekulare Aspekte der Implantation*. 9. Deutsches Gestose-Symposium, Freiburg
- Ezimokhai M, Aloamaka CP, Morrison J (1995) Effect of plasma from preeclamptic subjects on contractions of rat aorta. *Hypertension in Pregnancy*;14:39-48
- Ferrante A, Thong YH (1980) Optimal conditions for simultaneous purification of mononuclear and polymorphonuclear leucocytes from human blood by the hypaque-ficoll method. *J Immunol Methods*; 36(2):109-117
- Ferrara N, Henzel WJ (1989) Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells. *Biochem Biophys Res Commun*; 161:851-855
- Ferrara N, Carver-Moore K, Chen H et al. (1996) Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature*; 380:439-442
- Gargett CE, Zaitseva M, Bucak K (2002) 17Beta-estradiol up-regulates VEGF receptor-2 expression in human myometrial microvascular endothelial cells: role of estrogen receptor-alpha and -beta. *J Clin Endocrinol Metab*; 87(9):4341-9

Gerber HP, Condorelli F, Park J, Ferrara N (1997) Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes. FLT-1 but not FLK-1/KDR is up-regulated by hypoxia. *J Biol Chem*; 272:23659-23667

Graubert M, Ortega M, Kessel B (2001) Vascular repair after menstruation involves regulation of VEGF-receptor phosphorylation by sFLT-1. *Am J of Pathology*; 158:1399-1410

Guo D, Jia Q, Song H-Y, Warren RS, Donner DB (1995) Vascular endothelial growth factor promotes tyrosine phosphorylation of mediators of signal transduction that contain SH<sub>2</sub> domains. *J Biol Chem*; 270:6729-6733

Haeger M, Unander M, Norder-Hansson B, Tylman M, Bengtsson A (1992) Complement, neutrophil, and macrophage activation in women with severe preeclampsia and the syndrome of hemolysis, elevated liver enzymes, and low platelet count. *Obstet Gynecol*; 79:19-26

Hiratsuka S, Minowa O, Kuno J, Noda T, Shibuya M (1998) FLT-1 lacking the tyrosine kinase domain is sufficient for normal development and angiogenesis in mice. *Proc Natl Acad Sci*; 95:9349-9354

Hofbauer R, Hofbauer L, Moser D, Salfinger H, Speiser W, Frass M, Kapiotis S (1998) Leukocyte migration: a new triple migration chamber assay allows investigation of various cell interactions simultaneously. *Life Science*; 63:1725-1737

Hunter A, Aitkenhead M, Caldwell C, (2000) Serum levels of vascular endothelial growth factor in preeclamptic and normotensive pregnancy. *36:965-969*

Jakeman L, Armanini M, Phillips HS, Ferrara N (1993) Developmental expression of binding sites and messenger ribonucleic acid for vascular endothelial growth factor suggests a role for this protein in vasculogenesis and angiogenesis. *Endocrinology*; 133:848-859

Joukov V, Pajusola K, Kaipainen A (1996) A novel vascular endothelial growth factor, VEGF-C, is a ligand for the FLT-4 and KDR receptor tyrosine kinases. *EMBO J*; 15: 290-298

Keck, Neulen, Breckwolfdt u.a.: *Endokrinologie und Reproduktionsmedizin*, Stuttgart 1997  
S. 25

Kendall R, Thomas K (1993) Inhibition of vascular endothelial cell growth activity by an endogenously encoded soluble receptor. *Proc Natl Acad Sci*; 90:10705-10709

Kendall R, Wang G, Thomas K (1996) Identification of a natural soluble form of the vascular endothelial growth factor receptor FLT-1 and its heterodimerization with KDR. *Biochem Biophys Res Commun*; 226:324-328

Kroll J, Waltenberger J (1998) VEGF-A induces expression of eNOS and iNOS in endothelial cells via VEGF receptor-2 (KDR). *Biochem Biophys Res Commun*; 252:743-746

Kroll J, Waltenberger J (1999) A novel function of VEGF receptor-2 (KDR): rapid release of nitric oxide in response to VEGF-A stimulation in endothelial cells. *Biochem Biophys Res Commun*; 265:636-639

- Krüssel JS, Casan EM, Raga F, Hirchenhain J (1999) Expression of mRNA for VEGF transmembraneous receptors FLT-1 and KDR, and the soluble receptor sFLT in cycling human endometrium. *Molecular Human Reproduction*, Vol. 5, No. 5, 452-458
- Kupferminc MJ, Daniel Y, Englender T, Baram A, Many a, Jaffa AJ, Gull I, Lessing JB (1997) VEGF is increased in patients with PE. *Am J Reprod Immunol*; 38:302-306
- Leung DW, CachianesG, Kuang WJ, Goeddel DV, Ferrara N (1989) Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science*;246:1306-1309
- Levy AP, Levy NS, Wegner S, Goldberg MA (1995) Transcriptional regulation of the rat vascular endothelial growth factor gene by hypoxia. *J Biol Chem*; 270:13333-13340
- Levy NS, Chung S, Furneaux H, Levy AP (1998) Hypoxic stabilization of vascular endothelial growth factor mRNA by RNA-binding protein HuR. *J Biol Chem*;273:6417-6423
- Lichtenberger C, Zakeri S, Baier K, Willheim M, Holub M, Reinisch W (1999) A novel high-purity isolation method for human peripheral blood neutrophils permitting polymerase chain reaction-based mRNA studies. *J Immunol Methods*; 227:75-84
- Lincoln DW, Phillips PG, Bove K (2003) Estrogen-induced Ets-1 promotes capillary formation in *in vitro* tumor angiogenesis. *Breast Cancer Res Treat*, 78(2):167-78
- Lyall F, Greer IA, Boswell F, FlemingR (1997) Suppression of serum VEGF immunoreactivity in normal pregnancy and preeclampsia. *Brit J Obstet Gynecol*; 104:223-228.
- Maglione D, Guerriero V, Viglietto G, Delli-Bovi P, Persico MG (1991) Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc Natl Acad Sci*; 88:9267-9271
- Matsuura A, Kawashima S, Yamochi W, Hirata K, Yamaguchi T, Emoto N, Yokohama M (1997) VEGF increases endothelin-converting enzyme expression in vascular endothelial cells. *Biochem Biophys Res Commun*; 235: 713-716
- McCarthy AL, Woolfson RG, Raju SK, Poston L (1993) Abnormal endothelial cell function of resistance arteries in women with pre-eclampsia. *Am J Obstet Gynecol*; 168:1323-1330
- McCartney CP (1969) The acute hypertensive disorders of pregnancy, classified by renal histology. *Gynaecologia* 167:214-220
- Meduri G, Bausero P, Perrot-Applanat M (2000) Expression of VEGF receptors in the human endometrium: modulation during the menstrual cycle. *Biol Reprod* 62, 439-447
- Mellembakken JR, Aukrust P, Hetdal K, Ueland T, Abyholm T, Videm V (2001) Chemokines and leukocyte activation in the fetal circulation during preeclampsia. *Hypertension*; 38:394-398
- Mimeault M (2000) Structure-function studies of ets transcription factors. *Critical Reviews in Oncogenesis*; 11:227-253

- Mitola S, Sozzani S, et al. (1997) Tat-human immunodeficiency virus-1 induces human monocyte chemotaxis by activation of vascular endothelial growth factor receptor-1. *Blood*;90:1365-1372
- Morishita K, Johnson D, Williams L (1995) A novel promoter for vascular endothelial growth receptor (FLT-1) that confers endothelial-specific gene expression. *J Biol Chem*; 17:27948-27953
- Myatt L, Miodovnik M (1999) Prediction of preeclampsia. *Seminars in Perinatology*; Vol 23; 45-57
- Nadji P, Sommers SC (1973) Lesions of toxemia in first trimester pregnancies. *Am J Clin Pathol*; 59:344-348
- Neufeld G, Cohen T, Gengrinovitch S, Poltorak Z (1999) VEGF and its receptors. *FASEB J*; 13:9-22
- Nova A, Sibai BM, Barton JR (1991) Maternal plasma levels of endothelin is increased in preeclampsia. *Am J Obstet Gynecol*; 165:724-727
- Olofsson B, Pajusola K, Kaipainen A, von Euler G, Joukov V, Saksela O, Orpana A, Pettersson RF, Alitalo K, Eriksson U (1996) Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proc Natl Acad Sci USA*; 93:2576-2581
- Pfleiderer, Breckwolfdt u.a.: *Gynäkologie und Geburtshilfe*, Stuttgart 2002, S. 57
- Pirani BB, MacGillivray I (1975) The effect of plasma transfusion on the blood pressure in pregnancy. *Am J Obstet Gynecol*; 121:221-226
- Rizoli S, Kapus A, Parodo J, Fan J, Rotstein O (1999) Hypertonic immunomodulation is reversible and accompanied by changes in CD 11b expression. *J Surg Res*; 83:130-135
- Roberts J (1998) Endothelial dysfunction in preeclampsia. *Seminars in Reproductive Endocrinology*; Vol 16
- Sabatier F, Bretelle F, d'Ercole C, Boubli L, Sampol J, Dignat-George F (2000) Neutrophil activation in preeclampsia and isolated intrauterine growth restriction. *Am J Obstet Gynecol*; 183:1558-1563
- Sawano A, Iwai S, Sakurai Y, Ito M, Shitara K, Nakahata T, Shibuya M (2001) FLT-1, vascular endothelial growth factor receptor 1, is a novel cell surface marker for lineage of monocyte-macrophages in human. *Blood*; 97:785-791
- Shalaby F, Rossant J, Yamaguchi TP et al. (1995) Failure of blood-island formation and vasculogenesis in FLK-1-deficient mice. *Nature*; 376:62-66
- Shanklin DR, Sibai BM (1990) Ultrastructural aspects of preeclampsia. II. Mitochondrial changes. *Am J Obstet Gynecol*; 163:943-953

- Sharkey AM, Cooper JC, Balmforth JR, Mc Laren J (1996) Maternal plasma levels of VEGF in normotensive pregnancies and in pregnancies complicated by PE. *Eur J Clin Invest*; 26:1182-1185
- Shen H, Clauss M, Ryan J, Schmidt A-M, Tjiburg P et al. (1993) Characterisation of vascular permeability factor/ vascular endothelial growth factor receptors on mononuclear phagocytes. *Blood*; 10:2767-2773
- Shibuya M (1995) Role of VEGF-FLT receptor system in normal and tumor angiogenesis. *Adv. Cancer Res.*; 67:281-316
- Silver RK, Kupferminc MJ, Russel TL et al (1996) Evaluation of nitric oxide as a mediator of severe preeclampsia. *Am J Obstet Gynecol*; 175:1013-1017
- Smáráson AK, Sargent IL, Starkey PM et al. (1993) Effects of placental syncytiotrophoblast microvillous membranes from pre-eclamptic women on the growth of endothelial cells *in vitro*. *Br J Obstet Gynecol*; 100:943-949
- Spargo BH, Lichtig C, Luger AM (1976) The renal lesion in preeclampsia: Examination by light-, electron- and immunofluorescence-microscopy. *Hypertension in Pregnancy*: 129-137
- Sugino N, Kashida S, Karube-Harada A (2002) Expression of VEGF and its receptors in human endometrium throughout the menstrual cycle and in early pregnancy. *Reproduction*; 123(3): 379-87
- Suzuma I, Mandai M, Tagaki H (1999) 17 Beta-estradiol increases VEGF receptor-2 and promotes DNA synthesis in retinal microvascular endothelial cells. *Invest Ophthalmol Vis Sci*; 40(9): 2122-9
- Taylor A, Granger N (2000) Role of adhesion molecules in vascular regulation and damage. *Current Hypertension Reports*; 2:78-83
- Takahashi T, Shibuya M (1997) The 230 kDa mature form of KDR/FLK-1 activates the PLC-gamma pathway and partially induces mitotic signals in NIH3T3 fibroblasts. *Oncogene*; 14: 2079-2089
- Taylor RN, Varma M, Teng NNH (1990) Women with preeclampsia have higher plasma endothelin levels than women with normal pregnancies. *J Clin Endocrinol Metab*; 71:1675-1677
- Taylor RN, de Groot C, Cho YK, Lim K-H (1998) Circulating factors as markers and mediators of endothelial cell dysfunction in preeclampsia. *Seminars in Reproductive Endocrinology*; Vol 16
- Tsukimori K, Maeda H, Ishida K, Nagata H, Koyanagi T, Nakano H (1993) The superoxide generation of neutrophils in normal and preeclamptic pregnancies. *Obstet Gynecol*; 81:536-540
- Valter M, Hügel A, Huang H-J, Cavenee W, Wiestler O, Pietsch T, Wernert N (1999) Expression of the ets-1 transcription factor in human astrocytomas is associated with fms-like tyrosine kinase-1 (FLT-1) synthesis and neoangiogenesis. *Cancer Res*; 59:5608-5614

Vedernikov Y, Saade G, Garfield RE (1999) Vascular reactivity in preeclampsia. *Seminars of Perinatology*; Vol 23:34-44

von Dadelszen P, Sargent IL, Starkey PM, Redman C (1997) Increased intracellular calcium in maternal leukocytes in preeclampsia measured using fluo-3 by flow cytometry. *Hypertension*; 16:300 (abstract)

von Dadelszen P, Hurst G, Redman C (1999) Supernatants from co-cultured endothelial cells and syncytiotrophoblast microvillous membranes activate peripheral blood leukocytes *in vitro*. *Human Reproduction*;14:919-924

von Dadelszen P, Watson RWG, Noorwali F et al. (1999) Maternal neutrophil apoptosis in normal pregnancy, preeclampsia, and normotensive intrauterine growth restriction. *Am J Obstet Gynecol*; 181:408-414

Vuorela-Vepsäläinen P, Alfthan H, Orpana A, Alitalo K, Stenman U-H, Halmesmäki E (1998) Vascular endothelial growth factor is bound in amniotic fluid and maternal serum. *Human Reproduction*, 14:1346-1351

Wakiya K, Begue A, Stehelin D, Shibuya M (1996) A cAMP response element and an ets motif are involved in the transcriptional regulation of FLT-1 tyrosine kinase gene. *J Biol Chem*; 48:30823-30828

Waltenberger J, Claesson-Welsh L, Siegbahn A, Shibuya M, Heldin C-H (1994) Different signal transduction properties of KDR and FLT-1, two receptors for VEGF. *J Biochem Chem*; 269:26988-26995

Waltenberger J, Lange J, Kranz A (2000) Vascular endothelial growth factor-A-induced chemotaxis of monocytes is attenuated in patients with diabetes mellitus. *Circulation*; 102:185-190

Walzog B, Weinmann P, Jeblonski F, Scharfetter-Kochanek K, Bommert K, Gaehtgens P (1999) A role for  $\beta_2$  integrins (CD11/CD18) in the regulation of cytokine gene expression of polymorphnuclear neutrophils during the inflammatory response. *FASEB J*; 13:1855-1865

Webb N, Myers C, Watson C, Bottomley M, Brenchley P (1998) Activated human neutrophils express VEGF. *Cytokine*; 10:254-257

Zachary I (1989) VEGF: how it transmits its signal. *Exp Nephrol*; 6:480-487

Zeek PM, Assali NS (1950) Vascular changes in the decidua associated with eclamptogenic toxemia. *Am J Clin Pathol*; 20:1099-1109

Ziche M, Morbidelli L, Choudhuri R, Zhang H-T, Donnini S, Granger HJ, Bicknell R (1997) Nitric oxide synthase lies downstream from vascular endothelial growth factor-induced but not basic fibroblast growth factor-induced angiogenesis. *J Clin Invest*; 99:2625-2634