

## 8 References

1. Mathew R, Thilenius OG. The syndrome of familial defects of heart and upper extremities(Holt-Oram syndrome). Circulation 1996; 34:65
2. Romero T, Covell J, Friedman WF. A comparison of pressure-volume relations of the fetal newborn and adult heart. Amer. J. Physiol 1972; 222:1285
3. Tikoff G, Schmidt AM, Kuida H, et al. Heart failure in atrial septal defect. Amer.J.Med 1965; 39: 533
4. Arnfred E. Symptoms, signs and hemodynamics in one hundred cases of atrial septal defect confirmed by operation. J Cardior. Surg 1966; 7:349
5. Wahl A, Windecker S, Meier B. Patent foramen ovale: pathophysiology and therapeutic in symptomatic patients. Minerva Cardioangiologica 2001 Dec; 49(6): 403-11
6. Rodriguez CJ, Homma S. Patent Foramen Ovale and Stroke. Curr Treat Options Cardiovasc Med
7. Castro SD, Cartoni D, Fiorelli M, et al. Morphological and functional characteristics of patent foramen ovale and their embolic implications. Stroke 2000; 31: 2407-2413
8. Webster MWI, Chancellor AM, Smith HJ, et al. Patent foramen ovale in young stroke patients. Lancet 1988; 2: 11-12
9. Lechat P, Mas JL, Lascault G, et al. Prevalence of patent foramen ovale in patients with stroke. N Engl J Med 1988; 318: 1148-52
10. Di Tullio M, Sacco RL, Gopal A, et al. Patent foramen ovale as a risk factor for cryptogenic stroke. Ann Intern Med 1992; 117: 461-5
11. De Belder MA, Tourikis L, Leech G, et al. Risk of patent foramen ovale for thromboembolic events in all age groups. Am J Cardiol 1992; 69: 1316-20
12. Hausmann D, Mügge A, Becht I, et al. Diagnosis of patent foramen ovale by transesophageal echocardiography and association with cerebral and peripheral embolic events. Am J Cardiol 1992; 70: 668-72
13. Cohnheim J. Thrombose und embolie, Vorlesungen über allgemeine pathologie; ein handbuch fur aertze und studierende. Berlin, Hirschwald 1877; 1,134: 144-145

14. Chaikof EL, Campbell BE, Smith RB. Paradoxical embolism and acute arterial occlusion: rare or unsuspected? *J Vasc Surg* 1994; 20: 377-8
15. Mas JL, Zuber M et al. Recurrent cerebrovascular events in patients with patent foramen ovale, atrial septal aneurysm, or both and cryptogenic stroke or transient ischemic attack. *Am Heart J.* 1995 ; 130 : 1083-1088
16. Bogousslavsky J, Garazi S, Jeanrenaud X, et al. Stroke recurrence in patients with patent foramen ovale : the Lausanne Study. *Neurology.* 1996 ; 46 : 1301-1305
17. Cabanes L, Mas JL, Cohen A, et al. Atrial septal aneurysm and patent foramen ovale as risk factors for cryptogenic stroke in patients less than 55 years of age. *Stroke* 1993; 24: 1855-73
18. Herwig W. Schuchlenz, Wolfgang Weihs, Susanne Horner, et al. The association between the diameter of a patent foramen ovale and the risk of embolic cerebrovascular events. *Am J Med.*2000; 109: 456-462
19. Stein MM, Di Tullio MR, Rundek T, et al. Patent foramen ovale size and embolic brain imaging findings among patients with ischemic stroke. *Strke* 1998; 29: 944-948
20. Serena J, Segura T, Perez-Ayuso MJ, Bassaganyas J,et al. The need to quantify right-to-left shunt in acute ischemic stroke. *Stroke* 1998; 29: 1322-1328
21. Van Camp G, Schulze D, Cosyns B, et al. Relation between patent foramen ovale and unexplained stroke. *Am J Cardiol.* 1993; 71: 596-598
22. Stone DA, Godard J, Correti MC, et al. Patent foramen ovale: association between the degree of shunt by contrast transesophageal echocardiography and the risk of future ischemic neurologic events. *Am Heart J.* 1996; 131: 158-161
23. Natanzo A, Goldman ME. Patent foramen ovale: anatomy versus pathophysiology—which determines stroke risk? *J Am Soc Echocardiogr.* 2003; 16: 71-76
24. Agmon Y, Khandheira BK, Meissner I, et al. Frequency of atrial septal aneurysm in patients with cerebral ischemic events. *Circulation.* 1999; 99: 1942-1944
25. Caplan LR. Diagnosis and treatment of ischemic stroke. *JAMA.* 1991; 266: 2413-2418
26. Sherman DG, Dyken ML, Gent M, et al. Antithrombotic therapy for cerebrovascular disorders: an update. *Chest.* 1996; 104(suppl): 444S-456S

27. Shunichi Homma, Marco R.Di Tullio, Ralph L, et al. Surgical closure of patent foramen ovale in cryptogenic stroke patients. *Stroke* 1997; 28: 2376-2381
28. Ruchat P, Bogousslavsky J, Hurni M, et al. Systemic surgical closure of patent foramen ovale in selected patients with cerebral events due to paradoxical embolism. Early results of a preliminary study. *European Journal of Cardio-thoracic Surgery* 1997; 11: 824-827
29. Travis JA, Fuller SB, Ligush J Jr et al. Diagnosis and treatment of paradoxical embolus. *Vasc Surg.* 2001 Nov;34(5):860-5
30. Dearani JA, Ugurlu BS, Danielson GK et al. Surgical patent foramen ovale closure for prevention of paradoxical embolism-related cerebrovascular ischemic events. *Circulation* 1999 Nov 9;100(19 Suppl):II171-5
31. Joseph A. Dearani, Baran S, Gordon K, et al. Surgical patent foramen ovale closure for prevention of paradoxical embolism-related cerebrovascular ischemic events. *Circulation* 1999; 100(suppl II): II-171—II-175
32. Chaturvedi RR, Lincoln C, Gothard JW, et al. Left ventricular dysfunction after open repair of simple congenital heart defect in infants and children: quantitation with the use of a conductance catheter immediately after bypass. *J Thorac Cardiovasc Surg.* 1998; 115: 77-83
33. Hammarström E, Wranne B, Pinto FJ, et al Tricuspid annulus motion. *J Am Soc Echocardiogr* 1998; 4: 131-139
34. Jones CJL, Song GJ, Gibson DG. An echocardiographic assessment of atrial mechanical behaviour. *Br Heart J* 1991; 65: 215-220
35. Pinto FJ, Wranne B, St Goar FG, et al. Systemic venous flow during cardiac surgery examined by intraoperative transesophageal echocardiography. *Am J Cardiol* 1991; 17: 1493-1498
36. Wranne B, Pinto FJ, Siegel LC, et al. Abnormal postoperative interventricular motion: new intraoperative transesophageal echocardiographic evidence supports a novel hypothesis. *Am Heart J* 1993; 126: 161-167
37. Hauseus KC, Björkhem, Brodin et al. LA, Analysis of atrioventricular plane movements by Doppler tissue imaging and M-mode in children with atrial septal defects before and after surgical and device closure. *Pediatric Cardiology* 2002; 23: 152-159
38. King TD, Thompson SL, Steiner C, et al. Secundum atrial septal defects. Nonoperative closure during cardiac catheterization. *JAMA* 1976; 235: 2506-9

39. Mills NL, King TD. Nonoperative closure of left-to right shunts. *J Thorac Cardiovasc Surg* 1976; 72: 371-8
40. Sideris EB, Rao PS. Transcatheter closure of atrial septal defects: role of buttoned devices. *J Invasive Cardiol* 1996; 8: 289-96
41. Lock JE, Rome JJ, Davis R, et al. Transcatheter closure of atrial septal defects. Experimental studies. *Circulation* 1989; 79: 1091-9
42. Sideris EB, Sideris SE, Fowlkes JP, et al. Transvenous atrial septal defect occlusion in piglets with a buttoned double-disk device. *Circulation* 1990; 81: 312-8
43. Du ZD, Hijazi ZM, Kleinman CS, et al. Comparison between transcatheter and surgical closure of secundum atrial septal defect in children and adults. *J Am Coll Cardiol* 2002, June 5; 39(11) : 1836-44
44. Sievert H, Horvath K, Zadan E, et al. Patent foramen ovale closure in patients with transient ischemia attack/stroke. *J Interv Cardiol* 2001 April; 14(2): 261-6
45. Rao PS, Sideris EB, Hausdorf G, et al. International experience with secundum atrial septal defect occlusion by the button device, *Am Heart J* 1994; 128: 1022-1035
46. Worms AM, Rey C, Bourlon F, et al. French experience in the closure of atrial septal defects of the ostium secundum type with the Sideris button occluder. *Arch Mal Coeur Vaiss* 1996; 89: 509-515
47. Agarwal SK, Ghosh PK, Mittal PK, et al. Failure of devices used for closure of atrial septal defects: mechanisms and management. *J Thorac Cardiovasc Surg* 1996; 112: 21-26
48. Sievert H, Babic UU, Ensslen R, et al. Transcatheter closure of large atrial septal defects with the Babic system. *Cathet Cardiovasc Diagn* 1995; 36: 232-240
49. Bohm J, Bittigau K, Kohler F, et al. Surgical removal of atrial septal defect occlusion system-devices. *Eur J Cardiothorac Surg* 1997; 12: 869-872
50. Trepel T, Zeplin H, Sievert H, et al. Cardiac perforation following transcatheter PFO closure. *Cathet Cardiovasc Intervent* 2003; 58: 111-113
51. Onose Y, Oki T, Tabata T, et al. Assessment of the temporal relationship between left ventricular relaxation and filling during early diastole using pulsed Doppler echocardiography and tissue Doppler imaging. *Jpn Circ J* 1999; 63: 209- 215

52. Kitabatake A, Inoue M, Asao M, et al. Transmitral blood flow reflecting diastolic behaviour of the left ventricle in health and disease : a study by pulsed Doppler technique. *Jpn Circ J* 1982; 46: 92-102
53. Oki T, Fukuda N, Iuchi A, et al. Evaluation of left ventricular diastolic hemodynamics from the left ventricular inflow and pulmonary venous flow velocities in hypertrophic cardiomyopathy. *Jpn Heart J* 1995; 36: 617-627
54. Ishida Y, Meisner JS, Tsujioka K, Gallo JH, et al. Left ventricular filling dynamics: influence of left ventricular relaxation and left atrial pressure. *Circulation* 1986; 74: 187-196
55. Myreng Y, Smiseth OA. Assessment of left ventricular diastolic function by Doppler echocardiography: comparison of isovolumic relaxation. *Circulation* 1990; 81: 260-266
56. Thomas JD, Flachskampf FA, Chen C, et al. Isovolumic relaxation time varies predictably with its time constant and aortic and left atrial pressure: implications for the noninvasive evaluation of ventricular relaxation. *Am Heart J* 1992; 124:1305-1313
57. Issaz K, Thomson A, Palka P, et al. Doppler echocardiographic measurements of flow velocity motion of the left ventricular posterior wall. *Am J Cardiol* 1989; 64: 66-75
58. McDicken WN, Sutherland GR, Gordon LN. Color Doppler velocity imaging of the myocardium. *Ultrasound Med Biol* 1992; 18: 651-654
59. Miyatake K, Yamagishi M, Tanaka N, et al. New method for evaluating left ventricular wall motion by color-coded tissue Doppler imaging : in vitro and in vivo study. *J Am Coll Cardiol* 1995; 25: 717-724
60. Donovan CL, Armstrong WF, Bach DS. Quantitative Doppler tissue imaging of the left ventricular myocardium: validation in normal subjects. *Am Heart J* 1995; 130: 100-104
61. Galiuto L, Ignone G, Demarid AN. Contraction and relaxation velocities of the normal left ventricle using pulsed-wave tissue Doppler echocardiography. *Am J Cardiol* 1998; 81: 609-14
62. Palka P, Lange A, Fleming AD, et al. Age-related transmural peak mean velocities and peak velocity gradients by Doppler myocardial imaging in normal subjects. *Eur Heart J* 1996; 17: 940-50

63. Tasneem Z, Naqvi G, Neyman A, Broyde A, et al. Comparison of myocardial tissue Doppler with transmural flow Doppler in left ventricular hypertrophy. *J Am Soc Echocardiogr* 2001; 14(II): 1153-1160
64. Isaaz K, Munoz del Romeral L, Lee E, et al. Quantitation of the motion of the cardiac base in normal subjects by Doppler echocardiography. *J Am Soc Echocardiogr* 1993; 6:166-76
65. Garcia M, Thomas JD, Klein AL. New Doppler echocardiographic applications for the study of diastolic function. *J Am Coll Cardiol* 1998; 32: 865-75
66. Rodriguez L, Garcia M, Ares M, et al. Assessment of mitral annular dynamics during diastole by Doppler tissue imaging : comparison with mitral Doppler inflow in subjects without heart disease and in patients with left ventricular hypertrophy. *Am Heart J* 1996; 131: 982-7
67. Severino S, Caso P, Galderisi M, et al. Use of pulsed Doppler tissue imaging to assess regional left ventricular diastolic dysfunction in hypertrophic cardiomyopathy. *Am J Cardiol* 1998; 82: 1394-
68. Sohn DW, Chai IH, Lee DJ, et al. Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic function. *J Am Coll Cardiol* 1997; 30: 474-80
69. Pai GR, Gill KS. Amplitudes, durations, and timings of apical directed left ventricular myocardial velocities: II. Systolic and diastolic asynchrony in patients with left ventricular hypertrophy. *J Am Soc Echocardiogr* 1998; 11: 112-8
70. Heimdal A, Stoylen A, Torp H, et al. Real-time strain rate imaging of the left ventricle by ultrasound. *J Am Soc Echocardiogr* 1998; 11:1013-9.
71. Uematsu M, Miyatake K, Tanaka N, et al. Myocardial velocity gradient as a new indicator of regional left ventricular contraction: detection by a two-dimensional tissue Doppler imaging technique. *J Am Coll Cardiol* 1995; 26: 217-223
72. Jun K, Patricia RS, Rodney HF. Longitudinal myocardial function assessed by tissue velocity, strain, and strain rate tissue Doppler echocardiography in patients with AL(primary) cardiac amyloidosis. *Circulation* 2003;107(19): 2446-2452.
73. Greenberg NL, Firstenberg MS, Castro PL, et al. Doppler-derived myocardial systolic strain rate is a strong index of left ventricular contractility. *Circulation* 2002; 105: 99-105.

74. Zahn EM, Neil Wilson, Warren Cutright, et al. Development and testing of the Helex septal occluder, a new expanded polytetrafluoroethylene atrial septal defect occlusion system. *Circulation* 2001;104(6):711
75. Thompson T, Evans W. Paradoxical embolism. *Quart J Med* 1930; 23: 135-150
76. Hart R, Miller V. Cerebral infarction in young adults: a practical approach. *Stroke* 1983; 14: 100-114
77. Connheim J. Thrombose und Embolie: Vorlesung ueber allgemeine Pathologie, Berlin: Hirschwald; 1877. P175-177
78. King TD, Mills NL. Secundum atrial septal defect: nonoperative closure during cardiac catheterization. *JAMA* 1976; 235: 2506-2509
79. Windecker S, Wahl A, Chatterjee T, et al. Percutaneous closure of patent foramen ovale in patients with paradoxical embolism: long-term risk of recurrent thromboembolic events. *Circulation* 2000; 101: 893-898
80. Bogousslavsky J, Devuyst G, Nendaz M, et al. Prevention of stroke recurrence with presumed paradoxical embolism. *J Neurol* 1997; 244: 71-75
81. Sievert H. Interventional versus surgical closure of atrial and ventricular septal defects: advantages and limitations of the catheter-based approach. *J Intervent Cardiol* 2000; 13: 493-501
82. Formigari R, Di Donato RM, Mazzera E, et al. Minimally invasive or interventional repair of atrial septal defects in children: experience in 171 cases and comparison with conventional strategies. *J Am Coll Cardiol* 2001; 37: 1707-1712
83. Lee CH, Kwok OH, Fan K, et al. Transcatheter closure of atrial septal defect using Amplatzer septal occluder in Chinese adults. *Cathet Cardiovasc Intervent* 2001; 53: 373-377
84. Simonson JS, Schiller NB. Descent of the base of the left ventricle: an echocardiographic index of left ventricular function. *J Am Soc Echocardiogr* 1989; 2: 25-35
85. Nagueh SF, Middleton KJ, Kopelen HA, et al. Doppler tissue imaging : a technique for evaluation of left ventricular relaxation and estimation of filling pressure. *J Am Coll Cardiol* 1997; 30: 1527-33
86. Yamada H, Oki T, Tabata T, et al. Assessment of left ventricular systolic wall motion velocity with pulsed tissue Doppler imaging: comparison with peak dp/dt of the left ventricular pressure curve. *J Am Soc Echocardiogr* 1998; 11: 442-449

87. Oki T, Tabata T, Yamada H, et al. Clinical application of pulsed Doppler tissue imaging for assessment of abnormal left ventricular relaxation. *Am J Cardiol* 1997; 79: 921-8
88. Farias CA, Rodriguez L, Garcia MJ, et al. Assessment of diastolic function by tissue Doppler echocardiography: comparison with standard transmitral and pulmonary venous flow. *J Am Soc Echocardiogr* 1999; 12: 609-617
89. Ohte N, Narita H, Hashimoto T, et al. Evaluation of left ventricular early diastolic performance by color tissue Doppler imaging of the mitral annulus. *Am J Cardiol* 1998; 82: 1414-7
90. Rodriguez L, Garcia M, Ares M, et al. Assessment of mitral annular dynamics during diastole by Doppler tissue imaging: comparison with mitral Doppler inflow in subjects without heart disease and in patients with left ventricular hypertrophy. *Am Heart J* 1996; 77: 979-84
91. Rome JJ, Keane JF, Perry SB, et al. Double-umbrella closure of atrial defects: initial clinical application. *Circulation* 1990; 82: 751-8
92. Yip G, Wang M, Zhang Y, et al. Left ventricular long axis function in diastolic heart failure is reduced in both diastole and systole: time for a redefinition? *Heart* 2002; 87: 121-5
93. Mishiro Y, Oki T, Yamada H, et al. Evaluation of left ventricular contraction abnormalities in patients with dilated cardiomyopathy with the use of pulsed tissue Doppler imaging. *J Am Soc Echocardiogr* 1999; 13: 913-920
94. Carr-White GS, Kon M, Koh TW, et al. Right ventricular function after pulmonary autograft replacement of the aortic valve. *Circulation* 1999; 100(suppl II): II-36-41
95. Dhillon R, Henein M, Redington A. Transcatheter closure of septal defect preserves right ventricular function. *Heart* 2002; 87: 461-465
96. Frommelt, Peter C, Ballweg, Jean A, Whitstone, Beth N, et al. Usefulness of Doppler tissue imaging analysis of tricuspid annular motion for determination of right ventricular function in normal infants and children. *Am J Cardiol* 2002;89(5): 610-613
97. Selton-Suty C, Mock L, Piquemal R, et al. Doppler tissue imaging of the myocardium and right ventricle. *Arch Mal Coeur Vaiss* 2002 Oct;95(10):933-7
98. Mahbubul A, Johan W, Eva A, et al. Right ventricular function in patients with first inferior myocardial infarction: assessment by tricuspid annular motion and

- tricuspid annular velocity. Am Heart J 2000; 139(4): 710-715
99. Ali M, Marcy L, Sangeeta S, et al. Interrogation of the tricuspid annulus by a Doppler tissue imaging in patients with chronic pulmonary hypertension: implications for the assessment of right ventricular systolic and diastolic function. Cardiology 2001; 95: 101-104
100. Lange A, Coleman DM, Palka P, et al. Effect of catheter device of atrial septal defect on diastolic mitral annular motion. J Am Cardiol 2003; 91: 104-108
101. Greenberg NL, Firstenberg MS, Castro PL, et al. Doppler-derived myocardial systolic strain rate is a strong index of left ventricular contractility. Circulation 2002; 105: 99-105.
102. Abraham TP, Nishimura RA, Holmes DR , et al. Strain rate imaging for assessment of regional myocardial function-results from a clinical model of septal ablation. Circulation 2002; 105: 1403-1406.
103. Kiraly P, Kapusta L, Thijssen JM et al. Left ventricular myocardial function in congenital valvar aortic stenosis assed by ultrasound tissue velocity and strain rate techniques.Ultrasound in Med. & Biol. 2003; Vol. 29. No. 4. pp. 615-620.
104. Hoffmann R, Altio E, Nowak B, et al. Strain rate measurement by Doppler Echocardiography allows improved assessment of myocardial viability in patients with depressed left ventricular function. J Am Coll Cardiol 2002; 39: 443-9.