

Literaturliste

- Achenbach,T. (1991). *Manual for the Youth Self-Report and 1991 Profile*. Burlington,VT: University of Vermont, Department of Psychiatry.
- Adriani,W. & Laviola,G. (2004). Windows of vulnerability to psychopathology and therapeutic strategy in the adolescent rodent model. *Behav.Pharmacol*, 15(5-6), 341-352.
- American Psychiatric Association (1995). *Diagnostic and Statistical Manual of Mental Disorders*. Washington: APA.
- Andersen,S.L., Arvanitogiannis,A., Pliakas,A.M., LeBlanc,C., & Carlezon,W.A., Jr. (2002). Altered responsiveness to cocaine in rats exposed to methylphenidate during development. *Nat.Neurosci.*, 5(1), 13-14.
- Aquilino,W.S. (1997). Privacy Effects on Self-Reported Drug Use: Interactions With Survey Mode and Respondent Characteristics. *National Institute on Drug Abuse (NIDA) Research Monographs*, 167, 383-415.
- Arnold,L.E., Abikoff,H.B., Cantwell,D.P., Conners,C.K., Elliott,G., Greenhill,L.L., Hechtman,L., Hinshaw,S.P., Hoza,B., Jensen,P.S., Kraemer,H.C., March,J.S., Newcorn,J.H., Pelham,W.E., Richters,J.E., Schiller,E., Severe,J.B., Swanson,J.M., Vereen,D., & Wells,K.C. (1997). National Institute of Mental Health Collaborative Multimodal Treatment Study of Children with ADHD (the MTA). Design challenges and choices. *Arch.Gen.Psychiatry*, 54(9), 865-870.
- Baker,T.B. & Brandon,T.H. (1990). Validity of Self-Reports in Basic Research. *Behavioral Assessment*, 12, 33-51.
- Baumert,J. (1973). *Untersuchungen zur Valenz des HAWIK und die Entwicklung einer Kurzform (WIPKI)*. Stuttgart: Hogrefe.
- Benson,K. & Hartz,A. (2000). A comparison of observational studies and randomized, controlled trials. *N Engl.J Med.*, 342(25), 1879-1886.
- Bharucha-Reid,R., McCann,D., Schork,M.A., Foxman,B., Bass,A., Fraser,W., Cook,S., & Kaufman,R. (1995). A Comparison of Self-Reported Drug Use With a Urine Drug Screen in a Working Population. *Experimental and Clinical Psychopharmacology*, 3(3), 280-286.

- Biederman,J., Faraone,S.V., Monuteaux,M.C., & Feighner,J.A. (2000). Patterns of alcohol and drug use in adolescents can be predicted by parental substance use disorders. *Pediatrics*, 106(4), 792-797.
- Biederman,J., Wilens,T., Mick,E., Faraone,S., & Spencer,T. (1998). Does attention-deficit hyperactivity disorder impact the development course of drug and alcohol abuse and dependence? *Biological Psychiatry*, 44(4), 269-273.
- Biederman,J., Wilens,T., Mick,E., Spencer,T., & Faraone,S.V. (1999). Pharmacotherapy of attention-deficit/hyperactivity disorder reduces risk for substance use disorder. *Pediatrics*, 104(2), e20.
- Biederman,J., Wilens,T., Mick,E., Faraone,S., Weber,W., Curtis,S., Thornell,A., Pfister,K., Jetton,J., & Soriano,J. (1997). Is ADHD a Risk Factor for Psychoactive Substance Use Disorders? Findings From a Four-Year Prospective Follow-up Study. *J Am Acad Child Adolesc Psychiatry*, 36(1), 21-29.
- Borentein, M., Rothstein, H., and Cohen, J. Sample Power. [1.20]. 1997. SPSS Inc. Ref Type: Computer Program
- Bortz,J. & Döring,N. (1995). *Forschungsmethoden und Evaluation*. Heidelberg: Springer-Verlag.
- Brandon,C.L., Marinelli,M., Baker,L.K., & White,F.J. (2001). Enhanced reactivity and vulnerability to cocaine following methylphenidate treatment in adolescent rats. *Neuropsychopharmacology*, 25(5), 651-661.
- Brandon,C.L., Marinelli,M., & White,F.J. (2003). Adolescent exposure to methylphenidate alters the activity of rat midbrain dopamine neurons. *Biological Psychiatry*, 54(12), 1338-1344.
- Carelli,R.M. & Wightman,R.M. (2004). Functional microcircuitry in the accumbens underlying drug addiction: insights from real-time signaling during behavior. *Curr.Opin.Neurobiol.*, 14(6), 763-768.
- Carlezon,J., Mague,S.D., & Andersen,S.L. (2003). Enduring behavioral effects of early exposure to methylphenidate in rats. *Biological Psychiatry*, 54(12), 1330-1337.
- Carlezon,W.A., Jr., Thome,J., Olson,V.G., Lane-Ladd,S.B., Brodkin,E.S., Hiroi,N., Duman,R.S., Neve,R.L., & Nestler,E.J. (1998). Regulation of cocaine reward by CREB. *Science*, 282(5397), 2272-2275.
- Carr,K.D. (1984). The physiology of opiate hedonic effects and the role of opioids in motivated behavior. *Adv.Alcohol Subst.Abuse*, 3(3), 5-18.
- Carter,H.S. & Watson,W.A. (1994). IV pentazocine/methylphenidate abuse--the clinical toxicity of another Ts and blues combination. *J Toxicol.Clin.Toxicol.*, 32(5), 541-547.

- Chait,L.D. (1994). Reinforcing and subjective effects of methylphenidate in humans. *Behav Pharmacol*, 5(3), 281-288.
- Chau,D.T., Roth,R.M., & Green,A.I. (2004). The neural circuitry of reward and its relevance to psychiatric disorders. *Curr.Psychiatry Rep.*, 6(5), 391-399.
- Cierpka,M. & Frevert,G. (1994). *Die Familienbögen - Ein Inventar zur Einschätzung von Familienfunktionen*. Göttingen: Hogrefe.
- Cloninger,C.R., Przybeck,T.R., & Svarkic,D.M. (1994). *The Temperament and Character Inventory (TCI): A guide to its development and use*. St. Louis: Washington University.
- Cochran,W.G. (1968). The effectiveness of adjustment by subclassification in removing bias in observational studies. *Biometrics*, 24, 205-213.
- Collins,R.J., Weeks,J.R., Cooper,M.M., Good,P.I., & Russell,R.R. (1984). Prediction of abuse liability of drugs using IV self-administration by rats. *Psychopharmacology (Berl)*, 82(1-2), 6-13.
- Concato,J., Shah,N., & Horwitz,R. (2000). Randomized, controlled trials, observational studies, and the hierarchy of research designs *N Engl J Med*, 342(25), 1887-1892.
- Contet,C., Kieffer,B.L., & Befort,K. (2004). Mu opioid receptor: a gateway to drug addiction. *Curr.Opin.Neurobiol.*, 14(3), 370-378.
- Cook,R.C., Fradet,G., English,J.C., Soos,J., Muller,N.L., Connolly,T.P., & Levy,R.D. (1998). Recurrence of intravenous talc granulomatosis following single lung transplantation. *Can.Respir.J*, 5(6), 511-514.
- Curran,S., Newman,S., Taylor,E., & Asherson,P. (2000). Hypescheme: an operational criteria checklist and minimum data set for molecular genetic studies of attention deficit and hyperactivity disorders. *Am J Med.Genet*, 96(3), 244-250.
- D'Agostino,R.B., Jr. (1998). Propensity score methods for bias reduction in the comparison of a treatment to a non-randomized control group. *Stat Med*, 17(19), 2265-81.
- Debooy,V.D., Seshia,M.M., Tenenbein,M., & Casiro,O.G. (1993). Intravenous pentazocine and methylphenidate abuse during pregnancy. Maternal lifestyle and infant outcome. *Am J Dis.Child*, 147(10), 1062-1065.
- Ding,Y.S., Fowler,J.S., Volkow,N.D., Logan,J., Gatley,S.J., & Sugano,Y. (1995). Carbon-11-d-threo-methylphenidate binding to dopamine transporter in baboon brain. *J Nucl.Med.*, 36(12), 2298-2305.

- Disney,E.R., Elkins,I.J., McGue,M., & Iacono,W.G. (1999). Effects of ADHD, conduct disorder, and gender on substance use and abuse in adolescence. *Am J Psychiatry*, 156(10), 1515-1521.
- Faraone,S.V. & Wilens,T. (2003). Does stimulant treatment lead to substance use disorders? *J Clin.Psychiatry*, 64 Suppl 11, 9-13.
- Farone,S.V. (2001). Report from the second international meeting of the Attention Deficit Hyperactivity Disorder Molecular Genetics Network. *Am J Med.Genet*, 105(3), 255-258.
- Fergusson,D., LYNKEY,M., & Horwood,L. (1997). Attentional difficulties in middle childhood and psychosocial outcomes in young adulthood. *J Child Psychol Psychiatry*, 38, 633-644.
- Fergusson,D., LYNKEY,M., & Horwood,L. (2005). Attentional difficulties in middle childhood and psychosocial outcomes in young adulthood. *J Child Psychol Psychiatry*, 38, 633-644.
- Fergusson,D.M. & Horwood,L.J. (2000). Does cannabis use encourage other forms of illicit drug use? *Addiction*, 95(4), 505-520.
- Ferring,D.F. & Filipp,F.-H. (1989). Bewältigung kritischer Lebensereignisse: Erste Erfahrungen mit einer deutschsprachigen Version der "Ways of Coping Checklist". *Zeitschrift für Differentielle und Diagnostische Psychologie*, 10(4), 189-199.
- Garland,E.J. (1998). Intranasal abuse of prescribed methylphenidate. *J Am Acad.Child Adolesc Psychiatry*, 37(12), 1242-1243.
- Gillberg,C., Melander,H., von Knorring,A.L., Janols,L.O., Thernlund,G., Hagglof,B., Eidevall-Wallin,L., Gustafsson,P., & Kopp,S. (1997). Long-term stimulant treatment of children with attention-deficit hyperactivity disorder symptoms. A randomized, double-blind, placebo-controlled trial. *Arch Gen Psychiatry*, 54(9), 857-864.
- Haug,H. & Stieglitz,R. (1997). *Das AMDP-System in der klinischen Anwendung und Forschung*. Stuttgart: Hogrefe.
- Hechtman,L. (1985). Adolescent outcome of hyperactive children treated with stimulants in childhood: a review. *Psychopharmacol.Bull.*, 21(2), 178-191.
- Hoebel,B.G., Monaco,A.P., Hernandez,L., Aulisi,E.F., Stanley,B.G., & Lenard,L. (1983). Self-injection of amphetamine directly into the brain. *Psychopharmacology (Berl)*, 81(2), 158-163.
- Huss,M. (2004). Langzeiteffekte der Stimulanzietherapie. InM.Schulte-Markwort & A.Warnke (Eds.), *Methylphenidat* (pp. 34-42). Heidelberg: Springer.

- Huss,M. & Lehmkuhl,U. (1999). Trennung und Scheidung aus Sicht der Kinder und Jugendlichen: Chancen und Risiken für die psychische Entwicklung.
In J.Fegert (Ed.), *Kinder in Scheidungsverfahren nach der Kindschaftsreform* (pp. 31-44). Neuwied: Luchterhand-Verlag.
- Huss,M. & Lehmkuhl,U. (2002a). Methylphenidate and substance abuse: a review of pharmacology, animal, and clinical studies. *J. Atten. Disord.*, 6 Suppl 1, S65-S71.
- Huss,M. & Lehmkuhl,U. (2002b). Methylphenidate and substance abuse: a review of pharmacology, animal, and clinical studies. *J Atten. Disord.*, 6 Suppl 1, S65-S71.
- Huss, M. and Lehmkuhl, U. Sind mit Methylphenidat therapierte Menschen einem erhöhten Missbrauchs- und Abhängigkeitsrisiko ausgesetzt?
Forschungsbericht an das Bundesinstitut für Arzneimittel- und Medizinprodukte (BfArM) über die Ergebnisse der multizentrischen Studie Fo2.1-68503-200 ausgeschrieben im Deutschen Ärzteblatt Nr.30 vom 28.6.1995 . 2002c.
- Huss,M., Schmidt-Schulz,A., & Lehmkuhl,U. (2000). Wie gut stimmen subjektive Angaben ehemaliger Patienten mit Hyperkinetischen Syndrom mit objektiven Befunden des Drogenscreenings überein? Suchtmedizin in Forschung und Praxis. *Suchtmedizin in Forschung und Praxis*, 2(3), 135-140.
- Huss, Michael, Schmidt-Schulz, Andrea, Mendes, Ute, Völger, Margot, and Lehmkuhl, Ulrike. Discriminat Power of a New Diagnostic Tool for Retrospective Analysis of ADHD. 1998. Stockholm. 10-4-1998.
Ref Type: Conference Proceeding
- Ikegami,A. & Duvauchelle,C.L. (2004). Nucleus accumbens and medial prefrontal cortex dopaminergic response to self-administered cocaine in naive rats. *Neuroscience Letters*, 354(3), 205-208.
- Jaffe,R.B. & Koschmann,E.B. (1970). Intravenous drug abuse. Pulmonary, cardiac, and vascular complications. *Am J Roentgenol.Radium.Ther.Nucl.Med.*, 109(1), 107-120.
- Jaffe,S.L. (1991). Intranasal abuse of prescribed methylphenidate by an alcohol and drug abusing adolescent with ADHD. *J Am Acad.Child Adolesc Psychiatry*, 30(5), 773-775.
- Jaffe,S.L. (2002). Failed attempts at intranasal abuse of Concerta. *J Am Acad.Child Adolesc Psychiatry*, 41(1), 5.
- Jessup,M. (1996). Nicotine: a gateway drug? *J Am Med.Womens Assoc*, 51(1-2), 21.

- Joffe,M.M. & Rosenbaum,P.R. (1999). Invited commentary: propensity scores. *Am J Epidemiol*, 150(4), 327-33.
- Julien,R.M. (1997). *Drogen und Psychopharmaka*. Heidelberg: Springer-Verlag.
- Kandel,D.B., Yamaguchi,K., & Chen,K. (1992). Stages of progression in drug involvement from adolescence to adulthood: further evidence for the gateway theory. *J Stud.Alcohol*, 53(5), 447-457.
- Katz,J.L., Izenwasser,S., & Terry,P. (2000). Relationships among dopamine transporter affinities and cocaine-like discriminative-stimulus effects. *Psychopharmacology (Berl)*, 148(1), 90-98.
- Kenkel,D., Mathios,A.D., & Pacula,R.L. (2001). Economics of youth drug use, addiction and gateway effects. *Addiction*, 96(1), 151-164.
- Kleiber,D. & Soellner,R. (1998). *Cannabiskonsum: Entwicklungstendenzen, Konsummuster und Risiken*. Weinheim: Juventa.
- Kleining,G. & Moore,H. (1968). Soziale Selbsteinstufung (SSE). Ein Instrument zur Messung sozialer Schichten. *Kölner Z Soz Psychol* 20, 502-5. *Kölner Z Soz Psychol*, 20, 502-552.
- Kollins,S.H. (2003). Comparing the abuse potential of methylphenidate versus other stimulants: a review of available evidence and relevance to the ADHD patient. *J Clin.Psychiatry*, 64 Suppl 11, 14-18.
- Kollins,S.H., MacDonald,E.K., & Rush,C.R. (2001). Assessing the abuse potential of methylphenidate in nonhuman and human subjects: a review. *Pharmacol.Biochem.Behav.*, 68(3), 611-627.
- Koob,G.F. & Le Moal,M. (1997). Drug abuse: hedonic homeostatic dysregulation. *Science*, 278(5335), 52-58.
- Kupper,L.L., McMichael,A.J., & Spirtas,R. (1975). A hybrid epidemiologic study design useful in estimating relative risk. *J Am Stat Assoc*, 70, 524-528.
- Lambert,N.M. & Hartsough,C.S. (1998). Prospective study of tobacco smoking and substance dependencies among samples of ADHD and non-ADHD participants. *J Learn.Disabil.*, 31(6), 533-544.
- Larkin,M. & Griffiths,M.D. (1998). Response to Shaffer (1996): The Case for a 'Complex Systems' Conceptualisation of Addiction. *J Gambl.Stud.*, 14(1), 73-82.
- Lenton,S. (2001). Cannabis as a gateway drug: comments on Fergusson & Horwood (2000). *Addiction*, 96(3), 511-515.

- Lerman,C., Niaura,R., Collins,B.N., Wileyto,P., Audrain-McGovern,J., Pinto,A., Hawk,L., & Epstein,L.H. (2004). Effect of bupropion on depression symptoms in a smoking cessation clinical trial. *Psychol.Addict.Behav.*, 18(4), 362-366.
- Lindsay,G.B. & Rainey,J. (1997). Psychosocial and pharmacologic explanations of nicotine's "gateway drug" function. *J Sch Health*, 67(4), 123-126.
- Mague,S.D., Andersen,S.L., & Carlezon,W.A., Jr. (2005). Early developmental exposure to methylphenidate reduces cocaine-induced potentiation of brain stimulation reward in rats. *Biol.Psychiatry*, 57(2), 120-125.
- Massello,W., III & Carpenter,D.A. (1999). A fatality due to the intranasal abuse of methylphenidate (Ritalin). *J Forensic Sci.*, 44(1), 220-221.
- Mehta,H., Murray,B., & Loludice,T.A. (1984). Hepatic dysfunction due to intravenous abuse of methylphenidate hydrochloride. *J Clin.Gastroenterol.*, 6(2), 149-151.
- Meyer,L., Heijst,V., Geldmacher,A., & Mallinckrodt,M. (2005). *Cannabinoidnachweis im Harn. Mitteilungen der Senats-kommission der DFG für klinisch-toxikologische Analytik*. Weinheim: VCH-Verlagsgesellschaft.
- Milberger,S., Biederman,J., Faraone,S.V., Chen,L., & Jones,J. (1997). ADHD Is Associated With Early Initiation of Cigarette Smoking in Children and Adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(1), 37-44.
- Molina, B., Pelham, W., and Roth, J. Stimulant medication and substance use by adolescents with a childhood history of ADHD. 1999.
Ref Type: Conference Proceeding
- Molina,B.S. & Pelham,W.E. (2003). Childhood predictors of adolescent substance use in a longitudinal study of children with ADHD. *J.Abnorm.Psychol.*, 112(3), 497-507.
- Möllmann,C., Lenz,K., Naumann,A., Döpfner,M., Lehmkuhl,G., & Lehmkuhl,U. (1998). Klinikvergleich im Hinblick auf die Achse 5 des MAS: Unterschiedliche Inanspruchnahmepopulationen oder regionale Besonderheiten? *Z Kinder Jugendpsychiatr Psychother*, 26(2), 113-123.
- Nakano,H., Iwata,Y., Kanegae,H., Oshima,T., Aizawa,H., & Hara,N. (1998). [Talc pulmonary granulomatosis caused by intravenous administration of methylphenidate]. *Nihon Kokyuki Gakkai Zasshi*, 36(1), 111-115.
- Niederhofer,H. & Huber,M. (2004). Bupropion may support psychosocial treatment of nicotine-dependent adolescents: preliminary results. *Pharmacotherapy*, 24(11), 1524-1528.

- O'Donnell,D., Biederman,J., Jones,J., Wilens,T.E., Milberger,S., Mick,E., & Faraone,S.V. (1998). Informativeness of child and parent reports on substance use disorders in a sample of ADHD probands, control probands, and their siblings. *J Am Acad Child Adolesc Psychiatry*, 37(7), 752-758.
- Padley,S.P., Adler,B.D., Staples,C.A., Miller,R.R., & Muller,N.L. (1993). Pulmonary talcosis: CT findings in three cases. *Radiology*, 186(1), 125-127.
- Parran,T.V., Jr. & Jasinski,D.R. (1991). Intravenous methylphenidate abuse. Prototype for prescription drug abuse. *Arch Intern Med.*, 151(4), 781-783.
- Pelham, Bill. ADHD and Adolescent Drug Abuse. 3-8-2003.
Ref Type: Unpublished Work
- Remschmidt,H. & Schmidt,M. (1994). *Multiaxiales Klassifikationsschema für psychische Störungen des Kindes- und Jugendalters nach ICD-10 der Who*. Bern: Huber.
- Renz, Diane. Macht Ritalin süchtig? Frankfurter Allgemeine Sonntagszeitung [34]. 24-8-2003.
Ref Type: Newspaper
- Robbins,T.W. (2002). ADHD and addiction. *Nat Med.*, 8(1), 24-25.
- Robbins,T.W. & Everitt,B.J. (1999). Drug addiction: bad habits add up. *Nature*, 398(6728), 567-570.
- Roehrs,T., Papineau,K., Rosenthal,L., & Roth,T. (1999). Sleepiness and the reinforcing and subjective effects of methylphenidate. *Exp Clin Psychopharmacol*, 7(2), 145-150.
- Rosenbaum,P.R. & Rubin,D.B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70, 41-55.
- Rosenbaum,P.R. & Rubin,D.B. (1984). Reducing bias in observational studies using subclassification on the propensity score. *J Am Stat Association*, 79, 516-524.
- Sandoval,V., Riddle,E.L., Hanson,G.R., & Fleckenstein,A.E. (2002). Methylphenidate redistributes vesicular monoamine transporter-2: role of dopamine receptors. *J Neurosci.*, 22(19), 8705-8710.
- Sandoval,V., Riddle,E.L., Hanson,G.R., & Fleckenstein,A.E. (2003). Methylphenidate alters vesicular monoamine transport and prevents methamphetamine-induced dopaminergic deficits. *Journal of Pharmacology And Experimental Therapeutics*, 304(3), 1181-1187.
- Schenk,J.O. (2002). The functioning neuronal transporter for dopamine: kinetic mechanisms and effects of amphetamines, cocaine and methylphenidate. *Prog Drug Res.*, 59, 111-131.

- Shaffer,H.J. (1997). The most important unresolved issue in the addictions: conceptual chaos. *Subst.Use.Misuse.*, 32(11), 1573-1580.
- Shekim,W.O., Asarnow,R.F., Hess,E., Zaucha,K., & Wheeler,N. (1990). A clinical and demographic profile of a sample of adults with attention deficit hyperactivity disorder, residual state. *Compr.Psychiatry*, 31(5), 416-425.
- Shizgal,P. & Arvanitogiannis,A. (2003). Neuroscience. Gambling on dopamine. *Science*, 299(5614), 1856-1858.
- Simon,J.A., Duncan,C., Carmody,T.P., & Hudes,E.S. (2004). Bupropion for smoking cessation: a randomized trial. *Arch Intern.Med.*, 164(16), 1797-1803.
- Skinner,H.A., Steinhauer,P.D., & Santa-Barbara,J. (1983). The Fmily Assessment Measure. *Canadian Journal of Community Mental Health*, 2, 91-105.
- Stern,E.J., Frank,M.S., Schmutz,J.F., Glenny,R.W., Schmidt,R.A., & Godwin,J.D. (1994). Panlobular pulmonary emphysema caused by i.v. injection of methylphenidate (Ritalin): findings on chest radiographs and CT scans. *AJR Am J Roentgenol.*, 162(3), 555-560.
- Still,A., Gordon,M., Mercer,J., & Roake,J. (2001). Ritalin: drug of abuse. Two case reports of intra-arterial injection. *N.Z.Med.J*, 114(1144), 521-522.
- Swanson,J.M. & Volkow,N.D. (2002). Pharmacokinetic and pharmacodynamic properties of stimulants: implications for the design of new treatments for ADHD. *Behav.Brain Res.*, 130(1-2), 73-78.
- Swanson,J.M. & Volkow,N.D. (2003). Serum and brain concentrations of methylphenidate: implications for use and abuse. *Neuroscience & Biobehavioral Reviews*, 27(7), 615-621.
- Swanson,J.M., McBurnett,K., Christian,D.L., & Wigal,T. (1995). Stimulant medications and the treatment of children with ADHD. InT.H.Ollendick & R.J.Prinz (Eds.), *Advances in Clinical Child Psychology* (pp. 265-322). New York: Plenum.
- Swanson,J.M., McBurnett,K., Wigal,T., Pfiffner,L.J., Lerner,M.A., Williams,L., Christian,D.L., Tamm,L., Willcutt,E., Crowley,K., Clevenger,W., Khouzam,N., Woo,C., Crinella,F.M., & Fisher,T.D. (1993). Effect of stimulant medication on children with attention deficit disorder: A review of reviews. *Exceptional Children*, 60(2), 154-162.
- Talwar,A., Jain,M., & Vijayan,V.K. (2004). Pharmacotherapy of tobacco dependence. *Med.Clin.North Am*, 88(6), 1517-1534.
- Tannock,R. (1998). Attention deficit hyperactivity disorder: advances in cognitive, neurobiological, and genetic research. *J.Child Psychol.Psychiatry*, 39(1), 65-99.

- Taylor,E., Dopfner,M., Sergeant,J., Asherson,P., Banaschewski,T., Buitelaar,J., Coghill,D., Danckaerts,M., Rothenberger,A., Sonuga-Barke,E., Steinhausen,H.C., & Zuddas,A. (2004). European clinical guidelines for hyperkinetic disorder -- first upgrade. *Eur Child Adolesc Psychiatry*, 13 Suppl 1, I7-30.
- Taylor,E., Sergeant,J., Doeppner,M., Gunning,B., Overmeyer,S., Mobius,H.J., & Eisert,H.G. (1998). Clinical guidelines for hyperkinetic disorder. European Society for Child and Adolescent Psychiatry. *Eur Child Adolesc Psychiatry*, 7(4), 184-200.
- Taylor,E. (1900). Attention deficit and conduct disorder syndromes. In M.Rutter, A.H.Tuma, & I.S.Lann (Eds.), *Assessment and Diagnosis in Child Psychopathology* (pp. 377-407). New York: Guilford Press.
- Taylor,E. (1990). Attention deficit and conduct disorder syndromes. In M.Rutter, A.H.Tuma, & I.S.Lann (Eds.), *Assessment and Diagnosis in Child Psychopathology* (pp. 377-407). New York: Guilford Press.
- Terry,P. & Katz,J.L. (1997). Dopaminergic mediation of the discriminative stimulus effects of bupropion in rats. *Psychopharmacology (Berl)*, 134(2), 201-212.
- Tse,D.T. & Ober,R.R. (1980). Talc retinopathy. *Am J Ophthalmol.*, 90(5), 624-640.
- Unnewehr,S., Schneider,S., & Markgraf,J. (1995). *Kinder-DIPS - Diagnostisches Interview bei psychischen Störungen im Kindes- und Jugendalter*. Springer-Verlag.
- van Wolfswinkel,L., Seifert,W.F., & van Ree,J.M. (1988). Catecholamines and endogenous opioids in ventral tegmental self-stimulation reward. *Pharmacol Biochem Behav.*, 30(3), 589-595.
- Vastag,B. (2001). Pay attention: ritalin acts much like cocaine. *Jama*, 286(8), 905-906.
- Vitiello,B. (2001a). Long-term effects of stimulant medications on the brain: possible relevance to the treatment of attention deficit hyperactivity disorder. *J Child Adolesc Psychopharmacol.*, 11(1), 25-34.
- Vitiello,B. (2001b). Psychopharmacology for young children: clinical needs and research opportunities. *Pediatrics*, 108(4), 983-989.
- Vitiello,B., Severe,J.B., Greenhill,L.L., Arnold,L.E., Abikoff,H.B., Bukstein,O.G., Elliott,G.R., Hechtman,L., Jensen,P.S., Hinshaw,S.P., March,J.S., Newcorn,J.H., Swanson,J.M., & Cantwell,D.P. (2001). Methylphenidate dosage for children with ADHD over time under controlled conditions: lessons from the MTA. *J Am Acad Child Adolesc Psychiatry*, 40(2), 188-196.
- Volkow,N.D., Ding,Y.S., Fowler,J.S., Wang,G.J., Logan,J., Gatley,J.S., Dewey,S., Ashby,C., Liebermann,J., Hitzemann,R., & . (1995a). Is methylphenidate like

cocaine? Studies on their pharmacokinetics and distribution in the human brain. *Arch.Gen.Psychiatry*, 52(6), 456-463.

Volkow,N.D., Ding,Y.S., Fowler,J.S., Wang,G.J., Logan,J., Gatley,S.J., Schlyer,D.J., & Pappas,N. (1995b). A new PET ligand for the dopamine transporter: studies in the human brain. *J Nucl.Med.*, 36(12), 2162-2168.

Ward,S., Heyneman,L.E., Reittner,P., Kazerooni,E.A., Godwin,J.D., & Muller,N.L. (2000). Talcosis associated with IV abuse of oral medications: CT findings. *AJR Am J Roentgenol.*, 174(3), 789-793.

Weinman,S.A. & D'Agostino,J.L. (1993). Pyrogenic reaction after intravenous injection of pentazocine and methylphenidate. *J Emerg.Nurs.*, 19(1), 10-17.

Wilens,T.E., Biederman,J., & Spencer,T.J. (2002). Attention deficit/hyperactivity disorder across the lifespan. *Annu.Rev.Med.*, 53, 113-131.

Wilens,T.E., Faraone,S.V., Biederman,J., & Gunawardene,S. (2003). Does stimulant therapy of attention-deficit/hyperactivity disorder beget later substance abuse? A meta-analytic review of the literature. *Pediatrics*, 111(1), 179-185.

Yu,J. & Williford,W.R. (1992). The age of alcohol onset and alcohol, cigarette, and marijuana use patterns: an analysis of drug use progression of young adults in New York State. *Int.J Addict.*, 27(11), 1313-1323.