
11 Literaturverzeichnis

- Aletta JM, Lewis SA, Cowan NJ, Greene LA (1988) Nerve growth factor regulates both the phosphorylation and steady-state levels of microtubule-associated protein 1.2 (MAP1.2). *J Cell Biol* 106: 1573-1581.
- Arakawa T, Haniu M, Narhi LO, Miller JA, Talvenheimo J, Philo JS, Chute HT, Matheson C, Carnahan J, Louis JC, . (1994) Formation of heterodimers from three neurotrophins, nerve growth factor, neurotrophin-3, and brain-derived neurotrophic factor. *J Biol Chem* 269: 27833-27839.
- Bamji SX, Majdan M, Pozniak CD, Belliveau DJ, Aloyz R, Kohn J, Causing, CG, Miller FD (1998) The p75 neurotrophin receptor mediates neuronal apoptosis and is essential for naturally occurring sympathetic neuron death. *Journal of Cell Biology* 140: 911-923.
- Banfield MJ, Naylor RL, Robertson AG, Allen SJ, Dawbarn D, Brady RL (2001) Specificity in Trk receptor:neurotrophin interactions: the crystal structure of TrkB-d5 in complex with neurotrophin-4/5. *Structure (Camb)* 9: 1191-1199.
- Barde YA, Edgar D, Thoenen H (1982) Purification of a new neurotrophic factor from mammalian brain. *Embo J* 1: 549-553.
- Barker PA, Barbee G, Misko TP, Shooter EM (1994) The low affinity neurotrophin receptor, p75^{NTR}, is palmitoylated by thioester formation through cysteine 279. *J* 269: 30645-30650.
- Benzel I, Barde YA, Casademunt E (2001) Strain-specific complementation between NRIF1 and NRIF2, two zinc finger proteins sharing structural and biochemical properties. *Gene* 281: 19-30.
- Berkemeier LR, Winslow JW, Kaplan DR, Nikolics K, Goeddel DV, Rosenthal A (1991) Neurotrophin-5: a novel neurotrophic factor that activates trk and trkB. *Neuron* 7: 857-866.
- Bibel M, Hoppe E, Barde YA (1999) Biochemical and functional interactions between the neurotrophin receptors trk and p75^{NTR}. *Embo J* 18: 616-622.
- Bilderback TR, Grigsby RJ, Dobrowsky RT (1997) Association of p75^{NTR} with caveolin and localization of neurotrophin-induced sphingomyelin hydrolysis to caveolae. *J* 272: 10922-10927.
- Billger M, Wallin M, Karlsson JO (1988) Proteolysis of tubulin and microtubule-associated proteins 1 and 2 by calpain I and II. Difference in sensitivity of assembled and disassembled microtubules. *Cell Calcium* 9: 33-44.
- Blanquet PR (2000) Identification of two persistently activated neurotrophin-regulated pathways in rat hippocampus. *Neuroscience* 95: 705-719.

- Bloch A, Thoenen H (1995) Characterization of nerve growth factor (NGF) release from hippocampal neurons: evidence for a constitutive and an unconventional sodium- dependent regulated pathway. *Eur J Neurosci* 7: 1220-1228.
- Blom N, Gammeltoft S, Brunak S (1999) Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J Mol Biol* 294: 1351-1362.
- Bloom GS, Luca FC, Vallee RB (1985) Microtubule-associated protein 1B: identification of a major component of the neuronal cytoskeleton. *Proc Natl Acad Sci U S A* 82: 5404-5408.
- Botchkareva NV, Botchkarev VA, Chen LH, Lindner G, Paus R (1999) A role for p75 neurotrophin receptor in the control of hair follicle morphogenesis. *Dev Biol* 216: 135-153.
- Boyne LJ, Martin K, Hockfield S, Fischer I (1995) Expression and distribution of phosphorylated MAP1B in growing axons of cultured hippocampal neurons. *J Neurosci Res* 40: 439-450.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254.
- Brann AB, Scott R, Neuberger Y, Abulafia D, Boldin S, Fainzilber M, Futeran AH (1999) Ceramide signaling downstream of the p75 neurotrophin receptor mediates the effects of nerve growth factor on outgrowth of cultured hippocampal neurons. *J Neurosci* 19: 8199-8206.
- Brodski C, Schnurch H, Dechant G (2000) Neurotrophin-3 promotes the cholinergic differentiation of sympathetic neurons. *Proc Natl Acad Sci U S A* 97: 9683-9688.
- Brugg B, Matus A (1991) Phosphorylation determines the binding of microtubule-associated protein 2 (MAP2) to microtubules in living cells. *J Cell Biol* 114: 735-743.
- Brugg B, Reddy D, Matus A (1993) Attenuation of microtubule-associated protein 1B expression by antisense oligodeoxynucleotides inhibits initiation of neurite outgrowth. *Neuroscience* 52: 489-496.
- Bush MS, Goold RG, Moya F, Gordon-Weeks PR (1996) An analysis of an axonal gradient of phosphorylated MAP 1B in cultured rat sensory neurons. *Eur J Neurosci* 8: 235-248.
- Carlin RK, Grab DJ, Cohen RS, Siekevitz P (1980) Isolation and characterization of postsynaptic densities from various brain regions: enrichment of different types of postsynaptic densities. *Journal of Cell Biology* 86: 831-845.
- Carter BD, Kaltschmidt C, Kaltschmidt B, Offenhauser N, Bohm-Matthaei R, Baeuerle PA, Barde YA (1996) Selective activation of NF-kappa B by nerve growth factor through the neurotrophin receptor p75 [see comments]. *Science* 272: 542-545.
- Casaccia-Bonnel P, Carter BD, Dobrowsky RT, Chao MV (1996) Death of oligodendrocytes mediated by the interaction of nerve growth factor with its receptor p75. *Nature* 383: 716-719.

-
- Casademunt E, Carter BD, Benzel I, Frade JM, Dechant G, Barde YA (1999) The zinc finger protein NRIF interacts with the neurotrophin receptor p75(NTR) and participates in programmed cell death. *Embo J* 18: 6050-6061.
- Chan FK, Chun HJ, Zheng L, Siegel RM, Bui KL, Lenardo MJ (2000) A domain in TNF receptors that mediates ligand-independent receptor assembly and signaling. *Science* 288: 2351-2354.
- Chao MV, Hempstead BL (1995) p75 and Trk: a two-receptor system. *Trends Neurosci* 18: 321-326.
- Chapman BS (1995) A region of the 75 kDa neurotrophin receptor homologous to the death domains of TNFR-I and Fas. *FEBS Lett* 374: 216-220.
- Chen C, Okayama H (1987) High-efficiency transformation of mammalian cells by plasmid DNA. *Mol Cell Biol* 7: 2745-2752.
- Chen G, Kolbeck R, Barde YA, Bonhoeffer T, Kossel A (1999) Relative contribution of endogenous neurotrophins in hippocampal long-term potentiation. *J Neurosci* 19: 7983-7990.
- Chittka A, Chao MV (1999) Identification of a zinc finger protein whose subcellular distribution is regulated by serum and nerve growth factor. *Proc* 96: 10705-10710.
- Chuang HH, Prescott ED, Kong H, Shields S, Jordt SE, Basbaum AI, Chao MV, Julius D (2001) Bradykinin and nerve growth factor release the capsaicin receptor from PtdIns(4,5)P₂-mediated inhibition. *Nature* 411: 957-962.
- Cohen S (1960) Purification of a nerve-growth promoting protein from the mouse salivary gland and its neuro-cytotoxic antiserum. *Proc Natl Acad Sci U S A* 46: 302-311.
- Cosgaya JM, Shooter EM (2001) Binding of nerve growth factor to its p75 receptor in stressed cells induces selective I κ B β degradation and NF- κ B nuclear translocation. *J Neurochem* 79: 391-399.
- Coulson EJ, Reid K, Baca M, Shipham KA, Hulett SM, Kilpatrick TJ, Bartlett PF (2000) Chopper, a new death domain of the p75 neurotrophin receptor that mediates rapid neuronal cell death. *J Biol Chem* 275: 30537-30545.
- Craven SE, Bredt DS (1998) PDZ proteins organize synaptic signaling pathways [Review]. *Cell* 93: 495-498.
- Damerval C., le Guilloux M., Blaisonneau J., and de Vienne D. A simplification of Heukeshoven and Dernick's silver staining of proteins. *Electrophoresis* 8, 158-159. 1987.
Ref Type: Generic
- Davey F, Davies AM (1998) TrkB signalling inhibits p75-mediated apoptosis induced by nerve growth factor in embryonic proprioceptive neurons. *Current Biology* 8: 915-918.

- Diaz-Nido J, Serrano L, Mendez E, Avila J (1988) A casein kinase II-related activity is involved in phosphorylation of microtubule-associated protein MAP-1B during neuroblastoma cell differentiation. *J Cell Biol* 106: 2057-2065.
- DiTella MC, Feiguin F, Carri N, Kosik KS, Caceres A (1996) MAP-1B/TAU functional redundancy during laminin-enhanced axonal growth. *J Cell Sci* 109 (Pt 2): 467-477.
- Dobrowsky RT, Jenkins GM, Hannun YA (1995) Neurotrophins induce sphingomyelin hydrolysis. Modulation by co-expression of p75NTR with Trk receptors. *J* 270: 22135-22142.
- Dobrowsky RT, Werner MH, Castellino AM, Chao MV, Hannun YA (1994) Activation of the sphingomyelin cycle through the low-affinity neurotrophin receptor. *Science* 265: 1596-1599.
- Drubin D, Kobayashi S, Kellogg D, Kirschner M (1988) Regulation of microtubule protein levels during cellular morphogenesis in nerve growth factor-treated PC12 cells. *J Cell Biol* 106: 1583-1591.
- Edelmann W, Zervas M, Costello P, Roback L, Fischer I, Hammarback JA, Cowan N, Davies P, Wainer B, Kucherlapati R (1996) Neuronal abnormalities in microtubule-associated protein 1B mutant mice. *Proc Natl Acad Sci U S A* 93: 1270-1275.
- Ernfors P, Ibanez CF, Ebendal T, Olson L, Persson H (1990) Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain. *Proc Natl Acad Sci U S A* 87: 5454-5458.
- Esposito D, Patel P, Stephens RM, Perez P, Chao MV, Kaplan DR, Hempstead BL (2001) The cytoplasmic and transmembrane domains of the p75 and Trk A receptors regulate high affinity binding to nerve growth factor. *J Biol Chem* 276: 32687-32695.
- Fawcett JW, Mathews G, Housden E, Goedert M, Matus A (1994) Regenerating sciatic nerve axons contain the adult rather than the embryonic pattern of microtubule associated proteins. *Neuroscience* 61: 789-804.
- Feinstein DL, Larhammar D (1990) Identification of a conserved protein motif in a group of growth factor receptors. *FEBS Lett* 272: 7-11.
- Feinstein E, Kimchi A, Wallach D, Boldin M, Varfolomeev E (1995) The death domain: a module shared by proteins with diverse cellular functions [letter]. *Trends in Biochemical Sciences* 20: 342-344.
- Figurov A, Pozzo-Miller LD, Olafsson P, Wang T, Lu B (1996) Regulation of synaptic responses to high-frequency stimulation and LTP by neurotrophins in the hippocampus. *Nature* 381: 706-709.
- Forrer P, Jaussi R (1998) High-level expression of soluble heterologous proteins in the cytoplasm of *Escherichia coli* by fusion to the bacteriophage lambda head protein D. *Gene* 224: 45-52.
- Frade JM (2000) Unscheduled re-entry into the cell cycle induced by NGF precedes cell death in nascent retinal neurones. *J Cell Sci* 113 (Pt 7): 1139-1148.

- Frade JM, Rodriguez-Tebar A, Barde YA (1996) Induction of cell death by endogenous nerve growth factor through its p75 receptor. *Nature* 383: 166-168.
- Gaese F, Kolbeck R, Barde YA (1994) Sensory ganglia require neurotrophin-3 early in development. *Development* 120: 1613-1619.
- Gal V, Martin S, Bayley P (1988) Fast disassembly of microtubules induced by Mg²⁺ or Ca²⁺. *Biochem Biophys Res Commun* 155: 1464-1470.
- Galbiati F, Volonte D, Meani D, Milligan G, Lublin DM, Lisanti MP, Parenti M, marco (1999) The dually acylated NH₂-terminal domain of G(i1 alpha) is sufficient to target a green fluorescent protein reporter to caveolin-enriched plasma membrane domains - Palmitoylation of caveolin-1 is required for the recognition of dually acylated G-protein alpha subunits in vivo. *J* 274: 5843-5850.
- Gargano N, Levi A, Alema S (1997) Modulation of nerve growth factor internalization by direct interaction between p75 and TrkA receptors. *J Neurosci Res* 50: 1-12.
- Gentry, J. J., Rutkoski, N., Linggi, M. S., Musiek, E. S., Emeson, R., and Carter, B. D. A functional interaction between the p75 associated proteins NRIF and TRAF6. Annual Neuroscience Meeting 2001, San Diego, 695.14. 2001.
Ref Type: Abstract
- Giasson BI, Bruening W, Durham HD, Mushynski WE (1999) Activation of stress-activated protein kinases correlates with neurite outgrowth induced by protease inhibition in PC12 cells. *J Neurochem* 72: 1081-1087.
- Gogstad GO, Krutnes MB (1982) Measurement of protein in cell suspensions using the Coomassie brilliant blue dye-binding assay. *Anal Biochem* 126: 355-359.
- Goldowitz D, Hamre K (1998) The cells and molecules that make a cerebellum. *Trends Neurosci* 21: 375-382.
- Gong CX, Wegiel J, Lidsky T, Zuck L, Avila J, Wisniewski HM, Grundke-Iqbal I, Iqbal K (2000) Regulation of phosphorylation of neuronal microtubule-associated proteins MAP1b and MAP2 by protein phosphatase-2A and -2B in rat brain. *Brain Res* 853: 299-309.
- Goold RG, Owen R, Gordon-Weeks PR (1999) Glycogen synthase kinase 3beta phosphorylation of microtubule-associated protein 1B regulates the stability of microtubules in growth cones. *J Cell Sci* 112 (Pt 19): 3373-3384.
- Gotz R, Koster R, Winkler C, Raulf F, Lottspeich F, Scharf M, Thoenen H (1994) Neurotrophin-6 is a new member of the nerve growth factor family. *Nature* 372: 266-269.
- Graham FL, Smiley J, Russell WC, Nairn R (1977) Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J Gen Virol* 36: 59-74.

- Greene LA, Tischler AS (1976) Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc Natl Acad Sci U S A* 73: 2424-2428.
- Grob PM, Ross AH, Koprowski H, Bothwell M (1985) Characterization of the human melanoma nerve growth factor receptor. *J Biol Chem* 260: 8044-8049.
- Grundke-Iqbal I, Iqbal K, Tung YC, Quinlan M, Wisniewski HM, Binder LI (1986) Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A* 83: 4913-4917.
- Gundersen RW, Barrett JN (1979) Neuronal chemotaxis: chick dorsal-root axons turn toward high concentrations of nerve growth factor. *Science* 206: 1079-1080.
- Hallbook F, Ibanez CF, Persson H (1991) Evolutionary studies of the nerve growth factor family reveal a novel member abundantly expressed in *Xenopus* ovary. *Neuron* 6: 845-858.
- Hammarback JA, Obar RA, Hughes SM, Vallee RB (1991) MAP1B is encoded as a polyprotein that is processed to form a complex N-terminal microtubule-binding domain. *Neuron* 7: 129-139.
- Harlow E, Lane D (1988) *Using Antibodies: A Laboratory Manual [Illustrated]*. In: *Antibodies: A Laboratory Manual [Illustrated]* Plainview: Cold Spring Harbor Laboratory Press.
- Hart GW (1997) Dynamic O-linked glycosylation of nuclear and cytoskeletal proteins. [Review] [190 refs]. *Annual Review of Biochemistry* 66: 315-335.
- Hasegawa M, Arai T, Ihara Y (1990) Immunochemical evidence that fragments of phosphorylated MAP5 (MAP1B) are bound to neurofibrillary tangles in Alzheimer's disease. *Neuron* 4: 909-918.
- Higashijima T, Burnier J, Ross EM (1990) Regulation of Gi and Go by mastoparan, related amphiphilic peptides, and hydrophobic amines. Mechanism and structural determinants of activity. *J Biol Chem* 265: 14176-14186.
- Hofer MM, Barde YA (1988) Brain-derived neurotrophic factor prevents neuronal death in vivo. *Nature* 331: 261-262.
- Hohn A, Leibrock J, Bailey K, Barde YA (1990) Identification and characterization of a novel member of the nerve growth factor/brain-derived neurotrophic factor family. *Nature* 344: 339-341.
- Huber LJ, Chao MV (1995) A potential interaction of p75 and trkA NGF receptors revealed by affinity crosslinking and immunoprecipitation. *J Neurosci Res* 40: 557-563.
- Ip NY, Ibanez CF, Nye SH, McClain J, Jones PF, Gies DR, Belluscio L, Le Beau MM, Espinosa R, III, Squinto SP, . (1992) Mammalian neurotrophin-4: structure, chromosomal localization, tissue distribution, and receptor specificity. *Proc Natl Acad Sci U S A* 89: 3060-3064.

- Irie S, Hachiya T, Rabizadeh S, Maruyama W, Mukai J, Li Y, Reed JC, Bredesen DE, Sato TA (1999) Functional interaction of Fas-associated phosphatase-1 (FAP-1) with p75(NTR) and their effect on NF-kappaB activation. *FEBS Lett* 460: 191-198.
- Jiang H, Takeda K, Lazarovici P, Katagiri Y, Yu ZX, Dickens G, Chabuk A, Liu XW, Ferrans V, Guroff G (1999) Nerve growth factor (NGF)-induced calcium influx and intracellular calcium mobilization in 3T3 cells expressing NGF receptors. *J Biol Chem* 274: 26209-26216.
- Jing S, Tapley P, Barbacid M (1992) Nerve growth factor mediates signal transduction through trk homodimer receptors. *Neuron* 9: 1067-1079.
- Johnson D, Lanahan A, Buck CR, Sehgal A, Morgan C, Mercer E, Bothwell M, Chao M (1986) Expression and structure of the human NGF receptor. *Cell* 47: 545-554.
- Johnson EM, Jr., Gorin PD, Brandeis LD, Pearson J (1980) Dorsal root ganglion neurons are destroyed by exposure in utero to maternal antibody to nerve growth factor. *Science* 210: 916-918.
- Jungbluth S, Bailey K, Barde YA (1994) Purification and characterisation of a brain-derived neurotrophic factor/neurotrophin-3 (BDNF/NT-3) heterodimer. *Eur J Biochem* 221: 677-685.
- Kang H, Welcher AA, Shelton D, Schuman EM (1997) Neurotrophins and time: different roles for TrkB signaling in hippocampal long-term potentiation. *Neuron* 19: 653-664.
- Kaplan DR, Miller FD (2000) Neurotrophin signal transduction in the nervous system. *Curr Opin Neurobiol* 10: 381-391.
- Khursigara G, Orlicki JR, Chao MV (1999) Association of the p75 neurotrophin receptor with TRAF6. *J* 274: 2597-2600.
- Kischkel FC, Hellbardt S, Behrmann I, Germer M, Pawlita M, Krammer PH, Peter ME (1995) Cytotoxicity-dependent APO-1 (Fas/CD95)-associated proteins form a death-inducing signaling complex (DISC) with the receptor. *Embo J* 14: 5579-5588.
- Klein R (1994) Role of neurotrophins in mouse neuronal development. *FASEB J* 8: 738-744.
- Kong H, Boulter J, Weber JL, Lai C, Chao MV (2001) An evolutionarily conserved transmembrane protein that is a novel downstream target of neurotrophin and ephrin receptors. *J Neurosci* 21: 176-185.
- Korte M, Carroll P, Wolf E, Brem G, Thoenen H, Bonhoeffer T (1995) Hippocampal long-term potentiation is impaired in mice lacking brain-derived neurotrophic factor. *Proc Natl Acad Sci U S A* 92: 8856-8860.
- Large TH, Weskamp G, Helder JC, Radeke MJ, Misko TP, Shooter EM, Reichardt LF (1989) Structure and developmental expression of the nerve growth factor receptor in the chicken central nervous system. *Neuron* 2: 1123-1134.

- Lee KF, Li E, Huber LJ, Landis SC, Sharpe AH, Chao MV, Jaenisch R (1992) Targeted mutation of the gene encoding the low affinity NGF receptor p75 leads to deficits in the peripheral sensory nervous system. *Cell* 69: 737-749.
- Lee R, Kermani P, Teng KK, Hempstead BL (2001) Regulation of cell survival by secreted proneurotrophins. *Science* 294: 1945-1948.
- Leibrock J, Lottspeich F, Hohn A, Hofer M, Hengeler B, Masiakowski P, Thoenen H, Barde YA (1989) Molecular cloning and expression of brain-derived neurotrophic factor. *Nature* 341: 149-152.
- Levi-Montalcini R, Hamburger V (1953) A diffusible agent of mouse sarcoma producing hyperplasia of sympathetic ganglia and hyperneurotization of viscera in the chick embryo. *Journal of Experimental Zoology* 123: 233-278.
- Lien LL, Boyce FM, Kleyn P, Brzustowicz LM, Menninger J, Ward DC, Gilliam TC, Kunkel LM (1991) Mapping of human microtubule-associated protein 1B in proximity to the spinal muscular atrophy locus at 5q13. *Proc Natl Acad Sci U S A* 88: 7873-7876.
- Lien LL, Feener CA, Fischbach N, Kunkel LM (1994) Cloning of human microtubule-associated protein 1B and the identification of a related gene on chromosome 15. *Genomics* 22: 273-280.
- Liepinsh E, Ilag LL, Otting G, Ibanez CF (1997) NMR structure of the death domain of the p75 neurotrophin receptor. *Embo J* 16: 4999-5005.
- Lindholm D, Castren E, Berzaghi M, Blochl A, Thoenen H (1994) Activity-dependent and hormonal regulation of neurotrophin mRNA levels in the brain--implications for neuronal plasticity. *J Neurobiol* 25: 1362-1372.
- Locksley RM, Killeen N, Lenardo MJ (2001) The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 104: 487-501.
- Lovestone S, Reynolds CH (1997) The phosphorylation of tau: a critical stage in neurodevelopment and neurodegenerative processes. *Neuroscience* 78: 309-324.
- Lucas FR, Goold RG, Gordon-Weeks PR, Salinas PC (1998) Inhibition of GSK-3beta leading to the loss of phosphorylated MAP-1B is an early event in axonal remodelling induced by WNT-7a or lithium. *J Cell Sci* 111 (Pt 10): 1351-1361.
- Ma D, Chow S, Obrocka M, Connors T, Fischer I (1999) Induction of microtubule-associated protein 1B expression in Schwann cells during nerve regeneration. *Brain Res* 823: 141-153.
- Ma D, Nothias F, Boyne LJ, Fischer I (1997) Differential regulation of microtubule-associated protein 1B (MAP1B) in rat CNS and PNS during development. *J Neurosci Res* 49: 319-332.
- Maisonpierre PC, Belluscio L, Squinto S, Ip NY, Furth ME, Lindsay RM, Yancopoulos GD (1990) Neurotrophin-3: a neurotrophic factor related to NGF and BDNF. *Science* 247: 1446-1451.

- McDonald NQ, Chao MV (1995) Structural determinants of neurotrophin action. *J Biol Chem* 270: 19669-19672.
- McDonald NQ, Hendrickson WA (1993) A structural superfamily of growth factors containing a cystine knot motif. *Cell* 73: 421-424.
- McDonald NQ, Lapatto R, Murray-Rust J, Gunning J, Wlodawer A, Blundell TL (1991) New protein fold revealed by a 2.3-Å resolution crystal structure of nerve growth factor. *Nature* 354: 411-414.
- Meixner A, Haverkamp S, Wassle H, Fuhrer S, Thalhammer J, Kropf N, Bittner RE, Lassmann H, Wiche G, Probst F (2000) MAP1B is required for axon guidance and is involved in the development of the central and peripheral nervous system. *J Cell Biol* 151: 1169-1178.
- Memertino GN, Croall DE (1984) Purification and characterization of a protein inhibitor of calcium-dependent proteases from rat liver. *Arch Biochem Biophys* 232: 713-720.
- Moreno FJ, Diaz-Nido J, Jimenez JS, Avila J (1999) Distribution of CK2, its substrate MAP1B and phosphatases in neuronal cells. *Mol Cell Biochem* 191: 201-205.
- Mukai J, Hachiya T, Shoji-Hoshino S, Kimura MT, Nadano D, Suvanto P, Hanaoka T, Li Y, Irie S, Greene LA, Sato TA (2000) NADE, a p75^{NTR}-associated cell death executor, is involved in signal transduction mediated by the common neurotrophin receptor p75^{NTR}. *J Biol Chem* 275: 17566-17570.
- Muramoto K, Taniguchi H, Kawahara M, Kobayashi K, Nonomura Y, Kuroda Y (1994) A substrate of ecto-protein kinase is microtubule-associated protein 1B in cortical cell cultures undergoing synaptogenesis. *Biochem Biophys Res Commun* 205: 1467-1473.
- Muzio M, Chinnaiyan AM, Kischkel FC, O'Rourke K, Shevchenko A, Ni J, Scaffidi C, Bretz JD, Zhang M, Gentz R, Mann M, Krammer PH, Peter ME, Dixit VM (1996) FLICE, a novel FADD-homologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/APO-1) death-inducing signaling complex. *Cell* 85: 817-827.
- Neet KE, Campenot RB (2001) Receptor binding, internalization, and retrograde transport of neurotrophic factors. *Cell Mol Life Sci* 58: 1021-1035.
- Nguyen MD, Lariviere RC, Julien JP (2001) Deregulation of Cdk5 in a mouse model of ALS: toxicity alleviated by perikaryal neurofilament inclusions. *Neuron* 30: 135-147.
- Nicholas RS, Winter J, Wren P, Bergmann R, Woolf CJ (1999) Peripheral inflammation increases the capsaicin sensitivity of dorsal root ganglion neurons in a nerve growth factor-dependent manner. *Neuroscience* 91: 1425-1433.
- Nilsson AS, Fainzilber M, Falck P, Ibanez CF (1998) Neurotrophin-7: a novel member of the neurotrophin family from the zebrafish. *FEBS Lett* 424: 285-290.

- Noble M, Lewis SA, Cowan NJ (1989) The microtubule binding domain of microtubule-associated protein MAP1B contains a repeated sequence motif unrelated to that of MAP2 and tau. *J Cell Biol* 109: 3367-3376.
- O'Brien ET, Salmon ED, Erickson HP (1997) How calcium causes microtubule depolymerization. *Cell Motil Cytoskeleton* 36: 125-135.
- Okamoto T, Schlegel A, Scherer PE, Lisanti MP (1998) Caveolins, a family of scaffolding proteins for organizing "preassembled signaling complexes" at the plasma membrane. [Review] [72 refs]. *J* 273: 5419-5422.
- Oppenheim RW (1991) Cell death during development of the nervous system. *Annu Rev Neurosci* 14: 453-501.
- Oshima M, Koizumi S, Fujita K, Guroff G (1989) Nerve growth factor-induced decrease in the calpain activity of PC12 cells. *J Biol Chem* 264: 20811-20816.
- Patapoutian A, Reichardt LF (2001) Trk receptors: mediators of neurotrophin action. *Curr Opin Neurobiol* 11: 272-280.
- Patel TD, Jackman A, Rice FL, Kucera J, Snider WD (2000) Development of sensory neurons in the absence of NGF/TrkA signaling in vivo. *Neuron* 25: 345-357.
- Patterson SL, Abel T, Deuel TA, Martin KC, Rose JC, Kandel ER (1996) Recombinant BDNF rescues deficits in basal synaptic transmission and hippocampal LTP in BDNF knockout mice. *Neuron* 16: 1137-1145.
- Pearson WR, Wood T, Zhang Z, Miller W (1997) Comparison of DNA sequences with protein sequences. *Genomics* 46: 24-36.
- Pedrotti B, Ulloa L, Avila J, Islam K (1996) Characterization of microtubule-associated protein MAP1B: phosphorylation state, light chains, and binding to microtubules. *Biochemistry* 35: 3016-3023.
- Penna D, Muller S, Martinon F, Demotz S, Iwashima M, Valitutti S (1999) Degradation of ZAP-70 following antigenic stimulation in human T lymphocytes: role of calpain proteolytic pathway. *J Immunol* 163: 50-56.
- Petersen M, Segond von Banchet G, Heppelmann B, Koltzenburg M (1998) Nerve growth factor regulates the expression of bradykinin binding sites on adult sensory neurons via the neurotrophin receptor p75. *Neuroscience* 83: 161-168.
- Pigino G, Paglini G, Ulloa L, Avila J, Caceres A (1997) Analysis of the expression, distribution and function of cyclin dependent kinase 5 (cdk5) in developing cerebellar macroneurons. *J Cell Sci* 110 (Pt 2): 257-270.

- Pinter M, Aszodi A, Friedrich P, Ginzburg I (1994) Calpeptin, a calpain inhibitor, promotes neurite elongation in differentiating PC12 cells. *Neurosci Lett* 170: 91-93.
- Radeke MJ, Misko TP, Hsu C, Herzenberg LA, Shooter EM (1987) Gene transfer and molecular cloning of the rat nerve growth factor receptor. *Nature* 325: 593-597.
- Radziejewski C, Robinson RC (1993) Heterodimers of the neurotrophic factors: formation, isolation, and differential stability. *Biochemistry* 32: 13350-13356.
- Rechsteiner M, Rogers SW (1996) PEST SEQUENCES AND REGULATION BY PROTEOLYSIS [Review]. *Trends in Biochemical Sciences* 21: 267-271.
- Riederer B, Cohen R, Matus A (1986) MAP5: a novel brain microtubule-associated protein under strong developmental regulation. *J Neurocytol* 15: 763-775.
- Rohrer H, Hofer M, Hellweg R, Korsching S, Stehle AD, Saadat S, Thoenen H (1988) Antibodies against mouse nerve growth factor interfere in vivo with the development of avian sensory and sympathetic neurones. *Development* 103: 545-552.
- Ross AH, Daou MC, McKinnon CA, Condon PJ, Lachyankar MB, Stephens RM, Kaplan DR, Wolf DE (1996) The neurotrophin receptor, gp75, forms a complex with the receptor tyrosine kinase TrkA. *Journal of Cell Biology* 132: 945-953.
- Rothe M, Wong SC, Henzel WJ, Goeddel DV (1994) A novel family of putative signal transducers associated with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* 78: 681-692.
- Salehi AH, Roux PP, Kubu CJ, Zeindler C, Bhakar A, Tannis LL, Verdi JM, Barker PA (2000) NRAGE, a novel MAGE protein, interacts with the p75 neurotrophin receptor and facilitates nerve growth factor-dependent apoptosis. *Neuron* 27: 279-288.
- Sanchez S, Sayas CL, Lim F, Diaz-Nido J, Avila J, Wandosell F (2001) The inhibition of phosphatidylinositol-3-kinase induces neurite retraction and activates GSK3. *J Neurochem* 78: 468-481.
- Sato K, Kawashima S (2001) Calpain function in the modulation of signal transduction molecules. *Biol Chem* 382: 743-751.
- Schoenfeld TA, McKerracher L, Obar R, Vallee RB (1989) MAP 1A and MAP 1B are structurally related microtubule associated proteins with distinct developmental patterns in the CNS. *J Neurosci* 9: 1712-1730.
- Schoenfeld TA, Obar RA (1994) Diverse distribution and function of fibrous microtubule-associated proteins in the nervous system. *Int Rev Cytol* 151: 67-137.
- Seidl K, Erck C, Buchberger A (1998) Evidence for the participation of nerve growth factor and its low-affinity receptor (p75NTR) in the regulation of the myogenic program. *J Cell Physiol* 176: 10-21.

- Shea TB, Beermann ML, Nixon RA (1991) Multiple proteases regulate neurite outgrowth in NB2a/dl neuroblastoma cells. *J Neurochem* 56: 842-851.
- Shumway SD, Maki M, Miyamoto S (1999) The PEST domain of IkappaBalpha is necessary and sufficient for in vitro degradation by mu-calpain. *J Biol Chem* 274: 30874-30881.
- Slonimsky, J. D., Yang, B., and Birren, S. J. The p75 receptor mediates a rapid switch to cholinergic transmission in sympathetic neurons. Annual Neuroscience Meeting 2001, San Diego , 924.13. 2001.
Ref Type: Abstract
- Smart EJ, Ying YS, Mineo C, Anderson RG (1995) A detergent-free method for purifying caveolae membrane from tissue culture cells. *Proc Natl Acad Sci U S A* 92: 10104-10108.
- Snider WD (1994) Functions of the neurotrophins during nervous system development: what the knockouts are teaching us. *Cell* 77: 627-638.
- Song SK, Li Shengwe, Okamoto T, Quilliam LA, Sargiacomo M, Lisanti, MP. (1996) Co-purification and direct interaction of Ras with caveolin, an integral membrane protein of caveolae microdomains. Detergent- free purification of caveolae microdomains. *J* 271: 9690-9697.
- Suter U, Heymach JV, Jr., Shooter EM (1991) Two conserved domains in the NGF propeptide are necessary and sufficient for the biosynthesis of correctly processed and biologically active NGF. *Embo J* 10: 2395-2400.
- Takei Y, Kondo S, Harada A, Inomata S, Noda T, Hirokawa N (1997) Delayed development of nervous system in mice homozygous for disrupted microtubule-associated protein 1B (MAP1B) gene. *J Cell Biol* 137: 1615-1626.
- Taniuchi M, Johnson EM, Jr., Roach PJ, Lawrence JC, Jr. (1986) Phosphorylation of nerve growth factor receptor proteins in sympathetic neurons and PC12 cells. In vitro phosphorylation by the cAMP-independent protein kinase FA/GSK-3. *J Biol Chem* 261: 13342-13349.
- Togel M, Eichinger R, Wiche G, Propst F (1999) A 45 amino acid residue domain necessary and sufficient for proteolytic cleavage of the MAP1B polyprotein precursor. *FEBS Lett* 451: 15-18.
- Togel M, Wiche G, Propst F (1998) Novel features of the light chain of microtubule-associated protein MAP1B: microtubule stabilization, self interaction, actin filament binding, and regulation by the heavy chain. *J Cell Biol* 143: 695-707.
- Tokuoka H, Saito T, Yorifuji H, Wei F, Kishimoto T, Hisanaga S (2000) Brain-derived neurotrophic factor-induced phosphorylation of neurofilament-H subunit in primary cultures of embryo rat cortical neurons. *J Cell Sci* 113 (Pt 6): 1059-1068.
- Tran PT, Joshi P, Salmon ED (1997) How tubulin subunits are lost from the shortening ends of microtubules. *J Struct Biol* 118: 107-118.

- Tremblay R, Chakravarthy B, Hewitt K, Tauskela J, Morley P, Atkinson T, Durkin JP (2000) Transient NMDA receptor inactivation provides long-term protection to cultured cortical neurons from a variety of death signals. *J Neurosci* 20: 7183-7192.
- Tucker KL, Meyer M, Barde YA (2001) Neurotrophins are required for nerve growth during development. *Nat Neurosci* 4: 29-37.
- Tucker RP, Garner CC, Matus A (1989) In situ localization of microtubule-associated protein mRNA in the developing and adult rat brain. *Neuron* 2: 1245-1256.
- Tucker RP, Matus AI (1988) Microtubule-associated proteins characteristic of embryonic brain are found in the adult mammalian retina. *Dev Biol* 130: 423-434.
- Ulloa L, Avila J, Diaz-Nido J (1993a) Heterogeneity in the phosphorylation of microtubule-associated protein MAP1B during rat brain development. *J Neurochem* 61: 961-972.
- Ulloa L, Diaz-Nido J, Avila J (1993b) Depletion of casein kinase II by antisense oligonucleotide prevents neuritogenesis in neuroblastoma cells. *Embo J* 12: 1633-1640.
- Ulloa L, Diez-Guerra FJ, Avila J, Diaz-Nido J (1994a) Localization of differentially phosphorylated isoforms of microtubule-associated protein 1B in cultured rat hippocampal neurons. *Neuroscience* 61: 211-223.
- Ulloa L, Montejo dG, Gomez-Ramos P, Moran MA, Avila J (1994b) Microtubule-associated protein MAP1B showing a fetal phosphorylation pattern is present in sites of neurofibrillary degeneration in brains of Alzheimer's disease patients. *Brain Res Mol Brain Res* 26: 113-122.
- Vara JA, Portela A, Ortin J, Jimenez A (1986) Expression in mammalian cells of a gene from *Streptomyces alboniger* conferring puromycin resistance. *Nucleic Acids Res* 14: 4617-4624.
- Viereck C, Tucker RP, Matus A (1989) The adult rat olfactory system expresses microtubule-associated proteins found in the developing brain. *J Neurosci* 9: 3547-3557.
- von Schack D, Casademunt E, Schweigreiter R, Meyer M, Bibel M, Dechant G (2001) Complete ablation of the neurotrophin receptor p75NTR causes defects both in the nervous and the vascular system. *Nat Neurosci* 4: 977-978.
- Walsh GS, Krol KM, Kawaja MD, neurotrophin r, Transgenic, Nerve gf, Axon, growth, Sympathetic, Trigeminal (1999) Absence of the p75 neurotrophin receptor alters the pattern of sympathosensory sprouting in the trigeminal ganglia of mice overexpressing nerve growth factor. *Journal of Neuroscience* 19: 258-273.
- Wang JJ, Tasinato A, Ethell DW, Testa MP, Bredesen DE (2000) Phosphorylation of the common neurotrophin receptor p75 by p38beta2 kinase affects NF-kappaB and AP-1 activities. *J Mol Neurosci* 15: 19-29.

- White JV, Stultz CM, Smith TF (1994) Protein classification by stochastic modeling and optimal filtering of amino-acid sequences. *Math Biosci* 119: 35-75.
- Wiche G, Oberkanins C, Himmler A (1991) Molecular structure and function of microtubule-associated proteins. *Int Rev Cytol* 124: 217-273.
- Wiesmann C, Ultsch MH, Bass SH, de Vos AM (1999) Crystal structure of nerve growth factor in complex with the ligand-binding domain of the TrkA receptor. *Nature* 401: 184-188.
- Wilson MT, Kisaalita WS, Keith CH (2000) Glutamate-induced changes in the pattern of hippocampal dendrite outgrowth: a role for calcium-dependent pathways and the microtubule cytoskeleton. *J Neurobiol* 43: 159-172.
- Yamashita T, Tucker KL, Barde YA (1999) Neurotrophin binding to the p75 receptor modulates Rho activity and axonal outgrowth. *Neuron* 24: 585-593.
- Yamauchi E, Titani K, Taniguchi H (1997) Specific binding of acidic phospholipids to microtubule-associated protein MAP1B regulates its interaction with tubulin. *J Biol Chem* 272: 22948-22953.
- Yan GM, Lin SZ, Irwin RP, Paul SM (1995) Activation of G proteins bidirectionally affects apoptosis of cultured cerebellar granule neurons. *J Neurochem* 65 : 2425-2431.
- Yan Q, Johnson EM, Jr. (1987) A quantitative study of the developmental expression of nerve growth factor (NGF) receptor in rats. *Developmental Biology (Orlando)* 121: 139-148.
- Yan Q, Johnson EM, Jr. (1988) An immunohistochemical study of the nerve growth factor receptor in developing rats. *Journal of Neuroscience* 8: 3481-3498.
- Ye X, Mehlen P, Rabizadeh S, VanArsdale T, Zhang H, Shin H, Wang JJ, Leo E, Zapata J, Hauser CA, Reed JC, Bredesen DE (1999) TRAF family proteins interact with the common neurotrophin receptor and modulate apoptosis induction. *J Biol Chem* 274: 30202-30208.