

6 Literatur

Achard P, Cheng H, De Grauwé L, Decat J, Schoutteten H, Moritz T, Van Der Straeten D, Peng J, Hardberd NP (2006) Integration of plant responses to environmentally activated phytohormonal signals. *Science* **311**, 91 – 94

Achard P, Vriezen WH, Van Der Straeten D, Harberd NP (2003) Ethylene regulates *Arabidopsis* development via the modulation of DELLA protein growth repressor function. *Plant Cell* **15**, 2816 – 2825

Alonso-Peral MM, Candela H, del Pozo JC, Martínez-Laborda A, Ponce MR, Micó JL (2006) The HVE/CAND1 gene is required for the early patterning of leaf venation in *Arabidopsis*. *Development* **133**, 3755 – 3766

Arteca RN, Bachman JM (1987) Light inhibition of brassinosteroid-induced ethylene production, *J. Plant Physiol.* **129**, 13 – 18

Arteca RN, Bachman JM, Mandava NB (1988) Effects of indole-3-acetic acid and brassinosteroid on ethylene biosynthesis in etiolated mung bean hypocotyl segments. *J. Plant Physiol.* **133**, 430 – 435

Ashburner M, Bonner JJ (1979) The induction of gene activity in drosophila by heat shock. *Cell* **17**, 241 – 254

Bechtold N, Ellis J, Pelletier G (1993) A binary vector based on Basta resistance for in planta transformation of *Arabidopsis thaliana*. *C.R.Acad. Sci. Paris, Science de la vie/Life Sciences* **316** 1194 – 1199

Bernhardt A, Lechner E, Hano P, Schade V, Dieterle M, Anders M, Dubin MJ, Benvenuto G, Bowler C, Genschik P, Hellmann H (2006) CUL4 associates with DDB1 and DET1 and its downregulation affects diverse aspects of development in *Arabidopsis thaliana*. *Plant J.* **47** 591 – 603

Bewley JD (1997) Seed germination and dormancy. *Plant Cell* **9**, 1055 – 1066

Birnboim HC, Doly J (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucl. Acids Res.* **7** 1513 – 1523

Bishop GJ, Koncz C (2002) Brassinosteroids and plant steroid signaling. *Plant Cell* **14**, S97 – S110

Bouchez D, Camilleri C, Caboche M (1993) A binary vector based on Basta resistance for in planta transformation of *Arabidopsis thaliana*. *C.R.Acad. Sci. Paris, Science de la vie/Life Sciences* **316** 1188 – 1193

Bradford MM (1976) A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* **72** 248 – 254

Callis J, Carpenter T, Sun CW, Vierstra RD (1995) Structure and evolution of genes encoding polyubiquitin and ubiquitin-like proteins in *Arabidopsis thaliana* ecotype Columbia. *Genetics* **139**, 921 – 939

Cao H, Glazebrook J, Clarke JD, Volko S, Dong X (1997) The *Arabidopsis* NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell* **88**, 57 – 63

Carter RS, Pennington KN, Arrate P, Oltz EM, Ballard DW (2005) Site-specific monoubiquitination of IkappaB kinase IKKbeta regulates its phosphorylation and persistent activation. *J. Biol. Chem.* **280**, 43272 – 43279

Chae HS, Faure F, Kieber JJ (2003). The eto1, eto2, and eto3 mutations and cytokinin treatment increase ethylene biosynthesis in *Arabidopsis* by increasing the stability of ACS Protein. *Plant Cell* **15**, 545 – 559

Chau V, Tobias JW, Bachmair A, Marriott D, Ecker DJ, Gonda DK, Varshavsky A (1989) A multiubiquitin chain is confined to specific lysine in a targeted short-lived protein. *Science* **243** 1576 – 1583

Chen H, Shen Y, Tang X, Yu L, Wang J, Guo L, Zhang Y, Zhan H, Feng, S, Strickland E, Zheng N, Deng XW (2006) *Arabidopsis CULLIN4* Forms an E3 Ubiquitin Ligase with RBX1 and the CDD Komplex in Mediating Light Control of Development. *Plant Cell* **18**, 1991 – 2004

Chen Y-F, Etheridge N, Schaller G (2005) Ethylene signal transduction. *Ann. Bot.* **95**, 901 – 915

Chen Z, Sliva H, Klessig DF (1993) Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid. *Science* **262**, 1883 – 1868

Ciechanover A (1998) The ubiquitin-proteasome pathway: on protein death and cell life. *EMBO J.* **17**, 7151 – 7160

Clarke SM, Mur LA, Wood JE, Scott IM (2004) Salicylic acid dependent signaling promotes basal thermotolerance but is not essential for acquired thermotolerance in *Arabidopsis thaliana*. *Plant J.* **38**, 432 – 447

Coenen C, Lomax TL (1997). Auxin-cytokinin interactions in higher plants: Old problems and new tools. *Trends Plant Sci.* **2**, 351 – 356

Cope GA, Deshaies RJ (2003) COP9 signalosome: a multifunctional regulator of SCF and other cullin-based ubiquitin ligases. *Cell* **114**, 663 – 671

Craig EA (1985) The heat shock response. *CRC Crit. Rev. Biochem.* **18**, 239 – 280

Creelman RA, Mullet JE (1997) Biosynthesis and action of jasmonates in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**, 355 – 381

Davies PJ (2004) The plant hormones: their nature, occurrence, and functions. In: Davies PJ ed. *Plant hormones: biosynthesis, signal transduction, action!* Boston, MA: Kluwer Academic Publishers, 1 – 15

del Pozo JC, Dharmasiri S, Hellmann H, Walker L, Gray WM, Estelle M (2002) AXR1-ECR1-Dependent Conjugation of RUB1 to the *Arabidopsis* Cullin AtCUL1 Is Required for Auxin Response. *Plant Cell* **14**, 421 – 433

del Pozo JC, Estelle M (1999) The *Arabidopsis* cullin AtCUL1 is modified by the ubiquitin-related protein RUB1. *Proc. Natl. Acad. Sci. USA* **96**, 15342 – 15347

del Pozo JC, Timpte C, Tan S, Callis J, Estelle M (1998) The ubiquitin-related protein RUB1 and auxin response in *Arabidopsis*. *Science* **280**, 1760 – 1763

Deshaias RJ, (1999) SCF and Cullin/Ring H2-based ubiquitin ligases. *Annu. Rev. Cell Dev. Biol.* **15**, 435 – 467

Després C, DeLong C, Glaze S, Liu E, Fobert PR (2000) The *Arabidopsis* NPR1/NIM1 protein enhances the DNA binding activity of a subgroup of the TGA family of bZIP transcription factors. *Plant Cell* **12**, 279 – 290

Devoto A, Nieto-Rostro M, Xie D, Ellis C, Harmston R, Patrick E, Davis J, Sherratt L, Coleman M, Turner JG (2002) COI1 links jasmonate signalling and fertility to the SCF ubiquitin-ligase complex in *Arabidopsis*. *Plant J.* **32**, 457 – 466

Dharmasiri N, Dharmasiri S, Jones AM, Estelle M (2003b). Auxin action in a cell-free system. *Curr. Biol.* **13**, 1418 – 1422

Dharmasiri S, Dharmasiri N, Hellmann H, Estelle M (2003a) The RUB/ Nedd8 conjugation pathway is required for early development in *Arabidopsis*. *EMBO J.* **22**, 1762 – 1770

Dieterle M, Thomann A, Renou J-P, Parmentier Y, Cognat V, Lemonnier G, Müller R, Shen W-H, Kretsch T, Genschik P (2005) Molecular and functional characterization of *Arabidopsis* Cullin 3A. *Plant J.* **41**, 386 – 399

Dikic I (2003) Mechanisms controlling EGF receptor endocytosis and degradation. *Biochem. Soc. Trans.* **31**, 1178 – 1181

Dreher K, Callis J (2007) Ubiquitin, Hormones and Biotic Stress in Plants. *Ann. Bot.* **1** – 35

Dubiel W, Gordon C (1999) Ubiquitin pathway: Another link in the polyubiquitin chain? *Curr. Biol.* **9**, R554 – R557

Dunn R, Hicke L (2001) Multiple roles for Rsp5p-dependent ubiquitination at the internalization step of endocytosis. *J. Biol. Chem.* **276**, 25974 – 25981

Durner J, Klessig DF (1995) Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. *Proc. Natl. Acad. Sci.* **92**, 11312 – 11316

Durrant WE, Dong X (2004) Systemic acquired resistance. *Annu. Rev. Phytopathol.* **42**, 185 – 209

Estelle MA, Somerville C (1987) Auxin-resistant mutants of *Arabidopsis thaliana* with an altered morphology, *Mol. Gen. Genet.* **206**, 200 – 206

Feys BJ, Benedetti CE, Penfold CN, Turner JG (1994) *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen. *Plant Cell* **6**, 751 – 759

Feys BJ, Parker JE (2000) Interplay of signaling pathways in plant disease resistance. *Trends Genet.* **16**, 449 – 455

Figueroa P, Gusmaroli G, Serino G, Habashi J, Ma L, Shen Y, Feng S, Bostick M, Callis J, Hellmann H, Deng XW (2005) *Arabidopsis* has two redundant Cullin3 proteins that are essential for embryo development and that interact with RBX1 and BTB-Proteins to form multisubunit E3 ubiquitin ligase complexes in vivo. *Plant Cell* **17**, 1180 – 1195

Finkelstein R, Gampala S, Rock C (2002) Abscisic acid signaling in seeds and seedlings. *Plant Cell* **14**, S15 – S45

Finkelstein RR (2004) The Role of Hormones during Seed Development and Germination. In: Plant Hormones: Biosynthesis, Signal Transduction, Action! Edited by Davies PJ. Kluwer Academic Publishers, Netherlands

Finkelstein RR, Gibson SI (2002) ABA and sugar interactions regulating development: cross-talk or voices in a crowd? *Curr. Opin. Plant Biol.* **5**, 26 – 32

Finley D, Sadis S, Monia BP, Boucher P, Ecker DJ, Crooke ST, Chau V (1994) Inhibition of Proteolysis and cell cycle progression in a multi-ubiquitinationdeficient yeast mutant. *Mol. Cell Biol.* **14**, 5501 – 5509

Fu X, Harberd NP (2003) Auxin promotes *Arabidopsis* root growth by modulating gibberellin response. *Nature* **421**, 740 – 743

Fulton TM, Chunwongse J, Tanksley SD (1995) Microprep protocol for extraction of DNA from tomato and other herbaceous plants. *Plant Mol. Biol. Report.* **13**, 207 – 209

Furukawa M, He YJ, Borchers C, Xiong Y (2003) Targeting of protein ubiquitination by BTB-Cullin 3-Roc1 ubiquitin ligases. *Nat. Cell Biol.* **5**, 1001 – 1007

Gagne JM, Smalle J, Gingerich DJ, Walker JM, Yoo SD, Yanagisawa S, Vierstra RD (2004) *Arabidopsis* EIN3-binding F-box 1 and 2 form ubiquitin-protein ligases that repress ethylene action and promote growth by directing EIN3 degradation. *Proc. Natl. Acad. Sci. USA* **101**, 6803 – 6808

Gazzarrini S, McCourt P (2001) Genetic interaction between ABA, ethylene and sugar signaling pathways. *Curr. Opin. Plant Biol.* **4**, 387 – 391

Gingerich DJ, Gagne JM, Salter DW, Hellmann H, Estelle M, Vierstra RD (2005) Cullins 3a and 3b assemble with members of the broad complex/tramtrack/bric-a-brac (BTB) protein family to form essential ubiquitin-protein ligases (E3s) in *Arabidopsis*. *J. Biol. Chem.* **280**, 18810 – 18821

Glickman MH, Rubin DM, Coux O, Wefes I, Pfeifer G, Cjeka Z, Baumeister W, Fried VA, Finley D (1998). A subcomplex of the proteasome regulatory particle required for ubiquitin-conjugate degradation and related to the COP9-signalosome and eIF3. *Cell* **94**, 615 – 623

Gomez-Cadenas A, Zentella R, Walker-Simmons MK, Ho T- HD (2001) Gibberellin/abscisic acid antagonism in barley aleurone cells: site of action of the protein kinase PKABA1 in relation to gibberellin signaling molecules. *Plant Cell* **13**, 667 – 679

Gray WM, del Pozo JC, Walker L, Hobbie L, Risseeuw E, Banks T, Crosby WL, Yang M, Ma H, Estelle M (1999) Identification of an SCF ubiquitin-ligase complex required for auxin response in *Arabidopsis thaliana*. *Genes Dev.* **13**, 1678 – 1691

Gray WM, Hellmann H, Dharmasiri S, Estelle M, (2002) Role of the *Arabidopsis* RING-H2 Protein RBX1 in RUB Modification and SCF Function. *Plant Cell* **14**, 2137 – 2144

Gray WM, Kepinski S, Rouse D, Leyser O, Estelle M (2001). Auxin regulates SCFTIR1-dependent degradation of AUX/IAA proteins. *Nature* **414**, 271 – 276

Gubler F, Kalla R, Roberts JK, Jacobsen JV (1995) Gibberellin-regulated expression of a myb gene in barley aleurone cells: evidence for myb transactivation of a high-pl alpha-amylase gene promoter. *Plant Cell* **7**, 1879 – 1891

Guo H, Ecker JR (2004) The ethylene signaling pathway: new insights. *Curr. Opin. Plant Biol.* **7**, 40 – 49

Haglund K, Sigismund S, Polo S, Szymkiewicz I, Di Fiore PP, Dikic I (2003) Multiple monoubiquitination of RTKs is sufficient for their endocytosis and degradation. *Nat. Cell Biol.* **5**, 461 – 466

Hanahan D (1983) Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**, 557 – 560

Hanahan D (1985) Techniques for transformation of *Escherichia coli*. In: DNA Cloning, Volume I, a practical approach. Glover, D. M. (ed.). Oxford. pp. 109 – 135

He J-X, Gendron JM, Sun Y, Gampala SS, Gendron N, Sun CQ, Wang Z-Y (2005) BZR1 Is a Transcriptional Repressor with Dual Roles in Brassinosteroid Homeostasis and Growth Responses. *Science* **307**, 1634 – 1638

He JX, Gendron JM, Yang Y, Li J, Wang ZY (2002) The GSK3-like kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the brassinosteroid signaling pathway in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **99**, 10185 – 10190

Hellmann H, Estelle M., (2002) Plant development: regulation by protein degradation. *Science* **297**(5582), 793 – 797

Hellmann H, Hobbie L, Chapman A, Dharmasiri S, Dharmasiri N, del Pozo C, Reinhardt D, Estelle M (2003). *Arabidopsis AXR6* encodes CUL1 implicating SCF E3-Ligases in auxin regulation of embryogenesis. *EMBO J.* **22**, 3314 – 3325

Hershko A, Ciechanover A (1986) The ubiquitin pathway for the degradation of intracellular proteins *Prog. Nucleic Acid Res. Molec. Biol.* **33**, 19 – 56

Hershko A, Ciechanover A (1998) The ubiquitin system; *Annu. Rev. Biochem.* **67**, 425 – 479

Hershko A, Ciechanover A, Heller H, Haas AL, Rose IA (1980) Proposed Role of ATP in Protein Breakdown: Conjugation of Proteins with Multiple Chains of the Poly-peptide of ATP-Dependent Proteolysis. *Proc. Natl. Acad. Sci. U.S.A.* **77**, 1783 – 1786

Hochstrasser M (1998) There's the rub: A novel ubiquitin-like modification linked to cell cycle regulation. *Genes Dev.* **12**, 901 – 907

Hofmann RM, Pickart CM (2001) *In Vitro* assembly and recognition of Lys-63 polyubiquitin chains. *J. Biol. Chem.* **276**, 27936 – 27943

Inada S, Ohgishi M, Mayama T, Okada K, Sakai T (2004) RPT2 is a signal transducer involved in phototropic response and stomatal opening by association with phototropin 1 in *Arabidopsis thaliana*. *Plant Cell* **16**, 887 – 896

Iten M, Hoffmann T, Grill E (1999) Receptors and signalling components of plant hormones. *Journal of Receptor & Signal Transduction Research* **19**, 41 – 58

Jackson PK, Eldridge AG, Freed E, Furstenthal L, Hsu JL, Kaiser BK, Reimann JD (2000) The lore of the RINGs: substrate recognition and catalysis by ubiquitin ligases. *Trends Cell Biol.* **10**, 429 – 439

Jupin I, Chua NH (1996) Activation of the CaMV *as-1* cis-element by salicylic acid: differential DNA-binding of a factor related to TGA1a. *EMBO J.* **15**, 5679 – 5689

Kazuhiro I (2003) An Ubiquitin Ligase Recognizing a Protein Oxidized by Iron: Implications for the Turnover of Oxidatively Damaged Proteins. *J. Biochem.* **134**, 175 – 182

Kim S, Choi H-i, Ryu H-J, Park JH, Kim MD, Kim SY (2004a) ARIA, an *Arabidopsis* Arm repeat protein interacting with a transcriptional regulator of abscisic acid-responsive gene expression, is a novel abscisic acid signaling component. *Plant Phys.* **136**, 3639 – 3648

Kim S, Kang JY, Cho DI, Park JH, Kim SY (2004b) ABF2, an ABRE-binding bZIP factor, is an essential component of glucose signaling and its overexpression affects multiple stress tolerance. *Plant J.* **40**, 75 – 87

Klee HJ (2004) Ethylene signal transduction. Moving beyond *Arabidopsis*. *Plant Phys.* **135**, 660 – 667

Laemmli UK (1970) Cleavage of structural proteins during assembly of the heat of bacteriophage T4, *Nature* **227**, 680 – 685

Lam E, Benfey PN, Gilmartin PM, Fang RX, Chua NH (1989) Site-specific mutations alter in vitro factor binding and change promoter expression pattern in transgenic plants. *Proc. Natl. Acad. Sci. USA* **86**, 7890 – 7894

Lam YA, Lawson TG, Velayutham M, Zweier JL, Pickart CM (2002) A proteasomal ATPase subunit recognizes the polyubiquitin degradation signal. *Nature* **416**, 763 – 767

Larkindale J, Knight MR (2002) Protection against heat stress-induced oxidative damage in *Arabidopsis* involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol.* **128**, 682 – 695

Leung J, Giraudat J (1998) Abscisic acid signal transduction. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **49**, 199 – 222

Li J, Brader G, Kariola T, Palva ET (2006) WRKY70 modulates the selection of signaling pathways in plant defense. *Plant J.* **46**, 477 – 491

Li J, Brader G, Palva ET (2004) The WRKY70 Transcription Factor: A Node of Convergence for Jasmonate-Mediated and Salicylate-Mediated Signals in Plant Defense. *Plant Cell* **16**, 319 – 331

Li J, Wen J, Lease KA, Doke JT, Tax FE, Walker JC (2002) BAK1, an Arabidopsis LRR receptor-like protein kinase, interacts with BRI1 and modulates brassinosteroid signaling. *Cell* **110**, 213 – 222

Lincoln C, Britton JH, Estelle M, (1990) Growth and Development of the *axr1* Mutants of *Arabidopsis*. *Plant Cell* **2**, 1071 – 1081

Lindquist S (1986) The heat-shock response. *Ann. Rev. Biochem.* **55**, 1151 – 1191

Lo SC, Hannink M (2006) CAND1-mediated substrate adaptor recycling is required for efficient repression of Nrf2 by Keap1. *Mol. Cell Biol.* **26**, 1235 – 1244

Lopez-Delgado H, Dat, JF, Foyer CH, Scott IM (1998) Induction of thermotolerance in potato microplants by acetyl salicylic acid and H₂O₂. *J. Exp. Bot.* **49**, 713 – 730

Lopez-Molina L, Mongrand S, Chua NH (2001) A postgermination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **98**, 4782–4787

Lopez-Molina L, Mongrand S, Kinoshita N, Chua N-H (2003) AFP is a novel negative regulator of ABA signaling that promotes ABI5 protein degradation. *Genes Dev.* **17**, 410 – 418

Lorenzo O, Piqueras R, Sanchez-Serrano JJ, Solano R (2003) ETHYLENE RESPONSE FACTOR1 integrates signals from ethylene and jasmonate pathways in plant defense. *Plant Cell* **15**, 165 – 178

Lucero P, Penalver E, Vela L, Lagunas R (2000) Monoubiquitination is sufficient to signal internalization of the maltose transporter in *Saccharomyces cerevisiae*. *J. Bacteriol.* **182**, 241 – 243

Lyapina S, Cope G, Shevchenko A, Serino G, Tsuge T, Zhou C, Wolf DA, Wei N, Shevchenko A, Deshaies RJ (2001) Promotion of NEDD-CUL1 conjugate cleavage by COP9 signalosome. *Science* **292**, 1382 – 1385

Mandava NB, Sasse JM, Yopp JH (1981) Brassinolide, a growth-promoting steroid lactone. II. Activity in selected gibberellin and cytokinin bioassays. *Physiol. Plant.* **53**, 453 – 461

Masson N, Willam C, Maxwell PH, Pugh CW, Ratcliffe PJ (2001) Independent function of two destruction domains in hypoxia-inducible factor-alpha chains activated by prolyl hydroxylation. *EMBO J.* **20**, 5197 – 5206

Merrill CR (1981) Ultra sensitive stain for protein in polyacrylamide gels shows regional variation in cerebrospinal fluid proteins, *Science* **211**, 1437 – 1438

Miller JH (1972) Experiments in molecular genetics. Cold Spring Harbor Laboratory, New York

Mok DW, Mok MC (1994) Cytokinins: Chemistry, Activity and Function. Boca Raton, FL: CRC Press

Molnar G, Bancos S, Nagy F, Szekeres M (2002) Characterisation of BRH1, a brassinosteroid-responsive RING-H2 gene from *Arabidopsis thaliana*. *Planta* **215**, 127 – 133

Moon J, Zhao Y, Dai X, Zhang W, Gray WM, Huq E, Estelle M (2007) A New CULLIN 1 mutant has altered responses to hormones and light in *Arabidopsis*. *Plant Physiol.* **143**, 684 – 696

Mora-Garcia S, Vert G, Yin Y, Cano-Delgado A, Cheong H, Chory J (2004) Nuclear protein phosphatases with Kelch-repeat domains modulate the response to brassinosteroids in *Arabidopsis*. *Genes Dev.* **18**, 448 – 460

Morano KA, Thiele DJ (1999) Heat shock factor function and regulation in response to cellular stress, growth, and differentiation signals. *Gene Expr.* **7**, 271 – 282

- Mosesson Y, Shtiegman K, Katz M, Zwang Y, Vereb G, Szollosi J, Yarden Y** (2003) Endocytosis of receptor tyrosine kinases is driven by monoubiquitylation, not polyubiquitylation. *J. Biol. Chem.* **278**, 21323 – 21326.
- Motchoulski A, Liscum E** (1999) *Arabidopsis* NPH3: A NPH1 photoreceptor-interacting protein essential for phototropism. *Science* **286**, 961 – 964
- Mou Z, Fan WH, Dong XN** (2003) Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* **113**, 935 – 944
- Mur LA, Kenton P, Atzorn R, Miersch O, Wasternack C** (2006) The Outcomes of Concentration-Specific Interactions between Salicylate and Jasmonate Signaling Include Synergy, Antagonism, and Oxidative Stress Leading to Cell Death. *Plant Physiol.* **140**, 249 – 262
- Müssig C** (2005) Brassinosteroid-promoted growth. *Plant Biol.* **7**, 110 – 117
- Neuenschwander U, Vernooij B, Friedrich L, Uknes S, Kessman H, Ryals J** (1995) Is hydrogen peroxide a second messenger of salicylic acid in systemic acquired resistance? *Plant J.* **8**, 227 – 233
- Norberg M, Holmlund M, Nilsson O** (2005) The *BLADE ON PETIOLE* genes act redundantly to control the growth and development of lateral organs. *Development* **132**, 2203 – 2213
- Nover L** (1990) Molekulare Zellbiologie der Hitzeantwort Teil I. *Naturwissenschaften* **77**, 310 – 316
- Nover L** (1991) Heat Shock Response. CRC Press, Boca Raton
- Nover L, Bharti K, Döring P, Mishra S, Ganguli A, Scharf K-D** (2001) *Arabidopsis* and the Hsf world: How many heat stress transcription factors do we need? *Cell Stress Chap* **6**, 177 – 189
- Paciorek T, Friml J** (2006) Auxin signaling. *J. Cell Science* **119**, 1199 – 1202

- Pavri R, Zhu B, Li G, Trojer P, Mandal S, Shilatifard A, Reinberg D** (2006) Histone H2B monoubiquitination functions cooperatively with FACT to regulate elongation by RNA polymerase II. *Cell* **125**, 703 – 717
- Peters, JM** (2002) The anaphase-promoting complex: proteolysis in mitosis and beyond. *Mol. Cell* **9**, 931 – 943
- Pickart CM** (1997) Targeting of substrates to the 26S proteasome. *Faseb J.* **11**, 1055 – 1066
- Pickart CM** (2000) Ubiquitin in chains. *Trends Biochem. Sci.* **25**, 544 – 548
- Pieterse CM, Van Loon LC** (2004) NPR1: the spider in the web of induced resistance signaling pathways. *Curr. Opin. Plant Biol.* **7**, 456 – 464
- Pintard L, Willems A, Peter M** (2004) Cullin-based ubiquitin ligases: Cul3–BTB complexes join the family. *EMBO J.* **23**, 1681 – 1687
- Pintard L, Willis JH, Willems A, Johnson JL, Srivastava M, Kurz T, Glaser S, Mains PE, Tyers M, Bowerman B, Peter M** (2003) The BTB-Protein MEL-26 is a substrate-specific adaptor of the CUL-3 ubiquitin-ligase. *Nature* **425**, 311 – 316
- Potuschak T, Lechner E, Parmentier Y, Grava S, Koncz C, Genschik P** (2003) EIN3-dependent regulation of plant ethylene hormone signaling by two *Arabidopsis* F-Box proteins: EBF1 and EBF2. *Cell* **115**, 679 – 689
- Ren S, Mandadi KK, Boedeker AL, Rathore KS, McKnight TD** (2007) Regulation of telomerase in *Arabidopsis* by BT2, an apparent target of TELOMERASE ACTIVATOR1. *Plant Cell* **19**, 21 – 31
- Reymond P, Farmer EE** (1998) Jasmonate and salicylate as global signals for defense gene expression. *Curr. Op. Plant Biol.* **1**, 404 – 411
- Rochon A, Boyle P, Wignes T, Fobert PR, Després C** (2006) The coactivator function of *Arabidopsis* NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal cysteines. *Plant Cell* **18**, 3670 – 3685

Roddick JG (1994) Comparative root growth inhibitory activity of four brassinosteroids. *Phytochem.* **37**, 1277 – 1281

Rojo E, Solano R, Sánchez-Serrano JJ (2003) Interactions Between Signaling Compounds Involved in Plant Defense. *J. Plant Growth Regul.* **22**, 82 – 98

Ross AF (1961) Systemic acquired resistance induced by localized virus infections in plants. *Virology* **14**, 340 – 358

Ryals J, Weymann K, Lawton K, Friedrich L, Ellis D, Steiner HY, Johnson J, Delaney TP, Jesse T, Vos P, Uknes S (1997) The *Arabidopsis* NIM1 protein shows homology to the mammalian transcription factor inhibitor I κ B. *Plant Cell* **9**, 425 – 439

Sambrook J, Fritch EF, Maniatis T (1989) Molecular Cloning. A laboratory manual. Second Edition, Cold Spring Harbor Laboratory Press, New York

Sasaki A, Itoh H, Gomi K, Ueguchi-Tanaka M, Ishiyama K, Kobayashi M, Jeong DH, An G, Kitano H, Ashikari M, Matsuoka M (2003) Accumulation of phosphorylated repressor for gibberellin signaling in an F-box mutant. *Science* **299**, 1896 – 1898

Sasse JM (2003) Physiological actions of brassinosteroids: an update. *J. Plant Growth Regul.* **22**, 276 – 288

Schlagnhaufer CD, Arteca RN, Yopp JH (1984) A brassinosteroid-cytokinin interaction on ethylene production by etiolated mung bean segments. *Physiol. Plant.* **60**, 347 – 350

Schwechheimer C, Deng XW, (2001) COP9 signalosome revisited: a novel mediator of protein degradation. *Trends Cell Biol.* **11**, 420 – 426

Schwechheimer C, Serino G, Callis J, Crosby WL, Lyapina S, Deshaies RJ (2001) Interactions of the COP9 signalosome with the E3 ubiquitin ligase SCF^{TIR1} in mediating auxin response. *Science* **292**, 1379 – 1382

Senaratna T, Touchell D, Bunn E, Dixon K (2000) Acetyl salicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regul.* **30**, 157 – 161

Shen WH, Parmentier Y, Hellmann H, Lechner E, Dong A, Masson J, Granier F, Lepiniec L, Estelle M, Genschik P (2002) Null mutation of AtCUL1 causes arrest in early embryogenesis in *Arabidopsis*. *Mol. Biol. Cell* **13**, 1916 – 1928

Smalle J, Kurepa J, Yang P, Babiyukh E, Kushnir S, Durski A, Vierstra RD (2002) Cytokinin growth responses in *Arabidopsis* involve the 26S proteasome sub-unit RPN12. *Plant Cell* **14**, 17 – 32

Smalle J, Vierstra RD, (2004) The ubiquitin 26S proteasome proteolytic pathway. *Annu. Rev. Plant Biol.* **55**, 555 – 590

Solano R, Stepanova A, Chao QM, Ecker JR (1998) Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSEFACTOR1. *Genes Dev.* **12**, 3703 – 3714

Spoel SH, Koornneef A, Claessens SMC, Korzelius JP, Van Pelt JA, Mueller MJ, Buchala AJ, Me' traux J-P, Brown R, Kazan K, Van Loon LC, Dong X, Pieterse CM (2003) NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *Plant Cell* **15**, 760 – 770

Stogios JP, Downs GS, Jauhal JS, Nandra SK, Privé GG (2005) Sequence and structural analysis of BTB domain proteins. *Genome Biol.* **6**, R82.1 – R82.18

Strader LC, Ritchie S, Soule JD, McGinnis KM, Steber CM (2004) Recessive-interfering mutations in the gibberellin signaling gene SLEEPY1 are rescued by over-expression of its homologue, SNEEZY. *Proc. Natl. Acad. Sci. USA* **101**, 12771 – 12776

Swain SM, Singh DP (2005) Tall tales from sly dwarves: novel functions of gibberellins in plant development. *Trends Plant Sci.* **10**, 123 – 129

Tamura N, Yoshida T, Tanaka A, Sasaki R, Bando A, Toh S, Lepiniec L, Kawakami N (2006) Isolation and characterization of high temperature-resistant germination mutants of *Arabidopsis thaliana*. *Plant Cell Physiol.* **47**, 1081 – 1094

Thomann A, Brukhin V, Dieterle M, Gheyeselinck J, Vantard M, Grossniklaus U, Genschik P (2005) *Arabidopsis* CUL3A and CUL3B genes are essential for normal embryogenesis. *Plant J.* **43**, 437 – 448

Thomas PS (1980) Hybridization of denatured RNA and small DNA fragments transferred to nitrocellulose. *Proc. Natl. Acad. Sci. U.S.A.* **77**, 5201 – 5205

Thomas SG, Sun TP (2004) Update on gibberellin signaling. A tale of the tall and the short. *Plant Physiol.* **135**, 668 – 676

Thrower JS, Hoffman L, Rechsteiner M, Pickart, CM (2000) Recognition of the polyubiquitin proteolytic signal. *EMBO J.* **19**, 94 – 102

Tian Q, Nagpal P, Reed JW (2003). Regulation of *Arabidopsis* SHY2/IAA3 protein turnover. *Plant J.* **36**, 643 – 651

Towbin H, Staehelin T, Gordon, J (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedures and some applications. *Proc. Natl. Acad. Sci. USA* **76**, 4350 – 4354

van den Heuvel S (2004) Protein degradation: CUL-3 and BTB--partners in proteolysis. *Curr. Biol.* **14**, R59 – 61

Van Loon LC, Rep M, Pieterse CM (2006) Significance of inducible defense-related proteins in infected plants. *Annu. Rev. Phytopathol.* **44**, 135 – 162

Vogel JP, Woeste KE, Theologis A, Kieber JJ (1998) Recessive and dominant mutations in the ethylene biosynthetic gene ACS5 of *Arabidopsis* confer cytokinin insensitivity and ethylene overproduction, respectively. *Proc. Natl. Acad. Sci. USA* **95**, 4766 – 4771

Voges D, Zwickl P, Baumeister W (1999). The 26S proteasome: A molecular machine designed for controlled proteolysis. *Annu. Rev. Biochem.* **68**, 1015 – 1068

Wang KL, Yoshida H, Lurin C, Ecker JR (2004) Regulation of ethylene gas biosynthesis by the *Arabidopsis* ETO1 protein. *Nature* **428**, 945 – 950

Wang ZY, He JX (2004) Brassinosteroid signal transduction--choices of signals and receptors. *Trends Plant Sci.* **9**, 91 – 96

Weber H, Bernhardt A, Dieterle M, Hano P, Mutlu A, Estelle M, Genschik P, Hellmann H (2005) *Arabidopsis* AtCUL3a and AtCUL3b form complexes with members of the BTB/POZ-MATH protein family. *Plant Physiol.* **137**, 83 – 93

Weber H, Hano P, Hellmann H (2007) The Charming Complexity of CUL3. *Int. J. Plant Dev. Biol.* **1**, 178 – 184

Wimuttisuk W, Singer JD (2007) The Cullin3 ubiquitin ligase functions as a Nedd8-bound heterodimer. *Mol. Biol. Cell* **18**, 899 – 909

Woodcock DM, Crowther PJ, Doherty J, Jefferson S, DeCruz E, Noyer-Weidner M, Smith SS, Michael MZ, Graham MW (1989) Quantitative evaluation of *Escherichia coli* host strains for tolerance to cytosine methylation in plasmid and phage recombinants. *Nucleic Acids Res.* **17**, 3469 – 3478

Woodger FJ, Jacobsen JV, Gubler F (2004) GMPOZ, a BTB/POZ domain nuclear protein, is a regulator of hormone responsive gene expression in barley aleurone. *Plant Cell Phys.* **45**, 945 – 950

Wu J-T, Chan Y-R, Chien C-T (2006). Protection of cullin-RING E3-Ligases by CSN-UBP12. *Trends in Cell Biol.* **16**, 362 – 369

Xu L, Liu F, Lechner E, Genschik P, Crosby WL, Ma H, Peng W, Huang D, Xie D (2002) The SCF(CO1) ubiquitin-ligase complexes are required for jasmonate response in *Arabidopsis*. *Plant Cell* **14**, 1919 – 1935

Xu L, Wei Y, Reboul J, Vaglio P, Shin TH, Vidal M, Elledge SJ, Harper JW (2003) BTB-Proteins are substrate-specific adaptors in an SCF-like modular ubiquitin ligase containing CUL-3. *Nature* **425**, 316 – 321

Yokota T (1999) Brassinosteroids, in: P.J.J. Hooykaas, Hall MA, Libbenga KR (Eds.), Biochemistry and Molecular Biology of Plant Hormones, Elsevier Science, London, 277 – 293

Yopp JH, Mandava NB, Sasse JM (1981) Brassinolide, a growth-promoting steroid lactone. I. Activity in selected auxin bioassays. *Physiol. Plant.* **53** 445 – 452

Yoshida Y, Chiba T, Tokunaga F, Kawasaki H, Iwai K, Suzuki T, Ito Y, Matsuoka K, Yoshida M, Tanaka K, Tai T (2002) E3 ubiquitin ligase that recognizes sugar chains. *Nature* **418**, 438 – 442

Yu H, Peters J-P, King RW, Page A, Hieter P, Kirschner MW (1998) Identification of a cullin homology region in a subunit of the Anaphase-Promoting Complex. *Science* **279**, 1219 – 1222

Zachariae WA, Shevchenko PD, Andrews R, Ciosk M, Galova MJ, Stark Mann M, Nasmyth K, (1998) Mass spectrometric analysis of the anaphase-promoting complex from yeast: Identification of a subunit related to cullins. *Science* **279**, 1216 – 1219

Zhang X, Garreton V, Chua H-H (2005) The AIP E3 ligase acts as a novel negative regulator of ABA signalling by promoting ABI3 degradation. *Genes Dev.* **19**, 1535 – 1543

Zhang Y, Tessaro MJ, Lassner M, Li X (2003) Knockout analysis of Arabidopsis transcription factors TGA2, TGA5, and TGA6 reveals their redundant and essential roles in systemic acquired resistance. *Plant Cell* **15**, 2647 – 2653

Zhang YL, Fan WH, Kinkema M, Li X, Dong XN (1999): Interaction of NPR1 with basic leucine zipper protein transcription factors that bind sequences required for salicylic acid induction of the PR-1 gene. *Proc. Natl. Acad. Sci. USA*, **96**, 6523 – 6528

Zhou JM, Trifa Y, Silva H, Pontier D, Lam E, Shah J, Klessig DF (2000): NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the PR-1 gene required for induction by salicylic acid. *Mol. Plant-Microbe Interact.* **13**, 191 – 202