

13. Literature

1. Goedert, M.; Spillantini, M. G., A century of Alzheimer's disease. *Science* **2006**, 314, (5800), 777-781.
2. Sacchettini, J. C.; Kelly, J. W., Therapeutic strategies for human amyloid diseases. *Nat. Rev. Drug Discov.* **2002**, 1, (4), 267-275.
3. Dobson, C. M., Getting out of shape. *Nature* **2002**, 418, (6899), 729-730.
4. Taylor, J. P.; Hardy, J.; Fischbeck, K. H., Toxic proteins in neurodegenerative disease. *Science* **2002**, 296, (5575), 1991-1995.
5. Stefani, M.; Dobson, C. M., Protein aggregation and toxicity: New insights into protein folding, misfolding diseases and biological evolution. *J. Mol. Med.* **2003**, 81, 678-699.
6. Guijarro, J. I.; Sunde, M.; A. Jones, J.; Campell, I. D.; Dobson, C. M., Amyloid fibril formation by an SH3 domain. *Proc. Natl. Acad. Sci. U. S. A.* **1998**, 95, 4224-4228.
7. Dobson, C. M., Protein folding and misfolding. *Nature* **2003**, 426, (6968), 884-890.
8. Bucciantini, M.; Calloni, G.; Chiti, F.; Formigli, L.; Nosi, D.; Dobson, C. M.; Stefani, M., Prefibrillar amyloid protein aggregates share common features of cytotoxicity. *J. Biol. Chem.* **2004**, 279, (30), 31374-31382.
9. Bucciantini, M.; Giannoni, E.; Chiti, F.; Baroni, F.; Formigli, L.; Zurdo, J.; Taddei, N.; Ramponi, G.; Dobson, C. M.; Stefani, M., Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature* **2002**, 416, (6880), 507-511.
10. Rochet, J.-C.; Lansbury, P. T., Amyloid fibrillogenesis: Themes and variations. *Curr. Opin. Struct. Biol.* **2000**, 10, (1), 60-68.
11. Uversky, V. N.; Fink, A. L., Conformational constraints for amyloid fibrillation: The importance of being unfolded. *Biochim. Biophys. Acta. - Proteins and Proteomics* **2004**, 1698, (2), 131-153.
12. Selkoe, D. J., Alzheimer's disease: Genes, proteins, and therapy. *Physiol. Rev.* **2001**, 81, (2), 741-766.
13. Ghiso, J.; Frangione, B., Amyloidosis and Alzheimer's disease. *Adv. Drug Delivery. Rev.* **2002**, 54, (12), 1539-1551.
14. Monsellier, E.; Chiti, F., Prevention of amyloid-like aggregation as a driving force of protein evolution. *EMBO Rep.* **2007**, 8, (8), 737-742.
15. Citron, M., Alzheimer's disease: Treatments in discovery and development. *Nat. Neurosci.* **2002**, 5, 1055-1057.

16. Virchow, R., Ueber eine im Gehirn und Rueckenmark des Menschen aufgefundenen Substanz mit der chemischen Reaction der Cellulose. *Archiv f. pathol. Anat.* **1853**, 6, (4), 135-138.
17. Puchtler, H.; Sweat, F., A review of early concepts of amyloid in context with contemporary chemical literature from 1839 to 1859. *J. Histochem. Cytochem.* **1966**, 14, (2), 123-134.
18. Snipe, J. D.; Cohen, A. S., Review: History of the amyloid fibril. *J. Struct. Biol.* **2000**, 130, 88-98.
19. Ohnishi, S.; Takano, K., Amyloid fibrils from the viewpoint of protein folding. *Cell. Mol. Life Sci.* **2004**, 61, 511-524.
20. Divry, P.; Florkin, M., Sur les propriétés optiques de l'amyloïde. *C. R. Soc. Biol.* **1927**, 97, 1808-1810.
21. Cohen, A. S.; Calkins, E., Electron microscopic observations on a fibrous component in amyloid of diverse origins. *Nature* **1959**, 183, (4669), 1202-1203.
22. Eanes, E. D.; Glenner, G. G., X-ray diffraction studies on amyloid filaments. *J. Histochem. Cytochem.* **1968**, 16, 673-677.
23. Makin, O. S.; Serpell, L. C., Structures for amyloid fibrils. *FEBS J.* **2005**, 272, 5950-5961.
24. Stromer, T.; Serpell, L. C., Structure and morphology of the Alzheimer's amyloid fibril. *Microscopy Res. Technique* **2005**, 67, 210-217.
25. Jimenez, J. L.; Nettleton, E. J.; Bouchard, M.; Robinson, C. V.; Dobson, C. M.; Saibil, H. R., The protofilament structure of insulin amyloid fibrils. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, 9196-9201.
26. Bouchard, M.; Zurdo, J.; Nettleton, E. J.; Dobson, C. M.; Robinson, C. V., Formation of insulin amyloid fibrils followed by FTIR simultaneously with CD and electron microscopy. *Protein Sci.* **2000**, 9, 1960-1967.
27. Jiménez, J. L.; Guijarro, J. I.; Orlova, E.; Zurdo, J.; Dobson, C. M.; Sunde, M.; Saibil, H. R., Cryo-electron microscopy structure of an SH3 amyloid fibril and model of the molecular packing. *EMBO J.* **1999**, 18, 815-821.
28. Goldsbury, C. S.; Cooper, G. J. S.; Goldie, K. N.; Muller, S. A.; Saafi, E. L.; Gruijters, W. T. M.; Misur, M. P.; Engel, A.; Aeby, U.; Kistler, J., Polymorphic fibrillar assembly of human amylin. *J. Struct. Biol.* **1997**, 119, (1), 17-27.
29. Malinchik, S. B.; Inouye, H.; Szumowski, K. E.; Kirschner, D. A., Structural analysis of Alzheimer's β (1-40) amyloid: Protofilament assembly of tubular fibrils. *Biophys. J.* **1998**, 74, 537-545.
30. Goldsbury, C.; Frey, P.; Olivieri, V.; Aeby, U.; Muller, S. A., Multiple assembly pathways underlie amyloid- β fibril polymorphisms. *J. Mol. Biol.* **2005**, 352, (2), 282-298.

31. Siemer, A. B.; Arnold, A. A.; Ritter, C.; Westfeld, T.; Ernst, M.; Riek, R.; Meier, B. H., Observation of highly flexible residues in amyloid fibrils of the HET-s prion. *J. Am. Chem. Soc.* **2006**, 128, (40), 13224-13228.
32. Nelson, R.; Sawaya, M. R.; Balbirnie, M.; Madsen, A. O.; Riek, C.; Grothe, R.; Eisenberg, D., Structure of the cross- β spine of amyloid-like fibrils. *Nature* **2005**, 435, (7043), 773-778.
33. Eisenberg, D.; Nelson, R.; Sawaya, M. R.; Balbirnie, M.; Sambashivan, S.; Ivanova, M. I.; Madsen, A. O.; Riek, C., The structural biology of protein aggregation diseases: Fundamental questions and some answers. *Acc. Chem. Res.* **2006**, 39, 568-575.
34. Nelson, R.; Eisenberg, D., Recent atomic models of amyloid fibril structure. *Curr. Opin. Struct. Biol.* **2006**, 16, 260-265.
35. Perutz, M. F.; Johnson, T.; Suzuki, M.; Finch, J. T., Glutamine repeats as polar zippers: Their possible role in inherited neurodegenerative diseases. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, 91, 5355-5358.
36. Perutz, M. F.; Pope, B. J.; Owen, D.; Wanker, E. E.; Scherzinger, E., Aggregation of proteins with expanded glutamine and alanine repeats of the glutamine-rich and asparagine-rich domains of Sup35 and of the amyloid β -peptide of amyloid plaques. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (8), 5596-5600.
37. Azriel, R.; Gazit, E., Analysis of the minimal amyloid-forming fragment of the islet amyloid polypeptide. An experimental support for the key role of the phenylalanine residue in amyloid formation. *J. Biol. Chem.* **2001**, 276, (36), 34156-34161.
38. Iwata, K.; Fujiwara, T.; Matsuki, Y.; Akutsu, H.; Takahashi, S.; Naiki, H.; Goto, Y., 3D structure of amyloid protofilaments of β 2-microglobulin fragment probed by solid-state NMR. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (48), 18119-18124.
39. Ferguson, N.; Becker, J.; Tidow, H.; Tremmel, S.; Sharpe, T. D.; Krause, G.; Flinders, J.; Petrovich, M.; Berriman, J.; Oschkinat, H.; Fersht, A. R., General structural motifs of amyloid protofilaments. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (44), 16248-16253.
40. Petkova, A. T.; Ishii, Y.; Balbach, J. J.; Antzutkin, O. N.; Leapman, R. D.; Delaglio, F.; Tycko, R., A structural model for Alzheimer's β -amyloid fibrils based on experimental constraints from solid state NMR. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (26), 16742-16747.
41. Petkova, A. T.; Yau, W.-M.; Tycko, R., Experimental constraints on quaternary structure in Alzheimer's β -amyloid fibrils. *Biochemistry* **2006**, 45, 498-512.
42. Tycko, R., Progress towards a molecular-level structural understanding of amyloid fibrils. *Curr. Opin. Struct. Biol.* **2004**, 14, (1), 96-103.

43. Ritter, C.; Maddelein, M.-L.; Siemer, A. B.; Lührs, T.; Ernst, M.; Meier, B. H.; Sause, S. J.; Riek, R., Correlation of structural elements and infectivity of the HET-s prion. *Nature* **2005**, 435, (7043), 844-848.
44. Lührs, T.; Ritter, C.; Adrian, M.; Riek-Lohr, D.; Bohrmann, B.; Dobeli, H.; Schubert, D.; Riek, R., 3D structure of Alzheimer's amyloid- β (1-42) fibrils. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, 102, (48), 17342-17347.
45. Egnaczyk, G. F.; Greis, K. D.; Stimson, E. R.; Maggio, J. E., Photoaffinity cross-linking of alzheimer's disease amyloid fibrils reveals interstrand contact regions between assembled β -Amyloid peptide subunits. *Biochemistry* **2001**, 40, (39), 11706-11714.
46. Lazo, N. D.; Downing, D. T., Amyloid fibrils may be assembled from β -helical protofibrils. *Biochemistry* **1998**, 37, (7), 1731-1735.
47. Perutz, M. F.; Finch, J. T.; Berriman, J.; Lesk, A., Amyloid fibers are water-filled nanotubes. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (8), 5591-5595.
48. Govaerts, C.; Wille, H.; Prusiner, S. B.; Cohen, F. E., Evidence for assembly of prions with left-handed β -helices into trimers. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, 101, (22), 8342-8347.
49. Lazo, N. D.; Downing, D. T., Beta-helical fibrils from a model peptide. *Biochem. Biophys. Res. Commun.* **1997**, 235, 675-679.
50. Williams, A. D.; Portelius, E.; Kheterpal, I.; Guo, J.-T.; Cook, K. D.; Xu, Y.; Wetzel, R., Mapping A β amyloid fibril secondary structure using scanning proline mutagenesis. *J. Mol. Biol.* **2004**, 335, (3), 833-842.
51. Singer, S. J.; Dewji, N. N., Evidence that Perutz's double-beta-stranded subunit structure for β -amyloids also applies to their channel-forming structures in membranes. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (5), 1546-1550.
52. Lomakin, A.; Chung, D. S.; Benedek, G. B.; Kirschner, D. A.; Teplow, D. B., On the nucleation and growth of amyloid β -protein fibrils: Detection of nuclei and quantitation of rate constants. *Proc. Natl. Acad. Sci. U. S. A.* **1996**, 93, 1125-1129.
53. Walsh, D. M.; Lomakin, A.; Benedek, G. B.; Condron, M. M.; Teplow, D. B., Amyloid β -protein fibrillogenesis. *J. Biol. Chem.* **1997**, 272, (35), 22364-22372.
54. Gosal, W. S.; Morten, I. J.; Hewitt, E. W.; Smith, D. A.; Thomson, N. H.; Radford, S. E., Competing pathways determine fibril morphology in the self-assembly of β 2-microglobulin into amyloid. *J. Mol. Biol.* **2005**, 351, (4), 850-864.
55. Hwang, W.; Zhang, S.; Kamm, R. D.; Karplus, M., Kinetic control of dimer structure formation in amyloid fibrillogenesis. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, 101, (35), 12916-12921.
56. Kelly, J. W., The alternative conformations of amyloidogenic proteins and their multi-step assembly pathways. *Curr. Opin. Struct. Biol.* **1998**, 8, (1), 101-106.

57. Tjernberg, L. O.; Pramanik, A.; Björling, S.; Thyberg, P.; Thyberg, J.; Nordsted, C.; Berndt, K. D.; Terenius, L.; Rigler, R., Amyloid β -peptide polymerization studied using fluorescence correlation spectroscopy. *Chem. Biol.* **1999**, 6, 53-62.
58. Calamai, M.; Chiti, F.; Dobson, C. M., Amyloid fibril formation can proceed from different conformations of a partially unfolded protein. *Biophys. J.* **2005**, 89, (6), 4201–4210.
59. Baskakov, I. V.; Legname, G.; Baldwin, M. A.; Prusiner, S. B.; Cohen, F. E., Pathway complexity of protein assembly into amyloid. *J. Biol. Chem.* **2002**, 277, (24), 21140-21148.
60. Plakoutsi, G.; Bempord, F.; Calamai, M.; Taddei, N.; Dobson, C. M.; Chiti, F., Evidence for a mechanism of amyloid formation involving molecular reorganization with native-like precursor aggregates. *J. Mol. Biol.* **2005**, 351, 910-922.
61. Soldi, G.; Bemporad, F.; Torrassa, S.; Relini, A.; Ramazzotti, M.; Taddei, N.; Chiti, F., Amyloid formation of a protein in the absence of initial unfolding and destabilization of the native state. *Biophys. J.* **2005**, 89, 4234-4244.
62. Martsev, S. P.; Dubnovitsky, A. P.; Vlasov, A. P.; Hoshino, M.; Hasegawa, K.; Naiki, H.; Goto, Y., Amyloid fibril formation of the mouse V_L domain at acidic pH. *Biochemistry* **2002**, 41, (10), 3389-3395.
63. Lai, Z.; Colon, W.; Kelly, J. W., The acid-mediated denaturation pathway of transthyretin yields a conformational intermediate that can self-assemble into amyloid. *Biochemistry* **1996**, 35, (20), 6470-6482.
64. Klajnert, B.; Cladera, J.; Bryszewska, M., Molecular interactions of dendrimers with amyloid peptides: pH dependence. *Biomacromolecules* **2006**, 7, (7), 2186-2191.
65. Aggeli, A.; Bell, M.; Carrick, L. M.; Fishwick, C. W. G.; Harding, R.; Mawer, P. J.; Radford, S. E.; Strong, A. E.; Boden, N., pH as a trigger of peptide β -sheet self-assembly and reversible switching between nematic and isotropic phases. *J. Am. Chem. Soc.* **2003**, 125, (32), 9619-9628.
66. Dong, J.; Shokes, J. E.; Scott, R. A.; Lynn, D. G., Modulating amyloid self-assembly and fibril morphology with Zn(II). *J. Am. Chem. Soc.* **2006**, 128, (11), 3540-3542.
67. Cerasoli, E.; Sharpe, B. K.; Woolfson, D. N., Zico: A peptide designed to switch folded state upon binding zinc. *J. Am. Chem. Soc.* **2005**, 127, (43), 15008-15009.
68. Karr, J. W.; Kaupp, L. J.; Szalai, V. A., Amyloid- β binds Cu²⁺ in a mononuclear metal ion binding site. *J. Am. Chem. Soc.* **2004**, 126, (41), 13534-13538.
69. Bush, A. I., Metals and neuroscience. *Curr. Opin. Chem. Biol.* **2000**, 4, (2), 184-191.

70. Bush, A. I.; Masters, C. L.; Tanzi, R. E., Copper, β -amyloid, and Alzheimer's disease: Tapping a sensitive connection. *Proc. Natl. Acad. Sci. U. S. A.* **2003**, 100, (20), 11193-11194.
71. Brown, D. R.; Kozlowski, H., Biological inorganic and bioinorganic chemistry of neurodegeneration based on prion and Alzheimer diseases. *Dalton Trans.* **2004**, (13), 1907-1917.
72. McParland, V. J.; Kad, N. M.; Kalverda, A. P.; Brown, A.; Kirwin-Jones, P.; Hunter, M. G.; Sunde, M.; Radford, S. E., Partially unfolded states of β 2-microglobulin and amyloid formation in vitro. *Biochemistry* **2000**, 39, (30), 8735-8746.
73. Zhao, H.; Tuominen, E. K. J.; Kinnunen, P. K. J., Formation of amyloid fibers triggered by phosphatidylserine-containing membranes. *Biochemistry* **2004**, 43, (32), 10302-10307.
74. Liu, W.; Prausnitz, J. M.; Blanch, H. W., Amyloid fibril formation by peptide LYS (11-36) in aqueous trifluoroethanol. *Biomacromolecules* **2004**, 5, (5), 1818-1823.
75. Siegel, S. J.; Bieschke, J.; Powers, E. T.; Kelly, J. W., The oxidative stress metabolite 4-hydroxynonenal promotes Alzheimer protofibril formation. *Biochemistry* **2007**, 46, (6), 1503-1510.
76. Dobson, C. M., Protein misfolding, evolution and disease. *Trends Biochem. Sci.* **1999**, 24, (9), 329-332.
77. Carulla, N.; Caddy, G. L.; Hall, D. R.; Zurdo, J.; Gairi, M.; Feliz, M.; Giralt, E.; Robinson, C. V.; Dobson, C. M., Molecular recycling within amyloid fibrils. *Nature* **2005**, 436, (7050), 554-558.
78. Williams, A. D.; Shivaprasad, S.; Wetzel, R., Alanine scanning mutagenesis of $\text{A}\beta$ (1-40) amyloid fibril stability. *J. Mol. Biol.* **2006**, 357, (4), 1283-1294.
79. O'Nuallain, B.; Thakur, A. K.; Williams, A. D.; Bhattacharyya, A. M.; Chen, S.; Thiagarajan, G.; Wetzel, R.; Kheterpal, I.; Wetzel, R., Kinetics and thermodynamics of amyloid assembly using a high performance liquid chromatography-based sedimentation assay. In *Methods in Enzymology*, Academic Press: 2006; Vol. 413, pp 34-74.
80. Basu, A.; Schneider, J., Model systems. *Curr. Opin. Chem. Biol.* **2006**, 10, (6), 527-528.
81. Caflisch, A., Computational models for the prediction of polypeptide aggregation propensity. *Curr. Opin. Chem. Biol.* **2006**, 10, (5), 437-444.
82. Findeis, M. A., Peptide inhibitors of beta amyloid aggregation. *Curr. Top. Med. Chem.* **2002**, 2, 417-423.
83. Pagel, K.; Vagt, T.; Koksch, B., Directing the secondary structure of polypeptides at will: From helices to amyloids and back again? *Org. Biomol. Chem.* **2005**, 3, (21), 3843-3850.

84. Céline Adessi, C. S., Beta-sheet breaker strategy for the treatment of Alzheimer's disease. *Drug Dev. Res.* **2002**, 56, (2), 184-193.
85. Mart, R. J.; Osborne, R. D.; Stevens, M. M.; Ulijn, R. V., Peptide-based stimuli-responsive biomaterials. *Soft Matter* **2006**, 2, (10), 822-835.
86. Woolfson, D. N.; Ryadnov, M. G., Peptide-based fibrous biomaterials: Some things old, new and borrowed. *Curr. Opin. Chem. Biol.* **2006**, 10, (6), 559-567.
87. Gazit, E., Mechanisms of amyloid fibril self-assembly and inhibition: Model short peptides as a key research tool. *FEBS J.* **2005**, 272, (23), 5971-5978.
88. Esteras-Chopo, A.; Serrano, L.; de la Paz, M. L., The amyloid stretch hypothesis: Recruiting proteins toward the dark side. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, 102, (46), 16672-16677.
89. Hoang, T. X.; Marsella, L.; Trovato, A.; Seno, F.; Banavar, J. R.; Maritan, A., Common attributes of native-state structures of proteins, disordered proteins, and amyloid. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (18), 6883-6888.
90. Ma, B.; Nussinov, R., Simulations as analytical tools to understand protein aggregation and predict amyloid conformation. *Curr. Opin. Chem. Biol.* **2006**, 10, (5), 445-452.
91. Lopez de la Paz, M.; Goldie, K.; Zurdo, J.; Lacroix, E.; Dobson, C. M.; Hoenger, A.; Serrano, L., De novo designed peptide-based amyloid fibrils. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (25), 16052-16057.
92. Signarvic, R. S.; DeGrado, W. F., De novo design of a molecular switch: Phosphorylation-dependent association of designed peptides. *J. Mol. Biol.* **2003**, 334, (1), 1-12.
93. Ambroggio, X. I.; Kuhlman, B., Computational design of a single amino acid sequence that can switch between two distinct protein folds. *J. Am. Chem. Soc.* **2006**, 128, (4), 1154-1161.
94. Reches, M.; Porat, Y.; Gazit, E., Amyloid fibril formation by pentapeptide and tetrapeptide fragments of human calcitonin. *J. Biol. Chem.* **2002**, 277, (38), 35475-35480.
95. Lu, K.; Jacob, J.; Thiagarajan, P.; Conticello, V. P.; Lynn, D. G., Exploiting amyloid fibril lamination for nanotube self-assembly. *J. Am. Chem. Soc.* **2003**, 125, (21), 6391-6393.
96. Jones, S.; Manning, J.; Kad, N. M.; Radford, S. E., Amyloid-forming peptides from β 2-microglobulin--insights into the mechanism of fibril formation in vitro. *J. Mol. Biol.* **2003**, 325, (2), 249-257.
97. Morgan, D. M.; Dong, J.; Jacob, J.; Lu, K.; Apkarian, R. P.; Thiagarajan, P.; Lynn, D. G., Metal switch for amyloid formation: Insight into the structure of the nucleus. *J. Am. Chem. Soc.* **2002**, 124, (43), 12644-12645.

98. Zirah, S.; Kozin, S. A.; Mazur, A. K.; Blond, A.; Cheminant, M.; Segalas-Milazzo, I.; Debey, P.; Rebuffat, S., Structural changes of region 1-16 of the Alzheimer disease amyloid β -peptide upon zinc binding and in vitro aging. *J. Biol. Chem.* **2006**, 281, (4), 2151-2161.
99. Ricchelli, F.; Buggio, R.; Drago, D.; Salmona, M.; Forloni, G.; Negro, A.; Tognon, G.; Zatta, P., Aggregation/fibrillogenesis of recombinant human prion protein and Gerstmann-Sträussler-Scheinker disease peptides in the presence of metal ions. *Biochemistry* **2006**, 45, (21), 6724-6732.
100. Shivaraprasad, S.; Wetzel, R., Scanning cysteine mutagenesis analysis of A β -(1-40) amyloid fibrils. *J. Biol. Chem.* **2006**, 281, (2), 993-1000.
101. Chiti, F.; Calamai, M.; Taddei, N.; Stefani, M.; Ramponi, G.; Dobson, C. M., Studies of the aggregation of mutant proteins in vitro provide insights into the genetics of amyloid diseases. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (90004), 16419-16426.
102. Chiti, F.; Stefani, M.; Taddei, N.; Ramponi, G.; Dobson, C. M., Rationalization of the effects of mutations on peptide and protein aggregation rates. *Nature* **2003**, 424, 805-808.
103. Krishnan, R.; Lindquist, S. L., Structural insights into a yeast prion illuminate nucleation and strain diversity. *Nature* **2005**, 435, (7043), 765-772.
104. Baldwin, A. J.; Bader, R.; Christodoulou, J.; MacPhee, C. E.; Dobson, C. M.; Barker, P. D., Cytochrome display on amyloid fibrils. *J. Am. Chem. Soc.* **2006**, 128, (7), 2162-2163.
105. Dolphin, G. T.; Dumy, P.; Garcia, J., Control of amyloid β -peptide protofibril formation by a designed template assembly. *Angew. Chem. Int. Ed.* **2006**, 45, (17), 2699-2702.
106. Jack, E.; Newsome, M.; Stockley, P. G.; Radford, S. E.; Middleton, D. A., The organization of aromatic side groups in an amyloid fibril probed by solid-state ^2H and ^{19}F NMR spectroscopy. *J. Am. Chem. Soc.* **2006**, 128, (25), 8098-8099.
107. Lowik, D. W. P. M.; Garcia-Hartjes, J.; Meijer, J. T.; vanHest, J. C. M., Tuning secondary structure and self-assembly of amphiphilic peptides. *Langmuir* **2005**, 21, (2), 524-526.
108. DeGrado, W. F.; Summa, C. M.; Pavone, V.; Nastri, F.; Lombardi, A., De novo design and structural characterization of proteins and metalloproteins. *Annu. Rev. Biochem.* **1999**, 68, (1), 779-819.
109. Ambroggio, X. I.; Kuhlman, B., Design of protein conformational switches. *Curr. Opin. Struct. Biol.* **2006**, 16, (4), 525-530.
110. Zhang, S.; Rich, A., Direct conversion of an oligopeptide from a beta -sheet to an alpha-helix: A model for amyloid formation. *Proc. Natl. Acad. Sci. U. S. A.* **1997**, 94, (1), 23-28.
111. Rajagopal, K.; Schneider, J. P., Self-assembling peptides and proteins for nanotechnological applications. *Curr. Opin. Struct. Biol.* **2004**, 14, (4), 480-486.

112. Wang, W.; Hecht, M. H., Rationally designed mutations convert de novo amyloid-like fibrils into monomeric beta -sheet proteins. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (5), 2760-2765.
113. Xu, G.; Wang, W.; Groves, J. T.; Hecht, M. H., Self-assembled monolayers from a designed combinatorial library of de novo β -sheet proteins. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, 98, (7), 3652-3657.
114. West, M. W.; Wang, W.; Patterson, J.; Mancias, J. D.; Beasley, J. R.; Hecht, M. H., De novo amyloid proteins from designed combinatorial libraries. *Proc. Natl. Acad. Sci. U. S. A.* **1999**, 96, (20), 11211-11216.
115. Kamtekar, S.; Schiffer, J. M.; Xiong, H.; Babik, J. M.; Hecht, M. H., Protein design by binary patterning of polar and nonpolar amino acids. *Science* **1993**, 262, (5140), 1680-1685.
116. Brack, A.; Orgel, L. E., Beta structures of alternating polypeptides and their possible prebiotic significance. *Nature* **1975**, 256, (5516), 383-387.
117. Schneider, J. P.; Kelly, J. W., Synthesis and efficacy of square planar copper complexes designed to nucleate β -sheet structure. *J. Am. Chem. Soc.* **1995**, 117, (9), 2533-2546.
118. Lashuel, H. A.; LaBrenz, S. R.; Woo, L.; Serpell, L. C.; Kelly, J. W., Protofilaments, filaments, ribbons, and fibrils from peptidomimetic self-assembly: Implications for amyloid fibril formation and materials science. *J. Am. Chem. Soc.* **2000**, 122, (22), 5262-5277.
119. Deechongkit, S.; Powers, E. T.; You, S. L.; Kelly, J. W., Controlling the morphology of cross β -sheet assemblies by rational design. *J. Am. Chem. Soc.* **2005**, 127, (23), 8562-8570.
120. Lamm, M. S.; Rajagopal, K.; Schneider, J. P.; Pochan, D. J., Laminated morphology of nontwisting β -sheet fibrils constructed via peptide self-assembly. *J. Am. Chem. Soc.* **2005**, 127, (47), 16692-16700.
121. Pochan, D. J.; Schneider, J. P.; Kretsinger, J.; Ozbas, B.; Rajagopal, K.; Haines, L., Thermally reversible hydrogels via intramolecular folding and consequent self-assembly of a *de novo* designed peptide. *J. Am. Chem. Soc.* **2003**, 125, (39), 11802-11803.
122. Schneider, J. P.; Pochan, D. J.; Ozbas, B.; Rajagopal, K.; Pakstis, L.; Kretsinger, J., Responsive hydrogels from the intramolecular folding and self-assembly of a designed peptide. *J. Am. Chem. Soc.* **2002**, 124, (50), 15030-15037.
123. Aggeli, A.; Nyrkova, I. A.; Bell, M.; Harding, R.; Carrick, L.; McLeish, T. C. B.; Semenov, A. N.; Boden, N., Hierarchical self-assembly of chiral rod-like molecules as a model for peptide β -sheet tapes, ribbons, fibrils, and fibers. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, 98, (21), 11857-11862.
124. Mason, J. M.; Arndt, K. M., Coiled coil domains: Stability, specificity, and biological implications. *ChemBioChem* **2004**, 5, (2), 170-176.

125. Woolfson, D. N., The design of coiled-coil structures and assemblies. *Adv. Protein Chem.* **2005**, 70, 79-112
126. Yu, Y. B., Coiled-coils: Stability, specificity, and drug delivery potential. *Adv. Drug Delivery Rev.* **2002**, 54, (8), 1113-1129
127. Adamson, J. G.; Zhou, N. E.; Hodges, R. S., Structure, function and application of the coiled-coil protein folding motif. *Curr. Opin. Biotechnol.* **1993**, 4, (4), 428-437.
128. Liu, J.; Zheng, Q.; Deng, Y.; Cheng, C.-S.; Kallenbach, N. R.; Lu, M., A seven-helix coiled coil. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (42), 15457-15462.
129. Acharya, A.; Rishi, V.; Vinson, C., Stability of 100 homo and heterotypic coiled-coil α - α' pairs for ten amino acids (A, L, I, V, N, K, S, T, E, and R). *Biochemistry* **2006**, 45, (38), 11324-11332.
130. Monera, O. D.; Kay, C. M.; Hodges, R. S., Electrostatic interactions control the parallel and antiparallel orientation of α -helical chains in two-stranded .Alpha.-helical coiled-coils. *Biochemistry* **1994**, 33, (13), 3862-3871.
131. Mutter, M.; Gassmann, R.; Buttkus, U.; Altmann, K. H., Switch-Peptide: pH-induzierte Übergänge von der α -Helix-zur β -Faltblattstruktur in bisamphiphilen Oligopeptiden. *Angew. Chem.* **1991**, 103, (11), 1504-1506.
132. Mutter, M.; Gassmann, R.; Buttkus, U.; Altmann, K. H., Switch peptides: pH-induced α -helix to β -sheet transitions of bis-amphiphilic oligopeptides. *Angew. Chem. Int. Ed. Engl.* **1991**, 30, (11), 1514-1516.
133. Schenck, H. L.; Dado, G. P.; Gellman, S. H., Redox-triggered secondary structure changes in the aggregated states of a designed methionine-rich peptide. *J. Am. Chem. Soc.* **1996**, 118, (50), 12487-12494.
134. Mihara, H.; Takahashi, Y.; Ueno, A., Design of peptides undergoing self-catalytic α to β transition and amyloidogenesis. *Peptide Sci.* **1998**, 47, (1), 83-92.
135. Takahashi, Y.; Ueno, A.; Mihara, H., Design of a peptide undergoing α - β structural transition and amyloid fibrillogenesis by the introduction of a hydrophobic defect. *Chem. Eur. J.* **1998**, 4, (12), 2475-2484.
136. Takahashi, Y.; Mihara, H., Construction of a chemically and conformationally self-replicating system of amyloid-like fibrils. *Bioorg. Med. Chem.* **2004**, 12, (4), 693-699.
137. Takahashi, Y.; Ueno, A.; Mihara, H., Amyloid architecture: Complementary assembly of heterogeneous combinations of three or four peptides into amyloid fibrils. *ChemBioChem* **2002**, 3, (7), 637-642.
138. Takahashi, Y.; Ueno, A.; Mihara, H., Heterogeneous assembly of complementary peptide pairs into amyloid fibrils with α - β structural transition. *ChemBioChem* **2001**, 2, (1), 75-79.

139. Takahashi, Y.; Ueno, A.; Mihara, H., Optimization of hydrophobic domains in peptides that undergo transformation from α -helix to β -fibril. *Bioorg. Med. Chem.* **1999**, 7, (1), 177-185.
140. Yamashita, T.; Takahashi, Y.; Takahashi, T.; Mihara, H., Inhibition of peptide amyloidnext term formation by cationic peptides with homologous sequences. *Bioorg. Med. Chem. Lett.* **2003**, 13, (22), 4051-4054
141. Fezoui, Y.; Hartley, D. M.; Walsh, D. M.; Selkoe, D. J.; Osterhout, J. J.; Teplow, D. B., A de novo designed helix-turn-helix peptide forms nontoxic amyloid fibrils. *Nat. Struct. Mol. Biol.* **2000**, 7, (12), 1095-1099.
142. Pandya, M. J.; Cerasoli, E.; Joseph, A.; Stoneman, R. G.; Waite, E.; Woolfson, D. N., Sequence and structural duality: Designing peptides to adopt two stable conformations. *J. Am. Chem. Soc.* **2004**, 126, (51), 17016-17024.
143. Ciani, B.; Hutchinson, E. G.; Sessions, R. B.; Woolfson, D. N., A designed system for assessing how sequence affects alpha to beta conformational transitions in proteins. *J. Biol. Chem.* **2002**, 277, (12), 10150-10155.
144. Pandya, M. J.; Spooner, G. M.; Sunde, M.; Thorpe, J. R.; Rodger, A.; Woolfson, D. N., Sticky-end assembly of a designed peptide fiber provides insight into protein fibrillogenesis. *Biochemistry* **2000**, 39, (30), 8728-8734.
145. Smith, A. M.; Banwell, E. F.; Edwards, W. R.; Pandya, M. J.; Woolfson, D. N., Engineering increased stability into self-assembled protein fibers. *Adv. Funct. Mater.* **2006**, 16, (8), 1022-1030.
146. Smith, A. M.; Acquah, S. F. A.; Bone, N.; Kroto, H. W.; Ryadnov, M. G.; Stevens, M. S. P.; Walton, D. R. M.; Woolfson, D. N., Polar assembly in a designed protein fiber. *Angew. Chem. Int. Ed.* **2005**, 44, (2), 325-328.
147. Melnik, T. N.; Villard, V.; Vasiliev, V.; Corradin, G.; Kajava, A. V.; Potekhin, S. A., Shift of fibril-forming ability of the designed α -helical coiled-coil peptides into the physiological pH region. *Protein Eng.* **2003**, 16, (12), 1125-1130.
148. Potekhin, S. A.; Melnik, T. N.; Popov, V.; Lanina, N. F.; Vazina, A. A.; Rigler, P.; Verdini, A. S.; Corradin, G.; Kajava, A. V., De novo design of fibrils made of short alpha-helical coiled coil peptides. *Chem. Biol.* **2001**, 8, (11), 1025-1032.
149. Kammerer, R. A.; Steinmetz, M. O., De novo design of a two-stranded coiled-coil switch peptide. *J. Struct. Biol.* **2006**, 155, (2), 146-153
150. Kammerer, R. A.; Kostrewa, D.; Zurdo, J.; Detken, A.; Garcia-Echeverria, C.; Green, J. D.; Muller, S. A.; Meier, B. H.; Winkler, F. K.; Dobson, C. M.; Steinmetz, M. O., Exploring amyloid formation by a de novo design. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, 101, (13), 4435-4440.
151. Steinmetz, M. O.; García-Echeverría, C.; Kammerer, R. A., Design of a coiled-coil-based model peptide system to explore the fundamentals of amyloid fibril formation. *Int. J. Peptide Res. Therapeut.* **2005**, 11, (1), 43-52.

152. Dong, H.; Hartgerink, J. D., Role of hydrophobic clusters in the stability of α -helical coiled coils and their conversion to amyloid-like β -sheets. *Biomacromolecules* **2007**, 8, (2), 617-623.
153. Mutter, M.; Chandravarkar, A.; Boyat, C.; Lopez, J.; Santos, S. D.; Mandal, B.; Mimna, R.; Murat, K.; Patiny, L.; Saucède, L.; Tuchscherer, G., Switch peptides in statu nascendi: Induction of conformational transitions relevant to degenerative diseases. *Angew. Chem. Int. Ed.* **2004**, 43, (32), 4172-4178.
154. Mutter, M.; Chandravarkar, A.; Boyat, C.; Lopez, J.; Santos, S. D.; Mandal, B.; Mimna, R.; Murat, K.; Patiny, L.; Saucède, L.; Tuchscherer, G., Switch-Peptide in statu nascendi: Induktion von Konformationsübergängen und deren Bedeutung in degenerativen Erkrankungen. *Angew. Chem.* **2004**, 116, (32), 4267-4273.
155. DosSantos, S.; Chandravarkar, A.; Mandal, B.; Mimna, R.; Murat, K.; Saucedo, L.; Tella, P.; Tuchscherer, G.; Mutter, M., Switch-peptides: Controlling self-assembly of amyloid β -derived peptides in vitro by consecutive triggering of acyl migrations. *J. Am. Chem. Soc.* **2005**, 127, (34), 11888-11889.
156. Mimna, R.; Camus, M.-S.; Schmid, A.; Tuchscherer, G.; Lashuel, Hilal A.; Mutter, M., Disruption of amyloid-derived peptide assemblies through the controlled induction of a β -sheet to α -helix transformation: Application of the switch concept. *Angew. Chem. Int. Ed.* **2007**, 46, (15), 2681-2684.
157. Tuchscherer, G.; Chandravarkar, A.; Camus, M.-S.; Bérard, J.; Murat, K.; Schmid, A.; Lashuel, R. M. H. A.; Mutter, M., Switch-peptides as folding precursors in self-assembling peptides and amyloid fibrillogenesis. *Peptide Sci.* **2007**, 88, (2), 239-252.
158. Mimna, R.; Camus, M.-S.; Schmid, A.; Tuchscherer, G.; Lashuel, Hilal A.; Mutter, M., Auflösung von Amyloidaggregaten durch einen kontrolliert induzierten Übergang von einer β -Faltblatt- in eine α -Helix-Struktur: Eine Anwendung des Schalterkonzepts. *Angew. Chem.* **2007**, 119, (15), 2735-2738.
159. Sohma, Y.; Hayashi, Y.; Kimura, M.; Chiyomori, Y.; Taniguchi, A.; Sasaki, M.; Kimura, T.; Kiso, Y., The "O-acyl isopeptide method" For the synthesis of difficult sequence-containing peptides: Application to the synthesis of Alzheimer's disease-related amyloid β peptide (A β) 1-42. *J. Pept. Sci.* **2005**, 11, (8), 441-451.
160. Sohma, Y.; Kiso, Y., "Click peptides" - chemical biology-oriented synthesis of Alzheimer's disease-related amyloid β peptide (A β) analogues based on the "O-acyl isopeptide method". *ChemBioChem* **2006**, 7, (10), 1549-1557.
161. Sohma, Y.; Taniguchi, A.; Yoshiya, T.; Chiyomori, Y.; Fukao, F.; Nakamura, S.; Skwarczynski, M.; Okada, T.; Ikeda, K.; Hayashi, Y.; Kimura, T.; Hirota, S.; Matsuzaki, K.; Kiso, Y., "Click peptide": A novel "O-acyl isopeptide method" For peptide synthesis and chemical biology-oriented synthesis of amyloid β peptide analogues. *J. Pept. Sci.* **2006**, 12, (12), 823-828.
162. Sohma, Y.; Yoshiya, T.; Taniguchi, A.; Kimura, T.; Hayashi, Y.; Kiso, Y., Development of o-acyl isopeptide method. *Peptide Sci.* **2007**, 88, (2), 253-262.

163. Coin, I.; Dolling, R.; Krause, E.; Bienert, M.; Beyermann, M.; Sferdean, C. D.; Carpino, L. A., Depsi peptide methodology for solid-phase peptide synthesis: Circumventing side reactions and development of an automated technique via depsidipeptide units. *J. Org. Chem.* **2006**, 71, (16), 6171-6177.
164. Hentschel, J.; Borner, H. G., Peptide-directed microstructure formation of polymers in organic media. *J. Am. Chem. Soc.* **2006**, 128, (43), 14142-14149.
165. Hentschel, J.; Krause, E.; Borner, H. G., Switch-peptides to trigger the peptide guided assembly of poly(ethylene oxide)-peptide conjugates into tape structures. *J. Am. Chem. Soc.* **2006**, 128, (24), 7722-7723.
166. Fasman, G. D., *Circular dichroism and the conformational analysis of biomolecules*. Plenum Press: New York, 1996.
167. Berova, N.; Nakanishi, K.; Woody, R. W., *Circular dichroism - principles and applications*. 2nd ed.; Wiley & Sons: New York, 2000.
168. Szilágyi, A. **2002**, Electromagnetic waves and circular dichroism: An animated tutorial, <http://www.enzim.hu/~szia/cddemo/edemo0.htm> (10.06.2007).
169. Greenfield, N. J.; Fasman, G. D., Computed circular dichroism spectra for the evaluation of protein conformation. *Biochemistry* **1969**, 8, (10), 4108-4116.
170. Carter, D. **2003**, Proteininchemist.Com, www.proteininchemist.com (10.06.2007).
171. Reed, J.; Reed, T. A., A set of constructed type spectra for the practical estimation of peptide secondary structure from circular dichroism. *Anal. Biochem.* **1997**, 254, (1), 36-40.
172. Greenfield, N. J., Methods to estimate the conformation of proteins and polypeptides from circular dichroism data. *Anal. Biochem.* **1996**, 235, (1), 1-10.
173. Puchtler, H.; Sweat, F.; Levine, M., On the binding of Congo red by amyloid. *J. Histochem. Cytochem.* **1962**, 10, (3), 355-364.
174. Wu, C.; Wang, Z.; Lei, H.; Zhang, W.; Duan, Y., Dual binding modes of Congo red to amyloid protofibril surface observed in molecular dynamics simulations. *J. Am. Chem. Soc.* **2007**, 129, (5), 1225-1232.
175. Khurana, R.; Uversky, V. N.; Nielsen, L.; Fink, A. L., Is Congo red an amyloid specific dye. *J. Biol. Chem.* **2001**, 276, (25), 22715 - 22721.
176. Lorenzo, A.; Yankner, B. A., β -Amyloid neurotoxicity requires fibril formation and is inhibited by Congo red. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, 91, (25), 12243-12247.
177. Levine-III, H., Thioflavine T interaction with synthetic Alzheimer's disease β -amyloid peptides: Detection of amyloid aggregation in solution. *Protein Sci.* **1993**, 2, (3), 404-410.
178. Ban, T.; Hamada, D.; Hasegawa, K.; Naiki, H.; Goto, Y., Direct observation of amyloid fibril growth monitored by thioflavin T fluorescence. *J. Biol. Chem.* **2003**, 278, (19), 16462-16465.

179. Khurana, R.; Coleman, C.; Ionescu-Zanetti, C.; Carter, S. A.; Krishna, V.; Grover, R. K.; Roy, R.; Singh, S., Mechanism of thioflavin T binding to amyloid fibrils. *J. Struct. Biol.* **2005**, 151, (3), 229-238.
180. Krebs, M. R. H.; Bromley, E. H. C.; Donald, A. M., The binding of thioflavin-T to amyloid fibrils: Localisation and implications. *J. Struct. Biol.* **2006**, 149, (1), 30-37.
181. Klar, T. A.; Jakobs, S.; Dyba, M.; Egner, A.; Hell, S. W., Fluorescence microscopy with diffraction resolution barrier broken by stimulated emission. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, 97, (15), 8206-8210.
182. Donnert, G.; Eggeling, C.; Hell, S. W., Major signal increase in fluorescence microscopy through dark-state relaxation. *Nat. Meth.* **2007**, 4, (1), 81-86.
183. Donnert, G.; Keller, J.; Medda, R.; Andrei, M. A.; Rizzoli, S. O.; Luhrmann, R.; Jahn, R.; Eggeling, C.; Hell, S. W., Macromolecular-scale resolution in biological fluorescence microscopy. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, (31), 11440-11445.
184. Selby, C. C., Microscopy. II. Electron microscopy: A review. *Cancer Res.* **1953**, 13, (11), 753-775.
185. The Royal Swedish Academy of Science **1986**, Press release for the noble prize of physics 1986, http://nobelprize.org/nobel_prizes/physics/laureates/1986/press.html (12.06.2007).
186. Sengebusch, P. v. **2003**, Botany online - the internet hypertextbook, from: <Http://www.Biologie.Uni-hamburg.De/b-online/e00/contents.Htm>, (20.06.2007).
187. Lottspeich, F.; Zorbas, H., *Bioanalytik*. Spektrum Akademischer Verlag Heidelberg, Berlin, 1998.
188. Kiselev, N. A.; Sherman, M. B.; Tsuprun, V. L., Negative staining of proteins. *Electron Microsc. Rev.* **1990**, 3, (1), 43-72.
189. Unger, V. M., Electron cryomicroscopy methods. *Curr. Opin. Struct. Biol.* **2001**, 11, (5), 548-554.
190. Burkhard, P.; Stetefeld, J.; Strelkov, S. V., Coiled coils: A highly versatile protein folding motif. *Trends Cell Biol.* **2001**, 11, (2), 82-88.
191. Wagschal, K.; Tripet, B.; Lavigne, P.; Mant, C.; Hodges, R. S., The role of position a in determining the stability and oligomerization state of alpha-helical coiled coils: 20 amino acid stability coefficients in the hydrophobic core of proteins. *Protein Sci.* **1999**, 8, (11), 2312-2329.
192. Wagschal, K.; Tripet, B.; Hodges, R. S., De novo design of a model peptide sequence to examine the effects of single amino acid substitutions in the hydrophobic core on both stability and oligomerization state of coiled-coils. *J. Mol. Biol.* **1999**, 285, (2), 785-803.

193. Tripet, B.; Wagschal, K.; Lavigne, P.; Mant, C. T.; Hodges, R. S., Effects of side-chain characteristics on stability and oligomerization state of a de novo-designed model coiled-coil: 20 amino acid substitutions in position "D". *J. Mol. Biol.* **2000**, 300, (2), 377-402.
194. Monera, O. D.; Sönnichsen, F. D.; Hicks, L.; Kay, C. M.; Hodges, R. S., The relative positions of alanine residues in the hydrophobic core control the formation of two-stranded or four-stranded alpha-helical coiled-coils. *Protein Eng.* **1996**, 9, (4), 353-363.
195. Monera, O. D.; Zhou, N. E.; Lavigne, P.; Kay, C. M.; Hodges, R. S., Formation of parallel and antiparallel coiled-coils controlled by the relative positions of alanine residues in the hydrophobic core. *J. Biol. Chem.* **1996**, 271, (8), 3995-4001.
196. Oakley, M. G.; Kim, P. S., A buried polar interaction can direct the relative orientation of helices in a coiled coil. *Biochemistry* **1998**, 37, (36), 12603-12610.
197. Oakley, M. G.; Hollenbeck, J. J., The design of antiparallel coiled coils. *Curr. Opin. Struct. Biol.* **2001**, 11, (4), 450-447.
198. McClain, D. L.; Bin fet, J. P.; Oakley, M. G., Evaluation of the energetic contribution of interhelical coulombic interactions for coiled coil helix orientation specificity. *J. Mol. Biol.* **2001**, 313, (2), 371-383.
199. Gurnon, D. G.; Whitaker, J. A.; Oakley, M. G., Design and characterization of a homodimeric antiparallel coiled coil. *J. Am. Chem. Soc.* **2003**, 125, (25), 7518-7519.
200. Kohn, W. D.; Monera, O. D.; Kay, C. M.; Hodges, R. S., The effects of interhelical electrostatic repulsions between glutamic acid residues in controlling the dimerization and stability of two-stranded alpha-helical coiled-coils. *J. Biol. Chem.* **1995**, 270, (43), 25495-25506.
201. Kohn, W. D.; Monera, O. D.; Kay, C. M.; Hodges, R. S., The effects of interhelical electrostatic repulsions between glutamic acid residues in controlling the dimerization and stability of two-stranded alpha-helical coiled-coils. *J. Biol. Chem.* **1995**, 270, (43), 25495-506.
202. Pagel, K.; Seeger, K.; Seiwert, B.; Villa, A.; Mark, A. E.; Berger, S.; Koksch, B., Advanced approaches for the characterization of a de novo designed antiparallel coiled coil peptide. *Org. Biomol. Chem.* **2005**, 3, 1189 - 1194.
203. Burkhard, P.; Ivaninskii, S.; Lustig, A., Improving coiled-coil stability by optimizing ionic interactions. *J. Mol. Biol.* **2002**, 318, (3), 901-910.
204. Lacroix, E.; Kortemme, T.; Lopez de la Paz, M.; Serrano, L., The design of linear peptides that fold as monomeric beta-sheet structures. *Curr. Opin. Struct. Biol.* **1999**, 9, (4), 487-493.
205. Blanco, F. J.; Rivas, G.; Serrano, L., A short linear peptide that folds into a native stable β-hairpin in aqueous solution. *Nat. Struct. Mol. Biol.* **1994**, 1, (9), 584-590.

206. Yan, Y.; Erickson, B. W., Engineering of betabellin 14d: Disulfide-induced folding of a β -sheet protein. *Protein Sci.* **1994**, 3, (7), 1069-1073.
207. Quinn, T. P.; Tweedy, N. B.; Williams, R. W.; Richardson, J. S.; Richardson, D. C., Betadoublet: De novo design, synthesis, and characterization of a β -sandwich protein. *Proc. Natl. Acad. Sci. U. S. A.* **1994**, 91, (19), 8747-8751.
208. Ramirez-Alvarado, M.; Daragan, V. A.; Serrano, L.; Mayo, K. H., Motional dynamics of residues in a beta-hairpin peptide measured by ^{13}C -NMR relaxation. *Protein Sci.* **1998**, 7, (3), 720-729.
209. Kortemme, T.; Ram, I.; Rez-Alvarado, M.; Serrano, L., Design of a 20-amino acid, three-stranded β -sheet protein. *Science* **1998**, 281, (5374), 253-256.
210. Searle, M. S.; Ciani, B., Design of beta-sheet systems for understanding the thermodynamics and kinetics of protein folding. *Curr. Opin. Struct. Biol.* **2004**, 14, (4), 458-464.
211. Stanger, H. E.; Gellman, S. H., Rules for antiparallel β -sheet design: D-Pro-Gly is superior to L-Asn-Gly for β -hairpin nucleation. *J. Am. Chem. Soc.* **1998**, 120, (17), 4236-4237.
212. Tramontano, A., A brighter future for protein design. *Angew. Chem. Int. Ed.* **2004**, 43, (25), 3222-3223.
213. Pham, T. N.; Koide, A.; Koide, S., A stable single-layer beta-sheet without a hydrophobic core. *Nat. Struct. Biol.* **1998**, 5, (2), 115-119.
214. Koide, S.; Huang, X.; Link, K.; Koide, A.; Bu, Z.; Engelman, D. M., Design of single-layer beta-sheets without a hydrophobic core. *Nature* **2000**, 403, (6768), 456-460.
215. Balbirnie, M.; Grothe, R.; Eisenberg, D. S., An amyloid-forming peptide from the yeast prion Sup35 reveals a dehydrated β -sheet structure for amyloid. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, 98, (23), 2375-2380.
216. Perutz, M. F.; Pope, B. J.; Owen, D.; Wanker, E. E.; Scherzinger, E., Aggregation of proteins with expanded glutamine and alanine repeats of the glutamine-rich and asparagine-rich domains of Sup35 and of the amyloid beta-peptide of amyloid plaques. *Proc. Natl. Acad. Sci. U. S. A.* **2002**, 99, (8), 5596-5600.
217. Chou, P. Y.; Fasman, G. D., Prediction of protein conformation. *Biochemistry* **1974**, 13, (2), 222-245.
218. Chou, P. Y.; Fasman, G. D., Empirical predictions of protein conformation. *Annu. Rev. Biochem.* **1978**, 47, 251-276.
219. Chou, P. Y.; Fasman, G. D., Prediction of the secondary structure of proteins from their amino acid sequence. *Adv. Enzymol. Relat. Areas Mol. Biol.* **1978**, 47, 45-148.

220. Chou, P. Y.; Fasman, G. D., Conformational parameters for amino acids in helical, beta-sheet, and random coil regions calculated from proteins. *Biochemistry* **1974**, 13, (2), 211-222.
221. Samedov, K. Design und Synthese von switch-peptiden. Diploma Thesis, Freie Universität Berlin, Berlin, Germany, 2005.
222. Mühlberg, M. Untersuchungen zur Sequenz-Struktur-Beziehung amyloidbildender coiled coil-Peptide. Bachelor Thesis, Freie Universität Berlin, Berlin, Germany, 2006.
223. Araghi, R. R. Structural investigation on amyloid forming model peptides based on environmental factors. Master Thesis, Freie Universität Berlin, Berlin, Germany, 2006.
224. Zhou, N. E.; Kay, C. M.; Hodges, R. S., The net energetic contribution of interhelical electrostatic attractions to coiled-coil stability. *Protein Eng.* **1994**, 7, (11), 1365-1372.
225. Kohn, W. D.; Monera, O. D.; Kay, C. M.; Hodges, R. S., The effects of interhelical electrostatic repulsions between glutamic acid residues in controlling the dimerization and stability of two-stranded alpha-helical coiled-coils. *J. Biol. Chem.* **1995**, 270, (43), 25495-25506.
226. Kohn, W. D.; Kay, C. M.; Hodges, R. S., Protein destabilization by electrostatic repulsions in the two-stranded α -helical coiled-coil/leucine zipper. *Protein Sci.* **1995**, 4, (2), 237-250.
227. Pagel, K.; Wagner, S. C.; Samedov, K.; von Berlepsch, H.; Böttcher, C.; Koksch, B., Random coils, β -sheet ribbons, and α -helical fibers: One peptide adopting three different secondary structures at will. *J. Am. Chem. Soc.* **2006**, 128, (7), 2196-2197.
228. Wagner, S. C.; Araghi, R. R.; Pagel, K.; Berlepsch, H. v.; Böttcher, C.; Koksch, B., **2007**, submitted.
229. Wagner, S. C. Strukturelle Charakterisierung eines auf dem α -helikalen coiled coil-Motiv basierenden switch-Peptids. Diploma Thesis, Freie Universität Berlin, Berlin, Germany, 2005.
230. Klemba, M.; Gardner, K. H.; Marino, S.; Clarke, N. D.; Regan, L., Novel metal-binding proteins by design. *Nat. Struct. Mol. Biol.* **1995**, 2, (5), 368-373.
231. Schneider, J. P.; Kelly, J. W., Templates that induce α -helical, β -sheet, and loop conformations. *Chemical Reviews* **1995**, 95, (6), 2169-2187.
232. Ramadan, D.; Cline, D. J.; Bai, S.; Thorpe, C.; Schneider, J. P., Effects of As(III) binding on β -hairpin structure. *J. Am. Chem. Soc.* **2007**, 129 (10), 2981 - 2988.
233. Ambroggio, X. I.; Kuhlman, B., Computational design of a single amino acid sequence that can switch between two distinct protein folds. *J. Am. Chem. Soc.* **2006**, 128, (4), 1154-1161.

234. Doerr, A. J.; McLendon, G. L., Design, folding, and activities of metal-assembled coiled coil proteins. *Inorg. Chem.* **2004**, 43, (25), 7916-7925.
235. Tsurkan, M. V.; Ogawa, M. Y., Metal-mediated peptide assembly: Use of metal coordination to change the oligomerization state of an α -helical coiled-coil. *Inorg. Chem.* **2007**.
236. Dieckmann, G. R.; McRorie, D. K.; Lear, J. D.; Sharp, K. A.; DeGrado, W. F.; Pecoraro, V. L., The role of protonation and metal chelation preferences in defining the properties of mercury-binding coiled coils. *J. Mol. Biol.* **1998**, 280, (5), 897-912.
237. Iranzo, O.; Ghosh, D.; Pecoraro, V. L., Assessing the integrity of designed homomeric parallel three-stranded coiled coils in the presence of metal ions. *Inorg. Chem.* **2006**, 45, (25), 9959-9973.
238. Lee, K.-H.; Cabello, C.; Hemmingsen, L.; Marsh, E. N. G.; Pecoraro, V. L., Using nonnatural amino acids to control metal-coordination number in three-stranded coiled coils. *Angew. Chem. Int. Ed.* **2006**, 45, (18), 2864-2868.
239. Iranzo, O.; Cabello, C.; Pecoraro, V. L., Heterochromia in designed metallopeptides: Geometry-selective binding of Cd^{2+} in a de novo peptide. *Angew. Chem. Int. Ed.* **2007**, 9999, (9999), NA.
240. Kohn, W. D.; Kay, C. M.; Sykes, B. D.; Hodges, R. S., Metal ion induced folding of a de novo designed coiled-coil peptide. *J. Am. Chem. Soc.* **1998**, 120, (6), 1124-1132.
241. Futaki, S.; Kiwada, T.; Sugiura, Y., Control of peptide structure and recognition by Fe(III)-induced helix destabilization. *J. Am. Chem. Soc.* **2004**, 126, (48), 15762-15769.
242. Kiwada, T.; Sonomura, K.; Sugiura, Y.; Asami, K.; Futaki, S., Transmission of extramembrane conformational change into current: Construction of metal-gated ion channel. *J. Am. Chem. Soc.* **2006**, 128, (18), 6010-6011.
243. Krantz, B. A.; Sosnick, T. R., Engineered metal binding sites map the heterogeneous folding landscape of a coiled coil. *Nat. Struct. Mol. Biol.* **2001**, 8, (12), 1042-1047.
244. Pagel, K.; Vagt, T.; Kohajda, T.; Koksch, B., From α -helix to β -sheet - a reversible metal ion induced peptide secondary structure switch. *Org. Biomol. Chem.* **2005**, 3, (14), 2500-2502.
245. Sonnichsen, F. D.; Van Eyk, J. E.; Hodges, R. S.; Sykes, B. D., Effect of trifluoroethanol on protein secondary structure: An NMR and CD study using a synthetic actin peptide. *Biochemistry* **1992**, 31, (37), 8790-8798.
246. Seri, T. Influence of metal ions on the secondary structure of de novo designed peptides. Master Thesis, Freie Universität Berlin, Berlin, Germany, 2005.
247. Pagel, K.; Seri, T.; Berlepsch, H. v.; Griebel, J.; Kirmse, R.; Böttcher, C.; Koksch, B., How metal ions affect amyloid formation – Cu^{2+} and Zn^{2+} sensitive peptides. **2007**, in preparation.

248. Roe, R. R.; Pang, Y.-P., Zinc's exclusive tetrahedral coordination governed by its electronic structure. *Journal of Molecular Modeling* **1999**, 5, (7-8), 134-140.
249. Holm, R. H.; Kennepohl, P.; Solomon, E. I., Structural and functional aspects of metal sites in biology. *Chem. Rev.* **1996**, 96, (7), 2239-2314.
250. Herring, F. G.; Thompson, R. C.; Schwerdtfeger, C. F., *Can. J. Chem.* **1969**, 47, 555-558.
251. Preisach, J.; Blumberg, W. E., Structural implications derived from the analysis of epr spectra of natural and artificial copper proteins. *Arch. Biochem. Biophys.* **1974**, 165, 691-708.
252. Karr, J. W.; Akintoye, H.; Kaupp, L. J.; Szalai, V. A., N-terminal deletions modify the Cu²⁺ binding site in amyloid-β. *Biochemistry* **2005**, 44, (14), 5478-5487.
253. Jun, S.; Saxena, S., The aggregated state of amyloid-β peptide in vitro depends on Cu²⁺ ion concentration. *Angew. Chem. Int. Ed.* **2007**, 46, (21), 3959 - 3961.
254. Rasila, R. M.; Bertoncini, C. W.; Marsh, D.; Hoyer, W.; Cherny, D.; Zweckstetter, M.; Griesinger, C.; Jovin, T. M.; Fernandez, C. O., Structural characterization of copper(II) binding to α-synuclein: Insights into the bioinorganic chemistry of parkinson's disease. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, 102, (12), 4294-4299.
255. Viles, J. H.; Cohen, F. E.; Prusiner, S. B.; Goodin, D. B.; Wright, P. E.; Dyson, H. J., Copper binding to the prion protein: Structural implications of four identical cooperative binding sites. *Proc. Natl. Acad. Sci. U. S. A.* **1999**, 96, (5), 2042-2047.
256. Stellato, F.; Menestrina, G.; Serra, M. D.; Potrich, C.; Tomazzolli, R.; Meyer-Klaucke, W.; Morante, S., Metal binding in amyloid β-peptides shows intra- and inter-peptide coordination modes. *Eur. Biophys. J.* **2006**, 35, 340-351.
257. Aebersold, R.; Mann, M., Mass spectrometry-based proteomics. *Nature* **2003**, 422, (6928), 198-207.
258. Kaltahov, I. A.; Eyles, S. J., *Mass spectrometry in biophysics - conformation and dynamics of biomolecules*. John Wiley & Sons, Inc.: Hoboken, New Jersey, 2005.
259. Englander, S. W., Hydrogen exchange and mass spectrometry: A historical perspective. *J. Am. Soc. Mass Spectrom.* **2006**, 17, 1481-1489.
260. Thomas E. Wales, J. R. E., Hydrogen exchange mass spectrometry for the analysis of protein dynamics. *Mass Spectrom. Rev.* **2006**, 25, (1), 158-170.
261. Sinz, A., Chemical cross-linking and mass spectrometry for mapping three-dimensional structures of proteins and protein complexes. *J. Mass Spectrom.* **2003**, 38, (12), 1225-37.
262. Sinz, A., Investigation of protein-ligand interactions by mass spectrometry. *ChemMedChem* **2007**, 2, 425-431.

263. Loo, J. A., Studying noncovalent protein complexes by electrospray ionization mass spectrometry. *Mass Spectrom. Rev.* **1997**, 16, (1), 1-23.
264. Ashcroft, A. E., Recent developments in electrospray ionisation mass spectrometry: Noncovalently bound protein complexes. *Nat. Prod. Rep.* **2005**, 22, (4), 452-464.
265. Benesch, J. L. P.; Ruotolo, B. T.; Simmons, D. A.; Robinson, C. V., Protein complexes in the gas phase: Technology for structural genomics and proteomics. *Chem. Rev.* **2007**, 107, (8), 3544-3567.
266. Ruotolo, B. T.; Giles, K.; Campuzano, I.; Sandercock, A. M.; Bateman, R. H.; Robinson, C. V., Evidence for macromolecular protein rings in the absence of bulk water. *Science* **2005**, 310, (5754), 1658-1661.
267. Ruotolo, B. T.; Robinson, C. V., Aspects of native proteins are retained in vacuum. *Curr. Opin. Chem. Biol.* **2006**, 10, (5), 402-408.
268. Wolynes, P. G., Biomolecular folding in vacuo!!!(?). *Proc. Natl. Acad. Sci. U. S. A.* **1995**, 92, (7), 2426-2427.
269. Oomens, J.; Polfer, N.; Moore, D. T.; Meer, L. v. d.; Marshall, A. G.; Eyler, J. R.; Meijer, G.; Helden, G. v., Charge-state resolved mid-infrared spectroscopy of a gas-phase protein. *Phys. Chem. Chem. Phys.* **2005**, 7, (7), 1345-1348.
270. Shelimov, K. B.; Clemmer, D. E.; Hudgins, R. R.; Jarrold, M. F., Protein structure in vacuo: Gas-phase conformations of BPTI and Cytochrome C. *J. Am. Chem. Soc.* **1997**, 119, (9), 2240-2248.
271. Wendt, H.; Durr, E.; Thomas, R. M.; Przybylski, M.; Bosshard, H. R., Characterization of leucine zipper complexes by electrospray ionization mass spectrometry. *Protein Sci.* **1995**, 4, (8), 1563-1570.
272. Valle, J. J.; Eyler, J. R.; Oomens, J.; Moore, D. T.; Meer, A. F. G. v. d.; Helden, G. v.; Meijer, G.; Hendrickson, C. L.; Marshall, A. G.; Blakney, G. T., Free electron laser-fourier transform ion cyclotron resonance mass spectrometry facility for obtaining infrared multiphoton dissociation spectra of gaseous ions. *Rev. Sci. Instrum.* **2005**, 76, (2), 023103.
273. Polfer, N. C.; Oomens, J., Reaction products in mass spectrometry elucidated with infrared spectroscopy. *Phys. Chem. Chem. Phys.* **2007**, 9, (29), 3804-3817.
274. Oepts, D.; Meer, A. F. G. v. d.; Amersfoort, P. W. v., The free-electron-laser user facility FELIX. *Infrared Phys. Technol.* **1995**, 36, (1), 297-308.
275. Kupser, P. Konstruktion einer Elektrosprayquelle zur Erzeugung solvatisierter Biomoleküle – Nachweis von Sekundärstrukturen in der Gasphase. Diploma Thesis, Freie Universität Berlin, Berlin, Germany, 2007.