

## H Literaturverzeichnis

- Andrew, M., Carter, C., O`Brodovich, H., Heigenhauser, G.: Increases in factor VIII complex and fibrinolytic activity are dependent on exercise intensity. *J. Appl. Physiol.* 60(6): 1917-1922, 1986
- Arai, M., Yorifuji, H., Ikematsu, S. et al.: Influence of strenuous exercise (triathlon) on blood coagulation and fibrinolytic system. *Thromb. Res.* 57: 465-471, 1990
- Astrand, P.O. und Rodahl, K.: *Textbook of work physiology.* McGraw-Hill, New York, 1986
- Astrup, T.: Fibrinolysis in the organism. *Blood* 11: 781-806, 1956
- Bach-Gansmo, E.T., Godal, H.C., Skjonsberg, O.H.: Degradation of fibrinogen and cross-linked fibrin by human neutrophil elastase generates D-like fragments detected by ELISA but not latex D-dimer test. *Thromb. Res.* 92(3): 125-134, 1998
- Bärtsch, P., Welsch, B., Albert, M., Friedman, B., Levi, M., Kruithof, E.K.O.: Balanced activation of coagulation and fibrinolysis after a 2-h triathlon. *Med. Sci. Sports Exerc.* 27: 1465-1470, 1995
- Bärtsch, P., Haerberli, A., Straub, P.W.: Blood coagulation after long distance running: antithrombin III prevents fibrin formation. *Thromb. Hemost.* 63: 430-434, 1990
- Banfi, G., Marinelli, M., Roi, G.S., Giacometti, M.: Platelet indices in athletes performing a race in altitude environment. *J. Clin. Lab. Anal.* 9: 34-36, 1995
- Barold, S.S., Hannss, R., Falkoff, M.D., Sengupta, A.: Exercise-induced myocardial infarction due to coronary thrombosis. *Am. Heart J.* 109: 590-593, 1985
- Bar-Or, O., Dotan, R., Inbar, O.: A 30-second all-out ergometric test: Its reliability and validity for anaerobic capacity. *Israel J. Med. Sci.* 13: 326, 1977
- Bar-Or, O., Dotan, R., Inbar, O., Rotstein, A., Karlsson, J., Tesch, P.: Anaerobic capacity and muscle fibre type distribution in man. *Int. J. Sports Med.* 1: 89-92, 1980
- Barthels, M. und Poliwoda, H.: *Gerinnungsanalysen.* 5. Auflage, Thieme-Verlag, Stuttgart, 1997
- Bedrak, E., Beer, G., Furman, K.I.: Fibrinolytic activity and muscular exercise at high altitude. *J. Appl. Physiol.* 19: 469-471, 1964
- Beneke, R.: *Laktat bei Dauerleistung.* Habilitationsschrift, FU Berlin, 1999

- Beneke, R.: Maximal lactate steady state concentration (MLSS): experimental and modelling approaches. *Eur. J. Appl. Physiol.* 88: 361-369, 2003
- Beneke, R., Pollmann, Ch., Bleif, I., Leithäuser R.M., Hütler M.: How anaerobic is the wingate anaerobic test? *Eur. J. Appl. Physiol.* 87:388-392, 2002
- Berlin, J.A. und Colditz, G.A.: A meta-analysis of physical activity in the prevention of CHD. *Am. J. Epidemiol.* 132: 612-628, 1990
- Biggs, R., Mc Farlane, R.G., Pilling, J.: Observations on Fibrinolysis. Experimental activity produced by exercise or adrenaline. *Lancet* 1: 402-405, 1947
- Bourey, R.E. und Santoro, S.A.: Interactions of exercise, coagulation, platelets, and fibrinolysis: a brief review. *Med. Sci. Sports Exerc.* 20: 439-446, 1988
- Brommer, E.J.P., Barrett-Bergshoeff, M.M., Allen, R.A., Schicht, I., Bertina, R.M., Schalekamp, M.A.D.H.: The use of desmopressin (DDAVP) as a test of the fibrinolytic capacity of patients – analysis of responders and non-responders. *Thromb. Hemost.* 48: 156-161, 1982
- Brooks, G.A.: Anaerobic threshold: review of the concept and directions for future research. *Med. Sci. Sports Exerc.* 17(1): 22-31, 1985
- Cash, J.D., Woodfield, D.G., Allan, A.G.E.: Adrenergic mechanisms in the systemic plasminogen activator response to adrenalin in man. *Brit. J. Haematol.* 18: 487-494, 1970
- Chandler, W.L., Veith, R.C., et al.: Fibrinolytic response during exercise and epinephrine infusion in the same subjects. *JACC* 19(7): 1412-1420, 1992
- Chicharro, J.L., Sanchez, O., Brandes, F. et al.: Platelet aggregability in relation to the anaerobic threshold. *Thromb. Res.* 75: 251-257, 1994
- Ciampricotti, R., Gamal, M.E., Relik, T. et al.: Clinical characteristics and coronary angiographic findings of patients with unstable angina, acute myocardial infarction, and survivors of sudden ischemic death occurring during and after sport. *Am. Heart J.* 120: 1267-1278, 1990
- Cohen, R.J., Epstein, S.E., Cohen, S.L., Dennis, L.H.: Alterations in blood fibrinolysis and blood coagulation induced by exercise and the role of beta-adrenergic receptor stimulation. *Lancet* 2: 1264-1266, 1968
- Collen, D.: On the regulation and Control of Fibrinolysis. *Thromb. Haemost.* 43: 77-89, 1980

- Crowell, J.M., Houston, B.: Effect of acidity on blood coagulation. *Am. J. Physiol.* 201: 379-382, 1961
- Dahlbäck, B.: Resistance to Activated Protein C caused by Arg<sup>506</sup> to Gln Mutation in Factor V as a Pathogenic Risk Factor for Venous Thrombosis. 4. Würzburger Hämostaseologie-Symposium, 14.-15.09.1995, Abstractband, S.25, 1995
- Davis, G.L., Abidgaard, C.F., Bernauer, E.M., Britton, M.: Fibrinolytic and hemostatic changes during and after maximal exercise in males. *J. Appl. Physiol.* 40: 287-292, 1976
- De Paz, J.A., Lasierra, J., Villa, J.G., Vilades, E., et al.: Changes in the fibrinolytic system associated with physical conditioning. *Eur. J. Appl. Physiol.* 65: 388-393, 1992
- De Paz, J.A., Lasierra, J., Villa, J.G., Vilades, E., Gonzales-Gallego, J.: Effects of aerobic and anaerobic physical conditioning on fibrinolysis. *J. Sports Med. Phys. Fitness* 35(4): 263-267, 1995
- Deutsche Gesellschaft für Sportmedizin und Prävention e.V., Expertenkommission. Leitlinien zur Belastungsuntersuchung in der Sportmedizin. <http://www.dgsp.de/leitlinie.html>, 2002
- Dill, D.B. und Costill, D.L.: Calculation of percentage changes in volume of blood, plasma, and red cells in dehydration. *J. Appl. Physiol.* 37: 247-248, 1974
- Dost, F.H.: Grundlagen der Pharmakokinetik. Georg Thieme Verlag, Stuttgart, S. 29-62, 1968
- Drygas, W.K.: Changes in blood platelet function, coagulation, and fibrinolytic activity in response to moderate, exhaustive, and prolonged exercise. *Int. J. Sports Med.* 9(1): 67-72, 1988
- Dufaux, B., Order, U., Hollmann, W.: Can physical exercise induce an effective fibrinolysis? *Thromb. Res.* 36: 37-43, 1984
- Dufaux, B., Order, U., Liesen, H.: Effect of short maximal physical exercise on coagulation, fibrinolysis, and complement system. *Int. J. Sports Med.* 12 (Suppl.1): S38-42, 1991
- El-Sayed, M.S.: Fibrinolytic and hemostatic parameters response after resistance exercise. *Med. Sci. Sports Exerc.*: 25: 597-602, 1993
- El-Sayed, M.S., Davies, B., Morgan, D.B.: Vasopressin and plasma volume response to submaximal and maximal exercise in man. *J. Sports Med. Phys. Fitness* 30(4): 420-425, 1990

- El-Sayed, M.S., Lin, X., Rattu, A.J.M.: Blood coagulation and fibrinolysis at rest and in response to maximal exercise before and after a physical conditioning program. *Blood Coagul. Fibrinol.* 6: 747-752, 1996
- El-Sayed, M.S., Sale, C., Jones, P.G.W., Chester, M.: Blood hemostasis in exercise and training. *Med. Sci. Sports Exerc.* 32(5): 918-925, 2000
- Ernst, E.: Plasma fibrinogen. An independent cardiovascular risk factor. *Int. J. Med.* 227: 365-372, 1990
- Ferguson, E.W., Bernier, L.L., Banta, G.R., Yu-Yahiro, J., Schoemaker, E.B.: Effects of exercise and conditioning on clotting and fibrinolytic activity in men. *J. Appl. Physiol.* 62: 1416-1421, 1987
- Fowler, W.M., Showbury, S.R., Pearson, C.H., Gardner, G.: Changes in serum enzyme levels after exercise in trained and untrained subjects. *J. Appl. Physiol.* 17: 943, 1962
- Gawaz, M., Neumann, F., Schömig A.: Evaluation of platelet membrane glycoproteins in coronary artery disease. Consequences for diagnosis and therapy. *Circulation* 99: e1, 1999
- Gratas-Delamarche, A., Le Cam, R., Delamarche, P., Monnier, M., Koubi, H.: Lactate and catecholamine responses in male and female sprinters during a Wingate test. *Eur. J. Appl. Physiol.* 68: 362-366, 1994
- Hafner, G., Schinzel, H., Ehrental, W. et al.: Influence of blood sampling from venipunctures and catheter systems on serial determinations of prothrombin activation fragment 1+2 and thrombin-antithrombin III complex. *Ann. Hematol.* 67(3): 121-125, 1993
- Handa, K., Terao, Y., Mori, T. et al.: Different coagulability and fibrinolytic activity during exercise depending on exercise intensities. *Thromb. Res.* 66: 613-616, 1992
- Hebestreit, H., Meyer, F., Htay-Htay, Heigenhauser, G.J.F., Bar-Or, O.: Plasma metabolites, volume and electrolytes following 30-s high-intensity exercise in boys and men. *Eur. J. Appl. Physiol.* 72: 563-569, 1996
- Hedge, S.S., Goldfarb, A.H., Hedge, S.: Clotting and fibrinolytic activity change during the 1 h after a submaximal run. *Med. Sci. Sports Exerc.* 33(6): 887-892, 2001
- Hedlin, A.M., Milojewic, S., Korey, A.: Hemostatic changes induced by exercise during oral contraceptive use. *Can. J. Physiol. Pharmacol.* 56: 316-320, 1978

- Heffner, A. und Kline, J.: Role of the peripheral intravenous catheter in false-positive D-dimer testing. *Acad. Emerg. Med.* 8(2): 103-106, 2001
- Herren, T., Bärtsch, P., Haerberli, A., Straub, P.W.: Increased thrombin-antithrombin III complexes after 1h of physical exercise. *J. Appl. Physiol.* 73: 499-504, 1992
- Hilberg, T., Jeschke, D., Gabriel, H.H.W.: Hereditary thrombophilia in elite athletes. *Med. Sci. Sports Exerc.* 34(2): 218-221, 2002
- Hollmann, W. und Hettinger, Th. (Hrsg.): *Sportmedizin. Grundlagen für Arbeit, Training und Präventivmedizin.* 4. Auflage, Schattauer Verlag, Stuttgart, 2000
- Huisveld, I.A., Kluft, C., Hospers, A.J.H. et al.: The effect of exercise and oral contraceptive agents on fibrinolytic potential in trained females. *J. Appl. Physiol.* 56: 906-913, 1984
- Iatridis, S.G., Ferguson, J.H.: Effect of physical exercise on blood clotting and fibrinolysis. *J. Appl. Physiol.* 18: 337-344, 1963
- Ikarugi, H., Taka, T., Nakajima, S., Noguchi, T. et al.: Norepinephrine, but not epinephrine, enhances platelet reactivity and coagulation after exercise in humans. *J. Appl. Physiol.* 86(1): 133-138, 1999
- Ikkala, E., Myllylä, G., Sarajas, H.S.: Haemostatic changes associated with exercise. *Nature* 199: 459-461, 1963
- Inbar, O., Kaiser, P., Tesch, P.: Relationship between leg muscle fiber type distribution and leg exercise performance. *Int. J. Sports Med.* 2: 154-159, 1981
- Inbar, O., Bar-Or, O., Skinner, J.S.: *The Wingate Anaerobic Test.* Human Kinetics, Champaign, USA, 1996
- Jacobs, I., Bar-Or, O., Karlsson, J., Dotan, R., Tesch, P.A., Kaiser, P., Inbar, O.: Changes in muscle metabolites in females with 30-s exhaustive exercise. *Med. Sci. Sports Exerc.* 14 (6): 457-460, 1982
- Jern, C., Eriksson, E., Tengborn, L., et al.: Changes in plasma coagulation and fibrinolysis in response to mental stress. *Thromb. Haemost.* 62: 767-771, 1989
- Kaczkowski, W., Montgomery, D.L., Taylor, A.W., Klissouras, V.: The relationship between muscle fibre composition and maximal anaerobic power and capacity. *J. of Sports Medicine and Physical Fitness* 22: 407-413, 1982
- Kindermann, W. und Keul, J.: *Anaerobe Energiebereitstellung im Hochleistungssport.* Verlag Karl Hofmann, Schorndorf, 1977

- Kluft, C., Jie, A.F.H., Rijken, D.C., Verheijen, J.H.: Daytime fluctuation in blood of tissue-type plasminogen activator (t-PA) and its fast-acting inhibitor (PAI-1). *Thromb. Hemost.* 59: 329-332, 1988
- Löllgen, H., Winter, U.J., Erdmann, E. (Hrsg.): *Ergometrie – Belastungsuntersuchungen in Klinik und Praxis*. Springer Verlag, Berlin, Heidelberg, 1995
- Mader, A., Heck, H., Föhrenbach, R., Hollmann, W.: Das statische und dynamische Verhalten des Laktats und des Säure-Basen-Status im Bereich niedriger bis maximaler Azidosen bei 400- und 800-m Läufern bei beiden Geschlechtern nach Belastungsabbruch. *Dtsch. Z. Sportmed.* 30(7): 203-211, 30(8):249-261, 1979
- Mann, R.D.: Effect of age, sex, and diurnal variation on the human fibrinolytic system. *J. Clin. Path.* 20: 223-226, 1967
- Marsh, N.A. und Gaffney, P.J.: Exercise-induced fibrinolysis – fact or fiction? *Thromb. Hemost.* 48: 201-203, 1982
- Meyer zum Gottesberge, M., Fuchs, S., Kleyn, M.: Die Ermittlung des Trainingsfaktors bei der Herzfunktionsprüfung durch die Harnkreatininausscheidung und die Blutgerinnungszeit. *Z. Klin. Med.* 143: 388-395, 1943
- Mittleman, M.A., Maclure, M., Tofler, G.H., et al.: Triggering of acute myocardial infarction by heavy exertion – Protection against triggering by regular exertion. *New Engl. J. Med.* 329: 1677-1683, 1993
- Möckel, M., Ulrich, N.V., Röcker, L., et al.: Expression of CD 62p neoantigen on platelets and marked increase of new molecular markers of the hemostatic system during triathlon competition. *Circulation* 94 (Suppl.1): I-326, 1996
- Möckel, M., Ulrich, N.V., Röcker, L., et al.: Exhaustive Cycle Exercise Induces P-Selectin Expression, Coagulation, and Fibrinolysis Activation in Ultra-endurance Athletes. *Thromb. Research* 94: 263-269, 1999
- Möckel, M., Ulrich, N.V., Heller, G., et al.: Platelet Activation Through Triathlon Competition in Ultra-Endurance Trained Athletes: Impact of Thrombin and Plasmin Generation and Catecholamine Release. *Int. J. Sports Med.*, 22: 337-343, 2001
- Molz, A.B., Heyduck, B., Lill, H., Spanuth, E., Röcker, L.: The effect of different exercise intensities on the fibrinolytic system. *Eur. J. Appl. Physiol.* 67: 298-304, 1993
- Niewiarowski, S., Prou-Wartelle, O.: Role du Facteur Contact (Facteur Hageman) dans la Fibrinolyse. *Thromb. Diath. Haemorrh.* 3: 593-603, 1959

- Nilsson, I.M., Pandolfi, M.: Fibrinolytic response of the vascular wall. *Thromb. Diath. Haemorrh.* 40: 231-242, 1970
- Ohira, Y., Ito, A., Ikawa, S.: Correction of water content and solute concentration in blood during haemoconcentration. *J. Appl. Physiol.* 42(5): 739-743, 1977
- Prisco, D., Paniccia, R., Bandinelli, B. et al.: Evaluation of clotting and fibrinolytic activation after protracted exercise. *Thromb. Res.* 89: 73-78, 1998
- Prisco, D., Paniccia, R., Guarnaccia, V. et al.: Thrombin generation after physical exercise. *Thromb. Res.* 69: 159-164, 1993
- Prowse, C.V., Farrugia, A., Boulton, F.E., et al.: Comparative Study Using Immunological and Biological Assays of the Hemostatic Response to DDAVP Infusion, Venous Occlusion and Exercise in Normal Man. *Thromb. Haemost.* 110-114, 1983
- Prowse, C.V. und Cash, J.D.: Physiologic and pharmacologic enhancement of fibrinolysis. *Sem. In Thromb. And Haemost.* 10: 51-60, 1984
- Rankinen, T., Vaisanen, S., Penttila, I., Rauramaa, R.: Acute dynamic exercise increases fibrinolytic activity. *Thromb. Hemost.* 73: 281-286, 1995
- Robertson, B.R., Pandolfi, M., Nilsson, I.M.: „Fibrinolytic capacity“ in healthy volunteers as estimated from effect of venous occlusion of the arms. *Acta Chir. Scand.* 21: 1-8, 1972
- Röcker, L.: Das Verhalten von Plasmavolumen und Plasmaproteinen nach körperlicher Leistung, körperlichem Training und Hitzeeinwirkung. *Habilitationsschrift, FU Berlin*, 1978
- Röcker, L., Kirsch, K., Stoboy, H.: The Influence of Heat Stress on Plasma Volume and Intravascular Proteins in Sedentary Females. *Eur. J. Appl. Physiol.* 36: 187-192, 1977
- Röcker, L., Kirsch, K., Agraval, B.: Long-term observation on plasma antidiuretic hormone levels during and after heat stress. *Eur. J. Appl. Physiol.* 40: 59-62, 1982
- Röcker, L., Stiege-Quast, B., Schwandt, H.J., Quast, J.: Der Einfluß körperlicher Aktivität auf das plasmatische Gerinnungssystem. In: Franz, I.W., Mellerowicz, H., Noack, W. (Hrsg.): *Training und Sport zur Prävention und Rehabilitation in der technisierten Umwelt.* Springer-Verlag, S. 332-338, 1985
- Röcker, L., Drygas, W.K., Heyduck, B.: Blood platelet activation and increase in thrombin activity following a marathon race. *Eur. J. Appl. Physiol.* 55: 374-380, 1986

- Röcker, L., Kirsch, K.A., Heyduck, B., Altenkirch, H.-U.: Influence of Prolonged Physical Exercise on Plasma Volume, Plasma Proteins, Electrolytes, and Fluid-Regulating Hormones. *Int. J. Sports Med.* 10: 270-274, 1989
- Röcker, L., Taenzer, M., Drygas, W.K., Lill, H., Heyduck, B., Altenkirch, H.U.: Effect of prolonged physical exercise on the fibrinolytic system. *Eur. J. Appl. Physiol.* 60: 478-481, 1990
- Röcker, L., Möckel, M., Westpfahl, K.P., Gunga, H.C.: Influence of Maximal Ergometric Exercise on Endothelin Concentration in Relation to Molecular Markers of the Hemostatic System. *Thromb. Haemost.* 75(4): 612-616, 1996
- Röcker, L., Günay, S., Gunga, H.C., Hopfenmüller, W., et al.: Activation of Blood Platelets in Response to Maximal Isometric Exercise of the Dominant Arm. *Int. J. Sports Med.* 21: 191-194, 2000
- Scheele, K., Klaus, E.J., Walter, K.H., Burgdörfer, H.: Über das Verhalten der Blutgerinnung, der Fibrinolyse und der Transaminasen unter Ovulationshemmern bei und nach dosierter körperlicher Belastung. *Med. Welt* 23 (33/34): 1118-1120, 1972
- Schobersberger, W., Wirleitner, B., Puschendorf, B. et al.: Influence of an ultramarathon race at moderate altitude on coagulation and fibrinolysis. *Fibrinolysis* 10: 37-42, 1996
- Seitz, R., Lerch, L., Immel, A., Egbring, R.: D-dimer tests detect both plasmin and neutrophil elastase derived split products. *Ann. Clin. Biochem.* 32(Pt2): 193-195, 1995
- Serresse, O., Lortie, G., Bouchard, C., Boulay, M.R.: Estimation of the contribution of the various energy systems during maximal work of short duration. *Int. J. Sports Med.* 9 (6): 456-460, 1988
- Stevenson, E.T., Davy, K.P., Seals, D.R.: Hemostatic, metabolic androgenic risk factors for coronary heart disease in physically active and less active postmenopausal women. *Arterioscler. Thromb. Vasc. Biol.* 15: 669-677, 1995
- Strauss, M.B., Davis, R.K., Rosenbaum, J.D., Rossmeisl, E.C.: "Water Diuresis" produced during recumbency by the intravenous infusion of isotonic saline solution. *J. Clin. Invest.* 30: 862-868, 1951
- Streiff, M. und Bell, W.R.: Exercise and Hemostasis in Humans. *Semin. Hematol.* 31(2): 155-165, 1994
- Stützer, K.E., Amiral, J., Spanuth, E.: Neue Methoden zur spezifischen Bestimmung von Fibrinolyseprodukten (D-Dimere). *Lab. Med.* 12: 51-59, 1988



- Suzuki, T., Yamauchi, K., Yamada, Y. et al.: Blood coagulability and fibrinolytic activity before and after physical training during the recovery phase of acute myocardial infarction. *Clin. Cardiol.* 15: 358-364, 1992
- Szymanski, L.M., Pate, R.R.: Fibrinolytic response to moderate intensity exercise. Comparison of physically active and inactive men. *Arterioscler. Thromb.* 14(11): 1746-1750, 1994
- Szymanski, L.M. , Pate, R.R., Durstine, J.L.: Effects of maximal exercise and venous occlusion on fibrinolytic activity in physically active and inactive men. *J. Appl. Physiol.* 77:2305-2310, 1994
- Tamayo, M., Sucec, A., Phillips, W., Buon, M., Laubach, L., Frey, M.: The wingate anaerobic test, peak blood lactate, and maximal oxygen debt in elite volleyball players: A validation study. *Med. Sci. Sports Exerc.* 16: S126, 1984
- Thomas, L.: Labor und Diagnose. 5. Auflage, TH-Books Verlagsgesellschaft mbH, Frankfurt, 1998
- Tofler, G.H., Brezinski, D.A., Schafer, A.L., et al.: Concurrent morning increase in platelet aggregability and the risk of myocardial infarction and sudden cardiac death. *N. Engl. J. Med.* 316: 1514-1518, 1987
- Tofler, G.H., Muller, J.E., Stone, P.H., et al.: Modifiers of timing and possible triggers of acute myocardial infarction in the Thrombolysis in Myocardial Infarction Phase II (TIMI II) Study Group. *J. Am. Coll. Cardiol.* 20: 1049-1055, 1992
- Van den Burg, P.J.M., Hospers, J.E.H., Van Vliet, M. et al.: Effect of endurance training and seasonal fluctuation on coagulation and fibrinolysis in young sedentary men. *J. Appl. Physiol.* 82(2): 613-620, 1997
- Vind, J., Gleeurup, G., Nielsen, P.T., Winther, K.: The impact of static work on fibrinolysis and platelet function. *Thromb. Res.* 72(5): 441-446, 1993
- Vogt, A., Hofmann, V., Straub, P.W.: Lack of fibrin formation in exercise-induced activation of coagulation. *Am. J. Physiol.* 236: H577-HH579, 1979
- Wang, J.S., Jen, C.J., Kung, H.C., Lin, L.J., Hsiue, T.R., Chen, H.I.: Different effects of strenuous exercise and moderate exercise on platelet function in men. *Circulation* 90: 2877-2885, 1994
- Watson, R.D. und Sargeant, T.C.R.: Laboratory and on-ice test comparisons of anaerobic power of ice-hockey players. *Canad. J. Appl. Sport Sci.* 11: 218-224, 1986

- Watts, E.J.: Hemostatic changes in long-distance runners and their relevance to the prevention of ischemic heart disease. *Blood Coagul. Fibrinol.* 2: 221-225, 1991
- Weiss, C., Seitel, G., Bärtsch, P.: Coagulation and fibrinolysis after moderate and very heavy exercise in healthy male subjects. *Med. Sci. Sports Exerc.* 30(2): 246-251, 1998a
- Weiss, C., Welsch, B., Albert, M., Friedmann, B., Strobel, G., Jost, J., Nawroth, P., Bärtsch, P.: Coagulation and thrombomodulin in response to exercise of different type and duration. *Med. Sci. Sports Exerc.* 30(8): 1205-1210, 1998b
- Weiss, C., Bierhaus, A., Kinscherf, R., Hack, V., Luther, T., Nawroth, P.P., Bärtsch, P.: Tissue factor-dependent pathway is not involved in exercise-induced formation of thrombin and fibrin. *J. Appl. Physiol.* 92: 211-218, 2002
- Willich, S.N., Lewis, M., Lowel, H., Arntz, H.R., Schubert, F., Schroder, R.: Physical exertion as a trigger of acute myocardial infarction. *New Engl. J. Med.* 329: 1684-1690, 1993
- Winther, O.: Exercise and blood fibrinolysis. *Lancet* II: 1195-1196, 1966
- Winther, K. und Reine, E.: Exercise-induced platelet aggregation in angina, and its possible prevention by  $\beta$ 1-selective blockade. *Eur. Heart J.* 11: 819-823, 1990
- Witt, I., Beeser, H., Müller-Berghaus, G.: Minimalanforderungen zur Gewinnung von Zitratplasma für hämostaseologische Analysen. *Hämostaseologie* 15: 169-170, 1995
- Zivelin, A., Rosenberg, N., Faier, S.: A single genetic origin for the common prothrombotic G20210A polymorphism in the prothrombin gene. *Blood* 92: 1119-1124, 1998