

## 6. Literaturverzeichnis

- 1) Akal U K, Sayan N B, Aydogan S, Yaman Z:  
Evaluation of the neurosensory deficiencies of oral and maxillofacial region following surgery.  
Int J Oral Maxillofac Surg 29(5), 331-336 (2000)
- 2) Bailey P H, Bays R A:  
Evaluation of long-term sensory changes following mandibular augmentation procedures.  
J Oral Maxillofac Surg 42(11), 722-727 (1984)
- 3) Bell W H, Brammer J A, McBride K L, Finn R A:  
Reduction genioplasty: Surgical techniques and soft tissue changes.  
Oral Surg Oral Med Oral Pathol 51, 471-477 (1981)
- 4) Bull H G:  
Value of genioplasty as a profile improving intervention in esthetic facial surgery.  
Mund Kiefer Gesichtschir 1(1), 102-104 (1997)
- 5) Campbell R L, Shamaskin R G, Harkins S W:  
Assessment of recovery from injury to inferior alveolar and mental nerves.  
J Oral Surg 64, 519-523 (1987)
- 6) Cunningham L L, Tiner B D, Clark G M, Bays R A, Keeling S D, Rugh J D:  
A comparison of questionnaire versus monofilament assessment of neurosensory deficit.  
J Oral Maxillofac Surg 54(4), 454-459 (1996)
- 7) Davis W H, Davis C L, Daly B W, Taylor C:  
Long term bony and soft tissue stability following advancement genioplasty.  
J Oral Maxillofac Surg 46, 731-735 (1988)
- 8) Denecke H J, Meyer R:  
Plastische Operationen am Kopf und Hals  
Band 1: Korrigierende und rekonstruktive Nasenplastik;  
Berlin / Göttingen / Heidelberg (1964)
- 9) Driemel O, Kloss F, Roth B, Würzler C, Pistner H:  
Kinnosteotomie isoliert und in Kombination: Langzeitergebnisse unter besonderer Berücksichtigung der Sensibilität und der Fotoanalyse.  
Mund Kiefer Gesichtschir (2004 zur Publikation angenommen)
- 10) Ellis E, Dechow P C, McNamara J A, Carlson D S, Liskiewicz W E:  
Advancement genioplasty with and without soft tissue pedicle: an experimental investigation.  
J Oral Maxillofac Surg 42, 637-645 (1984)

- 11) Epker B M:  
Ästhetische Gesichtschirurgie;  
München (1996)
- 12) Ewing M, Ross R B:  
Soft tissue response to mandibular advancement and genioplasty.  
Am J Orthod Dentofac Orthop 101(6), 550-555 (1992)
- 13) Fridrich K L, Holton T J, Pansegrau K J, Buckley M J:  
Neurosensory recovery following the mandibular bilateral sagittal split osteotomy.  
J Oral Maxillofac Surg 53, 1300-1306 (1995)
- 14) Gallagher D M, Bell W H, Storum K A:  
Soft tissue changes associated with advancement genioplasty performed superior repositioning of the maxilla.  
J Oral Maxillofac Surg 42(4), 238-242 (1984)
- 15) Gattinger B:  
Konturkorrekturen im Kinnbereich.  
In: Fortschritte der Kiefer- und Gesichtschirurgie; Band 24  
Thieme-Verlag; Stuttgart; (1979) 105-108
- 16) Ghali G E, Epker B N:  
Clinical neurosensory testing: Practical applications.  
J Oral Maxillofac Surg 47, 1074-1078 (1989)
- 17) Goracy E S:  
Fracture of the mandibular body and ramus during horizontal osteotomy genioplasty.  
J Oral Surg 36(11), 893-894 (1978)
- 18) Gross B D, Moss R A:  
Cutaneous necrosis of the chin following mandibular advancement and genioplasty.  
J Maxillofac Surg 6(2), 140-142 (1978)
- 19) Guyuron B, Kadi J S:  
Problems following genioplasty. Diagnosis and treatment.  
Clin Plast Surg 24(3), 507-514 (1997)
- 20) Hofer O:  
Operation der Prognathie und Mikrogenie.  
Dtsch Zahn Mund Kieferheilkd 9, 121-132 (1942)
- 21) Hoffman G R, Moloney F B:  
The stability of facial osteotomies.3. Chin advancement.  
Aust Dent J 40(5), 289-295 (1995)

- 22) Hoffmann-Axthelm W:  
Die Geschichte der Zahnheilkunde; 2. Auflage  
Berlin (1985)
- 23) Hoffmann-Axthelm W:  
Die geschichtliche Entwicklung der Mund-, Kiefer- und Gesichtschirurgie  
Fortschritte der Mund-, Kiefer- und Gesichtschirurgie 21. 1976
- 24) Hoffmann-Axthelm W:  
Die Geschichte der Mund-, Kiefer- und Gesichtschirurgie;  
Berlin (1995)
- 25) Hohl T, Epker B M:  
Macrogenia: a study of treatment results, with surgical recommendations.  
Oral Surg Oral Med Oral Pathol 41(5), 545-567 (1976)
- 26) Kawamara P, Wessberg G A:  
Normal trigeminal neurosensory responses.  
Hawaii Dent J 16, 8-11 (1985)
- 27) Kelly J P, Malik S, Stucki-McCormick S U:  
Tender swelling of the chin 40 years after genioplasty.  
J Oral Maxillofac Surg 58(2), 203-206 (2000)
- 28) Kim S G, Lee J G, Lee Y C, Cho B O:  
Unusual complication after genioplasty.  
Plast Reconstr Surg 109(7), 2612-2613 (2002)
- 29) Kiyak H A, Hohl T, West R A:  
Psychologic changes in orthognathic surgery patients.  
J Oral Maxillofac Surg 42, 506-512 (1984)
- 30) Krekmanov L, Kahnberg K E:  
Soft tissue response to genioplasty procedures.  
Br J Oral Maxillofac Surg 30(2), 87-91 (1992)
- 31) Lanigan D T, Hey J, West R A:  
Haemorrhage following mandibular osteotomies: a reported 21 cases.  
J Oral Maxillofac Surg 49, 713-724 (1991)
- 32) Lässig H E, Müller R A:  
Die Zahnheilkunde in Kunst- und Kulturgeschichte  
Köln (1983)
- 33) Legan H L, Burstone C J:  
Soft tissue cephalometric analysis for orthognathic surgery.  
J Oral Surg 38, 744-751 (1980)

- 34) Linquist C C, Obeid G:  
Complications of genioplasty done alone or in combination with sagittal split ramus osteotomy.  
Oral Surg Oral Med Oral Pathol 66(1), 13-16 (1988)
- 35) Ludwig A, Merten H A, Jager A:  
Genioplasty. The long-term results after chin osteotomies with stable osteosynthesis.  
Schweiz Monatsschr Zahnmed 106(11), 1015-1022 (1996)
- 36) Luhr H G:  
Bedeutung des Kinns für die Ästhetik des Gesichts.  
In: Fortschritte der Kiefer- und Gesichtschirurgie; Band 34  
Thieme-Verlag; Stuttgart; (1989) 23-26
- 37) Mercuri L G, Laskin D M:  
Avascular necrosis after anterior horizontal augmentation genioplasty.  
J Oral Surg 35(4), 296-298 (1977)
- 38) Mulherin D, Schow C E Jr:  
Traumatic myositis ossificans after genioplasty.  
J Oral Surg 38(19), 786-789 (1980)
- 39) Nishioka G J, Zysset M K, Van Sickels J E:  
Neurosensory disturbance with rigid fixation of the bilateral sagittal split osteotomy.  
J Oral Maxillofac Surg 45(1), 20-26 (1987)
- 40) Omnell M L, Tong D C, Thomas T:  
Periodontal complications following orthognathic surgery and genioplasty 19-year-old: a case report.  
Int J Adult Orthod Orthognath Surg 9(2), 133-139 (1994)
- 41) Ousterhout D K:  
Sliding genioplasty, avoiding mental nerve injuries.  
J Craniofac Surg 7(4), 297-298 (1996)
- 42) Park H S, Ellis E, Fonseca R J, Reynolds S T, Mayo K H:  
A retrospective study of advancement genioplasty.  
Oral Surg Oral Med Oral Pathol 67, 481-489 (1989)
- 43) Posnick J C, Al-Qattan M M, Stepner N M:  
Alteration in facial sensibility in adolescents following sagittal split and correction of the mandible.  
Plast Reconstr Surg 97(5), 920-927 (1996)
- 44) Richard O, Ferrara J J, Cheynet F, Guyot L, Thiery G, Lanc J L:  
Complications of genioplasty.  
Rev Stomatol Chir Maxillofac 102, 34-39 (2001)

- 45) Ring M E:  
Dentistry. An Illustrated History;  
Toronto (1985)
- 46) Scheidemann G B, Legan H G, Bell W H:  
Soft tissue changes with combined mandibular setback and advancement genioplasty.  
J Oral Surg. 39, 505-509 (1981)
- 47) Schüle H:  
Ästhetische Gesichtspunkte bei der Planung von Kinnoperationen.  
In: Fortschritte der Kiefer- und Gesichtschirurgie; Band 18  
Thieme-Verlag; Stuttgart; (1974) 114-117
- 48) Segner D, Holtje W J:  
Long-term results after genioplasty.  
Fortschr Kieferorthop 52(5), 282-288 (1991)
- 49) Steinhäuser E W:  
Proportionen des ästhetischen Gesichts im Vergleich zur bildenden Kunst.  
In: Fortschritte der Kiefer- und Gesichtschirurgie; Band 34  
Thieme-Verlag; Stuttgart; (1989) 1-4
- 50) Storum K A, Bell W H, Nagura H:  
Microangiographic and histologic evaluation of revascularization and healing after genioplasty by osteotomy of the inferior border of the mandible.  
J Oral Maxillofac Surg 46(3), 210-216 (1988)
- 51) Talebzadeh N:  
Long-term hard and soft tissue relapse rate after genioplasty.  
Oral Surg Oral Med Oral Pathol 91(2), 153-156 (2001)
- 52) Tillmann B:  
Die anatomischen Zeichnungen des Leonardo da Vinci.  
In: Fortschritte der Kiefer- und Gesichtschirurgie; Band 40  
Thieme-Verlag; Stuttgart; (1995) 5-11
- 53) Trauner R, Obwegeser H:  
The surgical correction of mandibular prognathism and retrognathia with consideration of genioplasty.  
Oral Surg Oral Med Oral Pathol 10, 677-689 (1957)
- 54) Troulis J M, Kearns G J, Perrott D H, Kaban L B:  
Extended genioplasty: long-term cephalometric, morphometric and sensory results.  
Int J Oral Maxillofac Surg 29(3), 167-175 (2000)

- 55) Vedtofte P, Nattestad A., Hjorting-Hansen E, Svendsen H:  
Bone resorption after advancement genioplasty. Pedicled and non-pedicled.  
*J Craniomaxillofac Surg* 19(3), 102-107 (1991)
- 56) Wallace A F:  
The Progress of Plastic Surgery.  
Oxford (1982)
- 57) Westermark A, Bystedt H, von Konow L:  
Inferior alveolar nerve function after mandibular osteotomies.  
*Br J Oral Maxillofac Surg* 36(6), 425-428 (1998)
- 58) Westermark A, Englesson L, Bongenhielm U:  
Neurosensory function after sagittal split osteotomy of the mandible: a comparison  
between subjective evaluation and objective assessment.  
*Int J Adult Orthodon Orthognath Surg* 14(4), 268-275 (1999)
- 59) Wiese K G:  
Extreme chin advancement with tandem genioplasty.  
*Mund Kiefer Gesichts Chir* 1(1), 105-107 (1997)
- 60) Wittbjer J, Rune B:  
Changes of the profile after advancement genioplasty.  
*Scand J Plast Reconstr Surg Hand Surg* 23(1), 65-70 (1989)
- 61) Wolford L M, Fields R T:  
Surgical Planning.  
In: Booth P W, Schendel S A, Hausamen J E: *Maxillofacial Surgery, Band 2;*  
Churchill Livingstone; Edinburgh, London, New York; (1999) 1229
- 62) Wünsche P N:  
Die Geschichte der plastischen Chirurgie;  
Diss. med. Erlangen-Nürnberg (1990)
- 63) Ylikontiola L, Kinnunen J, Laukkanen P, Oikarinen K:  
Prediction of recovery from neurosensory deficit after bilateral split osteotomy.  
*Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 90(3), 275-281 (2000 / Nr.1)
- 64) Ylikontiola L, Kinnunen J, Oikarinen K:  
Factors affecting neurosensory disturbance after mandibular bilateral sagittal split os-  
teotomy.  
*J Oral Maxillofac Surg* 58(11), 1234-1239 (2000 / Nr.2)
- 65) Zuniga J R, Meyer R A, Gregg J M, Miloro M, Davis L F:  
The accuracy of clinical neurosensory testing for nerve injury diagnosis.  
*J Oral Maxillofac Surg* 56(1), 2-8 (1998)